

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

186,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Pre-Erythrocytic Vaccine Candidates in Malaria

Ken Tucker, Amy R. Noe, Vinayaka Kotraiah,
Timothy W. Phares, Moriya Tsuji,
Elizabeth H. Nardin and Gabriel M. Gutierrez

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/65592>

Abstract

A vaccine providing sterile immunity against malaria has been shown to be possible with antigens from the pre-erythrocytic stages of malaria. Therefore, it is reasonable to focus vaccine development efforts on the pre-erythrocytic stages, consisting of both sporozoites and liver stage parasites, where it is expected that sterile immunity against the parasite can be elicited to block the development of blood stage infection, clinical disease, and resulting parasite transmission. Accordingly, we will review the preclinical and clinical studies of malaria pre-erythrocytic efforts as well as highlight the advances, trends, and roadblocks encountered in these efforts.

Keywords: *Plasmodium falciparum*, malaria, vaccine, clinical trial, CSP, TRAP, CeTOS, LSA, EP1300, STARP, EXP-1, immune response, humoral, cellular, CD4+, CD8+

1. Introduction

Immunity to malaria correlates with age and develops only after years of repeated exposure to bites from infected mosquitoes [1, 2]. Most adolescents in malaria-endemic regions have developed a level of protective immunity that provides resistance to clinical disease but does not fully eliminate infection [1]. This lack of sterile immunity does not prevent further spread of the disease, as some level of parasitemia persists in these individuals. A complete understanding of how the malaria parasite avoids clearance remains unknown. However, mechanisms such as extreme allelic variation, strain diversity, as well as the unique complexity of the malaria parasite life cycle are undoubtedly involved. In this regard, identification of the conserved, critical epitopes of the parasite that are directly associated with eliciting

protection may provide the best viable target node for vaccine development. In addition, recent data from mouse models of malaria suggest that T cell exhaustion significantly impairs development of effective immune responses and contributes to chronic malaria infection [3–5]. Notwithstanding the fleeting immunity that individuals receive from frequent infection with the parasite, it suggests that there are targets for the immune system to attack.

Importantly, exposure to attenuated sporozoites that can infect hepatocytes but do not develop into the blood-phase of malaria can induce immunity that eliminates parasitic infection (i.e., sterile immunity). This has been demonstrated by conducting immunizations with irradiated sporozoites both in mouse models of malaria [6] as well as in humans with different strains of *Plasmodium falciparum* [7–10]. Similarly, induction of sterile immunity can occur in individuals undergoing chloroquine or mefloquine chemoprophylaxis. The resulting “chemically attenuated” *Plasmodium falciparum* sporozoites (CPS-CQ or CPS-MQ), which undergo full liver stage development but do not develop to clinical blood-stage infection [11, 12], can induce sterile immunity [12–15]. Of note is that sterile protection may not fully develop when sporozoites do not progress at least partially through liver stage development, potentially providing stage transcendent protection [16, 17]. These discoveries demonstrate that a vaccine providing sterile immunity against malaria is possible with antigens from the pre-erythrocytic stages of malaria. Therefore, it is reasonable to focus vaccine development efforts on the pre-erythrocytic stages, consisting of both sporozoites and liver stage parasites, where it is expected that sterile immunity against the parasite can be elicited to block the development of blood stage infection, clinical disease, and resulting parasite transmission.

1.1. Correlates of immunity

The correlates of immunity for sterile protection to malaria are not well defined [18, 19]. In both mice and humans immunized with irradiated sporozoites, sterile protective immunity is believed to be directed to both sporozoites and liver stages of malaria [1, 18]. Antibodies to sporozoites can opsonize sporozoites or neutralize invasion into hepatocytes, and potentially act synergistically with T cells to confer sterile immunity [20]. Additionally, several monoclonal antibodies against the major sporozoite surface antigen, the circumsporozoite protein (CSP), are capable of mediating protection upon passive transfer into animals [21].

Regarding cellular response, studies in mice involving depletion of CD4⁺ and CD8⁺ T cells indicate a T cell response is essential for protective immunity [22–24]. In mice immunized with irradiated sporozoites, IFN- γ produced by CD4⁺ and CD8⁺ T cells is believed to be the main antiparasitic effector produced by the immune response [18, 23, 25, 26]. IFN- γ may activate effector cells, such as natural killer cells and macrophages that can target and kill parasite-infected liver cells. CD8⁺ CTLs that recognize *Plasmodium* antigens presented by major histocompatibility complex (MHC) class I molecules on parasite-infected liver cells also may target those cells for destruction [25, 27–29]. This has led to identifying epitopes and vaccination approaches against malaria that elicit CD8⁺ T cell-mediated IFN- γ secretion and development of CTLs [25, 28, 30]. However, it is important to note that in studies of mice with incapacitated CD8⁺ T cell function, CD4⁺ T cell-dependent sterile immunity can be elicited upon immunization with irradiated sporozoites [31]. Additionally, studies evaluating peptide

vaccines in mice indicate that CD8⁺ T cells may not be necessary for sterile immunity but rather CD4⁺ T cells provide a critical role in immunity [18, 32–35]. Thus, in mice there is not a definitive marker for sterile immunity, and it is plausible that different immune responses may jointly contribute to protection.

In humans, immunization with irradiated sporozoites induces protective immunity consisting of both humoral and cell-mediated responses. Cell-mediated immunity comprises of a mixed population of CD4⁺ and CD8⁺ T cells producing IFN- γ and cytotoxic activity, both of which have been shown to possess antiliver stage activity [19, 25]. However, upregulation of CD8⁺ T cells does not always correlate with protective immunity in humans [10, 19, 25]. For example, immunity elicited in humans under chloroquine chemoprophylaxis with *Plasmodium falciparum* sporozoites is associated with CD4⁺ T cell production of IFN- γ , TNF, and IL-2 [12, 14], but not with upregulation of CD8⁺ T cells. Although the role of CD8⁺ T cells in mediating sterile protection in humans is still unclear, it appears that multiple factors including antibody, CD4⁺ and CD8⁺ T cells collectively contribute to overall protective immunity in humans.

1.1.1. Immunological challenges

Two genetic restrictions impact development of vaccines to malaria. The first genetic restriction is related to the extreme heterogeneity of the *Plasmodium* genome and resulting proteome, and existence of a large variety of different circulating strains (particularly for *P. falciparum*) that produce antigens that vary in sequence. As a result of this extreme heterogeneity, in many cases, efforts to identify antigens that can elicit protection against a broad range of malaria strains have been limited to the antigens that are the least variable. In this case, proteins selected for vaccine development are restricted to those relatively conserved across strains, which limits the available vaccine targets. The second restriction involves the population of HLA alleles on each person's cells that recognize these vaccine candidates and are involved in the induction of an immune response. Because of HLA polymorphism, individuals in a population will not generally recognize the same epitopes in an antigen. That is, a single individual in a given population will have HLAs that are capable of binding only a subset of the epitopes present in a vaccine. This means that a single epitope may only stimulate an effective immune response in a small percentage of any population, depending on the representation of the relevant HLA allele(s) in that population [28, 36–38]. The combined effect of these two restrictions is that an effective vaccine must provide many antigenic epitopes to ensure an effective immune response to the spectrum of circulating *Plasmodium* strains is elicited in a significant percentage of a population.

The need for a multiantigen vaccine has resulted in a number of studies identifying conserved epitopes in *Plasmodium* proteins that can stimulate CD4⁺ or CD8⁺ T cells from individuals either exposed to malaria or immunized with sporozoites [36–38]. The approach has been refined to identify promiscuous peptides capable of binding more than one type of HLA, resulting in antigens that can affect immune response in a larger percentage of the population. Incorporation of these epitopes into chimeric recombinant proteins for processing by immune cells to produce the peptides and stimulate T cells has been done [39, 40]. Examples of this approach are the multiepitope strings in ME-TRAP [41] and EP1300 (Section 4). Unfortunately,

in these constructs, the epitope strings produced low immune responses, even when other whole proteins in these vaccine compositions elicited a robust immune response [42–45].

Another confounding element to vaccine development, in general, is the elicitation of regulatory T cells (Tregs). It is understood that Tregs represent a “self-check” for providing immune tolerance to self and preventing the damaging effects of immune overreaction. For example, Tregs are involved in shutting down immune responses after they have successfully eliminated invading infectious agents, as well as their endogenous role of preventing autoimmunity [46]. However, some pathogens have been found to exploit development of Treg responses in order to evade elimination and persist in the human host. Tregs can also be induced by vaccination and some vaccine development efforts, such as those against tumors and chronic infection (e.g., tuberculosis), have included steps to overcome the constraints of such vaccination-induced Treg responses that result in suppression of immunity [47]. One somewhat successful approach pursued by biotechnology and pharmaceutical companies is development of adjuvant formulations, small molecules, and monoclonal antibodies to overcome the induction of and/or inhibit Treg function [48]. This area of study is also important regarding the development of malaria vaccines, as there is a growing set of data to support the conclusion that induction of Tregs during acute malaria infection limits the generation of immune memory and increases susceptibility to infection [49]. Of particular concern for malaria vaccine development efforts is that in the infant target population, functional Tregs are present in high numbers [49]. Therefore, tools to identify and remove Treg epitopes from malaria vaccines could be critical to the development of candidates that induce strong T cell responses.

1.1.2. Antigen discovery efforts

The understanding that infection with attenuated sporozoites can confer sterile immunity led to a new focus on discovery of liver-stage antigens [1]. Serological approaches to antigen discovery have been used in malaria research; however, sera from chronically infected individuals predominantly recognize the blood-phase of infection [50]. Therefore, discovery of liver-stages antigens has required methodologies that exploit recent discoveries of immunity targeting the liver-phase of infection. For example, immunological approaches using sera from individuals on antimalarial therapy (CPS-CQ or CPS-MQ) while in malaria-endemic regions were applied to screen libraries expressing *Plasmodium* peptides [51, 52]. Later, similar approaches were applied using blood from individuals immunized with irradiated sporozoites to evaluate expression libraries for reactivity with antibodies [53, 54], or T cell responses with peripheral blood mononuclear cells (PBMCs) [55]. These types of approaches continue to be a source of new antigens [56].

The availability of numerous genomic sequences from *Plasmodium* species has greatly facilitated such studies and made possible further large-scale genomic screening approaches that apply transcriptional analysis using isolated sporozoites with targeted comparisons to different phases in the *Plasmodium* lifecycle [57–61]. When applied to vaccine discovery, the genomic screens typically include a selection of proteins shown to react with antibodies or responsive T cells from individuals immune to *Plasmodium*. On a smaller scale, specific stages of the parasite’s lifecycle have been evaluated using proteomic analysis [54, 62, 63]. These

efforts have been augmented by extensive cross-comparison of results to prioritize those antigens identified by more than one approach. Many of these antigens, which show a relatively conserved primary protein structure, are being evaluated and developed as components of subunit vaccines against malaria (Section 5).

1.1.3. Surrogate models of protection

One of the challenges in vaccine development is the need for an animal model that clearly correlates to immunity and efficacy in humans. Mice are frequently used to evaluate antigens for both the resulting immune response and efficacy to preventing infection. Both inbred and outbred strains have been evaluated. While inbred strains (e.g., BALB/c and C57BL/6) provide more consistent results, outbred strains (e.g., CD1 and ICR) are believed to better represent the diversity of the immune response that is encountered in humans due to the polymorphism of the MHC (HLA in human). As *Plasmodium* strains that infect humans do not infect rodents, mouse models of malaria either require (1) use of rodent malaria species (e.g., *P. berghei*, and *P. yoelii*) and immunization with the rodent malaria ortholog of the target antigen, which is generally divergent from the corresponding vaccine target from a *Plasmodium* species that infects humans (e.g., *P. falciparum* and *P. vivax*); or (2) development of chimeric parasites where the rodent malaria ortholog of the target antigen has been replaced by the corresponding protein from a human malaria species. In some cases, orthologs of the vaccine targets do not exist in the rodent malaria species, making evaluation in animal models difficult (e.g., LSA-1 and SIAP2 [61], and SIAP2 [61, 64]). While chimeric rodent malaria strains that express the *P. falciparum* antigen have still been used in this type of scenario [33], the impact upon the physiology of the parasite, the host-pathogen interaction, and resulting response to vaccination is not clear. In rare cases, orthologs are sufficiently similar between the rodent and human malaria species to permit immunization using the human malaria antigen and challenge using a wild-type rodent malaria species [65]. While mouse models of malaria can be useful, immune response and level of protection to a target antigen can vary when different rodent malaria strains/species or chimeric models are used, making analysis difficult as it is not clear which of these datasets may be most applicable to or predictive of *P. falciparum* in humans [19, 28].

The use of nonhuman primate (NHP) models provides a more phylogenetically related model of human disease for both the host and parasite. However, cost and the regulations concerning research with NHPs have limited the use of these models. Even when used to evaluate immunity and safety of vaccines, NHPs are typically not challenged with sporozoites, so vaccine efficacy cannot be determined [66–68]. Further, interpretation of results using a NHP model must be balanced with the realization that old-world monkeys do not have the ortholog to the human HLA-C locus, which is one of the three classical human MHC class I loci [69]. In humans, HLA-A and HLA-B (the other two classical human MHC class I loci) exhibit extreme diversity through extensive allelic polymorphism in each class I gene [70]. Whereas in *Rhesus macaque* (the best characterized old-world monkey), there are very limited morphotypes for class I genes, but relative to humans there are more types of class I genes and extensive diversity of the combination of the genes resulting in a potential 10-fold greater diversity of proteins in NHPs [70, 71]. Therefore, the limited overlap of NHP and human MHC, with each species

containing unique MHC genes, and the much larger repertoire of MHCs in NHP makes it difficult to extrapolate results in NHP immune response to predict the response in humans. Thus, the immune response of NHPs to peptides developed for a human vaccine must be cautiously interpreted, and extrapolation of the results to define an appropriate dosage for use in humans may be problematic. Because of these limitations, many antigens have rapidly progressed to testing in humans with very limited efficacy testing in animals.

Lastly, it is noteworthy that several research groups have recently developed humanized mouse models that mimic human immune system for the purpose of testing human malaria vaccines. These humanized mice can mount both human T cell and antibody-mediated immunity in the context of HLA-restriction [72–74]. Other efforts have been made toward creating a mouse model having human liver [75–79]. These humanized liver mice can sustain development of human malaria parasites from sporozoites to blood stages.

2. Pre-erythrocytic vaccines based on CSP

2.1. Structure and antigenicity

One of the most studied of malaria proteins is the sporozoite surface antigen CSP, which forms a dense coat surrounding the parasite. This protein is involved in sporozoite motility and invasion where contact of the N-terminal region of CSP with hepatocytes triggers cleavage of the N-terminal region from the remainder of the protein [80–82]. This cleavage event is required for sporozoite infection of hepatocytes [80, 81, 83]. Apart from the central repeat region, CSP contains two regions of high conservation: Region 1 in the N-terminal third of the protein that includes the proteolytic cleavage site, and Region II in the C-terminal third of the protein; the latter has some hepatocyte binding activity [82, 84] and is located in a section of CSP enriched in T cell epitopes (**Figure 1**).




Figure 1. Schematic of full-length CSP identifying important epitopes.

Over 30 years ago, NANP repeats in the central repeat region of the protein were identified as the target of protective antibodies [85–88]. The T1 epitope, which is located in the *P. falciparum* CS minor repeat region, comprised of alternating NANPNVDP repeats, was originally identified by human CD4⁺ Th1-type T cells from a volunteer immunized with irradiated *P. falciparum* sporozoites [87]. Multiple CD4⁺ and CD8⁺ T cell epitopes have been identified in

the C-terminal region of CSP, and these are recognized by murine and human cells. Several CD4⁺ T cell epitopes in this region, labeled as T*, UTC, or TH2 in various publications, include one or more epitopes termed “universal” as they are recognized by a broad spectrum of HLA class II molecules. The T* epitope is present in the C-terminus of CSP amino acids 326–345 and promotes both cytotoxic and helper CD4⁺ T cell responses. The T* epitope was identified using CD4⁺ T cell clones derived from volunteers immunized by multiple exposures to the bites of irradiated *P. falciparum* (NF54 strain) infected mosquitos [89]. The T* epitope spans part of a polymorphic region as well as a portion of the conserved Region II. Different CD4⁺ T cell clones selected from these irradiated sporozoites volunteers recognized truncated peptides (8–18 amino acids long), all of which contained the RII sequence [89]. Importantly, the T* epitope, unlike the T1 epitope, is presented *in vitro* by multiple DR alleles reflecting a broad MHC class II binding pattern, thus suggesting a “universal CD4⁺ T cell epitope” for malaria [89, 90].

It has been speculated that CSP-specific T cells are required to provide protection through help for antibody production and CTL function, as well as direct cytokine-mediated antiparasite activity [22, 91]. However, the regions of CSP containing immunodominant T cell epitopes are also highly polymorphic among circulating *P. falciparum* strains [92], which is problematic regarding their use in vaccines.

2.1.1. RTS,S (Mosquirix)

The most advanced malaria vaccine (RTS,S) is a virus-like particle (VLP) consisting of a fragment (central repeat and C-terminal regions, amino acids 207–395) of the CSP fused to a hepatitis B virus surface antigen (**Figure 1**). Phase 3 clinical trials of RTS,S/AS01 (commercially known as Mosquirix), a liposomal adjuvant formulation AS01 which contains monophosphoryl lipid A (MPLA) and QS-21, have provided evidence for high levels of antiCSP antibody correlating with reduced clinical malaria episodes [93]. Interestingly, individuals immunized with irradiated sporozoites or RTS,S generate antibodies to the repeat region, which represents the immunodominant region of the protein under these circumstances; however, adults in endemic regions naturally infected with the parasite, and thus partially immune from disease, have high levels of antibodies against the C-terminus and other nonrepeat regions [94]. In malaria nonexposed individuals immunized with RTS,S vaccine, protection correlated with CD4⁺ and CD8⁺ T cells responses to the C-terminal region of CSP [95], while in other studies, sterile immunity was correlated with antirepeat antibodies [96]. In endemic areas, protection against clinical disease in RTS,S immunized infants also correlated with antiCSP antibodies [97]. Altogether, the protective efficacy of the CSP-based RTS,S/AS01 vaccine is a clear demonstration that a recombinant subunit vaccine containing a portion of the protein delivered on a heterologous VLP carrier can provide partial protection in humans. As 30–50% protection against clinical disease was obtained in RTS,S immunized infants and children in endemic areas, it is clear that there is room for improvement upon the RTS,S vaccine.

2.2. Vaccines containing CSP repeats

Whereas there are several multiantigen pre-erythrocytic vaccines that include CSP repeat units along with other CSP epitopes, the vaccines described in this section contain only CSP repeat

units or CSP repeat units combined with epitopes from other malaria antigens. One of the earliest vaccine trials with epitopes from CSP was a Phase I trial with controlled human malaria infection (CHMI) where 35 volunteers were immunized with a three-unit NANP repeat peptide (NANP)₃ conjugated to tetanus toxoid (TT) and adjuvanted with alum [98]. As the CSP repeats are a known B cell epitope, the three volunteers with the highest ELISA and immunofluorescence assay (IFA) titers were challenged with *P. falciparum* sporozoites (strain NF54); one of these volunteers was sterilely protected. A second trial with CHMI was conducted with this same vaccine where 202 volunteers were immunized and four volunteers with the highest titers were challenged; however, no sterile protection was seen in the challenged volunteers [99]. The results from these trials spurred development of vaccines containing minimal epitopes from CSP that contain CSP repeat units as well as the combination of CSP repeat units with epitopes from other malaria antigen vaccine targets.

One effort to explore combination of CSP repeat units with another malaria vaccine target includes development of a series of influenza virosome-based vaccines. However, virosome-based antigen developed contains a slight variation of the CSP repeat units, NPNA rather than NANP [100]. The resulting vaccine incorporating a constrained (NPNA)₃ repeat was termed PEV302. In a Phase I study, 16 volunteers received PEV302 and eight volunteers received a combination vaccine containing PEV302 as well as an influenza virosome containing a portion of domain III from AMA-1, termed PEV301 [101]. Sera collected from volunteers were used to assess ELISA and IFA titer to the constrained repeat peptide and *P. falciparum* (NF54) sporozoites, respectively. Of the 21 volunteers immunized with PEV302, 19 demonstrated titers $\geq 10^2$ after three immunizations [102]; however, a challenge was not conducted as part of this study. A second trial that included CHMI was conducted with these vaccines where the PEV301 and PEV302 epitopes were combined (known as PEV3A) and administered together to 12 volunteers [103]. In this same trial, 13 volunteers were co-immunized with PEV3A and ME-TRAP. Both groups were challenged; however, all volunteers developed parasitemia. The ELISA titers to UK-39 constrained repeat peptide and IFA titers *P. falciparum* (NF54) sporozoites elicited in both studies were similar with all individuals in the second study demonstrating titers $\geq 10^2$.

2.3. Vaccines containing minimal CSP epitopes

It has been suggested that malaria, along with other infectious diseases (e.g., HIV), has resisted classical vaccine strategies by utilizing a method of stimulating either strain-specific or incomplete, nonprotective immunity, which appears to have arisen from having co-evolved with humans for millions of years [104]. Further, studies with CSP in particular have suggested mechanisms other than allelic variation for parasite escape, even suggesting that the immune-dominant repeat region presents as an immunogenic “decoy” that elicits nonneutralizing antibody responses [105]. In general, the majority of vaccine developed strategies have targeted pathogens that display little antigenic variation, are highly conserved among isolates, and are readily inhibited by immune responses stimulated by native antigens, which is not the case for malaria. Furthermore, that only partially protective immune responses result after years of malaria exposure suggests that the parasites may be utilizing a “cloaking” strategy to hide

in plain sight of the host immune system such as incorporation of human-like protein sequences to induce Treg expansion [106]. Indeed, for vaccine targets, it has been shown that the degree of cross-conservation of predicted epitopes with the human genome inversely correlates with their immunogenicity [107], and in the case of malaria, Tregs (via PD1 upregulation) are responsible for a low frequency of precursor T cells [5]. Therefore, utilizing a minimal, well-designed epitope set for one or more targeted antigens may provide the most effective means to circumvent the parasites' avoidance mechanisms.

Efforts to combine the CSP repeat unit with other CSP epitopes include development of a series of synthesized peptide constructs containing (NANP)₃ (also known as B), the T1 epitope (NANPNVDP)₂, and in later studies, the T* epitope (**Figure 1**). A Phase I trial of a synthetic multiple antigenic peptide (MAP) immunogen, which includes four linked T1B peptides adjuvanted in alum with or without QS21, found that better ELISA and IFA titers were achieved with QS21 formulations [108]. This study also demonstrated that recognition of the T1 epitope by T cells is highly MHC-restricted with T cell responses limited to high responder MHC class II genotypes (present in only 25–35% of the population). None of the volunteers in this study were challenged. To address limitations of the T1 epitope, the T* epitope (which is recognized in the context of many human MHC class II molecules) was included in a branched synthetic peptide immunogen containing four linked T1BT* peptides as well as a linked Pam3Cys as endogenous adjuvant [109]. In a small Phase I trial of the tetra-branched T1BT* peptide in 10 volunteers of diverse HLA types, the majority of the volunteers (8/10) seroconverted following the first dose and reached peak antirepeat antibody titers of 10³–10⁴ following three immunizations [110]. The immunized volunteers developed T*-specific Th1- type CD4+ T cell responses. CD4+ T cell clones derived from PBMC up to 10 months after peptide immunization recognized the T* epitope in the context of multiple HLA DR and DQ molecules and secreted high levels of IFN- γ and variable levels of TNF- α [111]. The immunized volunteers in this study were not challenged. Note that up to 1 mg of immunogen was safely administered per dose in these clinical studies.

Additional clinical trials have investigated novel delivery platforms and adjuvants to enhance immunogenicity of CSP minimal T and B cell epitopes. A VLP based on the hepatitis B virus core antigen, ICC 1132, was engineered to express the T1, B (NANP)₃, and T* epitopes [112]. A total of four clinical studies were conducted with this antigen. Phase I trials using alum as adjuvant elicited suboptimal antibody and cellular responses [113, 114]. Use of a more potent water-in-oil emulsion-based adjuvant, Montanide ISA 720, enhanced not only immunogenicity, but also reactogenicity in a NHP model [115]. Two clinical studies were conducted with ICC 1132 formulated in Montanide ISA 720; however, due to the potential for increased reactogenicity, only a single administration was given. In the first study with ICC 1132 in ISA 720, escalating doses of antigen (from 20 to 50 μ g) were administered [116]. Both ELISA and IFA titers with a single dose of ICC 1132 in ISA 720 were improved as compared to those achieved with three doses of ICC 1132 in alum. A second clinical study was conducted where a single 50 μ g dose of ICC 132 in ISA 720 was administered to 11 volunteers that were subsequently challenged; however, none of the volunteers were protected [117]. A summary of Phase I clinical studies, with CSP vaccine candidates, is provided in **Table 1**.

Category	RTS,S				Pf CS 282-383	NANP-TT	(T1B)4	(T1BT*)4	ICC-1132			
Epitopes	NANP, aa 302–395 (207–395)				NANP, aa 302–383 (282–383)	(NANP)3	NANP, NVDP (T1)	NANP, NVDP (T1), (T*) aa 326–345	NANP, NVDP (T1), (T*) aa 326–345			
Carrier/platform	HBsAg				None	TT fusion	MAP	Polyoxime	HBcAg (truncated)			
Formulation/adjuvants	AS02	AS03	AS04	Alum	ISA 720	Alum	Alum +/- QS21	Pam3Cys	Alum, saline	Alum	ISA 720	ISA 720
Route	IM	IM	IM	IM	IM	IM	SC	SC	IM	IM	IM	IM
# Administrations	3	3	3	3	3	3	3	3	2–3	3	1	1
Dose	50µg 10µg, 25µg, 50µg	50µg	50µg	50µg	≥100µg	20µg, 50µg, 100µg, 160µg	500µg, 1000µg	1000µg	20µg, 50µg	20µg, 50µg	5µg, 20µg, 50µg	50µg
IFA: % titer >10 ² [group GMT]	100% [6400] 100% [3200–10500]	100% [4755]	75% [654] 100% [ND]	50% [ND]	100% [2364–2810]	23% [ND]	25–67% (+QS21) [5511]	70% [216]	25–63% [57–226]	33–71% [ND]	38–100% [ND]	ND
ELISA: % Seroconv. (titer >10 ²), coat Ag	100%, NANP 100%, NANP	100%, NANP	100%, NANP 100%, NANP	100%, NANP	100%, CS 282–383	33–57%, NANP	25–86% (+QS21), spz	100%, (T1BT*)4	50–75%, (T1B)4	40–75%, (T1B)4	63–100%, (T1B)4	100%, NANP
Cellular proliferation	ND 50–86%	ND	ND 10%	0%	63% (ISA)	0%	86% (+QS21)	60%	47–75%	ND	50%	ND
Protection (# protect- ed/# challenged)	86% (6/7) 50µg × 3 30% (3/10) 50µg × 1 50% (7/14) 50µg × 2 50% (3/6) 50µg × 3 57% (4/7) 25µg × 3 25% (1/4) 10 µg × 3	29% (2/7)	13% (1/8) 25% (2/8)	0% (0/6)	ND	33% (1/3)	ND	ND	ND	ND	ND	0% (0/11)
References	[121] [124]	[121]	[121] [122]	[122]	[119]	[98]	[123] [90]	[110] [90]	[113] [112]	[114]	[116]	[117]

Percentages indicate percent responders for IFA, ELISA, and cellular proliferation; ND, not determined; GMT, geometric mean titer.

Table 1. Summary of clinical trials of CSP vaccine candidates (including RTS,S CHMI studies).

2.4. Long synthetic CSP peptide vaccines

In addition to vaccine studies using minimal T and B cell epitopes, clinical studies have investigated immunogenicity of long synthetic peptides containing the entire N- or C-terminus of *P. falciparum* CSP [118]. Of interest is the Pf CSP 282-383 study where a long synthetic peptide including C-terminus amino acids 282–383 from CSP (PfCS102) was synthesized and administered to volunteers as a formulation in either alum or Montanide ISA 720 [119]. Elevated ELISA and IFA titers were seen with ISA 720 formulation as compared to alum. In a second clinical trial, PfCS102 formulated in ISA 720 or AS02A elicited IFN- γ secreting CD8⁺ T cells specific to malaria in some individuals [120]. Tenfold higher antibody and cellular responses were obtained with the AS02 adjuvant formulation. However, no volunteers were challenged in these studies. Note that synthesized peptide doses of $\geq 100\mu\text{g}$ were administered in this study.

2.5. Preclinical pipeline of novel CSP-based vaccines

Additional preclinical studies of VLP have utilized a woodchuck hepatitis core antigen engineered to express CSP repeats in the loop region and multiple universal T cell epitopes at the C-terminus, termed Mal-78-3T VLP [125]. Mice immunized intraperitoneally (i.p.) with Mal-78-3T VLP, formulated in alum/QS21 or ISA 720 adjuvant, developed high levels of antirepeat antibodies. Sterile immunity was obtained in Mal-78-3T immunized mice of three different strains following challenge by transgenic sporozoites expressing *P. falciparum* CSP repeats.

The minimal T and B cell epitopes have also been tested as a recombinant protein using a TLR5 agonist, flagellin, as adjuvant in murine studies [126, 127]. Chimeric flagellin protein containing multiple T1BT* modules, or nearly full-length *P. falciparum* CSP, elicited high levels of antirepeat antibody. The chimeric protein was immunogenic without addition of exogenous adjuvant when delivered subcutaneously or as a needle-free intranasal vaccine.

In addition to the TLR 5 agonist, a TLR-7 agonist, imiquimod, was also found to be a potent adjuvant for vaccines containing minimal T and B cell epitopes. Immunization of mice with the linear T1BT* peptide subcutaneously followed by application of topical TLR-7 agonist imiquimod induced protective antibody-mediated immunity as well as splenic CD4⁺ T cell responses [128]. Recent Phase I/II studies have demonstrated that topical imiquimod can also enhance immunogenicity and protective efficacy of a seasonal flu vaccine [129, 130], supporting the use of the murine model for identification of promising TLR agonist adjuvants for human trials.

Of interest are murine studies demonstrating that micro- and nanoparticle vaccines containing minimal T and B cell epitopes were immunogenic without the addition of exogenous adjuvants. Layer by Layer (LbL) microcapsules were constructed by sequential layering of positively charged poly-L-lysine and negatively charged poly-L-glutamic acid with derivatized peptides containing T1BT* epitopes added to the final layer [131]. Mice immunized with LbL microparticles in PBS elicited sporozoite neutralizing antibodies that blocked *in vitro* invasion of transgenic *P. berghei* sporozoites expressing *P. falciparum* CSP repeats into human hepatoma

cells and reduced parasite liver burden following challenge by exposure to the bites of mosquitoes infected with the transgenic rodent parasites. Protection was comparable to levels obtained following immunization with T1BT* peptide in Freund's adjuvant.

Self-assembling proteins that form nanoparticles (SAPNs) have also been examined in murine studies [132, 133]. The SAPN is comprised of 60 monomers of a recombinant linear protein containing trimeric and pentameric coiled-coil domains separated by a flexible linker with T and B cell epitopes expressed at N- or C-terminus. Following expression in *Escherichia coli*, monomers self-assemble to form spherical SAPN of ~40 nm in diameter. A *P. falciparum* SAPN was constructed comprised of monomer containing four copies of the NANP repeats at the C-terminus and three previously identified *P. falciparum* CSP CD8+ T cell epitopes (KPKDEL DY, MPNDPNRNV, and YLNKIQNSL) at the N-terminus, along with a designed pan-DR binding T cell epitope termed PADRE. All strains of mice immunized with two or three doses of SAPN developed high ELISA titers that persisted over 52 weeks and CD8+ T central memory cells secreting IL-2 and IFN- γ . Mice challenged by i.v. injection of transgenic rodent parasites expressing full length *P. falciparum* CSP protein showed 90–100% protection, with 50% of SAPN immunized mice still protected against sporozoite challenge at week 52. Protective immunity was mediated by both sporozoite neutralizing antirepeat antibodies and CD8+ T cells.

One notable target absent in most CSP-based constructs is the N-terminus of CSP, which contains both T and B cell epitopes as well as the highly conserved Region I (**Figure 2**). Further, within Region I, resides the proteolytic cleavage site and a ligand-binding domain which have been shown to be key requirements for sporozoite infection of hepatocytes [80, 81, 83]. Antibodies raised to this region have been shown to confer 90% protection in animal passive transfer studies [134]. Therefore, efforts are now being pursued to identify and incorporate this epitope into some of the strategies described in this section.




Figure 2. Diagram of LSA-1 (adapted from [164]).

3. Vaccines containing liver stage antigens

3.1. Cell-traversal protein for ookinetes and sporozoites (CelTOS)

3.1.1. Structure and antigenicity

As its name suggests, CelTOS is expressed in both the ookinetes and sporozoite stages of malaria, and functions in cell transversal [135]. This protein is highly conserved across multiple

species of *Plasmodium* including those that cause malaria in rodents (*P. berghei*) and primates (*P. falciparum* and *P. vivax*) [135]. Identification of CelTOS (182 AA—GenBank: AAN36249.1) as a potential protective antigen built upon a previous proteomic analysis identifying expressed proteins in specific phases of the *P. falciparum* lifecycle [62]. Using prototypical HLA supertypes, algorithms were developed to identify potential peptides that could bind to HLA supertypes, and peptides from 27 proteins were screened for stimulating PBMCs from volunteers vaccinated with irradiated sporozoites and subsequently challenged with infection by *P. falciparum* [55]. From these, CelTOS was identified as stimulating the highest number of effector T cells producing IFN- γ overall [55]. Subsequently, Kaiser et al. identified CelTOS using a suppression-subtractive hybridization approach in *P. yoelii* sporozoites versus merozoites [58].

In malaria-naive volunteers infected with irradiated sporozoites, peptides from CelTOS stimulate IFN- γ production from effector T cells in eight out of 12 volunteers [55]. Six out of 35 Ghanaian adults demonstrated effector T cells to CelTOS [136]. In mice immunized with irradiated sporozoites, low levels of IFN- γ producing CelTOS-specific CD8⁺ T cells are induced [137].

3.1.2. Preclinical and clinical trials of CelTOS vaccine

Vaccines comprised of CelTOS recombinant protein or expression vectors (i.e., DNA and viral vectors) have demonstrated protective efficacy against malaria in mouse models. CelTOS administered as a recombinant protein with the adjuvants Montanide ISA 720 or glucopyranosyl lipid A stable emulsion (GLA-SE) elicited CelTOS-specific antibodies, CD4⁺ and CD8⁺ T cells, and protection (10–76% protection, depending on dose and mouse strain) from infective challenge with *P. berghei* in BALB/c and CD1 mice [65, 138, 139]. This protection was demonstrated using homologous and heterologous challenge, demonstrating the unique level of conservation of CelTOS even across species of the parasite [65, 138]. Note that a challenge was not conducted with GLA-SE formulations. In mice vaccinated with recombinant CelTOS, protective immunity did not correlate with the level of antibody production, and passive immunization did not provide significant protection from infection [138]. In one study, self-adjuvanting bacterial vectors expressing CelTOS were used to vaccinate BALB/c mice and sterile immunity (40–60% protection, depending on dose) was demonstrated even though an antibody response to CelTOS was not detected [140]. In these studies, clearance of parasites infecting the liver plays a major role in protective immunity, which is likely potentiated by CD4⁺ and CD8⁺ T cells [138, 141]. Protective immunity in BALB/c mice was dependent on the mixed response of both the CD4⁺ and CD8⁺ T cells to CelTOS [138, 140, 141]. Studies using viral vectors encoding CelTOS did not demonstrate protective immunity in either BALB/c or CD1 mice, even though an antibody response and moderate levels of CelTOS-specific CD4⁺ and CD8⁺ T cells were induced, and control cohorts immunized with CSP in the same prime/boost protocol did demonstrate sterile immunity [33, 137].

Two Phase I trials with CHMI have been performed with a recombinant CelTOS protein produced in *E. coli* (FMP012) formulated with either GLA-SE or AS01B and tested for safety and efficacy (ClinicalTrials.gov: NCT01540474; ClinicalTrials.gov: NCT02174978). Malaria-

naïve adults (18–50 years of age) were vaccinated three times with FMP012 GLA-SE or four times with FMP012 AS01B, and adverse events were monitored. Antibody titers to FMP012 were monitored. Vaccinated volunteers along with nonvaccinated controls were challenged with *P. falciparum* to determine the efficacy of the immune response as determined by time to parasitemia evaluated by blood smear. Results for neither trial have been published.

3.2. Exported protein-1 (EXP-1)

3.2.1. Structure and antigenicity

Exported protein 1 (EXP-1) (162 AA—GenBank: AAN35808.1) was discovered by screening an *E. coli* expression library of blood stage antigens using a monoclonal antibody to *P. falciparum* and was designated antigen 5.1 (Ag5.1) [142]. It was subsequently rediscovered by other screening approaches and designated circumsporozoite-related antigen (CRA), EXP-1, QF116, and *P. yoelii* hepatocyte erythrocyte protein 17 (HEP17) [143–147]. Adults in malaria-endemic regions produced antibodies to the protein and EXP-1-dependent CD4+ and CD8+ T cells [37, 38, 143]. EXP-1 was incorporated in some of the earliest subunit vaccines for malaria tested in humans [148], and peptides from this protein continue to be incorporated into vaccines for malaria, such as ex23 used in ME-TRAP [44].

EXP-1 is conserved across strains of specific *Plasmodium* species [143, 147]. The demonstration that this protein elicited an immune response that provided protection across a diverse spectrum of murine malaria strains in challenged mice [149] led to analysis identifying supertype-peptides that elicited response from a variety of HLA types [37]. In addition to being a vaccine target, the protein is a glutathione S-transferase that has been associated with artesunate metabolism, making it a potential drug target for malaria [150].

The immune responses of people in malaria endemic regions demonstrate that EXP-1 elicits antibody and IFN- γ responses from CD4+ and CD8+ T cells. Indonesian and African adults had detectable antibodies that react with EXP-1 [151–153]. In African children in malaria endemic regions, the presence of antibody responding to an EXP-1 peptide encompassing amino acids 101–162 were associated with decreased infection [154]. Low level IFN- γ and CD4+ T cells were seen in response to EXP-1 in West Africans, although CD8+ T cells were not detected [151]. A similar limited response of CD8+ T cells and low level CD4+ T cells by Kenyans was noted with only one peptide, EXP-10, stimulating CD8+ T cells [37, 38]. Adults from Gambia and Tanzania with natural immunity to malaria develop EXP-1-dependent CTL and the EXP-1 peptide EX23 has also been shown to elicit a CTL response [155]. In another study, African children naturally exposed to malaria only developed low levels of EXP-1-dependent IFN- γ expression [156]. However, the peptides used for this study (EXP2, EXP80, and EXP91) were previously demonstrated to give no CD8+ T cell response in African adults [37], and may not reflect the potential for EXP-1 as a vaccine component. This potential is demonstrated in Caucasian adults, that when immunized with irradiated sporozoites developed EXP-1-dependent CD8+ T cells as well as CD4+ T cells [37, 55].

3.2.2. Preclinical and clinical trials of EXP-1

EXP-1 provided protection from malaria in a mouse model where protection was dependent on the development of CD8⁺ T cells and expression of IFN- γ or nitric oxide [149]. This study resulted in an early emphasis on CD8⁺-dependent IFN- γ production in the development of subunit vaccines against malaria.

5.1-(NANP)19 is a recombinant protein expressed in *E. coli* that contains EXP-1 fused with 19 repeats of the tetramer NANP from CSP [157]. Thirteen adults were vaccinated subcutaneously with 5.1-(NANP)19 using two or three immunizations containing 50 or 400 μ g of protein without adjuvant [148]. The vaccine was safe and only caused reactions at the site of injection. All volunteers developed antibody to the NANP peptide, and six developed 5.1-(NANP)19-specific effector T cells. Seven volunteers were subsequently challenged with *P. falciparum* by bites from infected mosquitoes. While all seven volunteers developed parasitemia, one did not develop symptoms of malaria.

In a study with 194 semiimmune children (6–12 years old) immunized with 5.1-(NANP)19, all but eight children had considerable levels of antibody to EXP-1 [158]. None of the nonvaccinated children developed malaria in this study over the 12 weeks of observation, and the protective efficacy of the vaccine could not be determined.

EXP-1 has been tested in a number of clinical trials as a component of multiantigen vaccines: EP1300, L3SEPTL, ME-TRAP, and MuStDP5. See the discussion in Section 5, **Table 3**.

3.3. Liver-stage antigen-1 (LSA-1)

3.3.1. Structure and antigenicity

Liver stage antigen 1 (LSA-1) is expressed after parasites have invaded hepatocytes and antigen accumulates in the parasitophorous vacuole [52, 159]. The function of this antigen is not known. LSA-1 was discovered using antisera from volunteers who remained on chloroquine while in a malaria endemic region to obtain antibodies to the liver stage of malaria. The antisera were used to screen recombinant DNA expression libraries [52]. Subsequently, a peptide ls94 from LSA-1 was found to bind to HLA-B53, which is MHC class I associated with resistance to severe malaria in West Africa. This indicated that class-I restricted CD8⁺ T cells are important for protective immunity against malaria in Africans [160, 161].

LSA-1 (UniProtKB/Swiss-Prot: Q25893) is a 1909 amino acid protein from the *P. falciparum* strain NF54 [162]. Even though strain 3D7 is derived from NF54 [163], the 3D7 strain produces a protein that is 1162 amino acids in length. The amino terminal and carboxyl terminal ends of the protein are nonrepetitive, whereas the middle of the protein contains multiple repeats (**Figure 2**). This repeat region results in significant variation of the protein between strains of *P. falciparum* [162].

LSA-1 antigen stimulates IFN- γ producing T cells, composed of CD4⁺ and CD8⁺ T cells, and CTL was demonstrated from volunteers in regions with endemic malaria [36, 37, 151, 164–166]. The immune response to LSA-1 typically increased with age and exposure to the parasite

[165–168]. In Gabonese children, LSA-1-dependent IFN- γ producing T cells were correlated with reduced disease severity [169]. Further, naturally infected individuals often express antibodies to LSA-1, targeting primarily the central repetitive region of the protein [170], and antibody to LSA-1 is associated with a reduction in disease severity [171–173].

Immunization of malaria-naive volunteers with irradiated sporozoites produced an immune response to LSA-1 that included CD4+ and CD8+ T cells, and a weak antibody response [37, 38, 55, 174]. However, when volunteers were challenged with infection, the volunteers who responded to LSA-1 were not protected from malaria [55]. Further, in malaria-naive volunteers immunized by exposure to bites from infected mosquitoes while receiving chloroquine, 74% of the volunteers produced antibodies to LSA-1; however, the humoral response did not provide sterile protection from malaria [175].

3.3.2. Preclinical and clinical trials of LSA-1 vaccine

Rodent strains of *Plasmodium* do not contain orthologs of LSA-1 [64, 176], and animal models evaluating LSA-1 are limited to NHPs. Analysis of immunity to peptides from LSA-1 in combination with LSA-3, SALSA, and STARP in NHPs has been the focus of several studies wherein immunity to the peptides is altered by lipidating the peptides or incorporating the peptides in Montanide ISA-51 [66, 177, 178]. High levels of B cells and CD4+ T cells producing IFN- γ in response to LSA-1 were demonstrated. One of the peptides for LSA-1 (LSA1-J) elicited CD8+ CTLs. The NHPs were not challenged with infection by *Plasmodium*, so the impact of this immune response is not known.

Analysis of LSA-1 vaccination using prime/boost approaches using DNA viral vehicles is limited. A prime boost of LSA-1 in chimpanzee adenovirus 63 (ChAd63) and modified vaccine Ankara (MVA) in BALB/c and CD-1 mice demonstrated sterile protection using a chimeric strain of *P. berghei* expressing the *P. falciparum* LSA-1 protein [33]. Seven out of eight mice, for both the BALB/c and CD-1 mice, demonstrated a significant delay in time to onset of parasitemia. LSA-1-dependent IFN- γ production by CD4+ and CD8+ T cells, and antibodies were detected; however, the mediators for protection were not clear. In BALB/c mice where CD4+ or CD8+ T cells were depleted, protection was not dependent on CD8+ T cells; however, results suggested that CD4+ T cells play a significant role. Analysis of the immune response developed in rhesus monkeys using DNA priming and pox virus boosting (using the nonreplicating canary pox virus ALVAC-Pf7) was carried out using an antigen cocktail containing LSA-1, CSP, TRAP, AMA1, and MSP1 [179]. This resulted in a predominant CD4+ T cell response with secretion of IL-2. Low-level IFN- γ production was attributed to CD4+ T cells and CD8+ T cells were low and only present in animals with high CD4+ T cell responses. No antibodies to LSA-1 were detected. The animals were not challenged with infection by *Plasmodium*, so the impact of this immune response is not known.

Recombinant protein LSA-NRC was developed to assess efficacy of an LSA-1 vaccine in the clinic. LSA-NRC incorporates *P. falciparum* NF54 LSA-1 nonrepetitive N-terminal and C-terminal regions, and includes two 17-amino-acid repeats from the repetitive central region of the protein (**Figure 3**). This antigen was administered with AS01B (a liposomal formulation with MPLA and QS-21) in rhesus monkeys and elicited high titers of antibodies and CD4+ T

cells producing IL-2 alone or IL-2 with IFN- γ , but CD8+ T cells were not detected [180]. The NHPs were not challenged with infection by *Plasmodium*, so the impact of this immune response is not known.




Figure 3. Diagram of LSA-NRC (adapted from [181]).

Phase I trials with CHMI were carried out in malaria naïve volunteers using a recombinant LSA 1 protein (FMP011). LSA-NRC adjuvanted with either AS01B or AS02A was well tolerated, safe, and immunogenic at both low and high doses of the antigen [64]. All volunteers developed antibodies after the first dose of the vaccine. Vaccinated volunteers developed LSA-1 specific CD4+ T cells following the second dose. The low dose of LSA-NRC, with either AS01B or AS02A, resulted in LSA-1-dependent CD4+ T cells that produced both IL-2 and IFN- γ . However, the LSA-1 high-dose groups had fewer CD4+ T cells and little to no IFN- γ response relative to the low-dose cohort. None of the groups produced LSA-1-dependent CD8+ T cells. These results were similar to the immune response seen in the preclinical studies in mice (BALB/c and A/J mice) and rhesus monkeys [180, 182]. Only the high-dose groups were challenged along with nonvaccinated volunteers using the homologous 3D7 strain of *P. falciparum*. All 22 volunteers that were challenged became parasitemic without delayed onset of the erythrocytic infection.

Several clinical trials of multiantigen pre-erythrocytic vaccines that included LSA-1 as an antigen (EP1300, L3SEPTL, ME-TRAP, MuStDP5, and NYVAC-Pf7) have been conducted and are discussed in Section 4, **Table 3**.

3.4. Liver-stage antigen-3 (LSA-3)

3.4.1. Structure and antigenicity of LSA-3

While LSA-3 is included with the liver-stage antigens, some report it is actually a blood stage antigen [183]. Using malaria-naïve volunteers immunized with irradiated *P. falciparum*, LSA-3 was identified by comparing antibody response of protected and nonprotected volunteers to a recombinant DNA expression library [53].

LSA-3 is composed of three nonrepeating regions (NR-A, NR-B, and NR-C) flanking two short repeat regions (R1 and R3) and one long repeat region (R2) (see **Figure 4**) [53]. The nonrepeat regions are well conserved across geographically diverse strains of *P. falciparum* [184]. The most significant variation is in the repeating regions, but this is due to the organization and the number of repeating subunits rather than composition of the repeating regions [184]. Vaccine development draws upon the protein from *P. falciparum* strains T9/96 and 3D7 (1586 AA GenBank: ACT22567.1).



Figure 4. Diagram of the domains of LSA-3 (adapted from [185]).

In individuals exposed to *P. falciparum*, LSA-3 elicits antibodies, and CD4⁺ and CD8⁺ T cells. West African adults demonstrated CD4⁺, and class I-LSA-3-dependent CTLs [151, 155]. Antibodies to LSA-3 were demonstrated in children from West Africa, and increased with age [171, 185], and malaria-naïve Europeans develop antibodies to LSA-3 when immunized with irradiated sporozoites [53].

3.4.2. Preclinical and clinical trials of LSA-3

Rodent strains of *Plasmodium* do not contain orthologs of LSA-3 [176], and animal models evaluating LSA-3 are limited to NHPs. A murine model with the infectious challenge using a strain of *P. berghei* that was genetically modified to contain the *P. falciparum* LSA-3 gene has been reported [33]. BALB/c and CD1 mice were primed with ChAd63 and boosted with MVA encoding LSA-3, but sterile immunity was demonstrated in only 13% of the BALB/c mice. The BALB/c mice developed antibodies and moderate levels of PBMCs producing LSA-3-specific IFN- γ were developed. Both CD4⁺ and CD8⁺ T cells were produced against the antigen, and CD8⁺ T cells predominated.

Vaccination with LSA-3 has been shown to elicit sterile immunity in chimpanzees [53]. This study demonstrated sterile immunity using various forms of LSA-3, albeit only one chimpanzee was used for each specific antigen. LSA-3 was expressed as GST-fused recombinant peptides in three segments: DG (encompassing AA 1–60), NN (encompassing AA 369–447), and PC (encompassing AA 869–1786). The mixture of these three peptides (designated as LSA-3 GST) elicited sterile immunity when administered with the adjuvant SBAS2 (SBAS2 is an adjuvant containing MPLA and QS21) or with Montanide ISA 51. The synthetic lapidated peptides CT1, NR1, NR2, and RE were also evaluated. Vaccination with NR2 alone elicited sterile immunity. Further, NR2 administered with NR1 and Montanide ISA 51, or with NR1 and CT1 with Montanide ISA 51 also elicited sterile immunity. Studies in chimpanzees using the

same LSA-3 peptides in combination with peptides derived from LSA-1, STARP, and SALSA demonstrated that the LSA-3 peptides elicited IFN- γ production with a mixed response of CD4⁺ and CD8⁺ T cells [66].

Later studies in chimpanzees immunized with plasmid expressing the nearly full length LSA-3 also demonstrated LSA-3 provided protection, with three of the four immunized chimpanzees demonstrating sterile immunity [186]. In this study, only low levels of IFN- γ were detected from a mixed class I and class II response, and no antibodies were detected. Studies in *Aotus* monkeys immunized with recombinant LSA-3 without adjuvant also developed low but sustained levels of antibodies to LSA-3, and LSA-3 dependent IFN- γ production; however, neither correlated with protection [187]. Subsequent challenge with *P. falciparum* demonstrated that three of the five immunized monkeys were sterilely protected. Interestingly, immunofluorescence assay titers to *P. falciparum* sporozoites were highest in monkeys that were protected. These results were repeated when *Aotus* monkeys were immunized using two peptides of LSA-3, amino acids 100–222 in the amino-terminal nonrepeat region, and amino acids 501–596 in the repeat-2 region of the protein from *P. falciparum* strain T9/96 [187]. In this later study, monkeys were immunized with the recombinant peptides adjuvanted with AS02, eliciting antibody and high IFN- γ responses with sterile immunity in four out of four immunized animals.

A Phase I clinical trial with challenge evaluating the safety and efficacy of recombinant LSA-3 protein adjuvanted with aluminum hydroxide or Montanide ISA 720 was carried out, but the results of the study have not been published [188].

Clinical trials for multistage pre-erythrocytic vaccines that contain LSA-3 include L3SEPTL, ME-TRAP, and MuStDP5 and are discussed in Section 4, **Table 3**.

3.5. Sporozoite threonine-asparagine-rich protein (STARP)

3.5.1. Structure and antigenicity

STARP (GenBank: CAA81224.1) was identified by screening a lambda library with antibodies developed in malaria exposed missionaries under chloroquine treatment [51]. It is a 604-amino acid protein with a central hydrophilic region (amino acids 85–489) that contains a complex repetitive structure that can vary in size between different strains of *P. falciparum*. Two peptides from this repetitive region (Rp10 and Rp45) have been evaluated in vaccines [51]. Even with variation in the repetitive structure, the protein is highly conserved across a variety of geographically separated strains [51, 189].

People in malaria endemic regions produce antibody to STARP [190–192], with the percentage of the population related to the rate of exposure to malaria [193]. The main STARP antibody response in Africans naturally exposed to malaria or volunteers immunized with irradiated sporozoites was to the Rp10 peptide, and purified antibody prevented 90% of the sporozoites from infecting human liver cells *in vitro* [193]. In West African children, the presence of antibody to STARP was associated with protection from malaria [171]. A conserved HLA class I epitope eliciting a CTL response in adults naturally infected with *P. falciparum* has been

identified and was incorporated in the multiple epitope used as part of the vaccine ME-TRAP [36, 41] (Section 4, **Table 3**).

3.5.2. Preclinical and clinical trials of STARP

A limited number of T cell epitopes have been identified in STARP. The conserved epitope st8 was demonstrated to elicit a CD8⁺ T cell CTL response in an African adult exposed to *P. falciparum* [36]. The Rp10 epitope is both a B cell and class I T cell epitope [189]. However, in a NHP model, the Rp10 epitope elicits low levels IFN- γ response but no detectable CTL; note that the NHPs were not challenged with infection in these studies [66, 178].

Homologues of STARP have been identified in *P. yoelii* (PY00217 and PY05105) and *P. reichenowi* (PRCDC_0700500) [176, 189]. However, no studies of animals vaccinated with STARP and challenged with infection are published.

Clinical trials for multiantigen vaccines that include STARP, L3SEPTL, and ME-TRAP, are discussed in Section 4, **Table 3**.

3.6. Thrombospondin-related anonymous protein (TRAP)

3.6.1. Structure and antigenicity of TRAP

The TRAP protein was initially identified based on peptide motifs in common with thrombospondin and properdin [194]. Subsequently, it was demonstrated that TRAP contributed to the binding of sporozoites to hepatic cells [195], and antibodies to the protein block binding of sporozoites to hepatic cells [196]. In addition to TRAP referring to “thrombospondin-related anonymous protein,” the literature refers to TRAP as “thrombospondin-related adhesion protein” [42] and as sporozoite surface protein-2 (SSP2) [196]. TRAP antigen from *P. falciparum* T9/96 (559 AA-PRF: 226137) differs by 6% from the amino acid sequence of TRAP in the 3D7 strain (574 AA-CAD52497.1). The major difference between the strains is the protein from the 3D7 strain contains hepta-repeat of the PPN sequence while T9/96 only contains a single PPN.

Early development of TRAP as a component in a malaria vaccine focused on identifying peptides that were recognized by MHC 1 and elicited a CTL response from individuals exposed to the parasite [36–38, 41, 197, 198]. Evaluation of the immune response to TRAP in naturally infected African donors demonstrated CD4⁺ and CD8⁺ T cells. Gambian and Kenyan adults demonstrated CD8⁺ T cell-dependent CTL activity against TRAP [36–38, 151, 197]. In a study in Kenya, TRAP-dependent IFN- γ producing T cells correlated with reduced disease severity in children [199]. Additionally, it has been demonstrated that a CD4⁺ T cell response dominated over a CD8⁺ T cell response to TRAP in Kenyan children and adults, and the CD4⁺ T cells reactive with TRAP were correlated with reduced risk of clinical malaria [200].

Analysis of the immune response to TRAP in malaria-naïve volunteers immunized with irradiated sporozoites (using the 3D7 clone of *P. falciparum* strain NF54) demonstrated CD8⁺ T cell-dependent CTL activity against TRAP [37, 198], and TRAP-specific CD4⁺ T cells [38]. In a separate study using 12 malaria-naïve Caucasian male volunteers immunized with irradiated

sporozoites, TRAP was recognized by five of the volunteers and resulted in the development of TRAP responsive CD4+ T cells, CD8+ T cells, and antibodies [55].

While antibodies are recognized as important in protection from the erythrocytic phase of malaria and reducing the severity of disease, their role in providing sterile protection in the liver-stage is controversial. *In vitro*, antibodies to TRAP can block binding of sporozoites to human hepatocytes, and it has been postulated that antibodies could help to prevent infection of the liver [201]. In a study using malaria-naive volunteers infected with *P. falciparum* while receiving chloroquine to provide sterile immunity, volunteers did not produce antibodies to TRAP or detectable memory B cells [175]. A recent study, using malaria-naive volunteers vaccinated with irradiated sporozoites and subsequently challenged with the homologous strain of *P. falciparum* (3D7 clone of NF54) correlated the magnitude of the antibody response to TRAP with sterile protective immunity [54]. In this study, both sterilely protected individuals (six volunteers) and individuals who developed blood-stage parasitemia (five volunteers) developed antibodies to TRAP and other *P. falciparum* antigens, but the magnitude of the immune response was significantly higher in the protected volunteers. In a similar but separate study using 12 volunteers immunized with irradiated sporozoites, while a subset of the volunteers responded to TRAP, the immune response to TRAP was associated with volunteers who did not develop sterile immunity [55].

3.6.2. Preclinical and clinical trials of TRAP vaccine

Recent studies have tested TRAP alone and in combination with RTS,S in malaria naive adults [202]. The expressed protein was a truncated form of TRAP consisting of amino acids 26–511 with an additional hepta-histidine at the carboxyl-terminus. Volunteers immunized with TRAP in AS02 developed antibodies and modest CD4+ and CD8+ T cell responses (46 SFU/10⁶ PBMCs). In an effort to determine if there was a synergistic effect of immune responses to TRAP and CSP, a clinical study with CHMI was conducted where volunteers were immunized with two 25 µg doses of TRAP or two doses of both TRAP and RTS,S (25 and 50 µg, respectively) in AS02. Following challenge, 1/11 volunteers (9%) developed sterile immunity in the cohort vaccinated with TRAP and RTS,S with no significant delayed patency in remaining volunteers. The level of protective efficacy was lower than was previously reported for immunization with RTS,S alone, suggesting potential antigenic competition.

Clinical trials of multiantigen vaccines containing TRAP, including EP1300, L3SEPTL, ME-TRAP, MuStDP5, and NYVAC-Pf7 are discussed in Section 4, **Table 3**.

4. Multiantigen pre-erythrocytic vaccines

Early studies with multiantigen vaccines were based on the rationale that a combination of antigens would elicit a broad immune response more comparable to that elicited by whole sporozoite vaccine. A number of multiantigen vaccines comprised of DNA, polyprotein, and recombinant viral vectors containing combinations of epitopes or entire pre-erythrocytic antigens have been tested in clinical trials (listed in alphabetical order).

4.1. EP1300

EP1300 is a multivalent DNA vaccine composed of a total of 38 CTL and 16 HTL epitopes derived from TRAP, CSP, EXP-1, and LSA-1 expressed as a single protein, and the NANP repeating epitope from CSP for antibody development (Figure 5). The epitopes are linked using spacers that facilitate proteolytic processing of the protein into individual epitopes in the body; while this worked in animals, processing may not have occurred correctly in humans.




Figure 5. Arrangement of epitopes from CSP, EXP-1, LSA-1, and TRAP in EP1300.

A Phase Ia dose-escalating study in healthy malaria-naive adults (18–40 years of age) was initiated in 2010 to evaluate the safety and immunogenicity of EP1300. EP1300 was delivered using electroporation. This trial was completed in 2015 [203]. The trial demonstrated the vaccine was safe and well tolerated but did not elicit a significant immune response in the volunteers (Unpublished Results, ClinicalTrials.gov: NCT01169077).

4.2. L3SEPTL polyprotein vaccine

L3SEPTL polyprotein is comprised of six pre-erythrocytic-stage antigens, LSA-3, STARP, EXP-1, Pfs16, TRAP, and LSA1-N/C, linked together to produce a 3240-aa-long polyprotein (Figure 6) [204]. In this construct, LSA-1 is modified so that only one repeat is included flanked by the conserved amino-end (amino acids 1–148) and the carboxyl-end (amino acids 1523–1909). The nucleic acid sequence expressing the recombinant fused protein is delivered as plasmid DNA using the human cytomegalovirus promoter, or in the viral vehicles MVA and Fowlpox (FP9), to provide the vaccines PP, MVA-PP, and FP9-PP, respectively.




Figure 6. Diagram of the L3SEPTL fusion protein (adapted from [204]).

The MVA-PP and FP9-PP vaccines were evaluated in a Phase I with challenge dose-escalation human clinical trial using malaria-naive volunteers [205]. Generally, the vaccine was well tolerated, but the immunity was lower than expected. Priming with MVA resulted in a stronger immune response than did FP9, and when FP9 was used for priming (FP9/FP9/MVA), the immune response was not significantly elicited until the MVA vehicle was used to deliver the antigens [205]. This was in contrast with the results seen in the preclinical study using FP9-PP for the priming. When delivered using MVA priming, all six proteins elicited IFN- γ -producing T cells, with stronger and more frequent responses seen for LSA-3, LSA-1, TRAP, and STARP. However, the number of IFN- γ -producing T cells was lower than had been reported in other studies [205]. Following challenge, none of the 15 vaccinated volunteers demonstrated protection or delayed onset of disease relative to the six nonvaccinated volunteers.

4.3. ME-TRAP

The most extensively tested multiantigen vaccine, ME (multi-epitope)-TRAP vaccine, was designed to elicit T cell responses to target the hepatic stages of the malaria parasite. Nucleic acid encoding an ME string was designed to include CD8+ and CD4+ T cell epitopes from TRAP, CSP, EXP-1, LSA-1, LSA-3, and STARP (Table 2) [36, 41, 197, 198]. The ME were fused in-frame to the nucleic acid sequence encoding the entire TRAP antigen from *P. falciparum* strain T9/96, encoding a polypeptide of 789 amino acids referred to as multi-epitope thrombospondin-related adhesion protein (ME-TRAP) [44].

Antigen	Designation	Amino acid sequence	Epitope
CSP	cp6	MNPNDPNRNV	HLA class I antigens
CSP	cp26	KPKDELDY	
CSP	cp39	YLNKIQNSL	
EXP-1	ex23	ATSVLAGL	
LSA1	ls6	KPIVQYDNF	
LSA1	ls8	KPNDKSLY	
LSA1	ls50	ISKYEDEI	
LSA1	ls53	KSLYDEHI	
LSA3	la72	MEKLKELEK	
STARP	st8	MINAYLDKL	
TRAP	tr26	HLGNVKYLV	
TRAP	tr29	LLMDCSGSI	
TRAP	tr39	GIAGGLALL	
TRAP	tr42/43	ASKNKEKALII	
CSP (<i>P. berghei</i>)	pb9	SYIPSAEKI	Mouse MHC class I antigen
CSP	NANP	NANPNANPNANPNANP	B cell epitopes
TRAP	TRAP-AM	DEWSPCSVTTCGKGTRSARKRE	Heparin binding motif
CSP	CSP	DPNANPNVDPNANPNV	Class II antigens (BCG and Tetanus toxin are not malarial antigens)
BCG		QVHFQPLPPAVVKL	
Tetanus toxin	QFIKANSKFIGITE		

Table 2. Peptides used in the multi-epitope domain of ME-TRAP [41].

Vaccine	Antigens (adjuvants)	Delivery regimen	Efficacy (# protected/# participants)	Immune response	Population	References
L3SEPTL (peptide)	LSA, STARP, Exp1, Pfs16, TRAP, and LSA1	FFM	0% protection (0/8)	85 SFU/10 ⁶ PBMCs	Malaria naïve adults	[205]
ME TRAP	TRAP and peptides from CSP, LSA-1, LSA-3, EXP-1, STARP expressed as a single protein.	MMF	0% protection (0/7)	96 SFU/10 ⁶ PBMCs		
		DDDMM	0% protection (8/8 delayed patency)	158-316 SFU/10 ⁶ PBMCs	Malaria Naïve adults	[44]
		DDMM	0% protection, (6/6 delayed patency)	234 SFU/10 ⁶ PBMCs		
		DDD	0% protection (0/5)	33 SFU/10 ⁶ PBMCs		
		MMM	0% protection (0/4)	44 SFU/10 ⁶ PBMCs		
		DDM	12.5% protection (1/8 sterile immunity, 7/8 delayed patency)	423 SFU/10 ⁶ PBMCs; low titer antibodies	Malaria naïve adults	[42]
		DDM	10% protective efficacy relative to control group	250 SFU/10 ⁶ PBMCs	Gambian adults	[219]
		FFM	12.5% protection (2/16 sterile immunity, 14/16 delayed patency)	430 SFU/10 ⁶ PBMCs, mixture of CD4+ and CD8+ T cells	Malaria naïve adults	[45, 207]
		MMM	0% protection (0/4)	0 SFU/10 ⁶ PBMCs		
		FM	0% protection (0/5)	380 SFU/10 ⁶ PBMCs, mixture of CD4+ and CD8+ T cells		
		MF	0% protection (0/5)	100 SFU/10 ⁶ PBMCs, mixture of CD4+ and CD8+ T cells		
		DDMF	0% protection (4/4 delayed patency)	200 SFU/10 ⁶ PBMCs, CD4+ T cells		
		DDFM	0% protection (0/3)	300 SFU/10 ⁶ PBMCs, CD4+ T cells		
		FFM	0% protection (73% infection in vaccines, 80% infection in controls; no statistically significant decrease)	107 SFU/10 ⁶ PBMCs	Kenyan children	[210]
		CM	21% protection (3/14 sterile immunity, 5/14 delayed patency)	1300 SFU/10 ⁶ PBMCs, CD4+ and CD8+ T cells producing IFN- γ , IL2, TNF- α (protection associated with CD8+ IFN- γ producing T cells)	Malaria naïve adults	[212]
		CM	13% protection (2/15 sterile immunity, 5/15 delayed patency)	2000 SFU/10 ⁶ PBMCs, and antibodies detected	Malaria naïve adults	[213]

Vaccine	Antigens (adjuvants)	Delivery regimen	Efficacy (# protected/# participants)	Immune response	Population	References
		CM	67% protection (18% infection in vaccines, 47% infection in controls)	1450 SFU/10 ⁶ PBMCs, CD4+ and CD8+ T cells producing IFN- γ , IL2, TNF- α	African adults	[105, 214]
MuStDO5	TRAP, CSP, EXP1, LSA1, and LSA3	DDD	0% protection (0/8)	For both regimens: class I— 41 SFU/10 ⁶ PBMCs; class II— 59 SFU/10 ⁶ PBMCs, no antibodies detected	Malaria naïve adults	[215, 216]
		DDD & plasmid with hGM-CSF	0% protection (0/23)			
NYVAC-Pf7	CSP, TRAP, LSA-1, MSP1, SERA, AMA1 and Pfs25	NNN	3% protection (1/35 sterile immunity, 34/35 delayed patency)	9/38 volunteers developed CTL to TRAP; 2/38 volunteers developed CTL to LSA1; 50% of volunteers developed antibody to TRAP and LSA1	Malaria naïve adults	[218]

*Regimens are the sequence of immunization using the vehicles D (plasmid DNA), F (attenuated fowlpox virus FP9), M (modified vaccinia virus Ankara), C (chimpanzee adenovirus 63), and N (attenuated vaccinia virus NYVAC) to provide gene sequences encoding the described antigens.

**T cells isolated by FAC, antigen-dependent T cell count not provided. SFU refers to antigen-specific IFN- γ producing T cells.

Table 3. Summary of the efficacy of multi-antigen pre-erythrocytic vaccines tested in clinical trials.

Expressing the ME-TRAP antigen *in situ* using DNA vaccines (plasmids) or viral vehicles has been employed to provide low-cost vaccination. While results in animal models using plasmids to deliver antigens were promising, the immunity elicited in humans was typically low. Heterologous prime/boost combinations, wherein the antigen remained constant but was delivered using different vehicles, reduced the development of immunity to the vehicle while provided a boost to the immune system with the target antigens [206]. In humans, the order of vehicle used for immunization was demonstrated to affect overall immune response [45, 207]. Thus, studies have focused on identifying the appropriate combination and order of vehicles that can be used to produce an elevated and expanded immune response. Plasmids encoding ME-TRAP only elicited a low-level immune response [44], and subsequent efforts have focused on viral vehicles for both priming and boosting (**Table 3**).

The MVA strain has proven to be a strong vehicle for boosting, and clinical trials using ME-TRAP subsequently focused on identifying an optimal priming vehicle and dosing regimen (see **Table 3**). Using an attenuated fowlpox virus vehicle (FP9) with MVA elicited strong immunity in malaria-naïve volunteers and limited protection [207], but provided only low-level immunity in people in malaria-endemic regions without providing significant protection [208–210] (see **Table 3**). The recent addition of a chimpanzee adenovirus 63 (ChAd63) for priming has provided encouraging results (see **Table 3**). AdCh63 priming elicits both CD4+ and CD8+ T cells at much higher levels than DNA or FP9 priming [211], and also results in a

much higher proportion of IFN- γ -secreting monofunctional CD8⁺ T cells that were correlated to sterile protection [212]. This combination has provided protection from infection in malaria-naive volunteers comparable to the immunity provided by CSP using the same delivery platform [212, 213]. Further, ME-TRAP using the ChAd63/MVA regimen elicited a strong immune response and provided significant protection in Africans who had prior exposure to malaria [43, 214]. Using a Cox regression analysis, vaccine efficacy was 67% during the 56 days of monitoring.

4.4. Multistage DNA vaccine operation five antigens (MuStDO5)

The MuStDO5 cocktail is composed of sequences encoding the antigens TRAP, CSP, EXP-1, LSA-1, and LSA-3 using sequences based on *P. falciparum* strain 3D7 [215, 216]. LSA-1 only encodes the nonrepetitive amino and carboxyl ends of the protein (462 amino acids), with the repetitive 727 amino acid middle domain removed. Each antigen is encoded on a separate plasmid using the same vector VCL-25 (Vical Inc., San Diego, CA, USA), with expression controlled by the promoter/enhancer of the human cytomegalovirus. The cocktail of plasmids is used for vaccination to provide each of the five antigens. In this study, the impact of including a plasmid expressing human granulocyte-macrophage colony-stimulating factor (hGM-CSF) as an immune enhancer was evaluated. This vaccine is safe and well tolerated. However, protective immunity was not demonstrated, and the addition of hGM-CSF decreased the response to class I epitopes.

4.5. NYVAC-Pf7 attenuated vaccinia virus

NYVAC-Pf7 is composed of an attenuated vaccinia virus NYVAC engineered to express seven antigens (Pf7) to target pre-erythrocytic, blood stages, and transmission stages of the parasite [217]. The seven antigens include three pre-erythrocytic antigens (CSP, TRAP, and LSA-1), three blood stage antigens (MSP1, SERA, and AMA1), and the Pfs25 antigen found on the sexual stage ookinete. CSP, TRAP, LSA-1, AMA1, and Pfs25 are derived from *P. falciparum* 3D7 strain. The *lsa-1* gene was modified to produce a protein linking amino acids 1–458 with 1630–1909, removing the sequence encoding the repeat region of the protein. Each gene is under a separate poxvirus promoter and expressed as a separate protein as follows: CSP under H6, TRAP under 42K, LSA-1 under C10LW, MSP1 under H6, SERA under 42K, AMA1 under I3L, and Pfs25 under I3L (**Figure 7**). Each gene was inserted into defined sites in the genome of the attenuated vaccinia strain NYVAC by *in vivo* recombination to produce NYVAC-Pf7.

A Phase I/II clinical trial was carried out with malaria-naive adults [218]. Volunteers were immunized with either high dose (1×10^8 pfu) or low dose (1×10^7 pfu) of NYVAC-Pf7 at 0, 4, and 26 weeks. Vaccination with NYVAC-Pf7 resulted in delayed onset of parasitemia following challenge in all volunteers who received the vaccine, as compared to nonvaccinated volunteers. The CTL response or antibody titer did not demonstrate a relationship to the time of delay to onset of parasitemia. One vaccinated volunteer from the low-dosage group demonstrated sterile immunity. About 50% of the volunteers in each of the dosage groups developed antibodies to TRAP and LSA-1, regardless of the dosage. Low levels of CTL were developed to TRAP, with no statistical significance due to the dosage. A CTL response to LSA-1 was only

seen in the low-dosage group. However, the CTL response did not correlate with the time to onset of parasitemia, and the one volunteer exhibiting sterile immunity did not develop a detected CTL response.




Figure 7. Genomic Organization of the NYVAC-Pf7 (adapted from [217]). The poxvirus promoters are represented with solid boxes.

5. New proteins for pre-erythrocytic malaria vaccines

New malarial proteins have been discovered through screens for pre-erythrocytic phase sporozoites, including identifying mRNA or proteins expressed in specific stages of the lifecycle of the parasite (e.g., sporozoites for the mosquito salivary glands), and typically included monitoring the immune response to the specific proteins using antibodies or T cells from volunteers vaccinated with sporozoites that were either irradiated or suppressed with chemotherapy, as discussed in the Introduction. Section 5 summarizes the preclinical studies carried out focusing on antigens analyzed beyond the initial identifying screen.

5.1. DNA vaccine encoding SAPI

Sporozoite asparagine-rich protein 1 (SAPI) was identified using suppressive-subtractive hybridization to identify *P. berghei* genes that are up-regulated in the sporozoite as it transitions from the mosquito midgut into the salivary gland [60]. In targeted deletion studies, *P. yoelii* SAPI knockout sporozoites can invade hepatocytes but arrest in early liver stage development [220]. The *P. yoelii* genome sequence encoding the SAPI protein domain corresponding to amino acids 3063–3227, and flanked by two CpG motifs to enhance the immune response, was cloned into pcDNA3.1(+) [221]. SAPI encoded in plasmid DNA was evaluated for protective immunity against challenge with a homologous strain of *P. yoelii*. BALB/c mice were immunized intramuscularly with 100 µg plasmid three times with 3 weeks between each vaccination. Controls included mice vaccinated with phosphate buffered saline. For the protection study, each group was composed of ten mice. BALB/c mice were challenged by bites from mosquitoes infected with *P. yoelii* 2 weeks after the final immunization. Onset of parasitemia was evaluated using blood smears. Mice without parasitemia 14 days after challenge were considered completely protected. The mice immunized with DNA encoding SAPI produced SAPI-

specific antibodies and moderate levels of T cells secreting IFN- γ . All 10 of the control mice vaccinated with phosphate buffered saline developed parasitemia. Two of 10 (20%) of the mice vaccinated with pcDNA3.1(+)/SAP1 did not develop parasitemia.

5.2. Adenovirus vectored pre-erythrocytic antigens

This was an exploratory study to evaluate the protective immunity of newly identified pre-erythrocytic-stage antigens, selected from sporozoite and liver stage libraries based on the predicted signal sequences [137]. The *P. yoelii* orthologues of P36p (PY01340; Py52), Ag5 (PY00410; hypothetical protein), Sporozoite Invasion-Associated Protein-1 (PY00455; SIAP1), Krueppel-like protein (PY00839; KLP), and TRAP-like protein (PY01499; TLP), CelTOS or CSP were delivered in 10^{10} virus particles of adenovirus encoding each individual antigen [137]. Protection was evaluated as the parasite burden in the liver following infection. Immunization of female BALB/c (H2-K^d) mice with adenovirus vector (Ad) expressing P36p, CelTOS, or Ag5 elicited a high IFN- γ T cell response that was comparable to the response to CSP. Following challenge, only mice immunized with Ad-CSP reduced the parasite burden. None of the other antigens reduced the burden of parasites as compared to nonvaccinated mice.

5.3. Heterologous prime boost with DNA and vaccinia viral vector

The protective immunity of the individual antigens CSP, Falstatin, PY03661, and UIS3, as well as combinations of the antigens, were evaluated using the *P. yoelii* model [222]. CD1 mice (14 per group) were vaccinated intramuscularly with a priming dose of the DNA vaccine, and boosted with the vaccinia vector vaccine. Parasitemia was evaluated by blood smears following challenge with 300 *P. yoelii* sporozoites. The combination of Falstatin and UIS3 and PY03661 provided protective immunity (57%) against challenge. Heterologous prime boost with CSP protected 36% of the mice. The addition of plasmid expressing murine GM-CSF to immunization scheme did not enhance protective immunity.

5.4. Heterologous prime boost with viral vectors ChAd63 and MVA

Humoral responses in mice were evaluated using recombinant ChAd63 and MVA viral vectors expressing eight individual *P. falciparum* 3D7 antigens [33]. Responses elicited following immunization with five newly identified antigens, upregulated in sporozoite 3 (UIS3), Falstatin, liver-stage associated protein 1 (LSAP1), liver-stage associated protein 2 (LSAP2), and the early transcribed membrane protein 5 (ETRAMP5) were compared to three previously characterized antigens, LSA-1, LSA-3, and CelTOS.

Protective efficacy was evaluated in both BALB/c (inbred) and CD1 (outbred) mice, with each antigen evaluated in a separate cohort of mice using heterologous prime boosts with ChAd63/MVA. Mice were challenged with *P. berghei* transgenic parasites that were modified to express the cognate *P. falciparum* antigen. The immunity and protective efficacy against infection was evaluated for each antigen using nonvaccinated controls for comparison.

The BALB/c mice developed antibodies to all the antigens except LSAP1. High levels of IFN- γ (>1000 SFU/10⁶ PBMCs) developed in mice immunized with UIS3 and LSA1. Moderate levels

of IFN- γ (>250 SFU/10⁶ PBMCs) were developed against ETRAMP5, CelTOS, LSAP2, Falstatin, and LSA-3. LSAP1 elicited very low levels of IFN- γ (<100 SFU/10⁶ PBMCs). CD8+ T cells predominated for UIS3, LSA-1, and LSA-3 cellular responses, while CD4+ T cells predominated for Falstatin. Highest levels of protection were found in BALB/c or CD1 mice immunized with LSAP2 (85 and 70%, respectively) or LSA-1 (87.5% for both murine strains). Mice immunized with CSP gave 37.5 and 33.3% sterile immunity in BALB/c and CD1 mice, respectively. Strain variation noted, with 30% of CD1, but none of the BALB/c, mice were protected following immunization with TRAP or LSAP1. Minimal or no protection was obtained in mice immunized with ETRAMPS, CelTOS, UIS3, Falstatin, or LSA-3.

6. Conclusions

Trends, advances, and roadblocks for malaria pre-erythrocytic vaccine development identified by the preclinical and clinical studies summarized in this chapter include:

- The hallmarks for protective immunity are not clear and the elements associated with sterile immunity vary in humans. Antibodies, IFN- γ , CD4+ T cells, CD8+ T cells, and cytolytic lymphocytes are believed to play essential roles in immunity, but are not always collectively demonstrated in immunity that protects from infection by *Plasmodium*.
- To achieve effective immunity that induces T cells across a wide population of people will require multiple and diverse T cell epitopes. Many of the studies that identified T cell epitopes in humans demonstrated that, within the limited number of volunteers in a Phase I clinical trial, only a limited subset of individuals respond to any single protein.
- Vaccines that string T cell epitopes from different proteins together in a composite recombinant protein (e.g., multiepitope strings) do not elicit significant T cell responses in humans. This is found even when other proteins in the vaccine do elicit strong immune response.
- For protective immunity in humans, the means and route of administration effect efficacy of the antigens. For example, irradiated sporozoites are more effective by intravenous verses intradermal administration, and TRAP delivered using heterologous vehicles of chimpanzee adenovirus followed by a modified vaccinia virus (heterologous prime boost regimen) provides more effective immunity than immunization using only one of these vehicles, or using other combinations of vehicles.
- Adjuvant formulations remain a critical factor for pre-erythrocytic vaccines, as highly purified subunit vaccine lack the pathogen-associated molecular patterns that trigger the innate immune responses required for initiation of adaptive immunity. Alum has been found to be a poor adjuvant for malaria subunit vaccines in Phase I/II trials. Oil-based adjuvants, e.g., ISA 50 or ISA 720, while enhancing immune responses, have been limited by increased reactogenicity. Precisely targeted adjuvants, such as TLR agonists, have been shown to enhance immunogenicity and protective efficacy of pre-erythrocytic vaccines, and hold promise as future adjuvant formulations.

- While cost competitiveness is a goal for malaria vaccines, it is unlikely that this has been a primary consideration in the early development plan for many candidates as much of the focus is on efficacy. Multiple platforms and approaches have been described in this chapter including DNA-based platforms, viral vehicles, recombinant proteins, and chimeric proteins composed of peptides. In consideration of vaccine design and development, production may be an initial focus to contain costs per dose. This includes consideration for adjuvants and processes to stabilize the vaccine for delivery to the clinic. However, it is not always obvious in early development which approaches may be the most cost effective for commercial vaccine production as efficiencies may be realized through optimizing processes and scaling of production so as to reduce costs significantly relative to the costs to produce research material. Further contributors to the direct cost for vaccination are transportation and storage requirements for the vaccine, the dosage, and number of boosters required to achieve protective immunity. This cost analysis may also need to consider compliance if complex or protracted boosting schedules are required. While development may strive to contain these direct costs, the vaccine still must be effective at preventing the morbidity and mortality caused by malaria.

Acknowledgements

This publication was made possible through support provided by the Office of Health, Infectious Diseases, and Nutrition, Bureau for Global Health, U.S. Agency for International Development, under the terms of the Malaria Vaccine Development Program (MVDP) Contract AID-OAA-C-15-00071, for which Leidos, Inc. is the prime contractor. The opinions expressed herein are those of the author(s) and do not necessarily reflect the views of the U.S. Agency for International Development.

Author details

Ken Tucker¹, Amy R. Noe¹, Vinayaka Kotraiah¹, Timothy W. Phares¹, Moriya Tsuji², Elizabeth H. Nardin³ and Gabriel M. Gutierrez^{1*}

*Address all correspondence to: gabriel.m.gutierrez@leidos.com

1 Leidos Life Sciences, Leidos Inc., Frederick, MD, USA

2 HIV and Malaria Vaccine Program, Aaron Diamond AIDS Research Center, Affiliate of the Rockefeller University, New York, NY, USA

3 Division of Parasitology, Department of Microbiology, New York University School of Medicine, New York, NY, USA

References

- [1] Crompton PD, Moebius J, Portugal S, Waisberg M, Hart G, Garver LS, et al. Malaria immunity in man and mosquito: insights into unsolved mysteries of a deadly infectious disease. *Annu Rev Immunol.* 2014;32:157–87. DOI: 10.1146/annurev-immunol-032713-120220
- [2] Gupta S, Snow RW, Donnelly CA, Marsh K, Newbold C. Immunity to non-cerebral severe malaria is acquired after one or two infections. *Nat Med.* 1999;5:340–3. DOI: 10.1038/6560
- [3] Horne-Debets JM, Faleiro R, Karunaratne DS, Liu XQ, Lineburg KE, Poh CM, et al. PD-1 dependent exhaustion of CD8+ T cells drives chronic malaria. *Cell Rep.* 2013;5:1204–13. DOI: 10.1016/j.celrep.2013.11.002
- [4] Ryg-Cornejo V, Ioannidis LJ, Ly A, Chiu CY, Tellier J, Hill DL, et al. Severe malaria infections impair germinal center responses by inhibiting T follicular helper cell differentiation. *Cell Rep.* 2016;14:68–81. DOI: 10.1016/j.celrep.2015.12.006
- [5] Wykes MN, Horne-Debets JM, Leow CY, Karunaratne DS. Malaria drives T cells to exhaustion. *Front Microbiol.* 2014;5:249. DOI: 10.3389/fmicb.2014.00249
- [6] Nussenzweig RS, Vanderberg J, Most H, Orton C. Protective immunity produced by the injection of x-irradiated sporozoites of *Plasmodium berghei*. *Nature.* 1967;216:160–2.
- [7] Clyde DF, McCarthy VC, Miller RM, Hornick RB. Specificity of protection of man immunized against sporozoite-induced falciparum malaria. *Am J Med Sci.* 1973;266:398–3.
- [8] Hoffman SL, Goh LM, Luke TC, Schneider I, Le TP, Doolan DL, et al. Protection of humans against malaria by immunization with radiation-attenuated *Plasmodium falciparum* sporozoites. *J Infect Dis.* 2002;185:1155–64. DOI: 10.1086/339409
- [9] Rieckmann KH, Carson PE, Beaudoin RL, Cassells JS, Sell KW. Letter: sporozoite induced immunity in man against an Ethiopian strain of *Plasmodium falciparum*. *Trans R Soc Trop Med Hyg.* 1974;68:258–9.
- [10] Seder RA, Chang LJ, Enama ME, Zephir KL, Sarwar UN, Gordon IJ, et al. Protection against malaria by intravenous immunization with a nonreplicating sporozoite vaccine. *Science.* 2013;341:1359–65. DOI: 10.1126/science.1241800
- [11] Bijker EM, Bastiaens GJ, Teirlinck AC, van Gemert GJ, Graumans W, van de Vegte-Bolmer M, et al. Protection against malaria after immunization by chloroquine prophylaxis and sporozoites is mediated by preerythrocytic immunity. *Proc Natl Acad Sci U S A.* 2013;110:7862–7. DOI: 10.1073/pnas.1220360110

- [12] Roestenberg M, McCall M, Hopman J, Wiersma J, Luty AJ, van Gemert GJ, et al. Protection against a malaria challenge by sporozoite inoculation. *N Engl J Med*. 2009;361:468–77. DOI: 10.1056/NEJMoa0805832
- [13] Behet MC, Foquet L, van Gemert GJ, Bijker EM, Meuleman P, Leroux-Roels G, et al. Sporozoite immunization of human volunteers under chemoprophylaxis induces functional antibodies against pre-erythrocytic stages of *Plasmodium falciparum*. *Malar J*. 2014;13:136. DOI: 10.1186/1475-2875-13-136
- [14] Bijker EM, Schats R, Obiero JM, Behet MC, van Gemert GJ, van de Vegte-Bolmer M, et al. Sporozoite immunization of human volunteers under mefloquine prophylaxis is safe, immunogenic and protective: a double-blind randomized controlled clinical trial. *PLoS One*. 2014;9:e112910. DOI: 10.1371/journal.pone.0112910
- [15] Schats R, Bijker EM, van Gemert GJ, Graumans W, van de Vegte-Bolmer M, van Lieshout L, et al. Heterologous protection against malaria after immunization with *Plasmodium falciparum* sporozoites. *PLoS One*. 2015;10:e0124243. DOI: 10.1371/journal.pone.0124243
- [16] Silvie O, Semblat JP, Franetich JF, Hannoun L, Eling W, Mazier D. Effects of irradiation on *Plasmodium falciparum* sporozoite hepatic development: implications for the design of pre-erythrocytic malaria vaccines. *Parasite Immunol*. 2002;24:221–3.
- [17] Suhrbier A, Winger LA, Castellano E, Sinden RE. Survival and antigenic profile of irradiated malarial sporozoites in infected liver cells. *Infect Immun*. 1990;58:2834–9.
- [18] Doolan DL, Hoffman SL. The complexity of protective immunity against liver-stage malaria. *J Immunol*. 2000;165:1453–62.
- [19] Perlaza BL, Sauzet JP, Brahimi K, BenMohamed L, Druilhe P. Interferon-gamma, a valuable surrogate marker of *Plasmodium falciparum* pre-erythrocytic stages protective immunity. *Malar J*. 2011;10:27. DOI: 10.1186/1475-2875-10-27
- [20] Dups JN, Pepper M, Cockburn IA. Antibody and B cell responses to *Plasmodium* sporozoites. *Front Microbiol*. 2014;5:625. DOI: 10.3389/fmicb.2014.00625
- [21] Fernandez-Arias C, Mashoof S, Huang J, Tsuji M. Circumsporozoite protein as a potential target for antimalarials. *Expert Rev Anti Infect Ther*. 2015;13:923–6. DOI: 10.1586/14787210.2015.1058709
- [22] Rodrigues M, Nussenzweig RS, Zavala F. The relative contribution of antibodies, CD4+ and CD8+ T cells to sporozoite-induced protection against malaria. *Immunology*. 1993;80:1–5.
- [23] Schofield L, Villaquiran J, Ferreira A, Schellekens H, Nussenzweig R, Nussenzweig V. Gamma interferon, CD8+ T cells and antibodies required for immunity to malaria sporozoites. *Nature*. 1987;330:664–6. DOI: 10.1038/330664a0

- [24] Weiss WR, Sedegah M, Beaudoin RL, Miller LH, Good MF. CD8+ T cells (cytotoxic/suppressors) are required for protection in mice immunized with malaria sporozoites. *Proc Natl Acad Sci U S A*. 1988;85:573–6.
- [25] Todryk SM, Walther M. Building better T-cell-inducing malaria vaccines. *Immunology*. 2005;115:163–9. DOI: 10.1111/j.1365-2567.2005.02154.x
- [26] Tsuji M, Miyahira Y, Nussenzweig RS, Aguet M, Reichel M, Zavala F. Development of antimalaria immunity in mice lacking IFN-gamma receptor. *J Immunol*. 1995;154:5338–44.
- [27] Huang J, Tsao T, Zhang M, Rai U, Tsuji M, Li X. A sufficient role of MHC class I molecules on hepatocytes in anti-plasmodial activity of CD8 (+) T cells in vivo. *Front Microbiol*. 2015;6:69. DOI: 10.3389/fmicb.2015.00069
- [28] Lalvani A, Aidoo M, Allsopp CE, Plebanski M, Whittle HC, Hill AV. An HLA-based approach to the design of a CTL-inducing vaccine against *Plasmodium falciparum*. *Res Immunol*. 1994;145:461–8.
- [29] Overstreet MG, Cockburn IA, Chen YC, Zavala F. Protective CD8 T cells against *Plasmodium* liver stages: immunobiology of an 'unnatural' immune response. *Immunol Rev*. 2008;225:272–83. DOI: 10.1111/j.1600-065X.2008.00671.x
- [30] Tsuji M. A retrospective evaluation of the role of T cells in the development of malaria vaccine. *Exp Parasitol*. 2010;126:421–5. DOI: 10.1016/j.exppara.2009.11.009
- [31] Oliveira GA, Kumar KA, Calvo-Calle JM, Othoro C, Altszuler D, Nussenzweig V, et al. Class II-restricted protective immunity induced by malaria sporozoites. *Infect Immun*. 2008;76:1200–6. DOI: 10.1128/iai.00566-07
- [32] Charoenvit Y, Majam VF, Corradin G, Sacci JB Jr, Wang R, Doolan DL, et al. CD4(+) T-cell- and gamma interferon-dependent protection against murine malaria by immunization with linear synthetic peptides from a *Plasmodium yoelii* 17-kilodalton hepatocyte erythrocyte protein. *Infect Immun*. 1999;67:5604–14.
- [33] Longley RJ, Salman AM, Cottingham MG, Ewer K, Janse CJ, Khan SM, et al. Comparative assessment of vaccine vectors encoding ten malaria antigens identifies two protective liver-stage candidates. *Sci Rep*. 2015;5:11820. DOI: 10.1038/srep11820
- [34] Tsuji M, Romero P, Nussenzweig RS, Zavala F. CD4+ cytolytic T cell clone confers protection against murine malaria. *J Exp Med*. 1990;172:1353–7.
- [35] Weiss WR, Sedegah M, Berzofsky JA, Hoffman SL. The role of CD4+ T cells in immunity to malaria sporozoites. *J Immunol*. 1993;151:2690–8.
- [36] Aidoo M, Lalvani A, Allsopp CE, Plebanski M, Meisner SJ, Krausa P, et al. Identification of conserved antigenic components for a cytotoxic T lymphocyte-inducing vaccine against malaria. *Lancet*. 1995;345:1003–7.

- [37] Doolan DL, Hoffman SL, Southwood S, Wentworth PA, Sidney J, Chesnut RW, et al. Degenerate cytotoxic T cell epitopes from *P. falciparum* restricted by multiple HLA-A and HLA-B supertype alleles. *Immunity*. 1997;7:97–112.
- [38] Doolan DL, Southwood S, Chesnut R, Appella E, Gomez E, Richards A, et al. HLA-DR-promiscuous T cell epitopes from *Plasmodium falciparum* pre-erythrocytic-stage antigens restricted by multiple HLA class II alleles. *J Immunol*. 2000;165:1123–37.
- [39] Thomson SA, Khanna R, Gardner J, Burrows SR, Coupar B, Moss DJ, et al. Minimal epitopes expressed in a recombinant polyepitope protein are processed and presented to CD8+ cytotoxic T cells: implications for vaccine design. *Proc Natl Acad Sci U S A*. 1995;92:5845–9.
- [40] Whitton JL, Sheng N, Oldstone MB, McKee TA. A “string-of-beads” vaccine, comprising linked minigenes, confers protection from lethal-dose virus challenge. *J Virol*. 1993;67:348–52.
- [41] Gilbert SC, Plebanski M, Harris SJ, Allsopp CE, Thomas R, Layton GT, et al. A protein particle vaccine containing multiple malaria epitopes. *Nat Biotechnol*. 1997;15:1280–4. DOI: 10.1038/nbt1197-1280
- [42] Dunachie SJ, Walther M, Epstein JE, Keating S, Berthoud T, Andrews L, et al. A DNA prime-modified vaccinia virus ankara boost vaccine encoding thrombospondin-related adhesion protein but not circumsporozoite protein partially protects healthy malaria-naive adults against *Plasmodium falciparum* sporozoite challenge. *Infect Immun*. 2006;74:5933–42. DOI: 10.1128/iai.00590-06
- [43] Kimani D, Jagne YJ, Cox M, Kimani E, Bliss CM, Gitau E, et al. Translating the immunogenicity of prime-boost immunization with ChAd63 and MVA ME-TRAP from malaria naive to malaria-endemic populations. *Mol Ther*. 2014;22:1992–03. DOI: 10.1038/mt.2014.109
- [44] McConkey SJ, Reece WH, Moorthy VS, Webster D, Dunachie S, Butcher G, et al. Enhanced T-cell immunogenicity of plasmid DNA vaccines boosted by recombinant modified vaccinia virus Ankara in humans. *Nat Med*. 2003;9:729–35. DOI: 10.1038/nm881
- [45] Vuola JM, Keating S, Webster DP, Berthoud T, Dunachie S, Gilbert SC, et al. Differential immunogenicity of various heterologous prime-boost vaccine regimens using DNA and viral vectors in healthy volunteers. *J Immunol*. 2005;174:449–55.
- [46] Shevach EM. Regulatory T cells in autoimmunity*. *Annu Rev Immunol*. 2000;18:423–49. DOI: 10.1146/annurev.immunol.18.1.423
- [47] Coleman MM, Keane J, Mills KH. Editorial: Tregs and BCG-dangerous liaisons in TB. *J Leukoc Biol*. 2010;88:1067–9. DOI: 10.1189/jlb.0710419

- [48] Mills KH. Designer adjuvants for enhancing the efficacy of infectious disease and cancer vaccines based on suppression of regulatory T cell induction. *Immunol Lett.* 2009;122:108–11. DOI: 10.1016/j.imlet.2008.11.007
- [49] Ndure J, Flanagan KL. Targeting regulatory T cells to improve vaccine immunogenicity in early life. *Front Microbiol.* 2014;5:477. DOI: 10.3389/fmicb.2014.00477
- [50] Chia WN, Goh YS, Renia L. Novel approaches to identify protective malaria vaccine candidates. *Front Microbiol.* 2014;5:586. DOI: 10.3389/fmicb.2014.00586
- [51] Fidock DA, Bottius E, Brahimi K, Moelans II, Aikawa M, Konings RN, et al. Cloning and characterization of a novel *Plasmodium falciparum* sporozoite surface antigen, STARP. *Mol Biochem Parasitol.* 1994;64:219–32.
- [52] Guerin-Marchand C, Druilhe P, Galey B, Londono A, Patarapotikul J, Beaudoin RL, et al. A liver-stage-specific antigen of *Plasmodium falciparum* characterized by gene cloning. *Nature.* 1987;329:164–7. DOI: 10.1038/329164a0
- [53] Daubersies P, Thomas AW, Millet P, Brahimi K, Langermans JA, Ollomo B, et al. Protection against *Plasmodium falciparum* malaria in chimpanzees by immunization with the conserved pre-erythrocytic liver-stage antigen 3. *Nat Med.* 2000;6:1258–63. DOI: 10.1038/81366
- [54] Trieu A, Kayala MA, Burk C, Molina DM, Freilich DA, Richie TL, et al. Sterile protective immunity to malaria is associated with a panel of novel *P. falciparum* antigens. *Mol Cell Proteomics.* 2011;10:M111.007948. DOI: 10.1074/mcp.M111.007948
- [55] Doolan DL, Southwood S, Freilich DA, Sidney J, Graber NL, Shatney L, et al. Identification of *Plasmodium falciparum* antigens by antigenic analysis of genomic and proteomic data. *Proc Natl Acad Sci U S A.* 2003;100:9952–7. DOI: 10.1073/pnas.1633254100
- [56] Aguiar JC, Bolton J, Wanga J, Sacci JB, Iriko H, Mazeika JK, et al. Discovery of novel *Plasmodium falciparum* pre-erythrocytic antigens for vaccine development. *PLoS One.* 2015;10:e0136109. DOI: 10.1371/journal.pone.0136109
- [57] Ishino T, Yano K, Chinzei Y, Yuda M. Cell-passage activity is required for the malarial parasite to cross the liver sinusoidal cell layer. *PLoS Biol.* 2004;2:E4. DOI: 10.1371/journal.pbio.0020004
- [58] Kaiser K, Matuschewski K, Camargo N, Ross J, Kappe SH. Differential transcriptome profiling identifies *Plasmodium* genes encoding pre-erythrocytic stage-specific proteins. *Mol Microbiol.* 2004;51:1221–32. DOI: 10.1046/j.1365-2958.2003.03909.x
- [59] Lasonder E, Janse CJ, van Gemert GJ, Mair GR, Vermunt AM, Douradinha BG, et al. Proteomic profiling of *Plasmodium* sporozoite maturation identifies new proteins essential for parasite development and infectivity. *PLoS Pathog.* 2008;4:e1000195. DOI: 10.1371/journal.ppat.1000195

- [60] Matuschewski K, Ross J, Brown SM, Kaiser K, Nussenzweig V, Kappe SH. Infectivity-associated changes in the transcriptional repertoire of the malaria parasite sporozoite stage. *J Biol Chem*. 2002;277:41948–53. DOI: 10.1074/jbc.M207315200
- [61] Siau A, Silvie O, Franetich JF, Yalaoui S, Marinach C, Hannoun L, et al. Temperature shift and host cell contact up-regulate sporozoite expression of *Plasmodium falciparum* genes involved in hepatocyte infection. *PLoS Pathog*. 2008;4:e1000121. DOI: 10.1371/journal.ppat.1000121
- [62] Florens L, Washburn MP, Raine JD, Anthony RM, Grainger M, Haynes JD, et al. A proteomic view of the *Plasmodium falciparum* life cycle. *Nature*. 2002;419:520–6. DOI: 10.1038/nature01107
- [63] Tarun AS, Peng X, Dumpit RF, Ogata Y, Silva-Rivera H, Camargo N, et al. A combined transcriptome and proteome survey of malaria parasite liver stages. *Proc Natl Acad Sci U S A*. 2008;105:305–10. DOI: 10.1073/pnas.0710780104
- [64] Cummings JF, Spring MD, Schwenk RJ, Ockenhouse CF, Kester KE, Polhemus ME, et al. Recombinant Liver Stage Antigen-1 (LSA-1) formulated with AS01 or AS02 is safe, elicits high titer antibody and induces IFN-gamma/IL-2 CD4+ T cells but does not protect against experimental *Plasmodium falciparum* infection. *Vaccine*. 2010;28:5135–44. DOI: 10.1016/j.vaccine.2009.08.046
- [65] Bergmann-Leitner ES, Mease RM, De La Vega P, Savranskaya T, Polhemus M, Ockenhouse C, et al. Immunization with pre-erythrocytic antigen CelTOS from *Plasmodium falciparum* elicits cross-species protection against heterologous challenge with *Plasmodium berghei*. *PLoS One*. 2010;5:e12294. DOI: 10.1371/journal.pone.0012294
- [66] BenMohamed L, Thomas A, Druilhe P. Long-term multi-epitopic cytotoxic-T-lymphocyte responses induced in chimpanzees by combinations of *Plasmodium falciparum* liver-stage peptides and lipopeptides. *Infect Immun*. 2004;72:4376–84. DOI: 10.1128/iai.72.8.4376-4384.2004
- [67] Capone S, Reyes-Sandoval A, Naddeo M, Siani L, Ammendola V, Rollier CS, et al. Immune responses against a liver-stage malaria antigen induced by simian adenoviral vector AdCh63 and MVA prime-boost immunisation in non-human primates. *Vaccine*. 2010;29:256–65. DOI: 10.1016/j.vaccine.2010.10.041
- [68] Ferraro B, Talbott KT, Balakrishnan A, Cisper N, Morrow MP, Hutnick NA, et al. Inducing humoral and cellular responses to multiple sporozoite and liver-stage malaria antigens using exogenous plasmid DNA. *Infect Immun*. 2013;81:3709–20. DOI: 10.1128/iai.00180-13
- [69] Adams EJ, Parham P. Species-specific evolution of MHC class I genes in the higher primates. *Immunol Rev*. 2001;183:41–64.

- [70] Otting N, Heijmans CM, Noort RC, de Groot NG, Doxiadis GG, van Rood JJ, et al. Unparalleled complexity of the MHC class I region in rhesus macaques. *Proc Natl Acad Sci U S A*. 2005;102:1626–31. DOI: 10.1073/pnas.0409084102
- [71] Shen S, Pyo CW, Vu Q, Wang R, Geraghty DE. The essential detail: the genetics and genomics of the primate immune response. *Ilar j*. 2013;54:181–95. DOI: 10.1093/ilar/ilt043
- [72] Danner R, Chaudhari SN, Rosenberger J, Surls J, Richie TL, Brumeanu TD, et al. Expression of HLA class II molecules in humanized NOD.Rag1KO.IL2RgcKO mice is critical for development and function of human T and B cells. *PLoS One*. 2011;6:e19826. DOI: 10.1371/journal.pone.0019826
- [73] Huang J, Li X, Coelho-dos-Reis JG, Wilson JM, Tsuji M. An AAV vector-mediated gene delivery approach facilitates reconstitution of functional human CD8+ T cells in mice. *PLoS One*. 2014;9:e88205. DOI: 10.1371/journal.pone.0088205
- [74] Huang J, Li X, Coelho-dos-Reis JG, Zhang M, Mitchell R, Nogueira RT, et al. Human immune system mice immunized with *Plasmodium falciparum* circumsporozoite protein induce protective human humoral immunity against malaria. *J Immunol Methods*. 2015;427:42–50. DOI: 10.1016/j.jim.2015.09.005
- [75] Coelho-dos-Reis JGA, Li X, Silveira ELV, Mandraju R, Velmurugan S, Chakravarty S, et al. *Plasmodium falciparum* infection in “humanised liver” mice. *MalariaWorld J*. 2013;4:1–4.
- [76] Foquet L, Meuleman P, Hermsen CC, Sauerwein R, Leroux-Roels G. Assessment of parasite liver-stage burden in human-liver chimeric mice. *Methods Mol Biol*. 2015;1325:59–68. DOI: 10.1007/978-1-4939-2815-6_5
- [77] Morosan S, Hez-Deroubaix S, Lunel F, Renia L, Giannini C, Van Rooijen N, et al. Liver-stage development of *Plasmodium falciparum*, in a humanized mouse model. *J Infect Dis*. 2006;193:996–04. DOI: 10.1086/500840
- [78] Sack BK, Miller JL, Vaughan AM, Douglass A, Kaushansky A, Mikolajczak S, et al. Model for in vivo assessment of humoral protection against malaria sporozoite challenge by passive transfer of monoclonal antibodies and immune serum. *Infect Immun*. 2014;82:808–17. DOI: 10.1128/iai.01249-13
- [79] Vaughan AM, Kappe SH, Ploss A, Mikolajczak SA. Development of humanized mouse models to study human malaria parasite infection. *Future Microbiol*. 2012;7:657–65. DOI: 10.2217/fmb.12.27
- [80] Coppi A, Natarajan R, Pradel G, Bennett BL, James ER, Roggero MA, et al. The malaria circumsporozoite protein has two functional domains, each with distinct roles as sporozoites journey from mosquito to mammalian host. *J Exp Med*. 2011;208:341–56. DOI: 10.1084/jem.20101488

- [81] Rathore D, Sacchi JB, de la Vega P, McCutchan TF. Binding and invasion of liver cells by *Plasmodium falciparum* sporozoites. Essential involvement of the amino terminus of circumsporozoite protein. *J Biol Chem*. 2002;277:7092–8. DOI: 10.1074/jbc.M106862200
- [82] Suarez JE, Urquiza M, Puentes A, Garcia JE, Curtidor H, Ocampo M, et al. *Plasmodium falciparum* circumsporozoite (CS) protein peptides specifically bind to HepG2 cells. *Vaccine*. 2001;19:4487–95.
- [83] Coppi A, Pinzon-Ortiz C, Hutter C, Sinnis P. The Plasmodium circumsporozoite protein is proteolytically processed during cell invasion. *J Exp Med*. 2005;201:27–33. DOI: 10.1084/jem.20040989
- [84] Sinnis P, Clavijo P, Fenyo D, Chait BT, Cerami C, Nussenzweig V. Structural and functional properties of region II-plus of the malaria circumsporozoite protein. *J Exp Med*. 1994;180:297–06.
- [85] Dame JB, Williams JL, McCutchan TF, Weber JL, Wirtz RA, Hockmeyer WT, et al. Structure of the gene encoding the immunodominant surface antigen on the sporozoite of the human malaria parasite *Plasmodium falciparum*. *Science*. 1984;225:593–9.
- [86] Enea V, Ellis J, Zavala F, Arnot DE, Asavanich A, Masuda A, et al. DNA cloning of *Plasmodium falciparum* circumsporozoite gene: amino acid sequence of repetitive epitope. *Science*. 1984;225:628–30.
- [87] Nardin EH, Nussenzweig V, Nussenzweig RS, Collins WE, Harinasuta KT, Tapchaisri P, et al. Circumsporozoite proteins of human malaria parasites *Plasmodium falciparum* and *Plasmodium vivax*. *J Exp Med*. 1982;156:20–30.
- [88] Zavala F, Tam JP, Hollingdale MR, Cochrane AH, Quakyi I, Nussenzweig RS, et al. Rationale for development of a synthetic vaccine against *Plasmodium falciparum* malaria. *Science*. 1985;228:1436–40.
- [89] Moreno A, Clavijo P, Edelman R, Davis J, Sztein M, Sinigaglia F, et al. CD4+ T cell clones obtained from *Plasmodium falciparum* sporozoite-immunized volunteers recognize polymorphic sequences of the circumsporozoite protein. *J Immunol*. 1993;151:489–99.
- [90] Calvo-Calle JM, Hammer J, Sinigaglia F, Clavijo P, Moya-Castro ZR, Nardin EH. Binding of malaria T cell epitopes to DR and DQ molecules in vitro correlates with immunogenicity in vivo: identification of a universal T cell epitope in the *Plasmodium falciparum* circumsporozoite protein. *J Immunol*. 1997;159:1362–73.
- [91] Huang J, Tsao T, Zhang M, Tsuji M. Circumsporozoite protein-specific K(d)-restricted CD8+ T cells mediate protective antimalaria immunity in sporozoite-immunized MHC-I-K(d) transgenic mice. *Mediators Inflamm*. 2014;2014:728939. DOI: 10.1155/2014/728939
- [92] Bailey JA, Mvalo T, Aragam N, Weiser M, Congdon S, Kamwendo D, et al. Use of massively parallel pyrosequencing to evaluate the diversity of and selection on

Plasmodium falciparum csp T-cell epitopes in Lilongwe, Malawi. J Infect Dis. 2012;206:580–7. DOI: 10.1093/infdis/jis329

- [93] Kester KE, Cummings JF, Ofori-Anyinam O, Ockenhouse CF, Krzych U, Moris P, et al. Randomized, double-blind, phase 2a trial of falciparum malaria vaccines RTS,S/AS01B and RTS,S/AS02A in malaria-naive adults: safety, efficacy, and immunologic associates of protection. J Infect Dis. 2009;200:337–46. DOI: 10.1086/600120
- [94] Calle JM, Nardin EH, Clavijo P, Boudin C, Stuber D, Takacs B, et al. Recognition of different domains of the *Plasmodium falciparum* CS protein by the sera of naturally infected individuals compared with those of sporozoite-immunized volunteers. J Immunol. 1992;149:2695–701.
- [95] Sun P, Schwenk R, White K, Stoute JA, Cohen J, Ballou WR, et al. Protective immunity induced with malaria vaccine, RTS,S, is linked to *Plasmodium falciparum* circumsporozoite protein-specific CD4+ and CD8+ T cells producing IFN-gamma. J Immunol. 2003;171:6961–7.
- [96] Stoute JA, Kester KE, Krzych U, Welde BT, Hall T, White K, et al. Long-term efficacy and immune responses following immunization with the RTS,S malaria vaccine. J Infect Dis. 1998;178:1139–44.
- [97] Olotu A, Lusingu J, Leach A, Lievens M, Vekemans J, Msham S, et al. Efficacy of RTS,S/AS01E malaria vaccine and exploratory analysis on anti-circumsporozoite antibody titres and protection in children aged 5–17 months in Kenya and Tanzania: a randomised controlled trial. Lancet Infect Dis. 2011;11:102–9. DOI: 10.1016/s1473-3099(10)70262-0
- [98] Herrington DA, Clyde DF, Losonsky G, Cortesia M, Murphy JR, Davis J, et al. Safety and immunogenicity in man of a synthetic peptide malaria vaccine against *Plasmodium falciparum* sporozoites. Nature. 1987;328:257–9. DOI: 10.1038/328257a0
- [99] Herrington DA, Clyde DF, Davis JR, Baqar S, Murphy JR, Cortese JF, et al. Human studies with synthetic peptide sporozoite vaccine (NANP)₃-TT and immunization with irradiated sporozoites. Bull World Health Organ. 1990;68 Suppl:33–7.
- [100] Moreno R, Jiang L, Moehle K, Zurbriggen R, Gluck R, Robinson JA, et al. Exploiting conformationally constrained peptidomimetics and an efficient human-compatible delivery system in synthetic vaccine design. Chembiochem. 2001;2:838–43.
- [101] Genton B, Pluschke G, Degen L, Kammer AR, Westerfeld N, Okitsu SL, et al. A randomized placebo-controlled phase Ia malaria vaccine trial of two virosome-formulated synthetic peptides in healthy adult volunteers. PLoS One. 2007;2:e1018. DOI: 10.1371/journal.pone.0001018
- [102] Okitsu SL, Silvie O, Westerfeld N, Curcic M, Kammer AR, Mueller MS, et al. A virosomal malaria peptide vaccine elicits a long-lasting sporozoite-inhibitory antibody

- response in a phase 1a clinical trial. *PLoS One*. 2007;2:e1278. DOI: 10.1371/journal.pone.0001278
- [103] Thompson FM, Porter DW, Okitsu SL, Westerfeld N, Vogel D, Todryk S, et al. Evidence of blood stage efficacy with a virosomal malaria vaccine in a phase IIa clinical trial. *PLoS One*. 2008;3:e1493. DOI: 10.1371/journal.pone.0001493
- [104] Liu W, Li Y, Learn GH, Rudicell RS, Robertson JD, Keele BF, et al. Origin of the human malaria parasite *Plasmodium falciparum* in gorillas. *Nature*. 2010;467:420–5. DOI: 10.1038/nature09442
- [105] Schofield L. The circumsporozoite protein of Plasmodium: a mechanism of immune evasion by the malaria parasite? *Bull World Health Organ*. 1990;68 Suppl:66–73.
- [106] Nelson RW, Beisang D, Tubo NJ, Dileepan T, Wiesner DL, Nielsen K, et al. T cell receptor cross-reactivity between similar foreign and self peptides influences naive cell population size and autoimmunity. *Immunity*. 2015;42:95–07. DOI: 10.1016/j.immuni.2014.12.022
- [107] Liu R, Moise L, Tassone R, Gutierrez AH, Terry FE, Sangare K, et al. H7N9 T-cell epitopes that mimic human sequences are less immunogenic and may induce Treg-mediated tolerance. *Hum Vaccin Immunother*. 2015;11:2241–52. DOI: 10.1080/21645515.2015.1052197
- [108] Nardin E. The past decade in malaria synthetic peptide vaccine clinical trials. *Hum Vaccin*. 2010;6:27–38.
- [109] Nardin EH, Calvo-Calle JM, Oliveira GA, Clavijo P, Nussenzweig R, Simon R, et al. *Plasmodium falciparum* polyoximes: highly immunogenic synthetic vaccines constructed by chemoselective ligation of repeat B-cell epitopes and a universal T-cell epitope of CS protein. *Vaccine*. 1998;16:590–600.
- [110] Nardin EH, Calvo-Calle JM, Oliveira GA, Nussenzweig RS, Schneider M, Tiercy JM, et al. A totally synthetic polyoxime malaria vaccine containing *Plasmodium falciparum* B cell and universal T cell epitopes elicits immune responses in volunteers of diverse HLA types. *J Immunol*. 2001;166:481–9.
- [111] Calvo-Calle JM, Oliveira GA, Nardin EH. Human CD4+ T cells induced by synthetic peptide malaria vaccine are comparable to cells elicited by attenuated *Plasmodium falciparum* sporozoites. *J Immunol*. 2005;175:7575–85.
- [112] Birkett A, Lyons K, Schmidt A, Boyd D, Oliveira GA, Siddique A, et al. A modified hepatitis B virus core particle containing multiple epitopes of the *Plasmodium falciparum* circumsporozoite protein provides a highly immunogenic malaria vaccine in preclinical analyses in rodent and primate hosts. *Infect Immun*. 2002;70:6860–70.
- [113] Gregson AL, Oliveira G, Othoro C, Calvo-Calle JM, Thorton GB, Nardin E, et al. Phase I trial of an alhydrogel adjuvanted hepatitis B core virus-like particle containing

- epitopes of *Plasmodium falciparum* circumsporozoite protein. PLoS One. 2008;3:e1556. DOI: 10.1371/journal.pone.0001556
- [114] Nardin EH, Oliveira GA, Calvo-Calle JM, Wetzel K, Maier C, Birkett AJ, et al. Phase I testing of a malaria vaccine composed of hepatitis B virus core particles expressing *Plasmodium falciparum* circumsporozoite epitopes. Infect Immun. 2004;72:6519–27. DOI: 10.1128/IAI.72.11.6519-6527.2004
- [115] Langermans JA, Schmidt A, Vervenne RA, Birkett AJ, Calvo-Calle JM, Hensmann M, et al. Effect of adjuvant on reactogenicity and long-term immunogenicity of the malaria Vaccine ICC-1132 in macaques. Vaccine. 2005;23:4935–43. DOI: 10.1016/j.vaccine.2005.05.036
- [116] Oliveira GA, Wetzel K, Calvo-Calle JM, Nussenzweig R, Schmidt A, Birkett A, et al. Safety and enhanced immunogenicity of a hepatitis B core particle *Plasmodium falciparum* malaria vaccine formulated in adjuvant Montanide ISA 720 in a phase I trial. Infect Immun. 2005;73:3587–97. DOI: 10.1128/IAI.73.6.3587-3597.2005
- [117] Walther M, Dunachie S, Keating S, Vuola JM, Berthoud T, Schmidt A, et al. Safety, immunogenicity and efficacy of a pre-erythrocytic malaria candidate vaccine, ICC-1132 formulated in Seppic ISA 720. Vaccine. 2005;23:857–64. DOI: 10.1016/j.vaccine.2004.08.020
- [118] Corradin G, Kajava AV, Verdini A. Long synthetic peptides for the production of vaccines and drugs: a technological platform coming of age. Sci Transl Med. 2010;2:50rv3. DOI: 10.1126/scitranslmed.3001387
- [119] Lopez JA, Weilenman C, Audran R, Roggero MA, Bonelo A, Tiercy JM, et al. A synthetic malaria vaccine elicits a potent CD8(+) and CD4(+) T lymphocyte immune response in humans. Implications for vaccination strategies. Eur J Immunol. 2001;31:1989–98. DOI: 10.1002/1521-4141(200107)31:7<1989::aid-immu1989>3.0.co;2-m
- [120] Audran R, Lurati-Ruiz F, Genton B, Blythman HE, Ofori-Anyinam O, Reymond C, et al. The synthetic *Plasmodium falciparum* circumsporozoite peptide PfCS102 as a malaria vaccine candidate: a randomized controlled phase I trial. PLoS One. 2009;4:e7304. DOI: 10.1371/journal.pone.0007304
- [121] Stoute JA, Slaoui M, Heppner DG, Momin P, Kester KE, Desmons P, et al. A preliminary evaluation of a recombinant circumsporozoite protein vaccine against *Plasmodium falciparum* malaria. RTS,S Malaria Vaccine Evaluation Group. N Engl J Med. 1997;336:86–91. DOI: 10.1056/NEJM199701093360202
- [122] Gordon DM, McGovern TW, Krzych U, Cohen JC, Schneider I, LaChance R, et al. Safety, immunogenicity, and efficacy of a recombinantly produced *Plasmodium falciparum* circumsporozoite protein-hepatitis B surface antigen subunit vaccine. J Infect Dis. 1995;171:1576–85.

- [123] Nardin EH, Oliveira GA, Calvo-Calle JM, Castro ZR, Nussenzweig RS, Schneckpeper B, et al. Synthetic malaria peptide vaccine elicits high levels of antibodies in vaccinees of defined HLA genotypes. *J Infect Dis.* 2000;182:1486–96. DOI: 10.1086/315871
- [124] Kester KE, McKinney DA, Tornieporth N, Ockenhouse CF, Heppner DG, Hall T, et al. Efficacy of recombinant circumsporozoite protein vaccine regimens against experimental *Plasmodium falciparum* malaria. *J Infect Dis.* 2001;183:640–7. DOI: 10.1086/318534
- [125] Whitacre DC, Espinosa DA, Peters CJ, Jones JE, Tucker AE, Peterson DL, et al. *P. falciparum* and *P. vivax* epitope-focused VLPs elicit sterile immunity to blood stage infections. *PLoS One.* 2015;10:e0124856. DOI: 10.1371/journal.pone.0124856
- [126] Carapau D, Mitchell R, Nacer A, Shaw A, Othoro C, Frevert U, et al. Protective humoral immunity elicited by a needle-free malaria vaccine comprised of a chimeric *Plasmodium falciparum* circumsporozoite protein and a Toll-like receptor 5 agonist, flagellin. *Infect Immun.* 2013;81:4350–62. DOI: 10.1128/iai.00263-13
- [127] Nacer A, Carapau D, Mitchell R, Meltzer A, Shaw A, Frevert U, et al. Imaging murine NALT following intranasal immunization with flagellin-modified circumsporozoite protein malaria vaccines. *Mucosal Immunol.* 2014;7:304–14. DOI: 10.1038/mi.2013.48
- [128] Othoro C, Johnston D, Lee R, Soverow J, Bystryn JC, Nardin E. Enhanced immunogenicity of *Plasmodium falciparum* peptide vaccines using a topical adjuvant containing a potent synthetic Toll-like receptor 7 agonist, imiquimod. *Infect Immun.* 2009;77:739–48. DOI: 10.1128/iai.00974-08
- [129] Hung IF, Zhang AJ, To KK, Chan JF, Li C, Zhu HS, et al. Immunogenicity of intradermal trivalent influenza vaccine with topical imiquimod: a double blind randomized controlled trial. *Clin Infect Dis.* 2014;59:1246–55. DOI: 10.1093/cid/ciu582
- [130] Hung IF, Zhang AJ, To KK, Chan JF, Li P, Wong TL, et al. Topical imiquimod before intradermal trivalent influenza vaccine for protection against heterologous non-vaccine and antigenically drifted viruses: a single-centre, double-blind, randomised, controlled phase 2b/3 trial. *Lancet Infect Dis.* 2016;16:209–18. DOI: 10.1016/s1473-3099(15)00354-0
- [131] Powell TJ, Tang J, Derome ME, Mitchell RA, Jacobs A, Deng Y, et al. *Plasmodium falciparum* synthetic LbL microparticle vaccine elicits protective neutralizing antibody and parasite-specific cellular immune responses. *Vaccine.* 2013;31:1898–904. DOI: 10.1016/j.vaccine.2013.02.027
- [132] Guo Q, Dasgupta D, Doll TA, Burkhard P, Lanar DE. Expression, purification and refolding of a self-assembling protein nanoparticle (SAPN) malaria vaccine. *Methods.* 2013;60:242–7. DOI: 10.1016/j.ymeth.2013.03.025
- [133] Kaba SA, McCoy ME, Doll TA, Brando C, Guo Q, Dasgupta D, et al. Protective antibody and CD8⁺ T-cell responses to the *Plasmodium falciparum* circumsporozoite protein induced by a nanoparticle vaccine. *PLoS One.* 2012;7:e48304. DOI: 10.1371/journal.pone.0048304

- [134] Espinosa DA, Gutierrez GM, Rojas-Lopez M, Noe AR, Shi L, Tse SW, et al. Proteolytic cleavage of the *Plasmodium falciparum* circumsporozoite protein is a target of protective antibodies. *J Infect Dis.* 2015;212:1111–9. DOI: 10.1093/infdis/jiv154
- [135] Kariu T, Ishino T, Yano K, Chinzei Y, Yuda M. CelTOS, a novel malarial protein that mediates transmission to mosquito and vertebrate hosts. *Mol Microbiol.* 2006;59:1369–79. DOI: 10.1111/j.1365-2958.2005.05024.x
- [136] Anum D, Kusi KA, Ganeshan H, Hollingdale MR, Ofori MF, Koram KA, et al. Measuring naturally acquired ex vivo IFN-gamma responses to *Plasmodium falciparum* cell-traversal protein for ookinetes and sporozoites (CelTOS) in Ghanaian adults. *Malar J.* 2015;14:20. DOI: 10.1186/s12936-014-0539-5
- [137] Mishra S, Rai U, Shiratsuchi T, Li X, Vanloubbeeck Y, Cohen J, et al. Identification of non-CSP antigens bearing CD8 epitopes in mice immunized with irradiated sporozoites. *Vaccine.* 2011;29:7335–42. DOI: 10.1016/j.vaccine.2011.07.081
- [138] Bergmann-Leitner ES, Legler PM, Savranskaya T, Ockenhouse CF, Angov E. Cellular and humoral immune effector mechanisms required for sterile protection against sporozoite challenge induced with the novel malaria vaccine candidate CelTOS. *Vaccine.* 2011;29:5940–9. DOI: 10.1016/j.vaccine.2011.06.053
- [139] Fox CB, Baldwin SL, Vedvick TS, Angov E, Reed SG. Effects on immunogenicity by formulations of emulsion-based adjuvants for malaria vaccines. *Clin Vaccine Immunol.* 2012;19:1633–40. DOI: 10.1128/cvi.00235-12
- [140] Bergmann-Leitner ES, Hosie H, Trichilo J, Deriso E, Ranallo RT, Alefantis T, et al. Self-adjuvanting bacterial vectors expressing pre-erythrocytic antigens induce sterile protection against malaria. *Front Immunol.* 2013;4:176. DOI: 10.3389/fimmu.2013.00176
- [141] Bergmann-Leitner ES, Li Q, Caridha D, O’Neil MT, Ockenhouse CF, Hickman M, et al. Protective immune mechanisms against pre-erythrocytic forms of *Plasmodium berghei* depend on the target antigen. *Trials Vaccinol.* 2014;3:6-10. DOI: 10.1016/j.trivac.2013.11.002
- [142] Hope IA, Mackay M, Hyde JE, Goman M, Scaife J. The gene for an exported antigen of the malaria parasite *Plasmodium falciparum* cloned and expressed in *Escherichia coli*. *Nucleic Acids Res.* 1985;13:369–79.
- [143] Coppel RL, Favaloro JM, Crewther PE, Burkot TR, Bianco AE, Stahl HD, et al. A blood stage antigen of *Plasmodium falciparum* shares determinants with the sporozoite coat protein. *Proc Natl Acad Sci U S A.* 1985;82:5121–5.
- [144] Doolan DL, Hedstrom RC, Rogers WO, Charoenvit Y, Rogers M, de la Vega P, et al. Identification and characterization of the protective hepatocyte erythrocyte protein 17 kDa gene of *Plasmodium yoelii*, homolog of *Plasmodium falciparum* exported protein 1. *J Biol Chem.* 1996;271:17861–8.

- [145] Kara U, Murray B, Pam C, Lahnstein J, Gould H, Kidson C, et al. Chemical characterization of the parasitophorous vacuole membrane antigen QF 116 from *Plasmodium falciparum*. *Mol Biochem Parasitol*. 1990;38:19–23.
- [146] Kara UA, Stenzel DJ, Ingram LT, Bushell GR, Lopez JA, Kidson C. Inhibitory monoclonal antibody against a (myristylated) small-molecular-weight antigen from *Plasmodium falciparum* associated with the parasitophorous vacuole membrane. *Infect Immun*. 1988;56:903–9.
- [147] Simmons D, Woollett G, Bergin-Cartwright M, Kay D, Scaife J. A malaria protein exported into a new compartment within the host erythrocyte. *Embo j*. 1987;6:485–91.
- [148] Sturchler D, Just M, Berger R, Reber-Liske R, Matile H, Etlinger H, et al. Evaluation of 5.1-[NANP]19, a recombinant *Plasmodium falciparum* vaccine candidate, in adults. *Trop Geogr Med*. 1992;44:9–14.
- [149] Doolan DL, Sedegah M, Hedstrom RC, Hobart P, Charoenvit Y, Hoffman SL. Circumventing genetic restriction of protection against malaria with multigene DNA immunization: CD8+ cell-, interferon gamma-, and nitric oxide-dependent immunity. *J Exp Med*. 1996;183:1739–46.
- [150] Lisewski AM, Quiros JP, Ng CL, Adikesavan AK, Miura K, Putluri N, et al. Super-genomic network compression and the discovery of EXP1 as a glutathione transferase inhibited by artesunate. *Cell*. 2014;158:916–28. DOI: 10.1016/j.cell.2014.07.011
- [151] Dodoo D, Hollingdale MR, Anum D, Koram KA, Gyan B, Akanmori BD, et al. Measuring naturally acquired immune responses to candidate malaria vaccine antigens in Ghanaian adults. *Malar J*. 2011;10:168. DOI: 10.1186/1475-2875-10-168
- [152] Meraldi V, Nebie I, Moret R, Cuzin-Ouattara N, Thiocone A, Doumbo O, et al. Recognition of synthetic polypeptides corresponding to the N- and C-terminal fragments of *Plasmodium falciparum* Exp-1 by T-cells and plasma from human donors from African endemic areas. *Parasite Immunol*. 2002;24:141–50.
- [153] Porter KR, Aguiar J, Richards A, Sandjaya B, Ignatias H, Hadiputranto H, et al. Immune response against the exp-1 protein of *Plasmodium falciparum* results in antibodies that cross-react with human T-cell lymphotropic virus type 1 proteins. *Clin Diagn Lab Immunol*. 1998;5:721–4.
- [154] Meraldi V, Nebie I, Tiono AB, Diallo D, Sanogo E, Theisen M, et al. Natural antibody response to *Plasmodium falciparum* Exp-1, MSP-3 and GLURP long synthetic peptides and association with protection. *Parasite Immunol*. 2004;26:265–72. DOI: 10.1111/j.0141-9838.2004.00705.x
- [155] Aidoo M, Lalvani A, Gilbert SC, Hu JT, Daubersies P, Hurt N, et al. Cytotoxic T-lymphocyte epitopes for HLA-B53 and other HLA types in the malaria vaccine candidate liver-stage antigen 3. *Infect Immun*. 2000;68:227–32.
- [156] Lyke KE, Burges RB, Cissoko Y, Sangare L, Kone A, Dao M, et al. HLA-A2 supertype-restricted cell-mediated immunity by peripheral blood mononuclear cells derived from

- Malian children with severe or uncomplicated *Plasmodium falciparum* malaria and healthy controls. *Infect Immun.* 2005;73:5799–808. DOI: 10.1128/iai.73.9.5799-5808.2005
- [157] Caspers P, Etlinger H, Matile H, Pink JR, Stuber D, Takacs B. A *Plasmodium falciparum* malaria vaccine candidate which contains epitopes from the circumsporozoite protein and a blood stage antigen, 5.1. *Mol Biochem Parasitol.* 1991;47:143–50.
- [158] Reber-Liske R, Salako LA, Matile H, Sowunmi A, Sturchler D. [NANP]19-5.1. A malaria vaccine field trial in Nigerian children. *Trop Geogr Med.* 1995;47:61–3.
- [159] Meis JF, Rijntjes PJ, Verhave JP, Ponnudurai T, Hollingdale MR, Smith JE, et al. Fine structure of the malaria parasite *Plasmodium falciparum* in human hepatocytes in vitro. *Cell Tissue Res.* 1986;244:345–50.
- [160] Hill AV, Allsopp CE, Kwiatkowski D, Anstey NM, Twumasi P, Rowe PA, et al. Common west African HLA antigens are associated with protection from severe malaria. *Nature.* 1991;352:595–600. DOI: 10.1038/352595a0
- [161] Hill AV, Elvin J, Willis AC, Aidoo M, Allsopp CE, Gotch FM, et al. Molecular analysis of the association of HLA-B53 and resistance to severe malaria. *Nature.* 1992;360:434–9. DOI: 10.1038/360434a0
- [162] Zhu J, Hollingdale MR. Structure of *Plasmodium falciparum* liver stage antigen-1. *Mol Biochem Parasitol.* 1991;48:223–6.
- [163] Walliker D, Quakyi IA, Wellems TE, McCutchan TF, Szarfman A, London WT, et al. Genetic analysis of the human malaria parasite *Plasmodium falciparum*. *Science.* 1987;236:1661–6.
- [164] Joshi SK, Bharadwaj A, Chatterjee S, Chauhan VS. Analysis of immune responses against T- and B-cell epitopes from *Plasmodium falciparum* liver-stage antigen 1 in rodent malaria models and malaria-exposed human subjects in India. *Infect Immun.* 2000;68:141–50.
- [165] Bucci K, Kastens W, Hollingdale MR, Shankar A, Alpers MP, King CL, et al. Influence of age and HLA type on interferon-gamma (IFN-gamma) responses to a naturally occurring polymorphic epitope of *Plasmodium falciparum* liver stage antigen-1 (LSA-1). *Clin Exp Immunol.* 2000;122:94–100.
- [166] Connelly M, King CL, Bucci K, Walters S, Genton B, Alpers MP, et al. T-cell immunity to peptide epitopes of liver-stage antigen 1 in an area of Papua New Guinea in which malaria is holoendemic. *Infect Immun.* 1997;65:5082–7.
- [167] John CC, Moormann AM, Sumba PO, Ofulla AV, Pregibon DC, Kazura JW. Gamma interferon responses to *Plasmodium falciparum* liver-stage antigen 1 and thrombospondin-related adhesive protein and their relationship to age, transmission intensity, and protection against malaria. *Infect Immun.* 2004;72:5135–42. DOI: 10.1128/iai.72.9.5135-5142.2004

- [168] John CC, Sumba PO, Ouma JH, Nahlen BL, King CL, Kazura JW. Cytokine responses to *Plasmodium falciparum* liver-stage antigen 1 vary in rainy and dry seasons in highland Kenya. *Infect Immun*. 2000;68:5198–204.
- [169] Luty AJ, Lell B, Schmidt-Ott R, Lehman LG, Luckner D, Greve B, et al. Interferon-gamma responses are associated with resistance to reinfection with *Plasmodium falciparum* in young African children. *J Infect Dis*. 1999;179:980–8. DOI: 10.1086/314689
- [170] Fidock DA, Gras-Masse H, Lepers JP, Brahim K, Benmohamed L, Mellouk S, et al. *Plasmodium falciparum* liver stage antigen-1 is well conserved and contains potent B and T cell determinants. *J Immunol*. 1994;153:190–204.
- [171] Crompton PD, Kayala MA, Traore B, Kayentao K, Ongoiba A, Weiss GE, et al. A prospective analysis of the Ab response to *Plasmodium falciparum* before and after a malaria season by protein microarray. *Proc Natl Acad Sci U S A*. 2010;107:6958–63. DOI: 10.1073/pnas.1001323107
- [172] John CC, Moormann AM, Pregibon DC, Sumba PO, McHugh MM, Narum DL, et al. Correlation of high levels of antibodies to multiple pre-erythrocytic *Plasmodium falciparum* antigens and protection from infection. *Am J Trop Med Hyg*. 2005;73:222–8.
- [173] John CC, Tande AJ, Moormann AM, Sumba PO, Lanar DE, Min XM, et al. Antibodies to pre-erythrocytic *Plasmodium falciparum* antigens and risk of clinical malaria in Kenyan children. *J Infect Dis*. 2008;197:519–26. DOI: 10.1086/526787
- [174] Krzych U, Lyon JA, Jareed T, Schneider I, Hollingdale MR, Gordon DM, et al. T lymphocytes from volunteers immunized with irradiated *Plasmodium falciparum* sporozoites recognize liver and blood stage malaria antigens. *J Immunol*. 1995;155:4072–7.
- [175] Nahrendorf W, Scholzen A, Bijker EM, Teirlinck AC, Bastiaens GJ, Schats R, et al. Memory B-cell and antibody responses induced by *Plasmodium falciparum* sporozoite immunization. *J Infect Dis*. 2014;210:1981–90. DOI: 10.1093/infdis/jiu354
- [176] Aurrecochea C, Brestelli J, Brunk BP, Dommer J, Fischer S, Gajria B, et al. PlasmoDB: a functional genomic database for malaria parasites. *Nucleic Acids Res*. 2009;37:D539–43. DOI: 10.1093/nar/gkn814
- [177] Benmohamed L, Thomas A, Bossus M, Brahim K, Wubben J, Gras-Masse H, et al. High immunogenicity in chimpanzees of peptides and lipopeptides derived from four new *Plasmodium falciparum* pre-erythrocytic molecules. *Vaccine*. 2000;18:2843–55.
- [178] Perlaza BL, Arevalo-Herrera M, Brahim K, Quintero G, Palomino JC, Gras-Masse H, et al. Immunogenicity of four *Plasmodium falciparum* preerythrocytic antigens in *Aotus lemurinus* monkeys. *Infect Immun*. 1998;66:3423–8.
- [179] Jiang G, Charoenvit Y, Moreno A, Baraceros MF, Banania G, Richie N, et al. Induction of multi-antigen multi-stage immune responses against *Plasmodium falciparum* in rhesus

- monkeys, in the absence of antigen interference, with heterologous DNA prime/poxvirus boost immunization. *Malar J.* 2007;6:135. DOI: 10.1186/1475-2875-6-135
- [180] Pichyangkul S, Kum-Arb U, Yongvanitchit K, Limsalakpetch A, Gettayacamin M, Lanar DE, et al. Preclinical evaluation of the safety and immunogenicity of a vaccine consisting of *Plasmodium falciparum* liver-stage antigen 1 with adjuvant AS01B administered alone or concurrently with the RTS,S/AS01B vaccine in rhesus primates. *Infect Immun.* 2008;76:229–38. DOI: 10.1128/iai.00977-07
- [181] Hillier CJ, Ware LA, Barbosa A, Angov E, Lyon JA, Heppner DG, et al. Process development and analysis of liver-stage antigen 1, a preerythrocyte-stage protein-based vaccine for *Plasmodium falciparum*. *Infect Immun.* 2005;73:2109–15. DOI: 10.1128/iai.73.4.2109-2115.2005
- [182] Brando C, Ware LA, Freyberger H, Kathcart A, Barbosa A, Cayphas S, et al. Murine immune responses to liver-stage antigen 1 protein FMP011, a malaria vaccine candidate, delivered with adjuvant AS01B or AS02A. *Infect Immun.* 2007;75:838–45. DOI: 10.1128/iai.01075-06
- [183] Doolan DL, Mu Y, Unal B, Sundaresh S, Hirst S, Valdez C, et al. Profiling humoral immune responses to *P. falciparum* infection with protein microarrays. *Proteomics.* 2008;8:4680–94. DOI: 10.1002/pmic.200800194
- [184] Prieur E, Druilhe P. The malaria candidate vaccine liver stage antigen-3 is highly conserved in *Plasmodium falciparum* isolates from diverse geographical areas. *Malar J.* 2009;8:247. DOI: 10.1186/1475-2875-8-247
- [185] Toure-Balde A, Perlaza BL, Sauzet JP, Ndiaye M, Aribot G, Tall A, et al. Evidence for multiple B- and T-cell epitopes in *Plasmodium falciparum* liver-stage antigen 3. *Infect Immun.* 2009;77:1189–96. DOI: 10.1128/iai.00780-07
- [186] Daubersies P, Ollomo B, Sauzet JP, Brahimi K, Perlaza BL, Eling W, et al. Genetic immunisation by liver stage antigen 3 protects chimpanzees against malaria despite low immune responses. *PLoS One.* 2008;3:e2659. DOI: 10.1371/journal.pone.0002659
- [187] Perlaza BL, Valencia AZ, Zapata C, Castellanos A, Sauzet JP, Blanc C, et al. Protection against *Plasmodium falciparum* challenge induced in Aotus monkeys by liver-stage antigen-3-derived long synthetic peptides. *Eur J Immunol.* 2008;38:2610–15. DOI: 10.1002/eji.200738055
- [188] Schwartz L, Brown GV, Genton B, Moorthy VS. A review of malaria vaccine clinical projects based on the WHO rainbow table. *Malar J.* 2012;11:11. DOI: 10.1186/1475-2875-11-11
- [189] Fidock DA, Sallenave-Sales S, Sherwood JA, Gachihi GS, Ferreira-da-Cruz MF, Thomas AW, et al. Conservation of the *Plasmodium falciparum* sporozoite surface protein gene, STARP, in field isolates and distinct species of Plasmodium. *Mol Biochem Parasitol.* 1994;67:255–67.

- [190] Ambrosino E, Dumoulin C, Orlandi-Pradines E, Remoue F, Toure-Balde A, Tall A, et al. A multiplex assay for the simultaneous detection of antibodies against 15 *Plasmodium falciparum* and *Anopheles gambiae* saliva antigens. *Malar J*. 2010;9:317. DOI: 10.1186/1475-2875-9-317
- [191] Sarr JB, Orlandi-Pradines E, Fortin S, Sow C, Cornelie S, Rogerie F, et al. Assessment of exposure to *Plasmodium falciparum* transmission in a low endemicity area by using multiplex fluorescent microsphere-based serological assays. *Parasit Vectors*. 2011;4:212. DOI: 10.1186/1756-3305-4-212
- [192] Suwancharoen C, Putaporntip C, Rungruang T, Jongwutiwes S. Naturally acquired IgG antibodies against the C-terminal part of *Plasmodium falciparum* sporozoite threonine-asparagine-rich protein in a low endemic area. *Parasitol Res*. 2011;109:315–20. DOI: 10.1007/s00436-011-2257-z
- [193] Fidock DA, Pasquetto V, Gras H, Badell E, Eling W, Ballou WR, et al. *Plasmodium falciparum* sporozoite invasion is inhibited by naturally acquired or experimentally induced polyclonal antibodies to the STARP antigen. *Eur J Immunol*. 1997;27:2502–13. DOI: 10.1002/eji.1830271007
- [194] Robson KJ, Hall JR, Jennings MW, Harris TJ, Marsh K, Newbold CI, et al. A highly conserved amino-acid sequence in thrombospondin, properdin and in proteins from sporozoites and blood stages of a human malaria parasite. *Nature*. 1988;335:79–82. DOI: 10.1038/335079a0
- [195] Robson KJ, Frevert U, Reckmann I, Cowan G, Beier J, Scragg IG, et al. Thrombospondin-related adhesive protein (TRAP) of *Plasmodium falciparum*: expression during sporozoite ontogeny and binding to human hepatocytes. *Embo j*. 1995;14:3883–94.
- [196] Rogers WO, Malik A, Mellouk S, Nakamura K, Rogers MD, Szarfman A, et al. Characterization of *Plasmodium falciparum* sporozoite surface protein 2. *Proc Natl Acad Sci U S A*. 1992;89:9176–80.
- [197] Plebanski M, Aidoo M, Whittle HC, Hill AV. Precursor frequency analysis of cytotoxic T lymphocytes to pre-erythrocytic antigens of *Plasmodium falciparum* in West Africa. *J Immunol*. 1997;158:2849–55.
- [198] Wizel B, Houghten RA, Parker KC, Coligan JE, Church P, Gordon DM, et al. Irradiated sporozoite vaccine induces HLA-B8-restricted cytotoxic T lymphocyte responses against two overlapping epitopes of the *Plasmodium falciparum* sporozoite surface protein 2. *J Exp Med*. 1995;182:1435–45.
- [199] Bejon P, Mwacharo J, Kai O, Todryk S, Keating S, Lowe B, et al. The induction and persistence of T cell IFN-gamma responses after vaccination or natural exposure is suppressed by *Plasmodium falciparum*. *J Immunol*. 2007;179:4193–01.
- [200] Todryk SM, Bejon P, Mwangi T, Plebanski M, Urban B, Marsh K, et al. Correlation of memory T cell responses against TRAP with protection from clinical malaria, and CD4

- CD25 high T cells with susceptibility in Kenyans. *PLoS One*. 2008;3:e2027. DOI: 10.1371/journal.pone.0002027
- [201] Muller HM, Reckmann I, Hollingdale MR, Bujard H, Robson KJ, Crisanti A. Thrombospondin related anonymous protein (TRAP) of *Plasmodium falciparum* binds specifically to sulfated glycoconjugates and to HepG2 hepatoma cells suggesting a role for this molecule in sporozoite invasion of hepatocytes. *EMBO J*. 1993;12:2881–9.
- [202] Kester KE, Gray Heppner D Jr, Moris P, Ofori-Anyinam O, Krzych U, Tornieporth N, et al. Sequential Phase 1 and Phase 2 randomized, controlled trials of the safety, immunogenicity and efficacy of combined pre-erythrocytic vaccine antigens RTS,S and TRAP formulated with AS02 adjuvant system in healthy, malaria naive adults. *Vaccine*. 2014;32:6683–91. DOI: 10.1016/j.vaccine.2014.06.033
- [203] WHO. Malaria Vaccine Rainbow Tables. 2016. Available from: http://www.who.int/vaccine_research/links/Rainbow/en/index.html. [Accessed: 2016-10-19]
- [204] Prieur E, Gilbert SC, Schneider J, Moore AC, Sheu EG, Goonetilleke N, et al. A *Plasmodium falciparum* candidate vaccine based on a six-antigen polyprotein encoded by recombinant poxviruses. *Proc Natl Acad Sci U S A*. 2004;101:290–5. DOI: 10.1073/pnas.0307158101
- [205] Porter DW, Thompson FM, Berthoud TK, Hutchings CL, Andrews L, Biswas S, et al. A human Phase I/IIa malaria challenge trial of a polyprotein malaria vaccine. *Vaccine*. 2011;29:7514–22. DOI: 10.1016/j.vaccine.2011.03.083
- [206] Hill AV, Reyes-Sandoval A, O'Hara G, Ewer K, Lawrie A, Goodman A, et al. Prime-boost vectored malaria vaccines: progress and prospects. *Hum Vaccin*. 2010;6:78–83.
- [207] Webster DP, Dunachie S, Vuola JM, Berthoud T, Keating S, Laidlaw SM, et al. Enhanced T cell-mediated protection against malaria in human challenges by using the recombinant poxviruses FP9 and modified vaccinia virus Ankara. *Proc Natl Acad Sci U S A*. 2005;102:4836–41. DOI: 10.1073/pnas.0406381102
- [208] Bejon P, Kai OK, Mwacharo J, Keating S, Lang T, Gilbert SC, et al. Alternating vector immunizations encoding pre-erythrocytic malaria antigens enhance memory responses in a malaria endemic area. *Eur J Immunol*. 2006;36:2264–72. DOI: 10.1002/eji.200636187
- [209] Bejon P, Mwacharo J, Kai O, Mwangi T, Milligan P, Todryk S, et al. A phase 2b randomised trial of the candidate malaria vaccines FP9 ME-TRAP and MVA ME-TRAP among children in Kenya. *PLoS Clin Trials*. 2006;1:e29. DOI: 10.1371/journal.pctr.0010029
- [210] Bejon P, Mwacharo J, Kai OK, Todryk S, Keating S, Lang T, et al. Immunogenicity of the candidate malaria vaccines FP9 and modified vaccinia virus Ankara encoding the pre-erythrocytic antigen ME-TRAP in 1–6 year old children in a malaria endemic area. *Vaccine*. 2006;24:4709–15. DOI: 10.1016/j.vaccine.2006.03.029

- [211] O'Hara GA, Duncan CJ, Ewer KJ, Collins KA, Elias SC, Halstead FD, et al. Clinical assessment of a recombinant simian adenovirus ChAd63: a potent new vaccine vector. *J Infect Dis.* 2012;205:772-81. DOI: 10.1093/infdis/jir850
- [212] Ewer KJ, O'Hara GA, Duncan CJ, Collins KA, Sheehy SH, Reyes-Sandoval A, et al. Protective CD8+ T-cell immunity to human malaria induced by chimpanzee adenovirus-MVA immunisation. *Nat Commun.* 2013;4:2836. DOI: 10.1038/ncomms3836
- [213] Hodgson SH, Ewer KJ, Bliss CM, Edwards NJ, Rampling T, Anagnostou NA, et al. Evaluation of the efficacy of ChAd63-MVA vectored vaccines expressing circumsporozoite protein and ME-TRAP against controlled human malaria infection in malaria-naive individuals. *J Infect Dis.* 2015;211:1076-86. DOI: 10.1093/infdis/jiu579
- [214] Ogwang C, Kimani D, Edwards NJ, Roberts R, Mwacharo J, Bowyer G, et al. Prime-boost vaccination with chimpanzee adenovirus and modified vaccinia Ankara encoding TRAP provides partial protection against *Plasmodium falciparum* infection in Kenyan adults. *Sci Transl Med.* 2015;7:286re5. DOI: 10.1126/scitranslmed.aaa2373
- [215] Richie TL, Charoenvit Y, Wang R, Epstein JE, Hedstrom RC, Kumar S, et al. Clinical trial in healthy malaria-naive adults to evaluate the safety, tolerability, immunogenicity and efficacy of MuStDO5, a five-gene, sporozoite/hepatic stage *Plasmodium falciparum* DNA vaccine combined with escalating dose human GM-CSF DNA. *Hum Vaccin Immunother.* 2012;8:1564-84. DOI: 10.4161/hv.22129
- [216] Wang R, Richie TL, Baraceros MF, Rahardjo N, Gay T, Banania JG, et al. Boosting of DNA vaccine-elicited gamma interferon responses in humans by exposure to malaria parasites. *Infect Immun.* 2005;73:2863-72. DOI: 10.1128/iai.73.5.2863-2872.2005
- [217] Tine JA, Lanar DE, Smith DM, Wellde BT, Schultheiss P, Ware LA, et al. NYVAC-Pf7: a poxvirus-vectored, multiantigen, multistage vaccine candidate for *Plasmodium falciparum* malaria. *Infect Immun.* 1996;64:3833-44.
- [218] Ockenhouse CF, Sun PF, Lanar DE, Wellde BT, Hall BT, Kester K, et al. Phase I/IIa safety, immunogenicity, and efficacy trial of NYVAC-Pf7, a pox-vectored, multiantigen, multistage vaccine candidate for *Plasmodium falciparum* malaria. *J Infect Dis.* 1998;177:1664-73.
- [219] Moorthy VS, Imoukhuede EB, Milligan P, Bojang K, Keating S, Kaye P, et al. A randomised, double-blind, controlled vaccine efficacy trial of DNA/MVA ME-TRAP against malaria infection in Gambian adults. *PLoS Med.* 2004;1:e33. DOI: 10.1371/journal.pmed.0010033
- [220] Aly AS, Mikolajczak SA, Rivera HS, Camargo N, Jacobs-Lorena V, Labaied M, et al. Targeted deletion of SAP1 abolishes the expression of infectivity factors necessary for successful malaria parasite liver infection. *Mol Microbiol.* 2008;69:152-63. DOI: 10.1111/j.1365-2958.2008.06271.x
- [221] Zhao J, Deng S, Liang J, Cao Y, Liu J, Du F, et al. Immunogenicity, protective efficacy and safety of a recombinant DNA vaccine encoding truncated *Plasmodium yoelii*

sporozoite asparagine-rich protein 1 (PySAP1). *Hum Vaccin Immunother.* 2013;9:1104–11. DOI: 10.4161/hv.23688

- [222] Limbach K, Aguiar J, Gowda K, Patterson N, Abot E, Sedegah M, et al. Identification of two new protective pre-erythrocytic malaria vaccine antigen candidates. *Malar J.* 2011;10:65. DOI: 10.1186/1475-2875-10-65

IntechOpen

IntechOpen

