

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

186,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Introductory Chapter: Advancements in the Management of Gallbladder Diseases

Hesham Abdeldayem

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/67832>

Gallbladder diseases are the most prevalent digestive diseases worldwide. They result in considerable amount of financial and social burden. At the same time, clinical studies on these diseases continue to advance at a rapid pace.

The surgical management of gallstones, the most common affliction of the biliary tree, has been parallel to the evolution of surgical techniques. The first surgical report on gallstones dates back to 1687 when Stal Pert Von Der Weil found gallstones while exploring a patient suffering from peritonitis [1]. Open cholecystectomy was first performed and reported by the German surgeon, Carl Johann Langenbuch since one century. Later, this technique became the gold standard for the treatment of symptomatic gallstones [2] and remained so for almost a century. Operative cholangiography was introduced by Mirizzi over 60 years ago for the detection of stones in the bile duct [3].

Dr Med Erich Mühe of Böblingen, Germany in 1985, while performing laparoscopy for gynecologic indication on a woman who was also suffering from symptomatic gallstones, moved the laparoscope to the subhepatic area and succeeded to remove the gallbladder laparoscopically and the patient recovered uneventfully. Once the safety of laparoscopic cholecystectomy was established, it became the treatment of choice for cholelithiasis [4] and one of the most commonly undertaken procedures in general surgery.

Since then, this procedure has undergone many refinements including reduction in the port size and number. Some surgeons tried two ports only; others described single port technique through the umbilical scar. No scar laparoscopy cholecystectomy has been also described, the so-called NOTES (natural orifice transluminal endoscopic surgery) [5]. In the later technique, the gallbladder is removed through transanal, transvaginal, transcolonic, and transgastric route. Percutaneous cholecystostomy is another option available for too ill patients who are not fit for the laparoscopic procedure. It seems that surgical management of gallstones is still open for innovation, and further advancement included robotic-assisted laparoscopic cholecystectomy [6].

Gallbladder cancer (GBC) is the most frequent type of cancer of the biliary tract. The most important risk factor is gallstones. The majority of GBCs are adenocarcinomas, followed by squamous cell, adenosquamous, and undifferentiated carcinomas [7].

Surgery is the only curative therapy for GBCs. Most of the resectable GBC cases are diagnosed incidentally after histopathological examination of the resected gallbladder after laparoscopic cholecystectomy performed for gallstones [8].

The aim of surgery is to get negative margins. The extent of resection varies depending on the extent of the disease. For locally advanced GBC, major hepatectomy and/or resection of the CBD would be mandatory to get R0 resection. On the other hand, the potential benefit of such major resections should be balanced against the high morbidity and the poor.

The roles of radiation, chemoradiation, and chemotherapy in the neoadjuvant and adjuvant settings remain to be defined. Chemotherapy has been used in advanced GBC with limited results. Molecularly targeted agents that inhibit angiogenesis and EGFR pathways are being investigated [9].

Advances in the understanding of the molecular pathways of and genetic profiling of gallbladder cancer patients together with integration and coordination of clinical research efforts are critical to improve the outcomes for GBC.

The articles in this book provide a state-of-the-art review of the current knowledge and advances in research and management of gallbladder diseases, as well as promote future research, and clinical studies on the biliary disorders worldwide.

The immunogenetic basis of cholecystitis including human leukocyte antigens, as well as single-nucleotide polymorphism, is discussed in a separate chapter. Other chapters also discuss the role of endoscopic ultrasound in the diagnosis of gallbladder diseases together with the diagnostic pitfalls of acute cholecystitis. Advances in laparoscopic cholecystectomy are reviewed, particularly those related to robot-assisted and laparoscopic cholecystectomy in special situations like pregnancy and left-sided gallbladder.

Advances in gallbladder cancer research including noncoding RNAs are reviewed. Topics related to incidental gallbladder cancer, including its incidence, management, and prognosis, are discussed in details. Recent advances in the diagnosis, staging, and management of gallbladder cancer whether surgical or non-surgical are reviewed as well.

This book focuses on basic science and current methods in the diagnosis and management of gallbladder diseases. It is written by recognized medical experts and expected to be of great value for researchers and practicing gastroenterologists, endoscopists, and surgeons.

Author details

Hesham Abdeldayem

Address all correspondence to: habdeldayem64@hotmail.com

National Liver Institute, Menoufia University, Egypt

References

- [1] Beal JM. Historical perspective of gallstone disease. *Surg Gynecol Obstet*. 1984;158(2):181–9.
- [2] Karam J, Roslyn JR. Cholelithiasis and cholecystectomy. *Maingot's Abdominal Operations*. 12th edn. Prentice Hall International Inc; 1997. Vol 2. pp. 1717–38.
- [3] Mirizzi PL. Operative cholangiography. *Lancet*. 1938;2:366–9.
- [4] Saleh JW. *Laparoscopy*. Philadelphia; WB Saunders Co: 1988. pp. 7–8.
- [5] Peng C, Ling Y, Ma C, Ma X, Fan W, Niu W, Niu J. Safety Outcomes of NOTES Cholecystectomy Versus Laparoscopic Cholecystectomy: A Systematic Review and Meta-Analysis. *Surg Laparosc Endosc Percutan Tech*. 2016 Oct;26(5):347–353.
- [6] Lim JH, Lee WJ, Park DW, Yea HJ, Kim SH, Kang CM. Robotic cholecystectomy using Revo-i Model MSR-5000, the newly developed Korean robotic surgical system: a pre-clinical study. *Surg Endosc*. 2016 Nov 21.
- [7] Zimmitti G, Manzoni A, Guerini F, Ramera M, Bertocchi P, Aroldi F, Zaniboni A, Rosso E. Current Role of Minimally Invasive Radical Cholecystectomy for Gallbladder Cancer. *Gastroenterol Res Pract*. 2016;2016:7684915.
- [8] Zou ZY, Yan J, Zhuge YZ, Chen J, Qian XP, Liu BR. Multidisciplinary collaboration in gallbladder carcinoma treatment: A case report and literature review. *Oncol Lett*. 2016 Oct;12(4):2696–2701.
- [9] Jang JY, Heo JS, Han Y, Chang J, Kim JR, Kim H, Kwon W, Kim SW, Choi SH, Choi DW, Lee K, Jang KT, Han SS, Park SJ. Impact of Type of Surgery on Survival Outcome in Patients With Early Gallbladder Cancer in the Era of Minimally Invasive Surgery: Oncologic Safety of Laparoscopic Surgery. *Medicine (Baltimore)*. 2016 May;95(22):e3675.

