

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

185,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.  
For more information visit [www.intechopen.com](http://www.intechopen.com)



---

# Discovery of the Phenomenon of Intracellular Development of Cardiac Stem Cell: A New Step in Understanding of Biology and Behavior of Tissue-Specific Stem Cells

---

Galina B. Belostotskaya, Tatyana A. Golovanova,  
Irina V. Nerubatskaya and Michael M. Galagudza

Additional information is available at the end of the chapter

---

## Abstract

In our experiments with an in vitro culture of rat cardiac cells, we identified and described for the first time the phenomenon of intracellular development of CSCs in mature CMs with formation of the “cell-in-cell structures” (CICs). Recently, we have confirmed the reproducibility of our results and existence of this phenomenon in rats of different age groups, 1-year-old bull, adult mice and humans. Moreover, we demonstrated the 5–10 times increase in the amount of CICs after exposure of in vitro cultures to hypoxia and acidosis, that is, these conditions stimulate intracellular development of CSCs. Our data strongly suggest that transitory amplifying cells (TACs), which release from CICs, are present as a very rare cell population in adult and old rats. Therefore, we assume that TACs are important for renewal of myocardium during ontogenesis. TACs should be considered as the major source of cells that can reduce myocardial damage in adult mammals with various pathologies of the cardiovascular system. In conclusion, precise and exhaustive analysis of the phenomenon of intracellular development of CSCs, CICs and TACs will pave the way for cell technologies of new generation in regenerative medicine.

**Keywords:** primary culture of myocardial cells, mature cardiac myocytes, resident cardiac stem cells, proliferation, differentiation, cell-in-cell structure, transitory amplifying cells

---

## 1. Introduction

The founder of the study of stem cells (SCs) in adult organism is A.A. Maximow, who in 1909 first used the term «stem cell» and proved the existence of pluripotent hematopoietic stem

cell publishing a paper «The Lymphocyte as a stem cell common to different blood elements in embryonic development and during the post-fetal life of mammals» [1]. However, the ideas of our fellow-countryman A.A. Maximow, a Professor at the Saint Petersburg Imperial Military Medical Academy and Head of the Department of Histology and Embryology, were developed only in the 1960s, when Canadian scientists E. McCulloch and J. Till found hematopoietic SCs in the bone marrow [2]. The next breakthrough in this direction was made by Russian scientist A.Ya. Friedenstein, a Corresponding Member of the Academy of Medical Sciences of the U.S.S.R. [3]. He found in the bone marrow not only hematopoietic, but also stromal SCs, that can turn into cells of bone, cartilage, fibrous and fatty tissue. Going back to the ideas of A.A. Maximow, it should be emphasized that he suggested the existence of progenitor cells in certain tissues, marking them as stem mesenchymal cells. In this concept, he not only put their embryonic origin, but also their differentiation potency, ability to progressive differentiation into all kinds of connective tissue in postnatal development, such as reparative histogenesis.

Basic elements of the stem cell concept, developed in the study of the hematopoietic system, have been further extended to other rapidly renewing organs and tissues. It was found that the undifferentiated tissue-specific SCs (progenitor cells) are located in various organs and tissues, such as skin [4], cornea [5], kidney [6], liver [7], tooth [8, 9] and others, and that they are responsible for renewal of their cell population, replacing dead cells.

Although the detailed discussion of the characteristics of tissue-specific SCs is beyond the scope of this review, the brief history of neuronal SC discovery is described below as an example of long-lasting debate between “classical” views and new original ideas. Despite the fact that a paper by J. Altman, titled «Are new neurons formed in the brains of adult mammals?» [10] and published in the Science Magazine in 1962, and a few other publications by him, have proved the existence of neurogenesis in the adult mammalian brain, a concept that «the nerve cells do not regenerate» still persisted. Only twenty years later neurogenesis was again «discovered», but in the brains of birds. In the mid-1980s, Professor F. Nottebohm at Rockefeller University (U.S.A.) was able to show that in the vocal center of adult male canaries, a neurogenesis process occurs constantly in varying degrees [11]. In the late 1980s and early 1990s neuronal SCs have also been found in adult amphibians at the laboratory of our Institute of Evolutionary Physiology and Biochemistry of the U.S.S.R. Academy of Sciences, led by physiologist, endocrinologist and morphologist, a Corresponding Member of the Academy of Sciences, Andrey L. Polenov [12, 13]. In the same time period, neuronal progenitor cells were identified and studied first in the embryos [14], and then in the adult vertebrate animals [15]. These studies, along with the following works, have led to recognition of the existence of neuronal SCs, that has become not only an important fundamental discovery in stem cell biology, but also open up the prospects for their use in the treatment of peripheral nerve injury [16], spinal cord trauma [17, 18], acute cerebral injury [19], and neurodegenerative diseases, such as Alzheimer disease and multiple sclerosis [20–22].

## **2. Which cells are involved in the renewal and regeneration of the mammalian myocardium?**

Paradoxically, mammalian heart was considered as an organ not capable of self-renewal until the beginning of the XXI century. It was widely believed that the cardiomyocytes formed in the first days after birth, persist throughout life and cannot be replaced in case of damage or destruction, and an increase in heart size from birth to adulthood is due to hypertrophy of cardiomyocytes but not to their proliferation [23].

The first signs of a possible myocardial recovery, manifested in the resumption of DNA synthesis, have been observed in the studies of Russian scientists in the late 1970s-80s [24–26], while the data on increased telomerase activity, stimulation of cyclins and cyclin-dependent kinases, and enhanced proliferation of myocytes in mammals, including humans, in the later stages of cardiac insufficiency after myocardial infarction began to appear at the turn of twentieth and twenty-first centuries [27–30].

Existence of the SCs in adult mammalian heart and their participation in remodeling of injured myocardium was revealed for the first time only in 2002 in the paper by Hierlihy et al. [31] and later papers [23, 32–35]. A number of studies performed on the culture of heart cells, yielded valuable information about types of the resident CSCs, their clonogenicity and ability to differentiate [32, 36–41]. Also, it was found that cardiac SCs are present in the myocardium in trace amounts: c-kit<sup>+</sup> – 1 per 10<sup>4</sup> myocytes [32], Islet-1<sup>+</sup> – 500-600 in the heart of a 1-5-day-old rats [38], SP<sup>+</sup> (Sca1) – 500-1000 in the adult heart [39].


Nevertheless, in spite the discovery of the resident CSCs in the heart of mammals, including humans, and accumulation of knowledge about their biology, there is no consensus in the scientific community about the regenerative potential of the myocardium. To date, there is no unifying theory that can explain the fact that the ability of mammalian myocardium to regenerate after injury is lost in a few days after birth [42], and also why body's own resident CSCs, as well as externally injected CSCs [43–46], or mesenchymal SCs [47–49], or bone marrow SCs [50] are unable to form mature cardiomyocytes in the infarct zone. Answers to these questions, on one hand, will establish a real picture of the self-renewal of myocardial cells in normal conditions and changes in homeostasis in injured heart in various cardiovascular diseases, and on the other hand, will make possible to develop a considered tactics to influence on regenerative processes using modulators of proliferation of the resident cardiac cells.

## **3. Intracellular development of stem cells: a new step in the understanding of biology and behavior of tissue-specific stem cells**

From 2005 to 2011, when we studied the behavior of neonatal myocardial cells in primary culture, we confirmed existing concept that after a burst of mitotic activity in the first 2-4




days of postnatal life in vivo [51], the division of neonatal CMs in culture also terminated in 4-5 days after seeding in vitro [52]. At the same time, just like in the body, a mitotic division of 60% of the cells is observed, and after the cessation of this division, a growth of their volume is occurred, that is also similar to the process of hypertrophy of the CMs in vivo. It has been shown that the volume of CMs increased from  $819\pm68\ \mu\text{m}^3$  on the 1st day to  $1532\pm212\ \mu\text{m}^3$  on the 3rd day and up to  $3246\pm190\ \mu\text{m}^3$  on the 6th day of culturing. Moreover, the growth rate of cultured cells is almost completely coincides with the speed of myocyte hypertrophy in the body. Just like in the body, hypertrophy was accompanied by formation of polyploid and multinucleate, mostly binucleate cells. Besides, it was first



**Figure 1.** The formation of cardiomyocyte colonies in the primary culture of rat neonatal myocardial cells. (A–C) The optical sections of colonies formed by Isl1<sup>+</sup>, c-kit<sup>+</sup>, and Sca1<sup>+</sup> CSCs on the 11th day in vitro (DIV). (D–F) Differentiation of c-kit<sup>+</sup> CSCs inside the colony on the 13th DIV. (G–I) Progressive increase in the contraction rate of the same colony during culturing. (G) The 18th DIV, the contraction rate: 25 beats/min. (H) The 25th DIV, the contraction rate: 46 beats/min. (I) The 30th DIV, the contraction rate: 58 beats/min.


shown that the resident CSCs of three types ( $c\text{-kit}^+$ ,  $Sca1^+$ ,  $Isl1^+$ ) in the primary culture of newborn rat myocardial cells form colonies of contracting cardiomyocytes by dividing and subsequent cardiac differentiation, simulating the cardiomyogenesis process from a single immature stem cell to functionally mature cardiomyocyte (**Figure 1**). Therefore, the formation of contractile cardiomyocyte colonies is a complete model for a basic research, testing of drugs and identification of the regenerative potential of CSCs for possible application of the resident self-renewing cells in the treatment of myocardial injury [53, 54]. In addition, immunocytochemical study of the freshly isolated suspension (*ex vivo*) of myocardial cells from rats of different age groups, adult mice, young bulls and humans, revealed the presence of colonies of different maturity formed by resident CSCs of all three types (**Figure 2**). This made it possible to suggest that the colony formation is a way of myocardial self-renewal that allows to replace the lost cells throughout the life of mammals, including humans.

By study of myocardial cells in monolayer culture from newborn, 20- and 40-day old rats with help of antibodies to CSC antigens and confocal microscopy technique we first revealed the presence of small immature cells (5-6  $\mu\text{m}$  in diameter) of all three types inside mature cardiomyocytes with formation of «cell-in-cell structures» (CICs). It was found that the cell that has penetrated inside, usually located near the nucleus, or between two nuclei, and stored in the vacuole, membrane of which protects this cell from the host cell cytoplasm (**Figure 3**). Intracellular regions of large cells expressing CSCs antigens increase in size during culturing. Expression of the Ki-67 protein that plays an important role in the regulation of cell division in such vacuoles indicates on proliferation inside located CSCs. As a result of increase in their number, the size of the vacuoles increases, and the membrane thickens and becomes more dense forming a capsule. As time goes by, a formation of two to five openings (pores) occurs in the capsule that appears to provide gas exchange and exchange of metabolites between CSCs and host cell (**Figure 3 C, D, F**). Expression of cardiac (**Figure 3**) proteins ( $\alpha$ -sarcomeric actin, sarcomeric  $\alpha$ -actinin and Troponin T) suggests that the host cells appear to be a mature cardiomyocytes [55].



**Figure 2.** CSC-derived colonies in freshly isolated myocardial cell suspensions (*ex vivo*) of different mammalian species. The nuclei of the cells have been stained with Hoechst. Transmitted light and fluorescent images are merged. Cardiomyogenic differentiation is verified with expression of sarcomeric  $\alpha$ -actinin and troponin T. (A) An undifferentiated small  $Sca1^+$ -colony inside a fragment of myocardium of 40-day-old rat. (B) An undifferentiated small  $c\text{-kit}^+$ -colony from myocardium of an adult mouse. (C) A large volume  $c\text{-kit}^+$ -colony inside a fragment of myocardium of one-year-old bull. (D) An undifferentiated small  $c\text{-kit}^+$ -colony inside a fragment of myocardium of 45-year-old woman.






**Figure 3.** The CSCs inside CMs and the formation of CSC-containing CICSs in the cultures obtained from newborn and 20- and 40-day-old rats. (A–C) Early stages of CICS formation. (A) c-kit<sup>+</sup> CSC inside a newborn rat CM obtained (day in vitro 6). (B) Isl<sup>+</sup> CSC inside a newborn rat CM (day in vitro 4). Both the CSC and the host cell exhibit proliferative ability, as documented by the expression of Ki67. (C) Sca<sup>+</sup> CSC encapsulated between the nuclei of the host cell (20-day-old rat, day in vitro 11). (D–F) The CICS capsule in detail. (D) c-kit<sup>+</sup> CSC-containing CICS with a prominent coating ("capsule") with three pores (white arrows) obtained from a newborn rat, day in vitro 6. (E) Erosion (black arrow) of the Isl<sup>+</sup> CSC-containing CICS capsule obtained from 40-day-old rat, day in vitro 8. The pores are also visualized (white arrows). The capsule interior is positive for sarcomeric  $\alpha$ -actinin. (F) Erosion (black arrow) of the c-kit<sup>+</sup> CSC-containing CICS capsule obtained from a newborn rat, day in vitro 20. The pores are seen (white arrows). The nuclei of the cells have been stained with Hoechst, transmitted light and fluorescent images are merged.

Long-term observation of CICSs in primary culture of newborn, 20- and 40-day old rats revealed that intracellular development of CSCs ended by rupture of CICS-capsule and release of large quantities (up to 200) of transient amplifying cells (TACs), positive against SC-proteins of one of the three CSC-subtypes investigated by us and cardiac proteins (**Figure 4**). In the process of further development of descendants of the original CSC, expression of the stem antigens is reduced, but at the same time, synthesis of cardiac proteins is increased which allowed to suggest a reasonable expectation that the TACs appears to be an intermediate stage between CSCs and cardiomyocytes.

The presence of the similar CICSs, intact and opened, and the TACs of varying degrees of maturity in suspension of myocardial cells (ex vivo) of rats in different ages from the 1st day of life (**Figure 4**) to old age (1.5 years), adult C57bl/6 mice, 1-year-old bull, and human (45 years old) allowed to suggest that self-renewal of the myocardium may occur by proliferation and initial differentiation of CSCs inside mature cardiomyocytes throughout the life of mammals including humans.



**Figure 4.** Identification of “cell-in-cell structures” (CICs) in freshly isolated suspension of myocardial cells (ex vivo) of mammals of different species. (A–D) Intact CICs. (A) Sca1<sup>+</sup> CICs from myocardium of 1.5-year-old rat. (B) Sca1<sup>+</sup> CICs from myocardium of 1-year-old bull. (C) c-kit<sup>+</sup> CICs from myocardium of an adult mouse. (D) Sca1<sup>+</sup> CICs from myocardium of 45-year-old woman. (E and F) Opened CICs. (E) c-kit<sup>+</sup> CICs from myocardium of an adult mouse. (F) c-kit<sup>+</sup> CICs from myocardium of 45-year-old woman. (G and H) The release of transitory amplifying cells (TACs) from opened CICs. (G) Opened CICs and TACs from myocardium of 4.5-month-old rat. (H) Opened CICs and TACs from myocardium of 7-month-old rat.

Because the colonies of contracting cardiomyocytes and intracellular structures are formed in the culture with almost the same frequency (1-2 cases per 100,000 cells plated), we have put forward a hypothesis that the formation of new cardiac cells from resident CSCs may occur by colony formation as well as by intracellular development in mature myocardial cells. It is suggested, therefore, that both processes are present in a healthy heart and, apparently, maintain homeostasis of the heart muscle by replacing the dead cells with new functionally active cardiomyocytes.

It has been found for the first time that in experiments in vitro under simulated conditions of myocardial ischemia zone like acidosis and hypoxia suppress the growth of colonies by blocking the differentiation of resident CSCs, but at the same time, stimulate their intracellular development, increasing the amount of intracellular structures by several times. Therefore, it has been suggested that Ca<sup>2+</sup>-overload, as well as acidosis and hypoxia, accompanying myocardial infarction, cause the death of not only the mature myocardial cells, but also of the resident CSCs that are forced to «hide» inside large cardiac cells, which allows us to understand their passive role during myocardial infarction. This also explains irrationality of stimulation of resident CSCs for proliferation, differentiation and cell therapy application (introduction of exogenous SCs into the injured region) during the acute phase of ischemia and infarction.



#### 4. Analysis of problems of mammalian myocardial regeneration in the light of the new experimental data

Penetration of one or more cells to another to form the «cell-in-cell structures» (CICSs) was revealed 90 years ago by W.H. Lewis [56] in blood cells. The following terms are used to identify the processes leading to the formation of «cell-in-cell structures»: entosis, emperitosis, cannibalism, emperipolesis and cytophagocytosis. Entosis, firstly described in 2007 [57], is the process of active invasion of one cell (effector cell) into another (host cell). Terms «cannibalism» and «cytophagocytosis» are used to describe the phenomenon of active ingestion by the host cell of another cell which in this case is more passive [58]. Emperipolesis was established in 1956 [59] to indicate the penetration, movement and existence processes of one cell inside the other and, according to Overholtzer and Brugge [58], is suitable to refer to all phenomena in which the formation of structures of «cell-in-cell» type occurs, regardless the fate of cell located inside.

Unlike cytophagocytosis in which dead or dying cells to be destroyed within 30-60 min are captured, if CICS is formed, cell trapped inside nonphagocytic cell can survive for several days or even weeks. But cells in the host cell cytoplasm are surrounded by vacuole membrane that is, presumably, a fragment of a host cell membrane formed after its invagination. Movement of the effector cells, their division, and sometimes exit outside of the host cell were monitored [58].

It is known that the formation of «cell-in-cell structures» occurs both in the lower and in the higher forms of organisms, as well as the penetration of cells of lower forms into cells of highly organized species (parasitism). It is also shown that this phenomenon is typical for immune system cells (cytophagocytosis and emperipolesis) and tumors (entosis), occurring in the inflammation regions [60]. However, there is no information in the scientific literature to date on the formation of such structures in the myocardium.

Discovery of previously unknown mechanism of CSC development within mature cardiac myocytes with subsequent formation of cardiac positive TACs is of great interest not only as another new variant of the «cell-in-cell structure» formation, but also, more significantly, as a phenomenon allowing to reveal the nature of the biological processes that underlie the behavior of CSCs.

Moreover, our in vitro and ex vivo data on the intracellular development of CSCs allow to offer a new perspective into consideration of myocardial infarction problems. The reason is that the current understanding of the physiology of cardiac muscle cells is highly contradictory. For example, data on the rate of formation of new cardiomyocytes range from less than 1% [61–63] to more than 40% per year [64]. In addition, parallel to accumulation of data on the participation of CSCs in cardiomyogenesis, some papers periodically appear arguing in favour of division of mature CMs in the adult heart [64–67], reviving debate about what cells, resident CSCs or mature CMs, participate in the renewal of adult myocardium. In addition, Omatsu-Kanbe et al. [68] found a small population of cells ( $d \sim 10 \mu\text{m}$ ) that can be differentiated in contracting CMs without preliminary division. In turn, Kimura et al. [69] showed the presence of small dividing cells with characteristics of neonatal CMs: small

size, mononuclearity and insignificant oxidative DNA lesions, in the myocardium during hypoxia. Similar neonatal-like cells have been described even after experimental ischemia [70], which allow to suggest that in the myocardium of adult animals these effects stimulate the proliferation of insignificant amount of small cardiac positive cells that are unable, however, to recover the injured myocardium. Therefore, next questions are brought to the fore: what are these cells, where are they located in a healthy heart, and why they are unable to regenerate a myocardium in case of injury? Since the division of large (with a volume of more than  $20,000 \mu\text{m}^3$ ) terminally differentiated CMs is contradicted [71], and small dividing cells exceed the size of resident CSCs, a question arises about the origin and the physiological role of these cells.

Proliferative activity of TACs, formed by intracellular proliferation of CSCs, gave us the reason to consider this category of cells as a main regenerative source in the myocardium of adult mammals. We suggest that these cells are able not only to renew the myocardium throughout life, but also partially participate in its regeneration, if not in the acute phase of ischemia and infarction, then possibly during the recovery period or in chronic heart failure.

Thus, discovery of the phenomenon of intracellular development of CSCs is opening further perspectives in the study and solution of the problems of myocardium, and can be seen as a new step in understanding the nature of resident CSCs and development of the approaches for use of cell technologies in regenerative medicine.

## Acknowledgements

This work was supported by the Government of Russian Federation (grant no 074-U01), the Russian Foundation of Basic Investigations (projects no 12-04-00941 and no 16-04-01424) and by the state assignment of FASO of Russia ("The mechanisms of development of neuropsychic, metabolic and hormonal dysfunctions in the nervous and endocrine diseases and the approaches for their correction").

## Author details

Galina B. Belostotskaya<sup>1,2\*</sup>, Tatyana A. Golovanova<sup>1,2</sup>, Irina V. Nerubatskaya<sup>1,2</sup> and Michael M. Galagudza<sup>2,3</sup>

\*Address all correspondence to: gbelost@mail.ru

1 Sechenov Institute of Evolutionary Physiology and Biochemistry of Russian Academy of Sciences, St. Petersburg, Russian Federation

2 Federal Almazov North-West Medical Research Centre, St. Petersburg, Russian Federation

3 ITMO University, St. Petersburg, Russian Federation

## References

- [1] Maximow A. Der Lymphozyt als gemeinsame Stammzelle der verschiedenen Blutelemente in der embryonalen Entwicklung und im postfetalen Leben der Säugetiere. Originally in: *Folia Haematologica*. 1909;8:125-134. *Cell Therapy and Transplantation*. 2009;1:e.000040.01. DOI: 10.3205/ctt-2008-en-000040.01
- [2] Till JE, McCulloch EA, Siminovitch L. A stochastic model of stem cell proliferation, based on the growth of spleen colony-forming cells. *Proceedings of the National Academy of Sciences of the United States of America*. 1964;51:29-36. PMID: 14104600
- [3] Friedenstein AJ, Petrakova KV, Kuralesova AI, Frolova OF. Heterotopic transplants of bone marrow. Analysis of precursor cells for osteogenic and hemopoietic tissues. *Transplantation*. 1968;6:230-247. PMID: 5654088
- [4] Potten CS, Hendry JH. Clonogenic cells and stem cells in epidermis. *International Journal of Radiation Biology and Related Studies in Physics, Chemistry, and Medicine*. 1973;24:537-540. PMID: 4543437
- [5] Tseng SCG. Concept and application of limbal stem cells. *Eye*. 1989;3:141-157. DOI: 10.1038/eye.1989.22
- [6] Dekel B, Zangi L, Shezen E, Reich-Zeliger S, Eventov-Friedman S, Katchman H, Jacob-Hirsch J, Amariglio N, Rechavi G, Margalit R, Reisner Y. Isolation and characterization of nontubular Sca-1+Lin- multipotent stem/progenitor cells from adult mouse kidney. *Journal of the American Society of Nephrology*. 2006;17:3300-3314. DOI: 10.1681/ASN.2005020195
- [7] Knight B, Matthews VB, Olynyk JK, Yeoh GC. Jekyll and Hyde: Evolving perspectives on the function and potential of the adult liver progenitor (oval) cell. *Bioessays*. 2005;27:1192-1202. DOI: 10.1002/bies.20311
- [8] Gronthos S, Mankani M, Brahimi J, Robey PG, Shi S. Postnatal human dental pulp stem cells (DPSCs) in vitro and in vivo. *PNAS*. 2000;97:13625-13630. DOI: 10.1073/pnas.240309797
- [9] Mao X, Liu Y, Chen C, Shi S. Mesenchymal stem cells and their role in dental medicine. *Dental Clinics of North America*. 2017;61:161-172. DOI: 10.1016/j.cden.2016.08.006
- [10] Altman J. Are new neurons formed in the brains of adult mammals? *Science*. 1962;135:1127-1128. PMID: 13860748
- [11] Nottebohm F. Neuronal replacement in adulthood. *Annals of the New York Academy of Sciences*. 1985;457:143-161. PMID: 3913361
- [12] Chetverukhin VK, Polenov AL. Ultrastructural radioautographic analysis of neurogenesis in the hypothalamus of the adult frog, *Rana temporaria*, with special reference to physiological regeneration of the preoptic nucleus. I. Ventricular zone cell proliferation. *Cell and Tissue Research*. 1993;271:341-350. PMID: 8453658



- [13] Polenov AL, Chetverukhin VK. Itrastructural radioautographic analysis of neurogenesis in the hypothalamus of the adult frog, *Rana temporaria*, with special reference to physiological regeneration of the preoptic nucleus. II. Types of neuronal cells produced. **Cell and Tissue Research**. 1993;**271**:351-362. PMID: 8453659
- [14] Frederiksen K, McKay RD. Proliferation and differentiation of rat neuroepithelial precursor cells in vivo. **Journal of Neuroscience**. 1988;**8**:1144-1151. PMID: 3357014
- [15] Alvarez-Buylla A, Lois C. Neuronal stem cells in the brain of adult vertebrates. **Stem Cells**. 1995;**13**:263-272. DOI: 10.1002/stem.5530130307
- [16] Sullivan R, Dailey T, Duncan K, Abel N, Borlongan CV. Peripheral nerve injury: Stem cell therapy and peripheral nerve transfer. **International Journal of Molecular Sciences**. 2016;**17**:2101-2112. DOI: 10.3390/ijms17122101
- [17] Oh SK, Jeon SR. Current concept of stem cell therapy for spinal cord injury: A review. **Korean Journal of Neurotrauma**. 2016;**12**:40-46. DOI: 10.13004/kjnt.2016.12.2.40
- [18] Duan H, Song W, Zhao W, Gao Y, Yang Z, Li X. Endogenous neurogenesis in adult mammals after spinal cord injury. **Science China Life Sciences**. 2016;**59**:1313-1318. DOI: 10.1007/s11427-016-0205-2
- [19] Jessberger S. Stem cell-mediated regeneration of the adult brain. **Transfusion Medicine and Hemotherapy**. 2016;**43**:321-326. DOI: 10.1159/000447646
- [20] Franklin RJ, French-Constant C. Remyelination in the CNS: From biology to therapy. **Nature Reviews Neuroscience**. 2008;**9**:839-855. DOI: 10.1038/nrn2480.
- [21] Lee JH, Oh IH, Lim HK. Stem cell therapy: A prospective treatment for Alzheimer's disease. **Psychiatry Investigation**. 2016;**13**:583-589. DOI: 10.4306/pi.2016.13.6.583
- [22] Kremer D, Gottle P, Hartung HP, Kury P. Pushing forward: Remyelination as the new frontier in CNS diseases. **Trends in Neurosciences**. 2016;**39**:246-263. DOI: 10.1016/j.tins.2016.02.004
- [23] Leri A, Kajstura J, Anversa P. Cardiac stem cells and mechanisms of myocardial regeneration. **Physiological Reviews**. 2005;**85**:1373-1416. DOI: 10.1152/physrev.00013.2005
- [24] Rumyantsev PP. Interrelations of the proliferation and differentiation processes during cardiac myogenesis and regeneration. **International Review of Cytology**. 1977;**51**:187-273. PMID: 338537
- [25] Rumyantsev PP, Borisov A. DNA synthesis in myocytes from different myocardial compartments of young rats in norm, after experimental infarction and in vitro. **Biomedica Biochimica Acta**. 1987;**46**:S610-S615. PMID: 3435518
- [26] Erokhina IL, Rumyantsev PP. Proliferation and biosynthetic activities of myocytes from conductive system and working myocardium of the developing mouse heart. Light microscopic autoradiographic study. **Acta Histochemica**. 1988;**84**:51-66. DOI: 10.1016/S0065-1281(88)80010-2

- [27] Kajstura J, Leri A, Finato N, et al. Myocyte proliferation in end-stage cardiac failure in humans. *Proceedings of the National Academy of Sciences of the United States of America*. 1998;**95**:8801-8805. PMID: 9671759
- [28] Beltrami AP, Urbanek K, Kajstura J, et al. Evidence that human cardiac myocytes divide after myocardial infarction. *New England Journal of Medicine*. 2001;**344**:1750-1757. DOI: 10.1056/NEJM200106073442303
- [29] Leri A, Barlucchi L, Limana F, et al. Telomerase expression and activity are coupled with myocyte proliferation and preservation of telomeric length in the failing heart. *Proceedings of the National Academy of Sciences of the United States of America*. 2001;**98**:8626-8631. DOI: 10.1073/pnas.151013298
- [30] Quaini F, Urbanek K, Beltrami AP, et al. Chimerism of the transplanted heart. *New England Journal of Medicine*. 2002;**346**:5-15. DOI: 10.1056/NEJMoa012081
- [31] Hierlihy AM, Seale P, Lobe CG, et al. The post-natal heart contains a myocardial stem cell population. *FEBS Letters*. 2002;**530**:239-243. PMID: 12387899
- [32] Beltrami AP, Barlucchi L, Torella D, et al. Adult cardiac stem cells are multipotent and support myocardial regeneration. *Cell*. 2003;**114**:763-776. PMID: 14505575
- [33] Urbanek K, Quaini F, Tasca G, et al. Intense myocyte formation from cardiac stem cells in human cardiac hypertrophy. *Proceedings of the National Academy of Sciences of the United States of America*. 2003;**100**:10440-10445. DOI: 10.1073/pnas.1832855100
- [34] Cai C-L, Liang X, Shi Y, et al. Isl1 identifies a cardiac progenitor population that proliferates prior to differentiation and contributes a majority of cells to the heart. *Developmental Cell*. 2003;**5**:877-889
- [35] Anversa P, Kajstura J, Leri A, Bolli R. Life and death of cardiac stem cells: A paradigm shift in cardiac biology. *Circulation*. 2006;**113**:1451-1463. DOI: 10.1161/CIRCULATIONAHA.105.595181
- [36] Oh H, Bradfute SB, Gallardo TD, et al. Cardiac progenitor cells from adult myocardium: Homing, differentiation, and fusion after infarction. *PNAS*. 2003;**100**:12313-12318. DOI: 10.1073/pnas.2132126100
- [37] Matsuura K, Nagai T, Nishigaki N, et al. Adult cardiac Sca-1-positive cells differentiate into beating cardiomyocytes. *Journal of Biological Chemistry*. 2004;**279**:11384-11391. DOI: 10.1074/jbc.M310822200
- [38] Laugwitz K-L, Moretti A, Lam J, et al. Postnatal isl1+ cardioblasts enter fully differentiated cardiomyocyte lineages. *Nature*. 2005;**433**:647-653. DOI: 10.1038/nature03215
- [39] Pfister O, Mouquet F, Jain M, et al. CD31- but not CD31+ cardiac side population cells exhibit functional cardiomyogenic differentiation. *Circulation Research*. 2005;**97**:52-61. DOI: 10.1161/01.RES.0000173297.53793.fa
- [40] van Vliet P, Roccio M, Smits AM, et al. Progenitor cells isolated from the human heart: A potential cell source for regenerative therapy. *Netherlands Heart Journal*. 2008;**16**:163-169. PMID: 18566670

- [41] Belostotskaya GB, Golovanova TA. Characterization of contracting cardiomyocyte colonies in the primary culture of neonatal rat myocardial cells: A model of in vitro cardiomyogenesis. *Cell Cycle*. 2014;**13**:910-918. DOI: 10.4161/cc.27768
- [42] Porrello ER, Mahmoud AI, Simpson E, et al. Transient regenerative potential of the neonatal mouse heart. *Science*. 2011;**331**:1078-1080. DOI: 10.1126/science.1200708
- [43] Centola M, Schuleri KH, Lardo AC, Hare JM. Stem cell therapy for myocardial regeneration: Mechanisms and current clinical applications. *Giornale Italiano Di Cardiologia (Rome)*. 2008;**9**:234-250. PMID: 18543793
- [44] Smits AM, van Laake LW, den Ouden K, et al. Human cardiomyocyte progenitor cell transplantation preserves long-term function of the infarcted mouse myocardium. *Cardiovascular Research*. 2009;**83**:527-535. DOI: 10.1093/cvr/cvp146
- [45] D'Alessandro DA, Kajstura J, Hosoda T, et al. Progenitor cells from the explanted heart generate immunocompatible myocardium within the transplanted donor heart. *Circulation Research*. 2009;**105**:1128-1140. DOI: 10.1161/CIRCRESAHA.109.207266
- [46] Welt FG, Gallegos R, Connell J, et al. Effect of cardiac stem cells on left ventricular remodeling in a canine model of chronic myocardial infarction. *Circulation: Heart Failure*. 2013;**6**:99-106. DOI: 10.1161/CIRCHEARTFAILURE.112.972273
- [47] Seth S, Narang R, Bhargava B, et al. Percutaneous intracoronary cellular cardiomyoplasty for nonischemic cardiomyopathy: Clinical and histopathological results: The first-in-man ABCD (Autologous Bone Marrow Cells in Dilated Cardiomyopathy) trial. *Journal of the American College of Cardiology*. 2006;**48**:2350-2351. DOI: 10.1016/j.jacc.2006.07.057
- [48] Vrtovc B, Poglajen G, Lezaic L, et al. Effects of intracoronary CD34+ stem cell transplantation in nonischemic dilated cardiomyopathy patients: 5-year follow-up. *Circulation Research*. 2013;**112**:165-173. DOI: 10.1161/CIRCRESAHA.112.276519
- [49] Fischer-Rasokat U, Assmus B, Seeger FH, et al. A pilot trial to assess potential effects of selective intracoronary bone marrow-derived progenitor cell infusion in patients with nonischemic dilated cardiomyopathy: Final 1-year results of the transplantation of progenitor cells and functional regeneration enhancement pilot trial in patients with nonischemic dilated cardiomyopathy. *Circulation: Heart Failure*. 2009;**2**:417-423. DOI: 10.1161/CIRCHEARTFAILURE.109.855023
- [50] Wei HM, Wong P, Hsu LF, Shim W. Human bone marrow-derived adult stem cells for post-myocardial infarction cardiac repair: Current status and future directions. *Singapore Medical Journal*. 2009;**50**:935-942
- [51] Golovanova TA, Belostotskaya GB. Regeneration of rat cardiac myocytes in vitro: The proliferative activity of neonatal cardiac myocytes. *Cell Transplantation and Tissue Engineering*. 2011;**6**:66-70 (in Russian)
- [52] Li E, Wang X, Capasso J, Gerdes A. Rapid transition of cardiac myocytes from hyperplasia to hypertrophy during postnatal development. *Journal of Molecular and Cellular Cardiology*. 1996;**28**:1737-1746. DOI: 10.1006/jmcc.1996.0163



- [53] Golovanova TA, Belostotskaya GB. Regeneration of rat cardiac myocytes in vitro: Contracting myocyte colonies in the primary culture of neonatal cardiac myocytes. *Cell Transplantation and Tissue Engineering*. 2012;**7**:67-72. (in Russian)
- [54] Belostotskaya GB, Golovanova TA. Characterization of contracting cardiomyocyte colonies in the primary culture of neonatal rat myocardial cells: A model of in vitro cardiomyogenesis. *Cell Cycle*. 2014;**13**:910-918. DOI: 10.4161/cc.27768
- [55] Belostotskaya G, Nevorotin A, Galagudza M. Identification of cardiac stem cells within mature cardiac myocytes. *Cell Cycle*. 2015;**14**:3155-3162. DOI: 10.1080/15384101.2015.1078037
- [56] Lewis WH. The exgulfment of living blood cells by others of the same type. *The Anatomical Record*. 1925;**3**:43-49
- [57] Overholtzer M, Mailleux AA, Mouneimne G, et al. A non-apoptotic cell death process, entosis, that occurs by cell-in-cell invasion. *Cell*. 2007;**131**:966-979. DOI: 10.1016/j.cell.2007.10.040
- [58] Overholtzer M, Brugge JS. The cell biology of cell-in-cell structures. *Nature Reviews Molecular Cell Biology*. 2008;**9**:796-809. DOI: 10.1038/nrm2504
- [59] Humble JG, Jayne WH, Pulvertaft R.J. Biological interaction between lymphocytes and other cells. *British Journal of Haematology*. 1956;**2**:283-294. DOI: 10.1111/j.1365-2141.1956.tb06700.x
- [60] He MF, Wang S, Wang Y, Wang XN. Modeling cell-in-cell structure into its biological significance. *Cell Death and Disease*. 2013;**4**:e630. DOI: 10.1038/cddis.2013.147
- [61] Soonpaa MH, Koh GY, Pajak L, et al. Cyclin D1 overexpression promotes cardiomyocyte DNA synthesis and multinucleation in transgenic mice. *Journal of Clinical Investigation*. 1997;**99**:2644-2654. DOI: 10.1172/JCI119453
- [62] Bergmann O, Bhardwaj RD, Bernard S, et al. Evidence for cardiomyocyte renewal in humans. *Science*. 2009;**324**:98-102. DOI: 10.1126/science.1164680
- [63] Walsh S, Ponten A, Fleischmann B, Jovinge S. Cardiomyocyte cell cycle control and growth estimation in vivo—An analysis based on cardiomyocyte nuclei. *Cardiovascular Research*. 2010;**86**:365-373. DOI: 10.1093/cvr/cvq005
- [64] Kajstura J, Gurusamy N, Ogórek B, et al. Myocyte turnover in the aging human heart. *Cardiovascular Research* 2010;**107**:1374-1386. DOI: 10.1161/CIRCRESAHA.110.231498
- [65] Leri A, Kajstura J, Anversa P. Mechanisms of myocardial regeneration. *Trends in Cardiovascular Medicine*. 2011;**21**:52-58. DOI: 10.1016/j.tcm.2012.02.006
- [66] Senyo SE, Steinhauser ML, Pizzimenti CL, et al. Mammalian heart renewal by pre-existing cardiomyocytes. *Nature*. 2013;**493**:433-436. DOI: 10.1038/nature11682
- [67] Mollova M, Bersell K, Walsh S, et al. Cardiomyocyte proliferation contributes to heart growth in young humans. *Proceedings of the National Academy of Sciences of the United States of America*. 2013;**110**:1446-1451. DOI: 10.1073/pnas.1214608110

- [68] Omatsu-Kanbe M, Matsuura H. A novel type of self-beating cardiomyocytes in adult mouse ventricles. *Biochemical and Biophysical Research Communications*. 2009;**381**:361-366. DOI: 10.1016/j.bbrc.2009.02.048
- [69] Kimura W, Xiao F, Canseco DC, et al. Hypoxia fate mapping identifies cycling cardiomyocytes in the adult heart. *Nature*. 2015;**523**:226-230. DOI: 10.1038/nature14582
- [70] Tang X-L, Li Q, Rokosh G, et al. Long-Term Outcome of administration of c-kit<sup>POS</sup> cardiac progenitor cells after acute myocardial infarction: Transplanted cells do not become cardiomyocytes, but structural and functional improvement and proliferation of endogenous cells persist for at least one year. *Circulation Research*. 2016;**118**:1091-1105. DOI: 10.1161/CIRCRESAHA.115.307647
- [71] Leri A, Rota M, Pasqualini FS, et al. Origin of cardiomyocytes in the adult heart. *Circulation Research*. 2015;**116**:150-166. DOI: 10.1161/CIRCRESAHA.116.303595