

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

186,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Sirtuins in Adipose Tissue Metabolism

Manu Sudhakar, Santhi Silambanan and
Ramya Ramakrishnan

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/intechopen.75704>

Abstract

Obesity, a complex metabolic disorder linked to the development of several diseases, is characterized by both hypertrophy and hyperplasia of adipocytes. While white adipose tissue (WAT) is an energy storage site, brown adipose tissue (BAT) activation generates heat from nutrients by non-shivering thermogenesis. The human orthologue of silencing information regulator 2 (Sir2) which was recognized as a regulator of life span in *S. cerevisiae*, includes seven sirtuins which are NAD⁺-dependent protein deacetylases distributed in different subcellular compartments. Sirtuins, particularly Sirt1, have emerged as important nutrient sensors and regulators of metabolism. Sirt1 has been shown to play a role in retarding the expansion of WAT while stimulating both differentiation and activation of brown adipose tissue as well as browning of WAT. This chapter focuses on the role of sirtuins in adipose tissue biology, their implications in obesity and potential as therapeutic targets.

Keywords: sirtuins, white adipose tissue, brown adipose tissue, adipogenesis, metabolic control, obesity

1. Introduction

Adipose tissue is a functionally diverse organ with remarkable plasticity to adapt to changes in energy balance and contribute to systemic regulation of metabolism. It is capable of expanding in response to over-nutrition preventing ectopic fat deposition in non-adipose tissues, as well as mobilizing stored lipids during starvation or energy demand. Apart from its storage function, its ability to produce multiple adipokines as an endocrine organ, influences functions of other metabolic tissues. The ability of adipose tissue to respond to changes in energy balance is finely regulated by molecular mechanisms linked to nutrient response and

the redox status of the tissue. Sirtuins are a group of NAD⁺-dependent protein-deacetylating enzymes that play a key role in metabolic homeostasis. They influence genome stability, transcription and activity of several enzymes contributing to epigenetic regulation and act as key nutrient sensors.

Acetylation and deacetylation of histones and other DNA associated proteins are key epigenetic processes that can influence chromatin structure regulating the transition between highly condensed and transcriptionally less active heterochromatin state to loose and transcriptionally active euchromatin structure [1]. The extent of acetylation of histones depends, on one hand, on the balance between the activity of acetyl transferases such as histone acetyl transferase and deacetylases such as histone deacetylases (HDACs) [2, 3] and on the other hand on acetyl CoA which is an important metabolite at the junction of several metabolic pathways. Phylogenetically, eukaryotic HDACs are ancient proteins comprising of two subfamilies of proteins with different structure and function [4, 5]. While the enzymes belonging to the classical HDAC family are generally Zn-dependent enzymes which remove acetyl group from lysine residues on acetylated protein substrates with the addition of a water molecule, those belonging to the sirtuin family remove acetyl moieties bound to protein substrates to another substrate viz. NAD⁺ cleaving it to nicotinamide and *O*-acetyl ADP-ribose. So far 11 members of the classical HDAC enzymes, subdivided into three classes (HDAC I, III and IV), and 7 members of sirtuins belonging to HDAC class III family have been identified.

Sirtuins, particularly SIRT1, play a key role in adipogenesis of white adipose tissue (WAT) and browning of WAT, metabolism of glucose and fat, insulin sensitivity, control of inflammation and energy homeostasis. Dysregulation of these physiological processes have major implications in the development of obesity and related metabolic diseases. The role of sirtuins in the regulation of adipose tissue biology and its implications in the development of obesity form the subject matter of this chapter. A number of reviews on the structure and functions of various sirtuins and their implications in aging and several pathological conditions are available [6–11].

2. Sirtuins in adipose tissue biology

2.1. Biochemistry and molecular biology of sirtuins

2.1.1. Classification and tissue distribution

Interest in understanding the role of sirtuins in mammalian system was driven by findings in calorie-restricted conditions in model organisms. The discovery of a protein in yeast, referred to as yeast-silencing information regulator-2 (Sir2) [12] followed by the demonstration of its localization in the nucleolus and telomeres [13] and low histone acetylation level in the genetic loci highlighted the importance of this group of proteins in regulating chromatin structure in specific loci. These findings were followed by demonstration of the importance of the protein in regulating yeast lifespan [14] and identification of Sir2 as an NAD-dependent histone deacetylase [15]. This revealed the connection between a molecule involved in gene

silencing and cellular metabolism and led to establishing sirtuins as important epigenetic regulators. This was followed by identification of *Sir2* orthologues in mammalian systems; these were demonstrated to be NAD⁺-dependent protein deacetylating enzymes that are highly conserved across bacteria to humans.

On the basis of sequence similarity, mammalian sirtuins are classified into four classes, class I–IV. Mammalian SIRT1–3 belong to class I, SIRT4 to class II, SIRT5 to class III and SIRT6 and 7 to class IV (**Table 1**) [9, 16]. All these enzymes share a conserved catalytic core structure, but they differ in their enzymic activity. While class I sirtuins (SIRT1–3) show robust deacetylase activity in vitro, SIRT4–7 show weak deacetylase activity; SIRT4 shows mono ADP-ribosyl transferase activity and SIRT2, 3 and 6 exhibit both deacetylase and ribosyl transferase activity. SIRT5 removes succinyl, malonyl and glutaryl groups from acylated protein-lysine residues. SIRT6 is more efficient in removing long chain fatty acyl groups such as myristoyl and palmitoyl groups. The enzymic reaction proceeds through the formation of a ternary complex involving acetyl protein substrate and NAD at the active site, which decomposes to form deacetylated protein, nicotinamide and 2-*O*-acetyl-ADP ribose. They also differ in tissue distribution and subcellular localization (**Table 1**) [6, 9]. SIRT1 is expressed in metabolic tissues such as liver, muscle, adipose tissue and other organs such as heart, brain, pancreas; it is localized mainly in the nucleus, but shuttles from nucleus to cytosol. SIRT2, primarily a cytosolic protein, is highly expressed in heart, brain and skeletal muscle. SIRT3–5 are expressed in mitochondria. SIRT3 and 5 are mainly expressed in kidney, brain, liver and heart. In contrast to white adipose tissue, brown adipose tissue (BAT) expresses SIRT3. SIRT4 is expressed in heart, liver, pancreas and vascular smooth muscle. SIRT6 and 7 are localized predominantly in the nucleus. While SIRT6 is expressed in brain, liver and muscle, SIRT7 is mainly found in liver and spleen.

2.1.2. Biological effects of sirtuins

Mammalian sirtuins influence various cellular processes such as chromatin silencing, cell cycle regulation, differentiation and survival, mitochondrial biogenesis, metabolism, inflammation and stress response. While most studies in model organisms indicate a role for sirtuins in mediating the increased longevity affected by calorie restriction [17], a similar role for mammalian sirtuins is debated [18]. Sirtuins deacetylate transcription factors and regulate their activities either by influencing their cytoplasmic-nuclear distribution, their binding to DNA or changing their interaction with regulatory proteins.

SIRT1 is by far the best characterized among all mammalian sirtuins. It has been linked to hypothalamic control of energy balance [19], has a role in adipogenesis and fat mobilization, as well as regulation of carbohydrate and lipid metabolism [6, 7]. SIRT1 promotes vasodilation and regenerative function in endothelial and smooth muscle cells of vascular wall by targeting eNOS for deacetylation [20]. In cardiomyocytes, SIRT1, 3 and 7 play a critical role in promoting cardiomyocyte resistance to stress and toxicity [21].

Important targets of SIRT1 include p53, forkhead box type O transcription factors (FOXO), PPAR γ co-activator-1 α (PGC1 α), NF κ B, androgen receptor and their co-regulatory molecules (**Table 1**). Apart from its effect on PGC1 α , a master regulator of mitochondrial biogenesis,

	Sirt 1	Sirt 2	Sirt 3	Sirt 4	Sirt 5	Sirt 6	Sirt 7
Class	I	I	I	II	III	IV	IV
Subcellular location	Nucleus, cytoplasm	Cytoplasm, nucleus	Mitochondria	Mitochondria	Mitochondria	Nucleus	Nucleolus
Tissue	Liver, heart, brain, pancreas, muscle, adipose tissue	Heart, brain, skeletal muscle	Brown adipose tissue, kidney brain, heart, liver	Vascular smooth muscle, skeletal muscle, heart, liver, pancreas	Brain, heart, muscle, liver, kidney	Brain, liver, muscle	Liver, spleen
Activity	Deacetylation	Deacetylation	Deacetylation	ADP-ribosylation	Deacetylation, demalonylation desuccinylation	Deacetylation, ADP-ribosylation	Deacetylation
Target	PGC1- α , PPAR γ , PPAR α , FOXO1, FOXO3, p53, notch, NF-kB, HIF-1 α , LXR, FXR, SREBP1c	Tubulin, PEPCK, FOXO-1, PAR-3	LCAD, HMGCS-2, SOD-2, GDH, IDH2	GDH, Malonyl CoA decarboxylase	CPS1	H3KK9, H3K56	SIRT 1
Biological effects (WAT)							
Adipogenesis	Inhibition	Inhibition					Stimulates
Lipogenesis	Inhibition (decreases AC1)			Stimulates malonyl decarboxylase			
Lipolysis	Stimulates (FOXO1, ATGL)	Stimulates					
β -oxidation							
Oxphos							
Glucose metabolism	Improves insulin sensitivity					Decreases insulin sensitivity	
Inflammation	Decreases					Decreases	Increases
Adipokines	Increases adiponectin Decreases leptin		Increases adiponectin			Increases adiponectin	Decreases leptin

	Sirt 1	Sirt 2	Sirt 3	Sirt 4	Sirt 5	Sirt 6	Sirt 7
Biological effects (BAT)							
Brown adipogenesis			Stimulates (expressed more in BAT than WAT)		Stimulates (expressed more in BAT than WAT)		Inhibits (expressed more in BAT than WAT)
Browning	Stimulates						
Whitening	Inhibits						
β-oxidation	Stimulates		Stimulates				
Oxphos	Stimulates mitochondrial biogenesis		Stimulates				
Thermogenesis	Stimulates		Stimulates				Inhibits
Glucose metabolism	Improves glucose tolerance			Increases on cold exposure		Increases glucose uptake	Decreases glucose tolerance
Effects on sirtuins							
Obesity	Decreased		Decreased	Increased	Decreased?	Decreased	Increased/decreased
CR	Increased		Increased	Decreased (?)		Increased	Increased
Gastric banding surgery	Increased			Increased			
References	[6, 7, 11, 31, 52, 53, 77, 78, 100, 102]	[6, 7, 11, 26, 31]	[6, 7, 11, 27, 31]	[6, 7, 11, 31, 65, 66]	[6, 7, 11, 31]	[6, 7, 11, 31]	[6, 7, 11, 31, 108]

Table 1. Sirtuins – targets, enzymic activity and biological effects.

SIRT1 acts on several transcription factors like estrogen like receptors, the nuclear respiratory factors 1 and 2 to induce mitochondrial gene expression [22].

SIRT1 appears to directly block lipid anabolism by interfering with PPAR γ and LXR signaling. The repressive effect of SIRT1 on PPAR γ activity requires the formation of a co-repressor complex that involves NCoR1 [23]. SIRT1 also has a role in reverse cholesterol transport; it stimulates cholesterol efflux from macrophages and the hepatic conversion of cholesterol to bile acids through LXR [24]. It is also present in the cytosol of many cell types and regulates major cytoplasmic enzymes such as acetyl CoA synthase and eNOS by deacetylation [20].

Compared to SIRT1, not much is known about the physiological effect of other mammalian sirtuins. SIRT2 shows similarity to SIRT1 in several of its biological effects. It appears to increase hepatic glucose which is beneficial in starving conditions where SIRT2 activity increases. While it suppresses glycolysis by deacetylating and destabilizing glucokinase, it activates gluconeogenesis by enhancing the action of a key gluconeogenic enzyme PEPCK by its deacetylation [25]. It also appears to have a role in the control of microtubule stability and cell cycle oscillations by deacetylating α -tubulin [26].

SIRT3 is a major regulator of mitochondrial function; it deacetylates several mitochondrial proteins which are critical in mitochondrial oxidative metabolism. Calorie restriction increases activity of SIRT3 which alters the mitochondrial acetylome [27, 28]. SIRT4 is a mitochondrial matrix protein with remarkable ADP-ribosyltransferase activity [29]. It is reported to regulate insulin secretion [30], the activity of glutamate dehydrogenase, and serves as a metabolic regulator by inhibiting the activity of several metabolic enzymes such as pyruvate dehydrogenase opposing the effect of SIRT3 [31]. Although recent studies have identified several SIRT5 target proteins, not much is known about its biological function. SIRT5 appears to play a role in energy homeostasis and free radical metabolism [32]. SIRT6 deficient mice aged prematurely and its overexpression increased life span of male mice apparently by altering IGF signaling [31].

2.1.3. Regulation of sirtuins

Sirtuins are regulated in response to nutritional and metabolic challenges, oxidative stress, and inflammation in a cell and tissue specific manner. They are subject to transcriptional control, post-transcriptional regulation by miRNA and post-translational modulation [31]. These regulatory events can either alter the levels of each sirtuin or their enzyme activity or both. Their activity can be modulated either directly by post-translational modifications (PTMs) such as phosphorylation and acetylation, protein interactions and by compounds that activate them, or indirectly by modulating NAD⁺ levels. The substrates themselves appear to regulate sirtuin expression indicating the possible formation of feedback regulatory loops.

Some of the key factors involved in metabolic homeostasis that cause upregulation of SIRT1 include CREB, FOXO1, FOX3a, C/EBP α , PPAR α and PPAR β/δ while the negative regulators include ChREBP, C/EBP. NF κ B, EGR1, APE1 positively regulate transcription of SIRT1 during stress conditions. SIRT1 in turn can modulate the activity of several of these transcription factors. The enzymatic activity of SIRT1 is enhanced by post-translational modification by phosphorylation [33] and SUMOylation (Lys734) [34]. The activity of SIRT1 can also be controlled

through interaction with different protein complexes such as DBC1 (nuclear protein deleted in breast cancer-1), AROS (active regulator of SIRT1) and NCoR1 [35, 36]. SIRT1 can also be activated indirectly by either increasing NAD⁺ synthesis from NAD⁺ precursors like nicotinic acid, NAM, nicotinamide riboside or decreasing NAD⁺ consumption by two alternate enzyme families, PARP and cADP ribose synthase [37]. SIRT1 can enhance its own activity by auto-deacetylation. This is inhibited by SIRT7 suggesting coordinated action of different sirtuins [38]. SIRT1 is inhibited by NADH which competes with NAD⁺ [39] and by nicotinamide [40]. *In vitro* studies using malignant cell lines, have reported the role of different miRNAs in the post-transcriptional regulation of SIRT1 [31]. Mir34a is reported to regulate SIRT1 at different levels; it inhibits translation of SIRT1 mRNA; it indirectly suppresses transcription and enzyme activity of SIRT1 by modulating the levels of NAD through regulation of the biosynthetic enzyme nicotinamide phosphoribosyl transferase [41].

Unlike SIRT1, information on the regulation of other sirtuins is limited. Cyclin(E-Cdk2 and A-Cdk2) mediated phosphorylation at S331 inhibits SIRT2 activity whereas Erk1/2 enhanced its activity [31]. Cyclic AMP–PKA pathway, which is activated in response to various stimuli, causes upregulation of SIRT3 expression through PGC1 α . cAMP may also activate SIRT3 by direct binding [31]. Activated AMPK also positively regulates SIRT3 by increasing NAD. Similar to SIRT3, expression of SIRT5 is also upregulated by PGC1 α , while AMPK appears to negatively regulate its expression [31]. Several transcription factors and miRNAs appear to regulate expression of SIRT6. While Nuclear Respiratory factor 1 (NRF-1) and co-activators induce SIRT6 expression during nutrient deprivation, cAMP signaling reduces SIRT6 expression. A relation between SIRT6 and mir122 has also been suggested [31].

2.2. Sirtuins regulate WAT development and metabolism

Adipose tissue is classified into white adipose tissue (WAT) which serves as the principal energy storage organ and brown adipose tissue (BAT) whose principal function is maintaining temperature by non-shivering thermogenesis. A third category includes beige or brite (brown in white) adipocytes within WAT which can potentially differentiate into cells of brown like phenotype.

2.2.1. Sirtuins and WAT development

The cellular and molecular mechanisms that govern the adipocyte life cycle have been extensively studied [42–45]. Both white and brown adipocytes develop by a highly regulated process of differentiation of mesenchymal stem cells (MSC). During the early phase of adipogenesis the pluripotent MSCs are committed to unipotent pre-adipocytes which in the latter phase undergo terminal differentiation acquiring the characteristic phenotype and functions of mature adipocytes. The complex process of adipocyte differentiation is coordinated by myriad factors. Isoforms of bone morphogenetic protein, BMP2 and BMP4 are the key positive regulators of commitment to white pre-adipocytes [46, 47]. This early phase of differentiation is also subject to negative regulation by several transcription factors including members of GATA and Forkhead family, Wnt and Notch signaling, Kruppel-like factors 2 and 7 (KLF2, KLF7) and CHOP proteins [48].

The differentiation of white pre-adipocytes to mature adipocytes is mediated by multiple transcription factors. The key transcription factors involved are CCAAT enhancer binding protein (C/EBP), PPAR γ and sterol-regulatory element binding protein 1 (SREBP1) [43]. C-EBP, which exists in six different isoforms, is activated and translocated to the nucleus in response to cAMP mediated signaling [43]. Hormones induce transient expression of C/EBP β and C/EBP δ which upregulate expression of PPAR γ and C/EBP α [43]. PPAR γ , (particularly the isoform PPAR γ 1) that belongs to the nuclear receptor superfamily, heterodimerises with retinoid receptor α (RXR α), another nuclear receptor, and binds to DNA to promote expression of adipocyte specific genes such as leptin, adiponectin, FABP4, and perilipin. Activation of C/EBP α leads to transcriptional activation of several genes encoding proteins such as GLUT4, SCD1, FABP4 which are critical in establishing adipocyte phenotype [43].

All seven sirtuin genes are expressed in human and rodent WAT. A difference in their level of expression, particularly a reduction in SIRT1, has been observed, in experimental obese models as well as obese human subjects [11]. Studies in experimental model systems suggest that sirtuins, particularly SIRT1, are negative modulators of WAT adipogenesis. Overexpression of SIRT1 decreased accumulation of fat while its knockdown increased fat accumulation in 3T3L1 cells undergoing differentiation [23]. Resveratrol, an activator of SIRT1 reduces osteoblastic differentiation of MSC to adipocytes [49]. SIRT1 is upregulated in WAT in calorie-restricted mice model in which there was significant reduction in fat mass [50, 51]. Further, transgenic mice overexpressing SIRT1 showed lower body weight and reduction of fat mass [52], while ablation of SIRT1 in WAT resulted in gain in body weight, increase in fat mass and an increase in the size of individual adipocytes [53]. Studies using bone marrow-derived MSC with SIRT1 deletion showed impaired self-renewal and differentiation to osteoblasts without significantly affecting their differentiation to adipocytes [54]. In the light of these observations it has been suggested that although SIRT1 inhibits adipocyte differentiation, its expression is critical for maintenance of MSC pool [11].

PPAR γ is an important substrate for SIRT1 [55]. SIRT1 dependent deacetylation of lysine residues (268 and 293 K) on PPAR γ is critical in the regulation of its transcriptional activity by its co-repressors NCoR and SMRT. Sirt1 can thus inhibit white adipogenesis by suppressing the transcriptional activity of PPAR γ by promoting the binding of its co-repressors NCoR and SMRT [23]. Further, C/EBP α , a PPAR γ dependent factor, regulates the expression of SIRT1 during adipogenesis [56]. miRNAs can also regulate effects of SIRT1 on adipogenesis. For example, mir 146b can promote adipogenesis by suppressing SIRT1-FOXO1 cascade [57]. SIRT2, the predominant sirtuin in adipose tissue, has also been shown to inhibit adipocyte differentiation by deacetylating FOXO1 and enhancing its repressive interaction with PPAR γ [58, 59]. Unlike SIRT1 and SIRT2, SIRT7 knockdown in human pre-adipocytes reduced lipid content and the number of FABP4⁺ differentiated adipocytes indicating a contrasting effect of SIRT7. Further, SIRT7 knockout mice had significantly reduced WAT and increased SIRT1 activity. This suggests that SIRT7 influences adipogenesis in mice by inhibiting autocatalytic activation of SIRT1, further highlighting the importance of cross talk among sirtuins in the control of adipose tissue maintenance [38].

2.2.2. Sirtuins and WAT metabolism

During times of positive energy balance, storage of lipids as triglycerides in lipid droplets in adipocytes leads to expansion of adipose tissue. It leads to changes in the levels of adipokines such as leptin and adiponectin that affect metabolic functions of other organs particularly liver, muscle and brain, which in turn can affect the adipose tissue function [43, 45]. The stored lipid is mobilized in response to hormones and systemic energy needs during periods of negative energy balance. Metabolism of adipose tissue is thus linked with that of other tissues. SIRT1 appears to serve as an important metabolic switch through which adipose tissue and other metabolic organs respond to energy needs [8].

When energy stores are high, excess nutrients particularly glucose and amino acids are used to synthesize fatty acids *de novo* primarily in liver and are exported to WAT where they are stored as TG [6]. SIRT1 deacetylates sterol-responsive element binding protein 1c (SREBP1c) causing reduction of its transcriptional ability and suppression of fatty acid synthesis [6]. Further, by increasing PGC1 α -mediated mitochondrial biogenesis, SIRT1 facilitates fatty acid oxidation. SIRT1 promotes gluconeogenesis in the liver through deacetylation and activation of PGC1 α and FOXO1, and increasing expression of gluconeogenic enzymes [22] and inhibits glycolysis [6]. This increases hepatic production of glucose during fasting. Apart from SIRT1, SIRT6 also has been shown to repress glycolysis in liver [31]. Further, SIRT1, by deacetylating STAT3, reduces its repressive effect on gluconeogenesis [60]. The hepatic effects of sirtuins can thus affect nutrient flux into the adipose tissue.

Sirtuins are not only critical in WAT adipogenesis, they also play a key role in maintaining the functions of differentiated adipocytes by regulating the expression of several PPAR γ -responsive genes involved in metabolism. Activation of SIRT1 reduces expression of acetyl coA carboxylase and other lipogenic genes involved in the *de novo* synthesis of FAs in pre-adipocytes. Multiple studies support the role of PPAR γ in regulating adipose tissue metabolism [61]. Over-expression of a dominant negative form of PPAR γ downregulates the expression of key genes involved in lipid metabolism, insulin signaling and decreases lipid content in 3T3L1 differentiated adipocytes [62]. Further, selective ablation of PPAR γ in mature white and brown adipocytes results in adipocyte death without any effect on pre-adipocyte differentiation indicating the requirement of PPAR γ for maintenance of differentiated functions of adipocytes [63]. Insulin dependent glucose uptake in the adipocyte takes place through the GLUT4 transporter. Expression of GLUT4 gene is regulated by PPAR γ . SIRT1 appears to regulate glucose-induced secretion of insulin by transcriptional repression of UCP2 which uncouples mitochondrial ATP production [64]. SIRT1 by deacetylating SREBP1, destabilizes and reduces its occupancy on the lipogenic gene promoters suppressing fatty acid synthesis [6]. Although SIRT1 inhibits lipogenesis in WAT, SIRT4 appears to have an opposite effect. Deletion of mitochondrial SIRT4 decreased lipogenesis apparently by regulating malonyl CoA decarboxylase, thereby altering the level of malonyl CoA that represses fatty acid oxidation [65, 66]. While an increase in SIRT4 in fed state results in deacetylation and inactivation of malonyl CoA decarboxylase resulting in enhanced FA synthesis, during fasting, reduction of SIRT4 expression

leads to activation of malonyl CoA decarboxylase resulting in enhanced lipid oxidation. It therefore appears that SIRT4 exhibits a dual effect regulating anabolism and catabolism of fatty acids [65]. SIRT6 also appears to regulate lipogenesis. In SIRT6 overexpressing mice, expression of diacylglycerol acyl transferase a key enzyme involved in TG synthesis was down regulated along with certain PPAR γ responsive genes involved in lipogenesis [31, 67].

2.2.3. Mobilization of depot fat

During periods of negative energy balance, the triglycerides stored in the lipid droplets, are hydrolysed to FFA which are released into circulation. In the basal state or during TG synthesis, perilipin, the structural protein on lipid droplet, is bound to protein CGI-58 (comparative gene identification-58) which is a co-activator of the adipose triglyceride lipase (ATGL). In response to cAMP dependent PKA-mediated phosphorylation of perilipin, CGI-58 is released leading to activation of ATGL [68]. Activated ATGL moves to the lipid droplet membrane surface to hydrolyse TG to diacyl glycerol. Further activation of hormone sensitive lipase (HSL) by PKA mediated phosphorylation causes binding of HSL to perilipin and continues lipolysis forming monoacyl glycerol which is acted upon by another specific monoacyl glycerol lipase (MGL) forming glycerol and FFA which are released to plasma for systemic utilization.

Unlike its suppressive effect on adipogenesis in WAT, SIRT1 appears to promote mobilization and utilization of depot fat. Overexpression of SIRT1 in differentiated 3T3L1 cells resulted in decreased triglyceride levels and increased release of FFAs [23], while knockdown of SIRT1 decreased basal and stimulated lipolysis in adipocytes in culture. In *in vivo* studies using mice receiving high fat diet, activators of SIRT1 such as resveratrol reduced fat mass [69]. Further, over expression of SIRT1 inhibited diet induced accumulation of fat [6, 31, 52]. SIRT1 regulates the expression of ATGL gene and thereby lipolysis in adipocytes through modulation of the acetylation and transcriptional activity of FOXO1 [70]. SIRT2 also appears to show similar effect on fat mobilization [58, 59]. Although sirtuins influence mitochondrial oxidative metabolism in other metabolic tissues such as liver, their role in WAT mitochondrial metabolism is poorly understood.

2.3. Role of sirtuins in BAT development and browning of WAT

Unlike white adipogenesis, BAT biogenesis involves BMP7-stimulated commitment of Myf 5⁺ cells to brown pre-adipocytes that mature into mitochondria-rich brown adipocytes. BMP7 stimulation of progenitor cells leads to down regulation of early adipogenic inhibitors such as Pref1, Nectin, Wnt signaling molecules [43]. This is followed by upregulation of transcription factors such as PPAR γ and C/EBP α which cause upregulation of expression of PRDM16, a key factor involved in adipocyte-myocyte switch through activation of expression of BAT specific genes (PGC1 α , UCP-1, ZIC1) and downregulation of myogenic genes such as Myf5 and MyoD or myogenin [43]. This leads to increased biogenesis of mitochondria. PRDM16 is a 140 kDa protein that binds and activates the transcriptional function of PPAR γ and PGC1 α . PGC1 α co-activates PPAR γ -RXR α heterodimer stimulating the expression of BAT-specific genes UCP1 and UCP3.

The ability of human white adipocytes to acquire brown fat-like phenotype, termed browning, in response to β -adrenergic stimulation, cold exposure [43] and by several molecules such as muscle derived irisin [71, 72] liver derived FGF21 [73] and small molecules such as β -aminoisobutyric acid [74] has been observed. These type of cells called beige or brite adipocytes express genes involved in thermogenesis such as UCP1, deiodinase type II and PGC1 α in response to stimulation of β 3-adrenergic receptors [43].

Sirtuins also appear to play a role in the differentiation and function of BAT. Like WAT, brown adipose tissue also has been shown to express all the members of the mammalian SIRT family [11]. While the relative level of expression of SIRT3 and 5 are higher, that of SIRT1 and 7 are lower in BAT than those in WAT. Calorie restriction (CR) and cold exposure upregulated the expression of SIRT3 present in the mitochondria in BAT [75]; SIRT1 and SIRT2 also showed an upregulation under such conditions. Conversely, SIRT3 is down regulated in BAT in high fat diet induced obese mice. SIRT1 also appears to have a role in differentiation of pre-adipocyte to brown adipocytes. It appears that SIRT1 influences BAT differentiation through repression of the MyoD-mediated myogenic gene expression signature and stimulation of PGC-1 α mediated mitochondrial gene expression [76]. SIRT1 appeared to improve glucose homeostasis in SIRT1 transgenic mice and brown adipocytes derived from them due to an enhanced response of brown adipocytes to β 3-adrenergic stimuli rather than differences in differentiation status [77]. Of the different sirtuins, SIRT3 appeared to be critical in the differentiation of brown adipocytes. SIRT3 has been shown to activate PGC1 α mediated thermogenic response in differentiating brown adipocytes.

Brown remodeling of white fat in response to cold exposure is shown to be regulated by SIRT1-dependent deacetylation of PPAR γ . SIRT1-dependent deacetylation of Lys 268 and Lys 293 of PPAR γ is required to recruit the co-activator PRDM16 to PPAR γ , leading to upregulation of BAT-specific genes and repression of WAT genes [78]. In response to different environmental stimuli, SIRT1 can differentially modulate PPAR γ in WAT. SIRT1 inhibits PPAR γ through local modulation of acetylation status of histones and recruitment of co-repressor NCoR in response to caloric restriction; but on cold exposure, it directly enhances PPAR γ signaling through deacetylation of PPAR γ itself [79]. SIRT1 deficiency in mice results in accumulation of lipid droplets and reduction of mitochondrial content in BAT indicating a role for SIRT1 in the white remodeling of BAT which appears to occur in obese conditions [80].

2.4. Sirtuins and obesity

Grossly elevated fat stores in adipose tissue with hypertrophic or hyperplastic adipocytes and concomitant development of blood vessels result in obesity. The relevance of sirtuins in adipose tissue development and metabolism and their effects on metabolism of glucose and lipids primarily in the liver, and insulin function suggest a possible link between sirtuins and obesity.

The level of expression and activity of SIRT1 decrease in adipose tissue in different obesity models. Expression of SIRT1 in adipose tissue of db/db leptin resistant obese mice and in mice fed on HFD was significantly low [53, 81]. Overexpression of SIRT1 in HFD-induced obese animals caused less inflammation and better glucose tolerance. SIRT1 expression in obese pigs

is reported to be lesser than that in lean pigs [82]. Apart from decrease in SIRT1 levels, its function is also affected by changes in its post translational modification in obesity. One of the important post-translational modifications of SIRT1 which has been shown to be affected in obesity leading to inhibition of its nuclear localization is casein kinase mediated phosphorylation of ser-164 which is enhanced in obese and not in lean animals [83]. Unlike SIRT1 which is decreased in WAT in obesity, there is no consensus on the changes in other sirtuins in obese WAT; while some reports show decrease in SIRT2–6, other reports do not show any significant differences between obese and respective controls. But in obese BAT, SIRT1 and SIRT3 are down regulated and SIRT7 is upregulated. It has been shown that mir34a, which regulates the expression and activity of Sirt1, is elevated in obesity [41]. A possible association of sirtuins with obesity and obesity-associated pathological conditions in humans has also been indicated mostly from observational studies [84–88]. There is significant reduction in sirtuins in adipose tissue and other metabolic tissues in obese subjects and that weight loss or long term fasting can result in increase in their expression.

2.4.1. Sirtuins, insulin response and energy homeostasis

Insulin resistance is a hallmark of obesity and a major factor contributing to obesity associated pathological conditions. *In vitro* and *in vivo* studies suggest that SIRT1 regulates insulin response. In insulin resistant cells where SIRT1 is down regulated, induction of SIRT1 expression increased insulin sensitivity [89]. SIRT1 regulated insulin-dependent glucose uptake in adipocytes. Increase in SIRT1 activity improved insulin sensitivity [90]. Adipose tissue-specific SIRT1 knockout mice were reported to be more prone to developing insulin resistance. In experimentally induced diabetic animals, overexpression of SIRT1 increased insulin sensitivity. Mechanistically, SIRT1 effect appears to involve transcriptional repression of protein tyrosine phosphatase 1B gene which is critical in insulin signaling [91]. Along with SIRT1 in WAT, SIRT3 and SIRT5 contribute to systemic glucose homeostasis. As indicated before, SIRT1 also regulates insulin secretion by β -cells of pancreas by repressing UCP-2 [64]. Inhibition of SIRT1 expression reduced insulin secretion in β -cell lines; conversely overexpression of SIRT1 increased it. *In vivo*, transgenic mice over expressing SIRT1 in pancreatic β -cells showed increase in glucose-stimulated insulin secretion [92]. Further, SIRT1 deficiency impaired insulin secretion apparently by disrupting glucose sensing and impairing response to fluctuations in glucose levels [93].

In addition to its effect on peripheral tissue metabolism, SIRT1 in hypothalamus appears to act as a key regulator of central control of energy homeostasis. Evidence in support of this include (a) increase in the expression and activity of SIRT1 in hypothalamus in both calorie restriction and fasting [19, 94] (b) inhibition of hypothalamic SIRT1 expression, specifically in anorexigenic POMC neurons, resulted in loss of response to leptin and reduced energy expenditure indicating requirement of SIRT1 in POMC neurons for homeostatic defense against diet-induced obesity [95] (c) deletion of SIRT1 expression specifically in orexigenic Agouti-related peptide (AgRP)-expressing neurons, which promotes feeding in response to fasting, decreased AgRP neuronal activity resulting in decreased food intake and body weight [96] (d) Central inhibition of SIRT1 in rodents on a high fat diet caused decreased body weight and increased energy expenditure. This is mediated through increased acetylated-FoxO1-mediated increased production of POMC and its active product α MSH which in turn augmented TRH and T3 levels suggesting a hypothalamic–pituitary–thyroid axis which stimulates energy expenditure [97].

2.4.2. *Sirtuins and inflammation in adipose tissue*

Inflammation of adipose tissue in obesity is a major contributor to insulin resistance and pathogenesis of the metabolic syndrome [98]. SIRT1 could act as a transcriptional regulator of inflammation in multiple tissues, particularly adipose tissue as well as macrophages and endothelial cells [90, 99]. The levels of SIRT1 is inversely related to inflammation in adipose tissue. SIRT1 expression in human subcutaneous adipose tissue was less in cases where macrophage infiltration was high [100–102]. The decrease in SIRT1 in obese conditions in adipose tissue is suggested to be due to its proteosomal degradation. Activation of C-jun N terminal kinase (JNK1), which is a key component in inflammation associated signaling pathway, leads to phosphorylation of SIRT1, followed by its degradation in proteasomes [53, 103]. The molecular basis of the beneficial effect of SIRT1 on inflammation is related to suppression of NFkB activation [104]. SIRT1 inhibits transcriptional activity of NFkB directly by deacetylating the RelA/p65 subunit of NFkB at Lys 310 [105]. Moreover mir34a dependent decrease in SIRT1 activity can increase NFkB activity [41]. This suggests that SIRT1 and inflammatory signals interact at various levels and that SIRT1 is an important molecular link between nutrients, inflammation and metabolic dysfunction of the tissue. Though not much data on the role of other sirtuins in inflammation in human subjects is available, SIRT5 expression levels also correlate inversely with markers of inflammation [106] Deletion of SIRT7 in mice reduced WAT inflammatory gene expression in HFD induced obesity, suggesting opposing functions for SIRT1 and 7 [107].

2.4.3. *Sirtuin activators for therapy*

Since sirtuins play an important role in regulation of adipogenesis, and adipose tissue metabolism, pharmacological activation of sirtuins could be a useful approach for the treatment of obesity and related metabolic disorders. Sirt1 is an allosteric enzyme which is regulated by ligand binding. Resveratrol, which is a naturally occurring polyphenol with anti-oxidant property, increased the enzyme activity of SIRT1 by binding to its allosteric site [108]. High throughput screening has identified several small molecular activators of SIRT1 [109]. The most potent of these is SRT1720 which, protects against diet induced obesity [6]. The administration of SRT1720 reduced expression of lipogenic enzymes and reduced hepatic lipid accumulation, it also enhanced oxidative metabolism in skeletal muscle, liver and BAT in mice, protecting from HFD induced obesity and insulin resistance [110]. However, increasing sirtuin activity could result in indiscriminate deacetylation of histones and several other key proteins in different tissues. Sirtuin activating therapies would therefore have to be target specific.

3. Conclusions

Epigenetic modifications have emerged as fundamental modulators of metabolic functions, and sirtuins, a group of class III histone deacetylases, play a key role in this context. In this chapter the regulatory effects of sirtuins on adipose tissue metabolism in both WAT and BAT, and implication of alterations in their expression and activity in obesity, inflammation, and insulin resistance have been highlighted.

In spite of considerable advances in molecular biology of mammalian sirtuins, many questions remain unanswered. Among the different sirtuins, the role of SIRT1 in development and metabolism of white adipose tissue is reasonably well known. Although all the other sirtuins are expressed in WAT, their role in WAT function is not clear. Similarly, the role of SIRT1 and 3 in brown remodeling of white fat has been elucidated, but the regulatory effects of the other sirtuins are still unknown.

Different sirtuins control similar cellular processes in adipose tissue. Unraveling the potential crosstalk and coordination between them will require further study. The significance of this in the possible gene regulatory network and coordinated action among sirtuins in metabolic regulation is evident from the antagonistic interaction between SIRT1 and 7 in adipose tissue metabolism.

Apart from functional differences between WAT and BAT, variations in the status of different depots of WAT are also related to a risk for obesity-associated diseases. It is not clear whether there is any depot dependent variation in sirtuin action.

As the expansion of a vascular tissue like adipose tissue is associated with neovascularization, adipogenesis and angiogenesis are interrelated. Understanding the role of sirtuins in adipose tissue angiogenesis is of paramount importance especially in brown tissue, where both the mitochondrial activity and oxygen demand are high.

Sirtuins appear to be an attractive target for the treatment of obesity and related metabolic disorders. Increasing sirt activity in adipose tissue by identifying natural compounds, or engineering small molecular activators is an area which needs intensive research; increasing intracellular levels of NAD⁺, a substrate for sirtuins, is an alternate approach.

Acknowledgements

Fellowship received from ICMR, New Delhi to MS under TSS Fellowship scheme is gratefully acknowledged.

Conflict of interest

The authors declare that there is no conflict of interest.

Author details

Manu Sudhakar^{1*}, Santhi Silambanan¹ and Ramya Ramakrishnan²

*Address all correspondence to: manu.sudhakar@gmail.com

1 Department of Biochemistry, Sri Ramachandra Medical College and Research Institute, Chennai, India

2 Department of Surgery, Sri Ramachandra Medical College and Research Institute, Chennai, India

References

- [1] Richards EJ, Elgin SCR. Epigenetic codes for heterochromatin formation and silencing: Rounding up the usual suspects. *Cell*. 2002;**108**:489-500
- [2] Grunstein M. Histone acetylation in chromatin structure and transcription. *Nature*. 1997;**389**:349-352
- [3] Lopez-Rodas G, Brosch G, Georgieva EI, et al. Histone deacetylase: A key enzyme for the binding of regulatory proteins to chromatin. *FEBS Letters*. 1993;**317**:175-180
- [4] de Ruijter A, van Gennip AH, Caron HN, et al. Histone deacetylases (HDACs): Characterization of the classical HDAC family. *The Biochemical Journal*. 2003;**370**:737-749
- [5] Gregoret I, Lee Y, Goodson HV. Molecular evolution of the histone deacetylase family: Functional implications of phylogenetic analysis. *Journal of Molecular Biology*. 2004;**338**:17-31
- [6] Houtkooper RH, Pirinen E, Auwerx J. Sirtuins as regulators of metabolism and healthspan. *Nature Reviews. Molecular Cell Biology*. 2012;**13**:225-238
- [7] Fiorino E, Giudici M, Ferrari A, et al. The Sirtuin class of histone deacetylases: Regulation and roles in lipid metabolism. *IUBMB Life*. 2014;**66**:89-99
- [8] Chang HC, Guarente L. SIRT1 and other sirtuins in metabolism. *Trends in Endocrinology & Metabolism*. 2014;**25**:138-145
- [9] Jing H, Lin H. Sirtuins in epigenetic regulation. *Chemical Reviews*. 2015;**115**:2350-2375
- [10] Poulou N, Raju R. Sirtuin regulation in aging and injury. *Biochimica et Biophysica Acta*. 1852;**2015**:2442-2455
- [11] Jokinen R, Pirnes-Karhu S, Pietilainen KH, et al. Adipose tissue NAD⁺ homeostasis, sirtuins and poly(ADP-ribose)polymerases-important players in mitochondrial metabolism and metabolic health. *Redox Biology*. 2017;**12**:246-263
- [12] Rine J, Herskowitz I. Four genes responsible for a position effect on expression from HML and HMR in *Saccharomyces cerevisiae*. *Genetics*. 1987;**116**:9-22
- [13] Gotta M, Strahl-Bolsinger S, Renauld H, et al. Localization of Sir2p: The nucleolus as a compartment for silent information regulators. *The EMBO Journal*. 1997;**16**:3243-3255
- [14] Kaeberlein M, McVey M, Guarente L. The SIR2/3/4 complex and SIR2 alone promote longevity in *Saccharomyces cerevisiae* by two different mechanisms. *Genes & Development*. 1999;**13**:2570-2580
- [15] Imai S, Armstrong CM, Kaeberlein M, et al. Transcriptional silencing and longevity protein Sir2 is an NAD-dependent histone deacetylase. *Nature*. 2000;**403**:795-800
- [16] Sauve AA, Wolberger C, Schramm VL, et al. The biochemistry of sirtuins. *Annual Review of Biochemistry*. 2006;**7**:435-465

- [17] Imai S, Guarente L. Ten years of NAD-dependent Sir2 family deacetylases: Implications for metabolic diseases. *Trends in Pharmacological Sciences*. 2010;**31**:212-220
- [18] Watroba M, Szukiewicz D. The role of sirtuins in aging and age-related diseases. *Advances in Medical Sciences*. 2016;**61**:52-62
- [19] Cakir I, Perello M, Lansari O, et al. Hypothalamic SIRT1 regulates food intake in a rodent model system. *PLoS One*. 2009;**4**(12):e8322
- [20] Mattagajasingh I, Cuk SK, Asma N, et al. SIRT1 promotes endothelium-dependent vascular relaxation by activating endothelial nitric oxide synthase. *Proceedings of the National Academy of Sciences*. 2007;**104**:14855-14860
- [21] Sundaresan NR, Samant SA, Pillai VB, et al. SIRT3 is a stress-responsive deacetylase in cardiomyocytes that protects cells from stress-mediated cell death by deacetylation of Ku70. *Molecular and Cellular Biology*. 2008;**28**:6384-6401
- [22] Rodgers JT, Lerin C, Haas W, et al. Nutrient control of glucose homeostasis through a complex of PGC-1 α and SIRT1. *Nature*. 2005;**434**:113-118
- [23] Picard F, Kurtev M, Chung N, et al. SIRT1 promotes fat mobilization in white adipocytes by repressing PPAR γ . *Nature*. 2004;**429**:771-776
- [24] Rodgers JT, Puigserver P. Fasting-dependent glucose and lipid metabolic response through hepatic sirtuin 1. *Proceedings of the National Academy of Sciences*. 2007;**104**:12861-12866
- [25] Jiang W, Wang S, Xiao M, et al. Acetylation regulates gluconeogenesis by promoting PEPCK1 degradation via recruiting the UBR5 ubiquitin ligase. *Molecular Cell*. 2011;**43**(1):33-44
- [26] North BJ, Marshall BL, Borra MT, et al. The human Sir2 ortholog, SIRT2, is an NAD⁺-dependent tubulin deacetylase. *Molecular Cell*. 2003;**11**:437-444
- [27] Hirschey MD, Shimazu T, Jing E, et al. SIRT3 deficiency and mitochondrial protein hyperacetylation accelerate the development of the metabolic syndrome. *Molecular Cell*. 2011;**44**:177-190
- [28] MJ1 R, Newman JC, Held JM, et al. Label-free quantitative proteomics of the lysine acetylome in mitochondria identifies substrates of SIRT3 in metabolic pathways. *Proceedings of the National Academy of Sciences of the United States of America*. 2013;**110**:6601-6606
- [29] Ahuja N, Schwer B, Carobbio S, et al. Regulation of insulin secretion by SIRT4, a mitochondrial ADP-ribosyltransferase. *The Journal of Biological Chemistry*. 2007;**282**:33583-33592
- [30] Haigis MC, Mostoslavsky R, Haigis KM, et al. SIRT4 inhibits glutamate dehydrogenase and opposes the effects of calorie restriction in pancreatic beta cells. *Cell*. 2006;**126**:941-954
- [31] Buler M, Andersson U, Hakkola J. Who watches the watchmen? Regulation of the expression and activity of sirtuins. *The FASEB Journal*. 2016;**30**:3942-3960

- [32] Kumar S, Lombard DB. Mitochondrial sirtuins and their relationships with metabolic disease and cancer. *Antioxidants & Redox Signaling*. 2015;**22**:1060-1077
- [33] Beausoleil SA, Jedrychowski M, Schwartz D, et al. Large-scale characterization of HeLa cell nuclear phosphoproteins. *Proceedings of the National Academy of Sciences of the United States of America*. 2004;**101**:12130-12135
- [34] Yang Y, Fu W, Chen J, et al. SIRT1 sumoylation regulates its deacetylase activity and cellular response to genotoxic stress. *Nature Cell Biology*. 2007;**9**:1253-1262
- [35] Zhao W, Kruse JP, Tang Y, et al. Negative regulation of the deacetylase SIRT1 by DBC1. *Nature*. 2008;**451**:587-590
- [36] Kim JE, Chen J, Lou Z. DBC1 is a negative regulator of SIRT1. *Nature*. 2008;**451**:583-586
- [37] Krishnakumar R, Kraus WL. The PARP side of the nucleus: Molecular actions, physiological outcomes, and clinical targets. *Molecular Cell*. 2010;**39**:8-24
- [38] Fang J, Ianni A, Smolka C, et al. SIRT7 promotes adipogenesis in the mouse by inhibiting autocatalytic activation of SIRT1. *Proceedings of the National Academy of Sciences of the United States of America*. 2017;**114**:E8352-E8361
- [39] Lin SJ, Ford E, Haigis M, et al. Calorie restriction extends yeast life span by lowering the level of NADH. *Genes & Development*. 2004;**18**:12-16
- [40] Bitterman KJ, Anderson RM, Cohen HY, et al. Inhibition of silencing and accelerated aging by nicotinamide, a putative negative regulator of yeast sir2 and human SIRT1. *The Journal of Biological Chemistry*. 2002;**277**:45099-45107
- [41] Choi SE, Fu T, Seok S, et al. Elevated microRNA-34a in obesity reduces NAD⁺ levels and SIRT1 activity by directly targeting NAMPT. *Aging Cell*. 2008;**28**:4439-4448
- [42] Rayalam S, Baile CA. Adipocyte growth and factors influencing adipocyte life cycle. In: Symonds ME, editor. *Adipose Tissue Biology*. New York: Springer; 2012. pp. 195-226
- [43] Rodriguez A, Ezquerro S, Mendez Gimenez L, et al. Revisiting the adipocyte: A model for integration of cytokine signalling in the regulation of energy metabolism. *American Journal of Physiology: Endocrinology and metabolism*. 2015;**309**:E691-E714
- [44] Rutkowski JM, Stern JH, Scherer PE. The cell biology of fat expansion. *The Journal of Cell Biology*. 2015;**208**:501-512
- [45] Luo L, Liu M. Adipose tissue in control of metabolism. *The Journal of Endocrinology*. 2016;**231**:R77-R99
- [46] Tang QQ, Otto TC, Lane MD. Commitment of C3H10T1/2 pluripotent stem cells to the adipocyte lineage. *Proceedings of the National Academy of Sciences of the United States of America*. 2004;**101**:9607-9611
- [47] Huang H, Song TJ, Li X, et al. BMP signaling pathway is required for commitment of C3H10T1/2 pluripotent stem cells to the adipocyte lineage. *Proceedings of the National Academy of Sciences of the United States of America*. 2009;**106**:12670-12675

- [48] Rosen ED, MacDougald OA. Adipocyte differentiation from the inside out. *Nature Reviews. Molecular Cell Biology*. 2006;**7**:885-896
- [49] Bäckesjö CM, Li Y, et al. Activation of SIRT1 decreases adipocyte formation during osteoblast differentiation of mesenchymal stem cells. *Journal of Bone and Mineral Research*. 2006;**21**:993-1002
- [50] Cohen HY et al. Calorie restriction promotes mammalian cell survival by inducing the sirt1 deacetylase. *Science*. 2004;**305**:390-392
- [51] Chen D et al. Tissue-specific regulation of SIRT1 by calorie restriction. *Genes & Development*. 2008;**22**:1753-1757
- [52] Bordone L, Cohen D, Robinson A, et al. SIRT1 transgenic mice show phenotype resembling calorie restriction. *Aging Cell*. 2007;**6**:759-767
- [53] Chalkiadaki A, Guarente L. High fat diet triggers inflammation induced cleavage of SIRT1 in adipose tissue to promote metabolic dysfunction. *Cell Metabolism*. 2012;**16**:180-188
- [54] Simic P, Zainabadi K, Bell E, et al. SIRT1 regulates differentiation of mesenchymal stem cells by deacetylating β -catenin. *EMBO Molecular Medicine*. 2013;**5**:430-440
- [55] Tontonoz P, Spiegelman BM. Fat and beyond: The diverse biology of PPAR γ . *Annual Review of Biochemistry*. 2008;**77**:289-312
- [56] Jin Q, Zhang F, Yan T, et al. C/EBP α regulates SIRT1 expression during adipogenesis. *Cell Research*. 2010;**20**:470-479
- [57] Ahn J, Lee H, Jung CH, et al. MicroRNA 146b promotes adipogenesis by suppressing the SIRT- FoxO1 cascade. *EMBO Molecular Medicine*. 2013;**5**:1602-1612
- [58] Jing E, Gesta S, Kahn CR. SIRT2 regulates adipocyte differentiation through FoxO1 acetylation/deacetylation. *Cell Metabolism*. 2007;**6**:105-114
- [59] Wang F, Tong Q. SIRT2 suppresses adipocyte differentiation by deacetylating FOXO1 and enhancing FOXO1's repressive interaction with PPAR γ . *Molecular Biology of the Cell*. 2009;**20**:801-808
- [60] Nie Y, Erion DM, Yuan Z, et al. STAT3 inhibition of gluconeogenesis is downregulated by SIRT1. *Nature Cell Biology*. 2009;**11**:492-500
- [61] Farmer SR. Transcriptional control of adipocyte formation. *Cell Metabolism*. 2006;**4**:263-273
- [62] Tamori Y, Masugi J, Nishino N, et al. Role of PPAR γ in maintenance of the characteristics of mature 3T3-L1 adipocytes. *Diabetes*. 2002;**51**:2045-2055
- [63] Imai T, Takakuwa R, Marchand S, et al. PPAR γ is required in mature white and brown adipocytes for their survival in the mouse. *Proceedings of the National Academy of Sciences of the United States of America*. 2004;**101**:4543-4547
- [64] Bordone L, Motta MC, Picard F, et al. SIRT1 regulates insulin secretion by repressing UCP2 in pancreatic beta cells. *PLoS Biology*. 2006;**4**:e31

- [65] Laurent G, German NJ, et al. SIRT4 coordinates the balance between lipid synthesis and catabolism by repressing malonyl CoA decarboxylase. *Molecular Cell*. 2013;**50**:686-698
- [66] Laurent G, de Boer VCJ, Finley WS, et al. SIRT4 represses PPAR α activity to suppress hepatic fat oxidation. *Molecular and Cellular Biology*. 2013;**33**:4552-4561
- [67] Kanfi Y, Peshti V, Gil R, et al. SIRT6 protects against pathological damage caused by diet-induced obesity. *Aging Cell*. 2010;**9**:162-173
- [68] Lass A, Zimmermann R, Oberer M, et al. Lipolysis – A highly regulated multi-enzyme complex mediates the catabolism of cellular fat stores. *Progress in Lipid Research*. 2011;**50**:14-27
- [69] Feige JN, Lagouge M, Canto C, et al. Specific SIRT1 activation mimics low energy levels and protects against diet-induced metabolic disorders by enhancing fat oxidation. *Cell Metabolism*. 2008;**8**:347-358
- [70] Chakrabarti P, English T, Karki S, et al. SIRT1 controls lipolysis in adipocytes via FOXO1-mediated expression of ATGL. *Journal of Lipid Research*. 2011;**52**:1693-1701
- [71] Böström P, Wu J, Jedrychowski MP, et al. A PGC1- α -dependent myokine that drives brown-fat-like development of white fat and thermogenesis. *Nature*. 2012;**481**:463-468
- [72] Rodríguez A, Becerril S, Méndez-Giménez L, et al. Leptin administration activates irisin-induced myogenesis via nitric oxide-dependent mechanisms, but reduces its effect on subcutaneous fat browning in mice. *International Journal of Obesity*. 2015;**39**:397-407
- [73] Fisher FM, Kleiner S, Douris N, et al. FGF21 regulates PGC-1 α and browning of white adipose tissues in adaptive thermogenesis. *Genes & Development*. 2012;**26**:271-281
- [74] Roberts LD, Bostrom P, O'Sullivan JF, et al. Beta-aminoisobutyric acid induces browning of white fat and hepatic beta-oxidation and is inversely correlated with cardiometabolic risk factors. *Cell Metabolism*. 2014;**19**:96-108
- [75] Shi T, Wang F, et al. SIRT3, a mitochondrial sirtuin deacetylase, regulates mitochondrial function and thermogenesis in brown adipocytes. *The Journal of Biological Chemistry*. 2005;**280**:13560-13567
- [76] Timmons JA, Wennmalm K, Larsson O, et al. Myogenic gene expression signature establishes that brown and white adipocytes originate from distinct cell lineages. *Proceedings of the National Academy of Sciences of the United States of America*. 2007;**104**:4401-4406
- [77] Boutant M, Joffraud M, Kulkarni SS, et al. SIRT1 enhances glucose tolerance by potentiating brown adipose tissue function. *Molecular Metabolism*. 2014;**18**:118-131
- [78] Qiang L, Wang L, Kon N, et al. Brown remodeling of white adipose tissue by SIRT1-dependent deacetylation of PPAR γ . *Cell*. 2012;**150**:620-632
- [79] Li X. SIRT1 and energy metabolism. *Acta Biochimica et Biophysica Sinica*. 2013;**45**:51-60
- [80] Fen X, Xiaobin Z, Lin B, et al. Diet-induced obesity and insulin resistance are associated with brown Fat degeneration in SIRT1 deficient mice. *Obesity*. 2016;**24**:634-642

- [81] Qiao L, Shao J. SIRT1 regulates adiponectin gene expression through Foxo1-C/EBP α transcriptional complex. *The Journal of Biological Chemistry*. 2006;**281**:39915-39924
- [82] Pang W, Wang Y, Wei N, et al. SIRT1 inhibits Akt2-mediated porcine adipogenesis potentially by direct protein-protein interaction. *PLoS One*. 2013;**8**:e71576
- [83] Choi SE, Kwon S, Seok S, et al. Obesity-linked phosphorylation of SIRT1 by casein kinase 2 inhibits its nuclear localization and promotes fatty liver. *Molecular and Cellular Biology*. 2017;**37**(15):e00006-17
- [84] Pedersen SB, Ølholm J, Paulsen SK, et al. Low SIRT1 expression, which is upregulated by fasting, in human adipose tissue from obese women. *International Journal of Obesity*. 2008;**32**:1250-1255
- [85] Rutanen J, Yaluri N, Modi S, et al. SIRT1 mRNA expression may be associated with energy expenditure and insulin sensitivity. *Diabetes*. 2010;**59**:829-835
- [86] de Kreutzenberg SV, Ceolotto G, Papparella I, et al. Downregulation of the longevity associated protein sirtuin1 in insulin resistance and metabolic syndrome: Potential biochemical mechanisms. *Diabetes*. 2010;**59**:1006-1015
- [87] Moschen AR, Wiese V, Gerner RR, et al. Adipose tissue and liver expression of SIRT1, 3 and 6 increase after extensive weight loss in morbid obesity. *Journal of Hepatology*. 2013;**59**:1315-1322
- [88] Zillikens MC, van Meurs JB, Rivadeneira F, et al. SIRT1 genetic variation is related to BMI and risk of obesity. *Diabetes*. 2009;**58**:2828-2834
- [89] Banks AS, Kon N, Knight C, et al. SIRT1 gain of function increases energy efficiency and prevents diabetes in mice. *Cell Metabolism*. 2008;**8**:333-341
- [90] Yoshizaki T, Milne JC, Imamura T, et al. SIRT1 exerts anti-inflammatory effects and improves insulin sensitivity in adipocytes. *Molecular and Cellular Biology*. 2009;**29**:1363-1374
- [91] Sun C, Zhang F, Ge X, et al. SIRT1 improves insulin sensitivity under insulin-resistant conditions by repressing PTP1B. *Cell Metabolism*. 2007;**6**:307-319
- [92] Ramsey KM, Mills KF, Satoh A, et al. Age-associated loss of SIRT1-mediated enhancement of glucose-stimulated insulin secretion in beta cell-specific SIRT1 - overexpressing (BESTO) mice. *Aging Cell*. 2008;**7**:78-88
- [93] Luu L, Dai FF, Prentice KJ, et al. The loss of SIRT1 in mouse pancreatic beta cells impairs insulin secretion by disrupting glucose sensing. *Diabetologia*. 2013;**56**:2010-2020
- [94] Satoh A, Brace CS, Ben-Josef G, et al. SIRT1 promotes the central adaptive response to diet restriction through activation of the dorsomedial and lateral nuclei of the hypothalamus. *The Journal of Neuroscience*. 2010;**30**:10220-10232
- [95] Ramadori G, Fujikawa T, Fukuda M, et al. SIRT1 deacetylase in POMC neurons is required for homeostatic defence against diet-induced obesity. *Cell Metabolism*. 2010;**12**:78-87

- [96] Dietrich MO, Antunes C, Geliang G, et al. AgRP neurons mediate sirt's action on the melanocortin system and energy balancing roles for SIRT1 neuronal firing and synaptic plasticity. *The Journal of Neuroscience*. 2010;**30**:11815-11825
- [97] Cyr NE, Sleger JS, Toorie AM, et al. Central SIRT1 regulates body weight and energy expenditure along with the OMC derived peptide α MSH and the processing enzyme CPE production in diet induced obese male rats. *Endocrinology*. 2014;**155**:2423-2435
- [98] Hotamisligil GS. Inflammation and metabolic disorders. *Nature*. 2006;**444**:860-867
- [99] Yoshizaki T, Schenk S, et al. SIRT1 inhibits inflammatory pathways in macrophages and modulates insulin sensitivity. *American Journal of Physiology. Endocrinology and Metabolism*. 2010;**298**:E419-E428
- [100] Xu F, Gao Z, Zhang J, et al. Lack of SIRT1 (mammalian Sirtuin 1) activity leads to liver steatosis in the SIRT1 +/-mice: A role of lipid mobilization and inflammation. *Endocrinology*. 2010;**151**:2504-2514
- [101] Purushotham A, Schug TT, Xu Q, et al. Hepatocyte specific deletion of SIRT1 alters fatty acid metabolism and results in hepatic steatosis and inflammation. *Cell Metabolism*. 2009;**9**:327-338
- [102] Gillum PM, Kotas EM, Erion DM, et al. SIRT1 regulates adipose tissue inflammation. *Diabetes*. 2011;**60**:3235-3245
- [103] Gao ZJ, Kheterpal I, et al. Sirtuin 1 (SIRT1) protein degradation in response to persistent c-Jun N-terminal kinase (JnK1) activation contributes to hepatic steatosis in obesity. *The Journal of Biological Chemistry*. 2011;**286**:22227-22234
- [104] Kauppinen A, Suuronen T, et al. Antagonistic crosstalk between NF- κ B and SIRT1 in the regulation of inflammation and metabolic disorders. *Cell. Signall*. 2013;**25**:1939-1948
- [105] Yeung F, Hoberg JE, Ramsey CS, et al. Modulation of NF-kappaB-dependent transcription and cell survival by the SIRT1 deacetylase. *The EMBO Journal*. 2004;**23**:2369-2380
- [106] Jukarainen S, Heinonen S, Rämö JT, et al. Obesity is associated with low NAD⁺/SIRT pathway expression in adipose tissue of BMI-discordant monozygotic twins. *The Journal of Clinical Endocrinology and Metabolism*. 2016;**101**:275-283
- [107] Yoshizawa T, Karim MF, Sato Y, et al. SIRT7 controls hepatic lipid metabolism by regulating the ubiquitin-proteasome pathway. *Cell Metabolism*. 2014;**19**:712-721
- [108] Hubbard BP, Gomes AP, Dai H, et al. Evidence for a common mechanism of SIRT1 regulation by allosteric activators. *Science*. 2013;**339**:1216-1219
- [109] Sinclair DA, Guarente L. Small-molecule allosteric activators of Sirtuins. *Annual Review of Pharmacology and Toxicology*. 2014;**54**:363-380
- [110] Yamazaki Y, Usui I, Kanatani Y, et al. Treatment with SRT1720 a SIRT1 activator ameliorates fatty liver with reduced expression of lipogenic enzymes in MSG mice. *American Journal of Physiology. Endocrinology and Metabolism*. 2009;**297**:E1179-E1186

