We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists



186,000

200M



Our authors are among the

TOP 1% most cited scientists





WEB OF SCIENCE

Selection of our books indexed in the Book Citation Index in Web of Science™ Core Collection (BKCI)

Interested in publishing with us? Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected. For more information visit www.intechopen.com



Progress in Second Generation Ethanol Production with Thermophilic Bacteria

Sean Michael Scully and Johann Orlygsson

Additional information is available at the end of the chapter

http://dx.doi.org/10.5772/intechopen.78020

Abstract

Thermophilic bacteria have gained increased attention as prospective organisms for bioethanol production from lignocellulosic biomass due to their broad substrate spectra including many of the hexoses pentoses, and disaccharides found in biomass and biomass hydrolysates, fast growth rates, and high tolerance for extreme cultivation conditions. Apart from optimizing the ethanol production by varying physiological parameters, genetic engineering methods have been applied. This review focuses upon those thermophilic anaerobes recognized as being highly ethanologenic, their metabolism, and the importance of various culture parameters affecting ethanol yields, such as the partial pressure of hydrogen, pH, substrate inhibition, and ethanol tolerance. Also, recent developments in evolutionary adaptation and genetic engineering of thermophilic anaerobes are addressed.

Keywords: thermophilic bacteria, biofuel, bioethanol, lignocellulosic biomass, bioprocessing, genetic engineering

1. Introduction

The production of sustainable biofuels have increased in recent years because of a driving need for highly renewable and environmentally friendly energy carriers with bioethanol, biobutanol, biomethane, and biohydrogen being the most widely investigated. In order to meet international obligations to address climatic and geopolitical issues, many governments have set production targets as a response to meet these mandates. Ethanol production has been the main aim of many authorities as a suitable biofuel, for instance, a target set by the EU necessitates that 20% of energy production must be from renewable sources and energy

IntechOpen

© 2018 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/3.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

efficiency must increase by 20% while greenhouse gases must decrease by 20% by 2020 [1]. This has led to a dramatic increase in the production of bioethanol from 48 billion liters in 2007 to 2.6 billion liters in 2017 [2]. Both the United States and Brazil are by far the largest producers of bioethanol although the vast majority of ethanol produced is from first generation biomasses such as sucrose-rich sugarcane and easily fermentable starch-rich crops such as corn. However, there is a growing concern over the use of these feedstocks because they are food and feed related and thus in a direct competition with food production [3–5]. In addition, increased concern has been regarding the negative impact on agricultural areas used for the production of this biomass.

Production of bioethanol by second-generation non-food (lignocellulosic) biomass, such as agricultural residues, addresses some of the above mentioned environmental concerns although poses several challenges as a raw material for bioprocessing. Second generation biomass requires extensive and costly chemical or physical pretreatment in addition to enzy-matic treatment processes which negatively impacts its industrial feasibility. Lignocellulosic biomass is often difficult to degrade due to the lignin sheath and the highly crystalline nature of cellulose [6]. In addition to cellulose, lignocellulose is also composed of lignin and hemicelluloses of which the latter contains a plethora of monosaccharides with various connectivities and varying degrees of branching. Therefore, processing lignocellulosic biomass and subsequent fermentation of the liberated sugars to ethanol has proven to be a major complication for large-scale production [3–5].

To address the challenges posed by lignocellulosic biomass, fermentative organisms that can meet these process needs will help improve the feasibility of bioethanol production from lignocellulosic biomass. At present, the majority of bioethanol is produced using well-established mesophilic organisms despite some of the inherent advantages to the use of thermophilic microorganisms such as higher operating temperatures and utilizing a non-glucose hexoses and pentoses such as xylose and arabinose. This work focuses on the physiology of ethanol-producing thermophiles with an emphasis on their salient features relevant to the utilization of lignocellulosic biomass as well as the use of genetic engineering to improve their potential for bioethanol utilization.

2. Selected aspects of ethanol production from lignocellulosic biomass

For the fermentative production of ethanol from biomass to be commercially successful, several key processes and organisms need to be considered [3, 4, 7–9]. These process requirements needed to simultaneously consider two viewpoints: the physiological properties of the ethanologen used and process requirements. Concerning organism requirements, an ideal strains should be homoethanologenic, with high productivity (> 1/g/L/h), have broad substrate spectra and high tolerance of ethanol, inhibitory compounds and high initial substrate concentrations. Other key factors include high cellulolytic activity, simple nutritional needs, low biomass production and ease of genetic manipulation. Ideally, a single organism that

is both highly ethanologenic and cellulolytic would be ideal for consolidated bioprocessing (CBP) although co-cultures of organisms together fulfilling these requirements may also be considered in a simultaneous saccharification and fermentation (SSF) setup. To adequately meet the process requirements, ethanol yields should reach a minimum of 90% of theoretical, achieve high ethanol titers (> 5% v/v), have a minimum number of process steps, and require minimal or no process cooling. Additionally, cells should be robust enough to be recyclable and substrates co-fermented and pretreatment should be limited or excluded.

No single wild type organism possesses all these features. Although genetic manipulation has yielded only modest improvements for ethanologens although transformants are not always stable [9–11]. Many thermophilic clostridia have much broader substrate spectrum as compared to standard ethanologens such as *Saccharomyces cerevisiae* and *Zymomonas mobilis*. Additionally, cultivations of thermoanaerobes do not require extensive agitation or temperature control of the fermentation vessels and these often tolerate extremes of pH and salt concentrations during fermentation with minimal need for nutrient supplementation [4]. The mixing of reaction vessels and pumping of liquids are easier at elevated temperatures due to reduced viscosity as well increased substrate solubility [12].

3. Ethanol producing thermophilic anaerobes

While more than 300 species of thermophilic anaerobic bacteria have described as of 2008 from a wide range of environs with new species being continuously discovered. Thermophilic anaerobes have been isolated from a diverse range of environments [13] including deep-sea vents [14], geothermal areas [15–17], compost piles [18], municipal solid waste or sewage sludge [19], oil wells [20], and canned goods [21]. Most thermophilic microorganisms are either obligatory or facultative anaerobic, likely due to the limited availability of oxygen and highly reducing nature of geothermal features [22]. The majority of the those that are highly ethanologenic that have been described in the literature are often strict anaerobes within the genera of *Clostridium, Caloramator, Caldanaerobacter, Thermoanaerobacter*, or *Thermoanaerobacterium* [3, 23].

The highly polyphyletic genus *Clostridium* within the class *Clostridia* (family *Clostridiaceae*, order *Clostridiales*) currently has greater than 200 species with standing in nomenclature although only about 15 are strains within the genus are thermophilic [24, 25] usually with temperature optima for growth between 45 and 65°C although several strains reportedly grow at temperatures as high at 75°C. All species within the genus are strictly anaerobic and can typically be isolated from a broad range of nutrient-rich environments [26]. Many members within the genus can hydrolyze cellulose and produce ethanol, making them target of extensive research on biofuel production from complex [27, 28].

C. thermocellum is a thermophilic species that degrades crystalline cellulose using a cellulosome which is comprised of a complex arrangement of glycohyldrolases attached to a scaffold-like matrix [6]. Several other members of *Clostridium* have glycohydrolases including *C. acetobutylicum* [29, 30] and *C. cellulovorans* [31]. Ethanol yields by *Clostridium* species are often moderate and vary depending on environmental conditions with other organic acids,

including lactic acid, being common end-products. Examples of ethanol production from sugars by members of the genus include *Clostridium thermocellum* [32, 33] and *Clostridium* strain AK1 with 1.5 mol ethanol/mol glucose [34].

The genus *Thermoanaerobacterium* is comprised of thermophilic anaerobes which fall within Cluster V of *Clostridia* [35]. Currently, the genus currently consists of nine species and *T. thermo-sulfurigenes* is the genus type species [36]. Species within *Thermoanaerobacterium* are usually amyloand xylanolytic with a T_{opt} between 55 and 65°C and have been isolated from a diverse range of environments including geothermal features and from heat-treated canned foods [21, 37, 38]. They catabolize a broad range of hexoses, pentoses, and disaccharides to a mixture of ethanol, acetate, lactate, hydrogen, and CO₂. One challenge for these organisms is achieving good ethanol yields from high initial substrate concentrations which considerably lower ethanol yields. Examples of ethanol production from sugars by members of the genus include *Thermoanaerobacterium saccharolyticum* with 1.18 mol ethanol/mol glucose [37] and *Thermoanaerobacterium* strain AK17 with 1.50 and 1.33 mol ethanol/mol glucose and xylose, respectively [39].

Thermoanaerobacter species have similar physiological characteristics as Thermoanaerobacterium species; all species within the genus are highly saccharolytic and produce end-products which include ethanol, acetate, lactate, alanine, CO₂, and H₂. Nineteen species and five subspecies belong to the genus [24, 25]. The main difference between Thermoanaerobacter and Thermoanaerobacterium, is that the majority of Thermoanaerobacter species produce H₂S from thiosulfate whereas Thermoanaerobacterium produces sulfur [37]. Additionally, the temperature optima for Thermoanaerobacter species (65-75°C) are higher as compared to Thermoanaerobacterium species (55–65°C). The type species, *Thermoanaerobacter ethanolicus* and several other species within the genus, have been extensively studied for ethanol production [40–43]. High ethanol yields have been observed by several members of the genus including T. pseudoethanolicus, T. mathranii, T. pentosaceus, Thermoanaerobacter strain AK5, and Thermoanaerobacter strain J1 [17, 38, 44-46]. The ethanol yields, however, vary extensively depending on culture conditions [17, 38]. Recently, Thermoanaerobacter subterraneous was moved to the genus Caldanaerobacter that currently comprises two species: Caldanaerobacter subterraneous (and its four subspecies) and Caldanaerobacter uzonensis [24, 25]. Other representative examples of thermophilic ethanologenic bacteria can be found within the genera of Caldicellulosiruptor [47], Caloramator [48], Geobacillus [49], Caloramator boliviensis, for example, produces 1.53 mol ethanol/mol xylose [50].

4. Culture parameters

Most saccharolytic thermophiles use the Embden-Meyerhof-Parnas (EMP) pathway [5, 51] but do not use pyruvate decarboxylase for converting pyruvate to acetaldehyde as do yeasts. The theoretical yields of ethanol from 1 mol of hexose and pentose are 2.0 and 1.66 mol, respectively [5]. There are several routes from pyruvate to other end-products than ethanol. The following equations show the most common end-products from glucose with anaerobic bacteria:

- **1.** 1 Glucose \rightarrow 2 Ethanol +2 CO₂
- **2.** 1 Glucose \rightarrow 2 Lactate
- 3. 1 Glucose \rightarrow 2 Acetate +2 CO₂ + 4H₂
- 4. 1 Glucose \rightarrow 1 Butyrate +2 CO₂ + 2 H₂
- 5. 1 Glucose +2 $NH_4^+ \rightarrow$ 2 Alanine

Butyrate is not a commonly observed end-product with thermophilic anaerobes and alanine is not commonly assayed. The distribution of end products are known to be influenced by various factors which can be of direct relevance for the production of ethanol; these conditions include the substrate types and concentrations, the partial pressure of hydrogen, pH, and temperature. Some of these factors are discussed in detail below.

4.1. Partial pressure of hydrogen

Early observations of the influence of hydrogen concentrations on the end-product formation of Thermoanaerobacter ethanolicus were reported in 1981 [15]. Higher partial pressure of hydrogen (pH₂) leads to increased ethanol production and less acetate production from glucose fermentations [15, 38, 46]. Strict anaerobes produce H₂ either via pyruvate ferredoxin oxidorecutase or NAD(P)-dependent oxidoreductase [52]. It has been well established that the high concentrations affects mesophilic bacteria more severely than thermophiles because the NADH ferredoxin oxidoreductase (FNOR) that converts NADH to Fd_{red} is more strongly inhibited. The reduction potential is -400 mV for the Fd_{red}/Fd_{ox} couple but -320 mV for the NADH/NAD⁺ couple [52, 53]. Therefore, at low temperatures, elevated hydrogen concentrations inhibit H₂ evolution at much lower concentrations as compared to at high temperatures. Microorganisms respond to this by directing their reducing equivalents to other more favorable electron acceptors and consequently produce reduced products such as ethanol and lactate. In nature, hydrogen accumulation usually does not occur because of hydrogen-utilizing organism such as methanogens and sulfate-reducers, allowing for a complete catabolism of glucose to end-products. However, batch fermentation with monocultures allows hydrogen to accumulate leading to a change in end production profile in some *Thermoanaerobacter* species [15, 38, 41]. For instance, during degradation of glucose and xylose, the major end-product for Thermoanaerobacter brockii was ethanol [54]. Under hydrogen scavenging conditions, however, the flow of electrons from glucose degradation was directed away from ethanol but towards acetate with extra ATP produced. Several experiments using different liquid-to-gas ratios have revealed that changes in end-product formation occur during hydrogen accumulation among species of *Clostridium*, *Thermoanaerobacter*, and *Caloramator*. Hydrogen accumulation in these cultures can either change the carbon flow to more reduced end-products or inhibit substrate degradation. The inhibition observed can be either direct, inhibiting the hydrogenases, or indirect by productions of acids, lowering the pH in a closed system, and thus stopping further degradation of the substrates.

4.2. Substrate loadings

In natural environments of thermophilic bacteria, the concentration of sugars is relatively low. It is thus not surprising that most thermophilic bacteria are inhibited at relatively low (often between 10 and 30 mM) initial substrate concentrations as compared to yeasts and *Z. mobilis* [4, 38, 39, 46]. This inhibition may be caused by accumulated hydrogen or by acid accumulation and pH drop, or it could also be an intriguing factor for thermophiles. *Thermoanaerobacter*, strain J1, has been shown to be very tolerant towards high sugar concentrations [17]. This high ethanol producing thermophile produces up to 1.7 mol ethanol/mol glucose at 100 mM initial glucose concentration. Recent work on *Thermoanaerobacter pentosaceus* showed a complete removal of xylose at 13.3 mM initial concentrations but only about 30% removal at 10 times higher concentrations [55]. Additionally, the ratio of ethanol to acetate and lactate decreased by a factor of more than six resulting in dramatic decrease in ethanol yields.

4.3. Ethanol tolerance

One of the most important traits for good ethanol producers is their ethanol tolerance. For an economic ethanol recovery to occur, using classical downstream processes, the microorganism should grow and produce ethanol in the presence of at least 4% (v/v) ethanol [56]. It is well known that growth rates of many organisms decrease markedly with increasing ethanol concentrations because of leaky membranes resulting in loss of energy during cellular metabolism and finally cell lysis. Yeasts and *Z. mobilis* tolerate much higher ethanol concentrations as compared to thermophiles mainly due to their composition of fatty acid in their cell membrane.

Studies on ethanol tolerance of wild-type species of thermophiles show tolerance between 0.5 and 3.0% (v/v) [4, 46, 57, 58]. Substantial efforts to increase ethanol tolerance of wild type thermophiles, have been done. The highest ethanol tolerance observed for a thermophile has been with a mutant strain of *Thermoanaerobacter ethanolicus* (12.7% v/v) [28]. However, later studies with one of its mutant derivatives, JW200 Fe 4, showed much less tolerance [59]. *Thermoanaerobacter* BG1L1 showed 8.3% (v/v) tolerance in continuous culture studies [43] on xylose. Increased ethanol tolerance was also observed with *Thermoanaerobacter thermohydrosulfuricus* 39E by successively sub-culturing the strain to higher ethanol concentrations [57]. The resulting mutant strain 39EA tolerated 10.1% (v/v ethanol) at 45°C but only 2.6% (v/v) at 68°C. Additionally, the ethanol yields dropped considerably.

4.4. Other culture parameters

Other environmental factors of importance for thermophilic bacteria is their pH and temperature growth optimum, their tolerance towards inhibitory compounds like furfuraldehyde and 5-hydroxymethyl-furfuraldehyde (5-HMF) and their need for trace elements and vitamins often originating from complex medium supplements like yeast extract.

5. Production of ethanol from lignocellulose

Production of bioethanol from lignocellulosic biomass by wild type thermophilic bacteria has been widely reported in the literature where the focus has been mostly on *Clostridium*

thermocellum and species within the genera *Thermoanaerobacterium* and *Thermoanaerobacter*. However, there is a large variation in the type and concentration of biomass used, fermentation processes (batch, semi-batch, continuous), pretreatment methods as well as whether pure or mixed cultures are used.

The theoretical maximum yield of ethanol obtained from glucose fermentation is 0.51 g ethanol/ g glucose (2 mol ethanol/mol glucose or 11.1 mM/g). Unsurprisingly, considering the complex structure of lignocellulosic biomass, ethanol yields are usually considerably lower from such substrates as seen in **Table 1**.

Organisms	Substrate	Fermentation mode	Pre- treatment	Ethanol yields (mM/g)	Temp (°C)	References
Clostridium thermocellum	Avicel (2.5 g/L)	Batch	А	5.00	60	[60]
Clostridium thermocellum	Whatman paper (8.0 g/L)	Batch	None	7.20-8.00	60	[61]
Clostridium thermocellum	Paddy straw (8.0 g/L)	Batch	None	6.10-8.00	60	[61]
<i>Clostridium</i> strain DBT-IOC-C19	Avicel (10.0 g/L)	Batch	None	3.26	60	[62]
Clostridium strain AK1	Hemp (5.0 g/L)	Batch	A/Alk	3.5	50	[34]
Thermoanaerobacter pentosaceus	Rapeseed straw (5 0 g/L)	Con	Alk	1.40	70	[55]
Thermoanaerobacter mathranii	Wheat straw (6.7 g/L)	Batch	WO/E	2.61	70	[63]
Thermoanaerobacter ethanolicus	Beet molasses (30.0 g/L)	Batch	None	4.81	65	[64]
Thermoanaerobacter BG1L1	Corn stover, wheat straw (25.0–150.0 g/L)	Batch	WO/E	8.50–9.20	70	[42]
Thermoanaerobacter BG1L1	Wheat straw (30.0–120.0 g/L)	Batch	WO/E	8.50–9.20	70	[65]
T. ethanolicus	Wood HL (8.0 g/L)	Batch	E	3.30-4.50	70	[66]
Thermoanaerobacter strain AK5	Grass (4.5 g/L)	Batch	A/E	4.31	65	[38]
Thermoanaerobacter strain J1	Hemp (4.5 g/L)	Batch	A/E	4.3	65	[17]
T. saccharoylticum	Xylan (10.0 g/L)	Batch	WO	6.30	60	[67]
<i>Thermoanaerobacterium</i> strain AK17	Grass (2.5 g/L)	Batch	A/Alk/E	5.5	60	[39]

Cultivation were either in batch or continuous (con). Ethanol yields given in mM/g substrate degraded as well as substrate concentrations and incubation temperature are also shown. A—acid; Alk—alkaline; E—enzymatic; and WO—wet oxidation.

Table 1. Examples of ethanol production from lignocellulosic biomass by thermophilic bacteria.

Early experiments on ethanol production from lignocellulose included as the ethanol-producing organisms Thermoanaerobacter ethanolicus and Clostridium thermocellum with hemicellulose from birch- and beechwood as a feedstock [66]. Clostridium thermocellum produced between 7.2 mM ethanol /g and 8.0 mM/g from avicel and Whatman paper, respectively. Studies of ethanol production from paddy straw, sorghum stover and corn stubs, pretreated with alkali showed similar results [68]. However, these results were obtained with relatively low substrate loadings (8.0 g/L) but later studies showed that increased substrate loadings lowered the ethanol yields considerable [69]. The highest ethanol yields reported from lignocellulose are by Thermoanaerobacter BG1L1 grown on corn stover and wheat straw [42, 43] that were pretreated with acid or wet oxidation, or 9.2 mM/g for biomass hydrolysates. Thermoanaerobacterium strain AK17 showed ethanol yields of 2.0 (paper) mM/g, 2.9 (grass) mM/g and 5.8 (cellulose) mM/g biomass [23]. Optimization experiments showed an increase in ethanol yields on grass and cellulose up to 4.0 and 8.6 mM·g⁻¹, respectively. The main culture factors increasing ethanol yields was obtained by lowering of the substrate concentration from 7.5 to 2.5 g/L [39]. Recent investigations on two Thermoanaerobacter strains, AK5 and J1, showed promising results from various types of hydrolysates made from chemically and enzymatically pretreated lignocellulosic biomass [17, 38] (Table 1).

6. Evolutionary adaptation and genetic engineering of thermophiles

The thermophilic anaerobes mentioned in the previous sections make logical targets for genetic improvement due to their ability to produce ethanol from a wide range of substrates as evidenced by acceptable yields on lignocellulosic biomass. There are two general strategies for enhancing characteristics for ethanol production by wild type microorganisms: evolutionary adaptation or genetically modify the organisms. Early work often used classical methods such as the selection of clones and nonspecific mutagenesis to improve ethanol production [70]. These methods are time-consuming, and genetic modification is not without drawbacks as modified strains can exhibit poor growth and unexpected shifts in end-product formation. More recent work has focused more on modern techniques in molecular biology discussed herein.

6.1. Evolutionary adaptation

One of the major drawbacks of using thermophiles for the production of ethanol is their low substrate and ethanol tolerance. The use of classical evolutionary adaptation methods, such as non-specific mutagenesis and artificial selection, to enhance specific traits of microorganisms for industrial bioethanol production have been applied to thermophilic anaerobes on a limited basis. Examples of adaptation methods on three new mutant strains of *Thermoanaerobacter ethanolicus* were obtained by selection of pyruvate and iron deprivation [51] leading to enhanced ethanol tolerance (10% v/v) at substrate concentrations above 10 g/L. *Clostridium thermocellum* showed increased ethanol tolerance (up to 5% v/v) by stepwise increasing and transferring cultures to increased ethanol concentrations [71]. *Thermoanaerobacter pentosaceus* has been gradually adapted

to higher substrate concentrations and demonstrated higher ethanol tolerance and substrate utilization [72]. Thus, evolutionary adaptation, may still be used for evolving of wild type strains and further improving GM strains to meet requirements for tolerance to high ethanol titers, improve substrate utilization, and potentially resistance to inhibitory compounds generated during biomass pretreatment such as 5-HMF and fufuraldehyde.

6.2. Genetic engineering

Despite other promising features, one of the main drawback of most wild type thermophiles is their production of mixed end-products resulting in lower ethanol yields and the fact that highly ethanologenic organisms are not natively cellulolytic and *vice versa*. Two main strategies have been used to metabolically engineer thermophilic organisms for consolidated bioprocessing (CBP). The first strategy is to increase the ethanol yields of cellulase-producing organisms while the other is to express enzymes for biomass deconstruction in highly ethanologenic microorganisms [73, 74]. The first approach involves increasing ethanol yields by redirecting the flow of carbon and electrons which involves eliminating other potential fermentation products. Obvious targets include knocking out acetate and lactate pathways. The second approach involves addition of cellulolytic genes to the genome of a good ethanol producing bacterium.

The first thermophilic bacterium to be genetically modified to increase ethanol yields was *Thermoanaerobacterium saccharolyticum* in 2004 [75]. Since then, several other ethanologenic thermophiles have been genetically modified to increase ethanol titers and minimize the formation of other end-products (**Table 2**).

Deletion of genes involved to the production of various end-products to increase ethanol production capacity is the most obvious way to increase ethanol titers. This has been done by knocking out lactate dehydrogenase in *Thermoanaerobacterium saccharolyticum* [73, 82], *Thermoanaerobacter mathranii* [79], *Clostridium thermocellum* [83] and *Geobacillus thermogluco-sidasius* [78].

Wild type *Clostridium thermocellum* produces a mixture of ethanol, acetate, lactate, hydrogen, and carbon dioxide [84] from cellulose and other substrates. The first successful transformation of the species was performed in 2006 [85], later on leading to the development of genetic systems to knock out the *pta* gene and thus acetate formation [85]. However, growth of the resultant strain was abnormal although cellulase active remained intact. Later work on *C. thermocellum* showed improved ethanol yields in an adapted strain (Δhpt , Δldh , Δpta) lacking acetate and lactate pathways and was successfully used in co-culture with *Thermoanaerobacterium saccharolyticum* [85].

Early work on *Thermoanaerobacterium saccharolyticum* were performed by using electroporation and shuttle vectors [86], but later on this strain has been further modified by inserting a cellobiohydrolase gene from *Clostridium thermocellum* into its genome [77]. Also a *ldh* gene knock out was done using *Thermoanaerobacterium saccharolyticum* [75] and then a double knock out of both *ldh* and *ak* [73]. The knocking out of acetate production led to less available energy,

Strain	Genotype	Substrate	Mode	Ethanol yields (mol/mol)	References
C. thermocellum	∆ <i>pyrF,</i> ∆ <i>pta</i> ::gapDHp-cat	Cellobiose (5.0 g/L)	Batch	0.59	[76]
C. thermocellum	∆ <i>pyr</i> F, ∆ <i>pta</i> ::gapDHp-cat	Avicel (5.0 g/L)	Batch	0.71	[76]
C. thermocellum $adhE^*(EA)$ Δldh	Δhpt , Δldh	Cellobiose (5.0 g/L)	Batch	0.37	[77]
C. thermocellum	Δhpt , Δldh , Δpta (evolved)	Avicel (19.5 g/L)	Batch	1.08	[77]
C. thermocellum/T. saccharolyticum	Δhpt , Δldh , Δpta (evolved) and Δpta , Δack , Δldh	Avicel (19.5 g/L)	Batch	1.26	[77]
T. saccharolyticum TD1	Δldh	Xylolse (5.0 g/L)	Batch	0.98	[77]
T. saccharolyticum ALK2	$\Delta pta, \Delta ack, \Delta ldh$	Cellobiose (70.0 g/L)	Con	ND	[73]
T. saccharolyticium HK07	Δldh , Δhfs	Cellobiose (1.8 g/L)	Batch	0.86	[74]
T. saccharolyticium M0355	Δldh , Δack , Δpta	Cellobiose (50.0 g/L)	Batch	1.73	[74]
T. saccharolyticum M1051	Δldh , Δack Δpta , ure	Cellobiose (27.5 g/L)	Batch	1.73	[74]
G. thermoglucosidasius TM242	Δldh -, pdh up, pflB-	Glucose (34.0 g/L)	Batch	1.73	[78]
G. thermoglucosidasius TM242	Δldh -, pdh up, $\Delta pflB$ -	Glucose (34.0 g/L)	Batch	1.84	[78]
G. thermoglucosidasius TM242	Δldh -, Δpdh up, $\Delta pflB$ -	Xylose (29.0 g/L)	Batch	1.37	[78]
T. mathranii BG1L1	Δldh	Wheat straw (30-120 g/L)	Con	1.53–1.67	[65]
T. mathranii BG1G1	$\Delta ldh, GldA$	Glucose + glycerol (5.0 g/L)	Batch	1.68	[79]
T. mathranii BG1G1	$\Delta ldh, GldA$	Xylose + glycerol (5.0 g/L)	Batch	1.57	[79]
T. mathranii BG1G1	$\Delta ldh, GldA$	Xylose + glycerol (12.8 and 7.2 g/L)	Con	1.53	[79]
<i>Thermoanarobacter</i> Pentocrobe 411	Δldh , Δack , Δpta	Wheat straw (65 g/L)	Con	1.84	[80]
C. bescii JWCB018	Δldh -	Celo (10 g/L)	Batch	0	[81]
C. bescii JWCB032	Δldh -, $adhE$ +	Celo (10 g/L)	Batch	0.66	[81]
C. bescii JWCB049	$\Delta pyrFA$, Δldh -	Celo (10 g/L)	Batch	0.54	[81]
C. bescii JWCB054	$\Delta pyrFA$, Δldh -	Celo (10 g/L)	Batch	0.28	[81]

ack—acetate kinase; *GldA*—glycerol dehydrogenase A; *hfs*—hydrogenase; *hpt*—hypoxanthine phosphoribosyl transferase; *pdh*—pyruvate decarboxylase; *pyrF*—orotidine-5-phoshate decarboxylase; *pfl*—pyruvate formate lyase; and *ure*—urease.

Table 2. Ethanol yields of genetically engineered thermophilic bacteria from different substrates and fermentation conditions.

thus less cell biomass and increased ethanol yields, both from glucose and xylose. Another double knock out of *Thermoanaerobacterium saccharolyticum* focused on the electron transfer system of the bacterium [74]. The *hfs* gene cluster, which codes for hydrogenase, and the *ldh* gene were knocked out resulting in a considerable increase in ethanol (44%) production as compared with the wild type.

Thermoanaerobacter mathranii has been modified and used in several investigations. The first mutant generated was BG1L1 by knocking out *ldh* resulting in a more than two-fold increase in ethanol production as compared with the wild type [87]. This strain showed good ethanol yields from undetoxified pretreated corn stover and wheat straw [42, 43]. Further manipulation of this strain involves overexpression of NAD(P)H-dependent alcohol dehydrogenase, resulting in the strain BG1E1. Clearly, this enzyme is of great importance for ethanol production and its overexpression resulted in higher ethanol yields [79]. The electron balance for sugar degradation was additionally focused upon with this strain when mannitol, which is more reduced than glucose and xylose, was used as a substrate [87] and this resulted in higher ethanol yields. The BG1G1 strain of *Thermoanaerobacter mathranii* was developed which included the insertion of a NAD⁺-dependent glycerol dehydrogenase which increased ethanol yields by 40% greater than the type strain. Additionally, the strain utilized the highly reduced glycerol and co-metabolism of glycerol and sugars.

Recently, the highly ethanologenic strain *Thermoanaerobacter* BG1 "Pentocrobe 411" was genetically engineered by knocking out lactate dehydrogenase, phosphotransacetylase, and acetate kinase [80]. Pentocrobe 411 achieved very high ethanol titers (1.84 to 1.92 mol ethanol/ mol hexose equivalent) nearing the maximum theoretical yield from hexoses and pentoses on various pretreated biomass in continuous culture.

A natural target for the strategy of converting a cellulolytic organism into a good ethanol producer would be members of the genus of *Caldicellulosiruptor* which has several cellulolytic members although none are good ethanol producers. Recent work with *Caldicellulosiruptor bescii*, a naturally cellulolytic organism, has produced ethanol producing strains [89–93].

The type strains of C. bescii typically yield a mixture of lactic and acetic acid in addition to hydrogen and CO₂ as end-products although other strains within the genus of *Caldicellulosiruptor* have been noted to produce low ethanol titers. Work by Cha [89] deleted the gene coding for lactate dehydrogenase by introducing a non-replicating plasmid via marker replacement. The resultant knockout strain did demonstrate increased biomass yield as well as acetate and hydrogen production with no lactate production when grown on cellobiose and lactose as well as switch grass hydrolysates. Subsequent work by Chung [81] inserted a NADHdependent adhE gene (from Clostridium thermocellum) into the ldh mutant (JWCB018) resulting in strain *C. bescii* JWCB032. The resultant *ldh⁻ adhE⁺* strain yielded less acetate (4.3 mM) but produced 14.8 mM of ethanol from 29.2 mM cellobiose or 12.7% of the theoretical yield. It should be noted that this strain only used a small portion (4.4 mM of 29.2 mM cellobiose) provided and not produce ethanol above 65°C. Work by Cha [89] and Chung [93] introduced the alcohol dehydrogenase genes (adhB and adhE) from Thermoanaerobacter pseudoethanolicus into the *ldh* deficient strain. The two resultant strains yielded ethanol at temperatures greater than 65°C although titers were lower than the aforementioned strain JWCB032 ($ldh^{-} adhE^{+}$). The *C. thermocellum* strain with *adhB* only produced 1.4 mM ethanol on avicel and 0.4 mM on switch grass while a strain with *adhE* gave 2.3 and 1.6 mM of ethanol on avicel and switch grass, respectively. One of the reasons for suggested for the low ethanol titers is the availability of cofactors and it should be noted that T. pseudoethanolicus ADHs utilize NADPH while the gene products from *C. thermocellum* use NADH as a source of reducing potential. Additional work is therefore needed to more carefully mimic the complex NAD(P)H system of multiple ADHs in *Thermoanaerobacter pseduoethanolicus*.

Overall, efforts to engineer thermophilic anaerobes to increase ethanol titers has resulted in modest gains in yields while minimizing or eliminating the formation of unwanted end products. Future targets for genetic manipulation might include the inclusion of the cellulolytic machinery of *C. thermocellum* into highly ethanologenic *Thermoanaerobacter* and *Thermoanaerobacterium* strains.

7. Conclusions

Bioethanol production from lignocellulosic biomass with thermophilic bacteria needs robust microbes regarding several aspects. One of the main advantages of thermophilic bacteria is their broad substrate spectra with many strains capable of simultaneous pentose and hexose degradations. Additionally, some thermophiles degrade complex carbohydrates like cellulose and hemicellulose although many of these strains are not highly ethanologenic. Recent advantages in genetic engineering have improved ethanol yields, mostly by knocking out pathways of undesired end-products. On the back side is the fact that yields and ethanol tolerance as well as low tolerance for high initial substrate concentrations still limits the use of thermophiles for large scale operations. The use of stable co-cultures where on microbe hydrolyses the sugar polymers and the other one ferments the sugars released to ethanol is an attractive way to go forward but warrants further investigations.

Author details

Sean Michael Scully and Johann Orlygsson*

*Address all correspondence to: jorlygs@unak.is

Faculty of Natural Resource Sciences, University of Akureyri, Akureyri, Iceland

References

- [1] European Commission. Directive 2009/28/EC of the European Parliament and of the Council of 23 April 2009 on the promotion of the use of energy from renewable sources and amending and subsequently repealing Directives 2001/77/EC and 2003/30/EC; 2009
- [2] RFA Renewable fuels association. 2013. World fuel ethanol production. http://ethanolrfa.org/pages/World-Fuel-Ethanol-Production (Accessed: February 27, 2018)
- [3] Sánchez ÓJ, Cardona CA. Trends in biotechnological production of fuel ethanol from different feedstocks. Bioresource Technology. 2008;**99**:5270-5295
- [4] Taylor MP, Eley KL, Martin S, Tuffin MI, Burton SG, Cowan DA. Thermophilic ethanologenesis: Future prospects for second-generation bioethanol production. Trends in Biotechnology. 2009;27:398-405
- [5] Scully SM, Orlygsson J. Recent advantages in second generation ethanol production by thermophilic bacteria. Energies. 2015;8:1-30
- [6] Demain AL, Newcomb M, Wu JHD. Cellulase, *Clostridia*, and ethanol. Microbiology and Molecular Biology Reviews. 2005;**69**:124-154
- [7] Gnansounou E, Dauriat A. Techno-economic analysis of lignocellulosic ethanol: A review. Bioresource Technology. 2010;**101**:4980-4991
- [8] Chang T, Yao S. Thermophilic, lignocellulolytic bacteria for ethanol production: Current state and persepectives. Applied Microbiology and Biotechnology. 2011;**92**:13-27
- [9] Ostergaard S, Olsson L, Nielsen J. Metabolic engineering of *Saccharomyces cerevisae*. Microbiology and Molecular Biology Reviews. 2000;**64**:34-50
- [10] Jeffries TW. Engineering yeasts for xylose metabolism. Current Opinion in Biotechnology. 2006;17:320-326
- [11] He MX, Wu B, Qin H, Ruan ZY, Tan FR, Wang JL, Shui ZX, Dai LC, Zhu QL, Pan K, Tang XY, Wang WG, Hu QC. *Zymomonas mobilis*: A novel platform for future biorefineries. Biotechnology for Biofuels. 2014;7:101
- [12] Turner P, Mamo G, Karlsson EN. Potential and utilization of thermophiles and thermostable enzymes in biorefining. Microbial Cell Factories. 2007;6:9

- [13] Wagner ID, Wiegel J. Diversity of thermophilic anaerobes. In: Incredible anaerobes: From physiology to genomics fuels. Annals of the New York Academy of Sciences. 2008; 1125:1-43
- [14] Slobodkin AI, Tourova TP, Kuznetsov BB, Kostrikina NA, Chernyh NA, Bonch-Osmolovskaya EA. *Thermoanaerobacter siderophilus* sp. nov., a novel dis-similatory Fe(III)-reducing, anaerobic, thermophilic bacterium. International Journal of Systematic Bacteriology. 1999;49:1471-1478
- [15] Wiegel J, Ljungdahl LG. *Thermoanaerobacter ethanolicus* gen. Nov., spec. Nov., a new, extreme thermophilic, anaerobic bacterium. Archives of Microbiology. 1981;**128**:343-348
- [16] Larsen L, Nielsen P, Ahring BK. *Thermoanaerobacter mathranii* sp. nov, an ethanolproducing, extremely thermophilic anaerobic bacterium from a hot spring in Iceland. Archives of Microbiology. 1997;168:114-119
- [17] Jessen JE, Orlygsson J. Production of ethanol from sugars and lignocellulosic biomass by *Thermoanaerobacter* J1 isolated from a hot spring in Iceland. Journal of Biomedicine and Biotechnology. 2012;186982. DOI: 10.1155/2012/186982
- [18] Fong JCN, Svenson CJ, Nakasugi K, Leong CTC, Bowman JP, Chen B, Glenn DR, Neilan BA, Rogers PL. Isolation and characterization of two novel ethanol-tolerant facultative-anaerobic thermophilic bacteria strains from waste compost. Extremophiles. 2006;10:363-372
- [19] Sekiguchi Y, Imachi H, Susilorukmi A, Muramatsu M, Ohashi A, Harada H, Hanada S, Kamagata Y. *Tepidanaerobacter syntrophicus* gen. Nov., sp. nov., an anaerobic, moderately thermophilic, syntrophic alcohol- and lactate-degrading bacterium isolated from thermophilic digested sludges. International Journal of Systematic and Evolutionary Microbiology. 2006;56:1621-1629
- [20] Cayol JL, Ollivier B, Patel BKC, Ravot G, Magot M, Ageron E, Grimont PAD, Garcia JL. Description of *Thermoanaerobacter brockii* subsp. *lactiethylicus* subsp. nov., isolated from a deep subsurface French oil well, a proposal to reclassify *Thermoanaerobacter finnii* as *Thermoanaerobacter brockii* subsp.*finnii* comb. nov., and an emended description of *Thermoanaerobacter brockii*. International Journal of Systematic Bacteriology. 1995; 45:783-789
- [21] Cann IK, Stroot PG, Mackie KR, White BA, Mackie RI. Characterization of two novel saccharolytic, anaerobic thermophiles, *Thermoanaerobacterium polysaccharolyticum* sp. nov. and *Thermoanaerobacterium zeae* sp. nov., and emendation of the genus *Thermoanaerobacterium*. International Journal of Systematic and Evolutionary Microbiology. 2001;51: 293-302
- [22] Amend JP, Shock EL. Energetics of overall metabolic reactions of thermophilic and hyperthermophilic *Archaea* and bacteria. FEMS Microbiology Reviews. 2001;25:175-243
- [23] Sveinsdottir M, Baldursson SRB, Orlygsson J. Ethanol production from monosugars and lignocellulosic biomass by thermophilic bacteria isolated from Icelandic hot springs. Icelandic Agricultural Sciences. 2009;22:45-58

- [24] Euzéby JP. List of bacterial names with standing in nomenclature: A folder available on the. International Journal of Systematic Bacteriology. 1997;47:590-592
- [25] Parte AC. LPSN-list of prokaryotic names with standing in nomenclature. Nucleic Acids Research. 2014;42:D613-D616
- [26] Wiegel J, Tanner R, Rainey FA. An introduction to the family clostridae. In: Dworkin M, Falkow S, Rosenberg E, Schleifer K-H, Stackebrandt E, editors. The Prokaryotes, 3rd ed. Springer: New York, NY, USA; 2006; part 1. pp. 654-678
- [27] Canganella F, Wiegel J. The potential of thermophilic clostridia in biotechnology. In: Woods DR, editor. The Clostridia and Biotechnology. Vol. 23. Freepost, England: Butterworth-Heinemann; 1993. pp. 394-429
- [28] Carreira LH, Ljungdahl LG. Production of ethanol from biomass using anaerobic thermophilic bacteria. In: Wise DL, editor. Liquid Fuel Developments. Boca Raton, Flordia, USA: CRC Press, ISBN 0849360943; 1993. pp. 1-28
- [29] Nölling J, Breton G, Omelchenko MV, Makarova KS, Zeng Q, Gibson R, Lee HM, Dubois J, Qiu D, Hitti J, Wolf YI, Tatusov RL, Sabathe F, Doucette-Stamm L, Soucaille P, Daly MJ, Bennett GN, Koonin EV, Smith DR. Genome sequence and comparative analysis of the solvent-producing bacterium *Clostridium acetobutylicum*. Journal of Bacteriology. 2001;**183**:4823-4838
- [30] Sabathe F, Belaich A, Soucaille P. Characterization of the cellulolytic complex (cellulosome) of *Clostridium acetobutylicum*. FEMS Microbiology Letters. 2002;217:15-22
- [31] Han SO, Yukawa H, Inui M, Doi RH. Transcription of *Clostridium cellulovorans* cellulosomal cellulase and hemicellulase genes. Journal of Bacteriology. 2003;185:2520-2527
- [32] Balusu R, Paduru RMR, Seenyya G, Reddy G. Production of ethanol from cellulosic biomass by *Clostridium thermocellum* SS19 in submerged fermentation: Screening of nutrients using Plackett-Burman design. Applied Biochemistry and Biotechnology. 2004;117:133-141
- [33] Rani KS, Seenayya G. High ethanol tolerance of new isolates of *Clostridium thermocellum* strains SS21 and SS22. World Journal of Microbiology & Biotechnology. 1999;15:173-178
- [34] Orlygsson J. Ethanol production from biomass by a moderate thermophile. *Clostridium* AK1. Icelandic Agricultural Science. 2012;**25**:25-35
- [35] Collins MD, Lawson PA, Willems A, Cordoba JJ, Fernandez-Garayzabal J, Garcia P, Cai J, Hippe H, Farrow JA. The phylogeny of the genus *Clostridium*: Proposal of five new genera and eleven new species combinations. International Journal of Systematic Bacteriology. 1994;44:812-826
- [36] Schink B, Zeikus JG. *Clostridium thermosulfurogenes* sp. nov, a new thermophile that produces elementar sulfur from thiosulfate. Journal of General Microbiology. 1983;**129**: 1145-1158
- [37] Lee YE, Jain MK, Lee C, Lowe SE, Zeikus JG. Taxonomic distinction of saccharolytic thermophilic anaerobes: Description of *Thermoanaerobacterium xylanolyticum* gen. Nov., sp. nov., and *Thermoanaerobacterium saccharolyticum* gen. Nov., sp. nov.; reclassification

of Thermoanaerobium brockii, Clostridium thermosulfurogenes, and Clostridium thermohydrosulfuricum E100-69 as Thermoanaerobacter brockii comb. nov., Thermoanaerobacterium thermosulfurigenes comb. nov., and Thermoanaerobacter thermohydrosulfuricus comb. nov., respectively; and transfer of Clostridium thermohydrosulfuricum 39E to Thermoanaerobacter ethanolicus. International Journal of Systematic Bacteriology. 1993;**43**:41-51

- [38] Brynjarsdottir H, Wawiernia B, Orlygsson J. Ethanol production from sugars and complex biomass by *Thermoanaerobacter* AK₅: The effect of electron-scavenging systems on end-product formation. Energy and Fuels. 2012;26:4568-4574
- [39] Almarsdottir AR, Sigurbjornsdottir MA, Orlygsson J. Effects of various factors on ethanol yields from lignocellulosic biomass by *Thermoanaerobacterium* AK17. Biotechnology and Bioengineering. 2012;**109**:686-694
- [40] Lacis LS, Lawford HG. Ethanol-production from xylose by *Thermoanaerobacter ethanolicus* in batch and continuous culture. Archives of Microbiology. 1988;150:48-55
- [41] Lee Y-J, Dashti M, Prange A, Rainey FA, Rohde M, Whitman WB, Wiegel J. *Thermoan-aerobacter sulfurigignens* sp. nov., an anaerobic thermophilic bacerium that reduces 1 M thiosulfate to elemental sulfur and tolerates 90 mM sulfite. International Journal of Systematic and Evolutionary Microbiology. 2007;57:1429-1434
- [42] Georgieva TI, Ahring BK. Evaluation of continuous ethanol fermentation of dilute-acid corn Stover hydrolysate using thermophilic anaerobic bacterium *Thermoanaerobacter* BG1L1. Applied Microbiology and Biotechnology. 2007;77:61-68
- [43] Georgieva TI, Mikkelsen MJ, Ahring BK. High ethanol tolerance of the thermophilic anaerobic ethanol producer *Thermoanaerobacter* BG1L1. Cent. Europ. Journal of Biology. 2007;2:364-377
- [44] Lovitt RW, Shen GJ, Zeikus JG. Ethanol-production by thermophilic bacteria biochemical basis for ethanol and hydrogen tolerance in *Clostridium thermohydrosulfuricum*. Journal of Bacteriology. 1988;170:2809-2815
- [45] Larsen L, Nielsen P. *Thermoanaerobacter mathranii* sp. nov., an ethanol-producing extremely thermophilic bacterium from hot spring in Iceland. Archives of Microbiology. 1997;168: 114-119
- [46] Tomás AF, Karagöz P, Karakashev D, Angelidaki I. Extreme thermophilic ethanol production from rapeseed straw: Using the newly isolated *Thermoanaerobacter pentosaceus* and combining it with *Saccharomyces cerevisiae* in a two-step process. Biotechnology and Bioengineering. 2013;**110**:1574-1582
- [47] Svetlitchnyi VA, Kensch O, Falkenhan DA, Korseska SG, Lippert N, Prinz M, Sassi J, Schickor A, Curvers S. Single-step ethanol production from lignocellulose using novel extremely thermophilic bacteria. Biotechnology for Biofuels. 2013;6:31
- [48] Crespo C, Pozzo T, Karlsson EN, Alvarez MP, Mattiasson B. Caloramator boliviensis sp. nov., a thermophilic, ethanol-producing bacterium isolated from a hot spring. International Journal of Systematic and Evolutionary Microbiology. 2012;62:1679-1686

- [49] Zambare V, Bhalla A, Muthukumarappan K, Sani RK, Christopher L. Bioprocessing of agricultural waste to ethanol utilizing a cellulolytic extremophile. Extremophiles. 2011;**15**:611-618
- [50] Crespo RE, Badshah M, Alvarez MT, Mattiasson B. Ethanol production by continuous fermentation of d-(+)-cellobiose, d-(+)-xylose and sugarcane bagasse hydrolysate using the thermoanaerobe *Caloramator boliviensis*. Bioresource Technology. 2012; 103:186-191
- [51] He Q, Lokken PM, Chen S, Zhou J. Characterization of the impact of acetate and lactate on ethanolic fermentation by *Thermoanaerobacter ethanolicus*. Bioresource and Technology. 2009;**100**:5955-5965
- [52] Jones P. Improving fermentative biomass-derived H₂-production by engineered microbial metabolism. International Journal of Hydrogen Energy. 2008;33:5122-5130
- [53] Hallenbeck PC. Fermentative hydrogen production: Principles, progress and prognosis. International Journal of Hydrogen Energy. 2009;**34**:7379-7389
- [54] Fardeau ML, Patel BKC, Magot M, Ollivier B. Utilization of serine, leucine, isoleucine and valine by *Thermoanaerobacter brockii* in the presence of thiosulfate or *Methanobacterium* sp as electron acceptors. Anaerobe. 1997;3:405-410
- [55] Tomás AF, Karakashev D, Angelidaki I. *Thermoanaerobacter pentosaceus* sp. nov., an anaerobic, extreme thermophilic, high ethanol-yielding bacterium isolated from household waste. International Journal of Systematic and Evolutionary Microbiology. 2012;63:2396-2404
- [56] Hahn Hagerdahl B, Galbe M, Gorwa-Grauslund MF, Liden G, Zacchi G. Bio-ethanol the fuel of tomorrow from residues today. Trends in Biotechnology. 2006;24:549-556
- [57] Lovitt RW, Longin R, Zeikus JG. Ethanol production by thermophilic bacteria: Physiological comparison of solvent effects on parent and alcohol-tolerant strains of *Clostridium thermohydrosulfuricum*. Applied and Environmental Microbiology. 1984;48:171-177
- [58] Wang DIC, Avgerinos GC, Biocic I, Wang SD, Fang HY. Ethanol from cellulosic biomass. Philosophical transactions of the Royal Society of London Series B-Biological Sciences. 1983;300:323-333
- [59] Hild HM, Stuckey DC, Leak DJ. Effect of nutrient limitation on product formation during continuous fermentation of xylose with *Thermoanaerobacter ethanolicus* JW200 Fe(7). Applied Microbiology and Biotechnology. 2003;60:679-686
- [60] Lynd LR, Grethlein HE, Wolkin RH. Fermentation of cellulosic substrates in batch and continuous culture by *Clostridium thermocellum*. Applied and Environemental Microbiology. 1989;55:3131-3139
- [61] Rani KS, Swamy MV, Seenayya G. Increased ethanol production by metabolic modulation of cellulose fermentation in *Clostridium thermocellum*. Biotechnology Letters. 1997; 19:819-823

- [62] Singh N, Mathur AS, Tuli DK, Gupta RP, Barrow CJ, Puri M. Cellulosic ethanol production via consolidated bioprocessing by a novel thermophilic anaerobic bacterium isolated from a Himalayan hot spring. Biotechnology for Biofuels. 2017;**10**:73
- [63] Ahring BK, Licht D, Schmidt AS, Sommer P, Thomsen AB. Production of ethanol from wet oxidised wheat straw by *Thermoanaerobacter mathranii*. Bioresource Technology. 1999;
 68:3-9
- [64] Avci A, Donmez S. Effect of zinc on ethanol production by two *Thermoanaerobacter* strains. Process Biochemistry. 2006;**41**:984-989
- [65] Georgieva TI, Mikkelsen MJ, Ahring BK. Ethanol production from wet-exploded wheat straw hydrolysate by thermophilic anaerobic bacterium *Thermoanaerobacter* BG1L1 in a continuous immobilized reactor. Applied Biochemistry and Biotechnology. 2008;145: 99-110
- [66] Wiegel J, Carreira LH, Mothershed CP, Puls J. Production of ethanol from bio-polymers by anaerobic, thermophilic, and extreme thermophilic bacteria. II. *Thermoanaerobacter ethanolicus* JW200 and its mutants in batch cultures and resting cell experiments. Biotechnology and Bioengineering. 1983;13:193-205
- [67] Ahring BK, Jensen K, Nielsen P, Bjerre AB, Schmidt AS. Pretreatment of wheat straw and conversion of xylose and xylan to ethanol by thermophilic anaerobic bacteria. Bioresource Technology. 1996;58:107-113
- [68] Rani KS, Swamy MV, Seenayya G. Production of ethanol from various pure and natural cellulosic biomass by *Clostridium thermocellum* strains SS21 and SS22. Process Biochemistry. 1988;**33**:435-440
- [69] Lin CW, Wu CH, Tran DT, Shih MC, Li WH, Wu CF. Mixed culture fermentation from lignocellulosic materials using thermophilic lignocellulose-degrading anaerobes. Process Biochemistry. 2010;46:489-493
- [70] Lynd LR, Weimer PJ, van Zyl WH, Pretorius LS. Microbial cellulose utilization fundamentals and biotechnology. Microbiology Molecular Biology Reviews. 2002;66:506-577
- [71] Shao X, Raman B, Zhu M, Mielenz JR, Brown SD, Guss AM, Lynd LR. Mutant selection and phenotypic and genetic characterization of ethanol-tolerant strains of *Clostridium thermocellum*. Applied Microbiology and Biotechnology. 2011;92:641-652
- [72] Sittijunda S, Tomas AF, Reungsang A, O-Thong S, Angelidaki I. Ethanol production from glucose and xylose by immobiliezed *Thermoanaerobacter pentosaceus* at 70°C in an up-flow anaerobic sludge blanket (UASB) reactor. Bioresource Technology. 2013;143:598-607
- [73] Shaw AJ, Podkaminer KK, Desai SG, Bardsley JS, Rogers SR, Thorne PG, Hogsett DA, Lynd LR. Metabolic engineering of a thermophilic bacterium to produce ethanol at high yield. Proceedings of the National Academy of Sciences of the United States of America. 2008;105:13769-13774

- [74] Shaw AJ, Hogsett DA, Lynd LR. Identification of the [FeFe]-hydrogenase responsible for hydrogen generation in *Thermoanaerobacterium saccharolyticum* and demonstration of increased ethanol yield via hydrogenase knockout. Journal of Bacteriology. 2009;191: 6457-6464
- [75] Desai SG, Guerinot ML, Lynd LR. Cloning of L-lactate dehydrogenase and elimination of lactic acid production via gene knockout in *Thermoanaerobacterium saccharolyticum* JW/ SL-YS485. Applied Microbiology and Biotechnology. 2004;65:600-605
- [76] Tripathi SA, Olson DG, Argyros DA, Miller BB, Barrett TF, Murphy DM, Mccool JD, Warner AK, Rajgarhia VB, Lynd LR, Hogsett DA, Caiazza NC. Development of pyrFbased genetic system for targeted gene deletion in *Clostridium thermocellum* and creation of a pta mutant. Applied and Environmental Microbiology. 2010;76:6591-6599
- [77] Biswas R, Prabhu S, Lynd LR, Guss AM. Increase in ethanol yield via elimination of lactate production in an ethanol-tolerant mutant of *Clostridium thermocellum*. PLoS One. 2014:9. DOI: 10.1371/journal.pone.0086389
- [78] Cripps RE, Eley K, Leak DJ, Rudd B, Taylor M, Todd M, Biakes S, Martin S, Atkinson T. Metabolic engineering of *Geobacillus thermoglucosidasius* for high yields ethanol production. Metabolic Engineering. 2009;11:398-408
- [79] Yao S, Mikkelsen MJ. Metabolic engineering to improve ethanol production in *Thermoanaerobacter mathranii*. Applied Microbiology and Biotechnology. 2010;**88**:199-208
- [80] Andersen RL, Jensen KM, Mikkelsen MJ. Continuous ethanol fermentation of pretreated lignocellulosic biomasses, waste biomasses, molasses and syrup using the anaerobic, thermophilic bacterium *Thermoanaerobacter italicus* Pentocrobe 411. PLoS One. 2015;**10**:8
- [81] Chung D, Cha M, Guss AM, Westpheling J. Direct conversion of plant biomass to ethanol by engineered *Caldicellulosiruptor bescii*. Proceedings of the National Academy of Sciences of the United States of America. 2014;111:8931-8936
- [82] Shaw AJ, Hogsett DA, Lynd LR. Natural competence in *Thermoanaerobacter* and *Thermoanaerobacterium* species. Applied and Environmental Microbiology. 2010;**76**:4713-4719
- [83] Argyros DA, Tripathi SA, Barrett TF, Rogers SR, Feinberg LF, Olson DG, Foden JM, Miller BB, Lynd LR, Hogsett DA, Caiazza NC. High ethanol titers from cellulose by using metabolically engineered thermophilic, anaerobic microbes. Applied and Environmental Microbiology. 2011;77:8288-8294
- [84] Xu L, Tschirner U. Immobilized anaerobic fermentation for bio-fuel production by *Clostridium* co-culture. Bioprocess and Biosystems Engineering. 2014;**37**:1551-1559
- [85] Tyurin MV, Lynd LR, Wiegel J. Gene transfer systems for obligately anaerobic thermophilic bacteria. In: Rainey FA, Oren A, editors. Methods in Microbiology. Vol. 35. London, England: Academic Press Ltd. Elsvier Science Ltd; 2006. pp. 309-330

- [86] Mai V, Lorenz WW, Wiegel J. Transformation of *Thermoanaerobacterium* sp. strain JW/ SL-YS485 with plasmid pIKM1 conferring kanamycin resistance. FEMS Microbiology Letters. 1997;148:163-167
- [87] Yao S, Mikkelsen MJ. Identification and overexpression of a bifunctional aldehyde/alcohol dehydrogenase responsible for ethanol production in *Thermoanaerobacter mathranii*. Journal of Molecular Microbiology and Biotechnology. 2010;**19**:123-133
- [88] Van Zyl LJ, Taylor MP, Eley K, Tuffin M, Cowan DA. Engineering pyruvate decarboxylase-mediated ethanol production in the thermophilic host *Geobacillus thermoglucosidasius*. Applied Microbiology and Biotechnology. 2014;98:1247-1259
- [89] Cha M, Chung D, Elkins JG, Guss AM, Westpheling J. Metabolic engineering of *Caldi-cellulosiruptor bescii* yields increased hydrogen production from lignocellulosic biomass. Biotechnology for Biofuels. 2013;6:85
- [90] Chung D, Cha M, Farkas J, Westpeling J. Construction of a stable replicating shuttle vector for *Caldicellulosiruptor* species: Use of extending genetic methodologies to other members of this genus. PLoS One. 2013a;8:1-10
- [91] Chung D, Farkas J, Westpheling J. Overcoming restriction as a barrier to DNA transformation in *Caldicellulosiruptor* species results in efficient marker replacement. Biotechnology for Biofuels. 2013b;**6**:82
- [92] Chung D, Cha M, Snyder EN, Elkins JG, Guss AM, Westpheling J. Cellulosic ethanol production via consolidated bioprocessing at 75°C by engineered *Caldicellulosiruptor bescii*. Biotechnology for Biofuels. 2015a;8
- [93] Chung D, Verbeke TJ, Cross KL, Wespheling J, Elkins JG. Expression of heat-stable NADPH-dependent alcohol dehydrogenase in *Caldicellulosiruptor bescii* result in furan aldehyde detoxification. Biotechnology for Biofuels. 2015b;102

