

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

185,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.  
For more information visit [www.intechopen.com](http://www.intechopen.com)



---

## Apicomplexa and Histone Variants: What's New?

---

Laura Vanagas, Susana Marisol Contreras and  
Sergio Oscar Angel

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/intechopen.81409>

---

### Abstract

*Plasmodium* spp. and *Toxoplasma gondii* present a conserved nucleosome composition based on canonical H3 and variants, H4, canonical H2A and variants, and H2B. One-off, the phylum has also a variant H2B, named H2B.Z, which was shown to form a double variant nucleosome H2A.Z/H2B.Z. These histones also present conserved and unique post-translational modifications (PTMs). Histone variants have shown particular genomic localization and PTMs along euchromatin and heterochromatin, including telomere-associated sequences (TAS), suggesting fine-grained chromatin structure modulation. Several other nonhistone proteins present remarkable participation in controlling chromatin state, especially at TAS. Based on that, we discuss the role of epigenetics (PTMs and histone variants) in *Plasmodium* and *Toxoplasma* gene expression, replication, and DNA repair. We also discuss TAS structures and chromatin composition and its impact on antigenic variant expression in *Plasmodium*.

**Keywords:** *Plasmodium*, *Toxoplasma*, epigenetics, histone variants, H2B.Z, chromatin, antigenic variation, telomere-associated region

---

### 1. Introduction

Apicomplexa is a large phylum of unicellular obligate intracellular protozoan parasites responsible for a range of human and animal diseases with considerable medical and economic impact worldwide [1]. The phylum comprises several well-known genera such as *Cryptosporidium*, *Eimeria*, *Babesia*, and *Theileria*, but the most studied genera are *Plasmodium* and *Toxoplasma*.


*Plasmodium* genus is comprised by several species of which five infect humans: *P. falciparum*, *P. ovale*, *P. malariae*, *P. vivax*, and *P. knowlesi*. The infection due to *Plasmodium* genus is known

as Malaria, a mosquito-borne infectious disease endemic in the tropical and subtropical zones of Asia, Africa, South, and Central America. Malaria also constitutes a serious problem for travelers as well as for people working in endemic regions. In 2016, an infection rate of 216 million cases was reported, causing some 445,000 deaths globally. Data show a stalling in declining burden of *Plasmodium* observed over the last decade (<http://apps.who.int/iris/bitstream/10665/259492/1/9789241565523-eng.pdf?ua=1>). In addition, the mass drug administration program has shown a limit due to the spreading of multidrug-resistant malaria [2]. The severe disease is mainly caused by *P. falciparum*, whereas *P. vivax* causes the majority of malaria morbidity outside Africa. The clinical manifestations of malaria include fever, shivering, arthralgia (joint pain), vomiting, jaundice, hemoglobinuria, convulsions, metabolic alterations, renal failure, liver and lung dysfunctions, anemia, and cerebral malaria (coma).

*Toxoplasma gondii* is the only one species of the *Toxoplasma* genus, and it is able to infect birds and mammals, including human, and cause toxoplasmosis. The infection occurs worldwide and the chronic stage reaches more than 500 million people [3]. During the first few weeks of infection, toxoplasmosis is either asymptomatic or causes a mild flu-like illness. However, those with a weakened immune system, such as AIDS patients, infected fetus during gestation or newborns with a congenital infection, may become seriously ill, and occasionally die. The parasite can cause encephalitis (inflammation of the brain) and neurologic diseases, and can affect the heart, liver, inner ears, and eyes (chorioretinitis). Recent research has also linked toxoplasmosis with neuropsychiatric symptoms such as attention-deficit hyperactivity disorder, obsessive compulsive disorder, bipolar disease, and schizophrenia [4–8]. The present chemotherapy for toxoplasmosis is efficient but, sometimes, it is not well tolerated by individuals with AIDS, and it is effective against the acute or active stage, but not against the chronic/latent stage.

## 2. Genome and nucleus

Both *Plasmodium* protozoan parasites and *T. gondii* present a highly complex life cycle, involving several stages along the cycle (**Figure 1**). The genome sizes are 23.3 Mb for *Plasmodium* and 80 Mb for *Toxoplasma*, being haploid (1 N) almost all their life cycle but diploid during sexual replicative stages (2 N) (**Figure 1**). *Plasmodium* genus and *T. gondii* were the first apicomplexan parasites to be included in genome projects [9, 10]. Since then, several other apicomplexan parasites genome projects were taken forward and the data uploaded at EuPathDB (<https://eupathdb.org/eupathdb/>). From these databases, it could be observed that *T. gondii* and *P. falciparum* present about 5300 ([https://protists.ensembl.org/Plasmodium\\_falciparum/Info/Index](https://protists.ensembl.org/Plasmodium_falciparum/Info/Index)) and 8172 ([http://protists.ensembl.org/Toxoplasma\\_gondii/Info/Annotation/#assembly](http://protists.ensembl.org/Toxoplasma_gondii/Info/Annotation/#assembly)) gene transcripts, respectively, organized as single copy genes along the chromosomes, and in general, they are not clustered by function, pathway, or stage of expression. All of these data suggest that these genes are finely regulated throughout the cell cycle and the life cycle by transcriptional regulators and chromatin. Regarding transcription factors, they are poorly represented in apicomplexan, with only one large family of transcriptional regulators, with 24 AP2 family factors in *P. falciparum* and 68 in *T. gondii* [11, 12]. In this context, epigenetic



**Figure 1.** (A) Life cycles of *T. gondii* and *P. falciparum*. Haploid and diploid DNA content is referred to as 1 and 2 N. Black arrows and letters represent common parasite stages, blue arrows and letters represent *T. gondii* specific stages, and red arrows and letters represent *P. falciparum*-specific stages. (B) Epichromatin labeling in *T. gondii*. Epichromatin is a conformational epitope formed by DNA and histones H2A and H2B localized only at the exterior chromatin surface. During tachyzoite replication, the 2 N nucleus is divided, entering one genome to each budding cell, forming a typical U shape. The nuclear envelope does not disappear, and chromosomes do not present high level of condensation. Epichromatin labels predominantly one side of the nucleus, suggesting a nonhomogeneous organization of chromosomes and nuclear envelope interaction. IMC1: inner membrane complex 1.

control was proposed as a key element to facilitate parasite gene expression, DNA replication, and DNA repair [13–15].

An interesting aspect of apicomplexan parasites is that they never lose the nuclear envelope during cell division, and their chromosomes do not present the higher order level of condensation observed in metaphase chromosomes of higher eukaryotes [16]. So, the nucleus presents

the same aspect along the cell cycle. However, it seems to be not homogenous: *Toxoplasma gondii* nuclear envelope and chromosomes seem to have a dynamic relocalization and/or rotation inside the nucleus during parasite budding as observed by epichromatin localization (**Figure 1B**). Epichromatin is a conformational epitope formed by DNA and histones H2A and H2B localized only at the exterior chromatin surface [17, 18]. More recently, it was observed that epichromatin forms superbead domains associated to DNA-A at the nuclear envelope [19]. A 3D analysis also shows that *P. falciparum* nucleus presents a polarization of the nuclear pore complex: in the early multinucleated schizont, it clusters in the nucleus region facing the mother plasma membrane, whereas in the late stages, when prepared for budding, it clusters toward the cytoplasm of the incipient merozoite [20].

In addition to putative polarization of the genome inside the nucleus of Apicomplexan parasites, in *T. gondii*, it was observed that the centromeres (CenH3, see below) are localized at a single spot at the apical region of the nucleus, indicating that all of them are attached to the centrocone, a structure associated to the nuclear envelope, which is traversed by microtubules coordinating the cell division [21]. Similarly, Chromo1, a *T. gondii* protein that binds to the telomere, presents a focalized localization in the nucleus, also suggesting a certain degree of chromosome organization within the parasite nucleus [22]. In *P. falciparum* prior to replication, in late ring stages and young trophozoites, CenH3 localizes to a single nuclear focus suggesting that centromeres are clustered in a single spot that most likely continues to be attached to the mitotic spindle until the end of schizogony and the intraerythrocytic developmental cycle, similar to that observed in *T. gondii* [23].

### 3. H3 histones: a multivariant family

H3 histone family presents canonical forms: H3, H3.1, H3.2 and variants: H3.3 and cenH3 [24]. H3.3 differs from canonical H3s in various aspects. Canonical H3s are expressed and associate to chromatin during the S-phase of cell cycle. Canonical H3s and H3.3 are highly identical differing in only four to five amino acids. The CAF-1 complex is involved in the incorporation of canonical H3s whereas CHD1/ATRX remodelers as well as HIRA chaperone complex are involved in the incorporation of H3.3 [25–30]. In addition, H3.3 is enriched in transcribed genes, enhancers, regulatory elements, and also heterochromatic repeats, including telomeres and pericentromeric regions [31–34]. In general, H3.3 is linked to gene activation or open chromatin. Moreover, it has been found to be methylated at K4, K36, and K79 and acetylated at K9 and K14, all being marks of active chromatin [32, 35]. H3.3 and H2A.Z were detected at active promoters generating nucleosomes that promote gene transcription [36–38]. Recently, it was found that H3.3 plays an essential biological role during mammal development since mice that lack H3.3 presented developmental retardation and early embryonic lethality [39]. Rather than gene expression troubles, H3.3 depletion causes genome instability due to dysfunction of heterochromatin structures at telomeres, centromeres, and pericentromeric regions of chromosomes, leading to mitotic defects.

There is little information regarding H3 histone family and the variants H3.3 in Apicomplexan parasites. The first approach is from W.J Sullivan [40] who was able to clone the entire ORFs




encoding H3 and H3.3 in *Toxoplasma gondii* and also in *Plasmodium falciparum*. In this work, it was confirmed that, like in most other organisms, there is not much difference between the two variants: only four amino acids in *T. gondii* and eight between the *P. falciparum* variants. In most other species, the critical residues that differ between H3 and H3.3, resulting in different roles of these histones, are a motif, which contains SAVM in H3 canonical histone, but changes to AAIG in H3.3 [24]. However, while PfH3 has the typical SAVM motif, it changes to QAVL in PfH3.3, whereas in TgH3, the motif is SAVL and changes to QAIL in TgH3.3 [40]. Besides, there is another difference in Apicomplexa, which seems to be exclusive: KF changes for RY at position 54–55 in H3.3 [40].

In *Plasmodium*, H3.3 had a similar expression pattern to another important histone variant, H2A.Z, namely localization to active chromatin [41] (see **Figure 2**). As observed in other eukaryotic cells, it has been recently demonstrated by ChIP-seq experiments that euchromatic regions in the genome are demarcated by the presence of the H3.3 variant histone [42]. However, in *P. falciparum*, there is a particular AT versus GC content along the genome with euchromatic intergenic regions richer in AT-content compared to coding sequences with less AT content [23]. Fraschka et al. [42] have seen a particular correlation between enrichment in PfH3.3 histone variant and GC content, with this variant mainly located not only in euchromatic GC-rich sequences, but also in subtelomeric GC-rich repetitive regions. Interestingly, this correlation with the nucleotide composition is also observed with the double-variant nucleosome H2A.Z-H2B.Z (see below), but in this case, it is just the contrary: the regions with more AT content show abundance of this nucleosome [42]. However, GC-poor intergenic regions show the lowest H3.3 coverage, but the authors still argue that the incorporation of this variant to coding regions is more dependent on GC content than transcriptional activity.

It is well documented that *P. falciparum* depends on the *var* multigene family, encoding for a highly variable cytoadherence protein called *P. falciparum* erythrocyte membrane protein 1 (PfEMP1) to avoid host immunity [43–46]. This is due to the expression of only one of the ~60 *var* gene family members in any given parasite.

Regarding this important gene family, H3.3 stably occupies the promoter region and coding sequence of the active *var* gene but is evidently less incorporated into the promoter and coding sequence of silenced *var* genes [42] (see **Figure 3**). Additionally, it has been demonstrated that the PTMs affecting histone H3 are extremely important in the regulation of *var* expression. Data from fluorescence *in situ* hybridization (FISH) suggest that the *P. falciparum* SETvs (*P. falciparum* variant-silencing SET gene), which encodes an ortholog of *Drosophila melanogaster* ASH1 and controls histone H3 lysine 36 trimethylation (H3K36me3) on *var* genes, is specifically involved in *var* gene silencing, and its *knock-out* results in the transcription of virtually all *var* genes in the single parasite nuclei [47]. Besides, ChIP-qPCR analysis showed that the TSS occupancy of H3K36me3 is considerably higher in the silent *var* genes compared to the active one (see **Figure 3**) [47].

A detailed mass spectrometry study has been accomplished for *P. falciparum* histone PTMs by Trelle et al. [48], and it has been established that lysines 4, 9, 14, 18, 23, and 27 of both H3 and H3.3 are capable of being modified by acetylations and/or methylations. Also, arginine in position 17 may be mono or bimethylated. Some of these modifications had already been




**Figure 2.** (A) Schematic model for double-variant nucleosome in *T. gondii* and *P. falciparum*. Hypothetical active (or poised) and silenced genes are represented with the nucleosome composition in each promoter and also downstream the genes. PTM marks are represented as well and listed below with the differences found in the literature between both the parasites. Specific HATs and HDACs are speculated to acetylate/deacetylate the unique H2B.Z histone variant. (B) Sequence alignment of histone variant H2B.Z in Apicomplexa. Pf: *Plasmodium falciparum*, Tg: *Toxoplasma gondii*, Et: *Eimeria tenella*, Vb: *Vitrella brassicaformis*, Bv: *Babesia bovis*, Cf: *Cytauxzoon felis*, Gn: *Gregarina niphandrodes* and Cp: *Cryptosporidium parvum*. All sequences can be found in EuPathDB (<https://eupathdb.org/eupathdb/>).

identified for H3 and H3.3 also by Miao et al. [41]. More recently, a lysine residue in the core of H3, K56, was also indicated as a site of acetylation [49, 50]. In the same way, *T. gondii* histone H3 has many lysines and also arginines capable of being modified: lysines in the positions 4, 9, 14, 23, 27 and also 36, 37, 56, 115 and 122 can be acetylated, methylated and besides some of them receive formylation, ubiquitination, or succinylation [51]. Besides, arginines 2, 17, 26, 40, and 83 can be methylated [51].

But not only acetylations and methylations are marking histones; with the development of improved acid and high-salt purification methods for *P. falciparum* histone phosphoprotein analysis, multiple phosphorylation sites have been found mostly at the N-terminal region of

## Telomere Telomeric associated sequences



**Figure 3.** Hypothetical telomeric and subtelomeric structure in *T. gondii* (upper panel) and *P. falciparum* (lower panel). Here, we show the TgTAS and PFTAS composition and the possible protein interactors listed in **Table 1** and described in Section 6. In *P. falciparum*, we only illustrate three of the six TAREs associated. The punctuated lines define proteins described in other organisms, but some of them present an ortholog in any of those parasites. The proteins represented in continuous lines have already been described for those parasites. Proteins illustrated in grey are unknown proteins. We also show the possible histone variant exchange on silenced or active *tsf* or *var* genes. Specific PTMs on silenced or active genes are described in the figure. The questions are open because these proteins and PTMs have not been confirmed yet in these TAS regions.

most histones, including H3 and H3.3 [52]. These marks are frequently seen in combination with neighboring lysine acetylation (and methylation). In this work, they also described a Pf14-3-3 as a phosphohistone mark binding protein.

In parasites, among the most conserved modifications is histone 3 trimethylation of lysine 4 (H3K4me3), a marker of potentially active promoters. Opposed to that is H3K9 methylation, associated with silent genes and densely packed heterochromatin, although protozoan parasite histones are more highly enriched in the activation marks associated with euchromatin with



lower abundance of histone modifications associated with heterochromatin [53]. However, it has been shown that the epigenome in *P. falciparum* is highly dynamic, and dependent on the stage, and, for example, H3K4me3 and H3K9ac are cycle regulated at *P. falciparum* genes [49]. This could also probably be true for *T. gondii*, where the tachyzoite to bradyzoite conversion is regulated at an epigenetic level. In this sense, it has been speculated that the H3R17me2 mark may have significance during the tachyzoite to bradyzoite differentiation process, as it was found only restricted to a subset of promoters, and taking into account the importance of arginine methylation during early development of mouse embryo [54]. In this study, using ChIP-on-chip technique, they found that H3K9ac, H4ac, and H3K4me3 modifications co-localize at focused loci in the *T. gondii* genome and correlate with significant gene expression, while the H3K4me1 and the H3K4me2 modifications were found at equal amounts in active and inactive chromatin [54].

#### 4. Centromeric H3

CenH3, the centromere-specific H3, has been observed in animals, fungi, and plants [24] and also in Apicomplexa, including *T. gondii*, *Plasmodium spp.*, and *N. caninum* [55]. This fact was recently confirmed by Fräschka et al. [42] who found the centromeres depleted of PfH3.3 and PfH3, but occupied by PfCenH3. In *T. gondii*, this histone variant was characterized with the aim to understand the way in which chromosomes are delivered to the daughter cells after mitosis, a process that is still intriguing [21]. In this work, the authors labeled all the histone H3 variants, and used TgCenH3 as a marker of centromeres, to perform ChIP and microarray assays [21]. They found a particular combination of histone PTMs surrounding centromeres; this region had a huge concentration of H3K9 di- and trimethylation, marks usually associated to heterochromatin and found in subtelomeric regions in *P. falciparum* but not in *T. gondii*. In this parasite, these modifications concentrate in two peaks directly flanking the center of the centromere in each chromosome, while H3K4me3 or H3K9ac are not present [21, 52]. In contrast, H3K9me3 and heterochromatin protein 1 (HP1, chromodomain protein that binds to H3K9me3) were not associated with centromeres in *P. falciparum* [23], but rather found in islands of the genome that contain transcriptionally silent members of multigene families [56]. In this parasite, the enrichment of PfCenH3 on centromeres of all the chromosomes has also been demonstrated by genome-wide ChIP-seq analysis [23]. Besides, it has been characterized that a region within the carboxy-terminal histone fold domain, which is also named CENP-A targeting domain (CATD), is essential for mediating centromere targeting, while the N-terminus is not [57].

#### 5. H2A.Z-H2B.Z: the double variant nucleosome

H2A family also has a canonical H2A and several variants: H2A.Z, H2A.X, both exchangeable by H2A.Z-H2B or H2A.X-H2B, allowing the modulation of gene transcription, DNA replication, and/or DNA damage repair [58, 59]. In vertebrates, the H2A family has two more

variants: H2Abd and macro-H2A. When talking about H2A-H2B and the incorporation of variants into such nucleosomes, there are vast differences if we take a glance at Apicomplexan parasites compared to most other eukaryotes. One of the most surprising discoveries in these parasites was the presence of a novel H2B variant (formerly named H2Bv, but recently reclassified as H2B.Z [60]), a histone, which is known to be deficient in variants, similar to H4 [58, 61]. Variants of this histone family, though, are not only found in these parasites, but also in *Trypanosomatids* (even though they are not evolutionary related), and some rare testis-specific variants in human and other mammalian species (reviewed in [55]).

Different studies performed in *Toxoplasma* have shown a nucleosome composition in which H2A.Z, but not H2A.X, dimerizes with H2B.Z, while H2A.X dimerizes with canonical H2B (H2Ba in *T. gondii*), but never with H2B.Z [62, 63]. This fact is also seen in *P. falciparum*, although this parasite lacks H2A.X variant [41] and has driven the hypothesis of a new double variant nucleosome exclusive of parasites with particular characteristics that will be described in this section [64, 65] (**Figure 2A**). As it can be observed in the sequence alignment of H2B.Z in many Apicomplexan species, this histone variant is quite conserved (**Figure 2B**), suggesting that this histone, and likely the double variant nucleosome H2A.Z-H2B.Z, may have had an important role in the expansion of the phylum.

Since H2B.Z is not represented in yeast, insects, or mammals, almost all the current knowledge about the double-variant nucleosome relies on H2A.Z studies. H2A.Z is so widespread that has been catalogued as “universal” because of its origin before the divergence of eukaryotes [66]. The first observation that appears is the hyperacetylation of its N-terminal tail in most species [48–50, 67–69]. It is thought that this possibility gives H2A.Z the faculty of mediating responsiveness to the environmental changes, with so varied and seemingly contradictory effects as gene activation, heterochromatic silencing, transcriptional memory, and others, depending on the binding of activating or repressive complexes [66]. H2A.Z containing nucleosomes mark active and bivalent promoters as well as enhancers, correlating with open chromatin [70, 71]. However, acetylation of H2A.Z is necessary for gene induction and is most often associated with active gene transcription [67, 68, 70, 71], whereas ubiquitylation, which can occur at the C-terminal tail, is linked to transcriptional repression and polycomb silencing [72–75]. Acetylated H2A.Z composes nucleosomes flanking the nucleosome-depleted regions [76]. Regulation of gene expression by acetylation of H2A.Z histone tail may be a result of the participation of other proteins as “readers” in the histone code; for example, the SWR-C chromatin remodeling enzyme and related INO80 family are well characterized to catalyze chromatin incorporation of the histone variant from yeast to human, and the acetylation of histone H3 on lysine 56 (H3-K56Ac) was said to lead to promiscuous dimer exchange in which either H2A.Z or H2A can be exchanged from nucleosomes, although this is in discussion [77–82]. NuA4 acetylation activity, which is homologous to the TIP60/p400 complex, was found to be associated with SWR1-driven incorporation of H2A.Z into chromatin [83]. Besides, bromodomain-containing proteins are known to be implicated in “reading” the acetylation patterns of H2A.Z: acetylated lysines in histones, and other proteins are recognized by this motif, common in remodelers [77, 78, 84, 85]. In fact, for SWR1, bromodomains have been studied to recognize a pattern of acetylation (including H3K14ac), which may influence the deposition of H2A.Z-H2B variant dimers into the appropriate nucleosome [77, 78]. By using *Tetrahymena* as a model, it

could be observed that these protozoa cannot survive with all acetyltable lysines replaced by arginines, indicating that H2A.Z acetylation modulates a charge patch with an essential function in chromatin regulation [69, 75]. Unlike the histone code, these changes need not to be site-specific. If this hypothesis is true, modulation of the charge at any one of a number of clustered sites could inhibit nucleosome condensation, facilitating transcription [86].

*T. gondii* H2A.Z, together with H2B.Z, was enriched in the promoters of active genes in tachyzoites, while repressed genes were enriched with H2A.X-H2Ba nucleosomes [63] (**Figure 2A**). In addition, H2A.Z-H2B.Z was also recruited within the coding region of silent bradyzoite-specific genes and within promoter regions but not coding regions of actively expressed genes [87]. It is tempting to speculate that the enrichment at active promoters or poised regions could be ruled by different PTM stages of these histone variants. In agreement with this, H2A.Z and H2B.Z have shown to be highly acetylated at the amino-terminal tail, in contrast to canonical H2A and H2B histones and the H2A.X variant [51]. Considering that H2A.Z has shown to be essential in regulating the changing gene expression program during differentiation [79–81, 88–90], and recently, it was observed that overexpression of mutated version of H2A.Z, where all five potential acetyltable lysines on H2A.Z-GFP (K4, 7, 11, 13, and 15) were mutated to arginines, blocked myoblast differentiation through disruption of myoD expression [91], it may be that the H2B variant is involved in the *T. gondii* cell differentiation process as part of H2A.Z-H2B.Z nucleosome. Whether through a patch charge modulation and/or histone code remains an open question, considering that *T. gondii* presents several bromodomain-containing proteins that can recognize some of the acetylated lysine [92].

As stated above, the sequence alignment of H2B.Z in many Apicomplexan species reveals a high degree of conservation for this histone variant (**Figure 2B**). Interestingly, every lysine that has been proved to be acetylated in *T. gondii* and *P. falciparum*, H2B.Z was detected in the other Apicomplexan species, which is also true for H2A.Z [50, 53]. Maybe, the double-variant nucleosome is present in the phylum with same PTMs and similar biological role.

### 5.1. Double-variant nucleosome in var genes

In *P. falciparum*, H2A.Z-containing nucleosomes were proposed to demarcate intergenic/regulatory regions of the genome, serving as a scaffold for stage specific as well as transcription-coupled recruitment of histone modifying enzymes [93]. H3K9ac and H3K4me3 were found preferentially placed/retained on or next to H2A.Z-containing nucleosomes [49]. However, it was observed that *P. falciparum* intergenic regions, including promoters, display a global nucleosome depletion, while telomeres harbored the highest nucleosomal occupancy, except for the *var* gene with the highest expression level, which again showed the lowest nucleosomal occupancy [94]. Apparently, the little amount of nucleosomes in these areas is composed largely of variant nucleosomes. Petter et al. [95] also showed an enrichment of PfH2A.Z in the promoter of a set of developmentally regulated genes in the euchromatin compartment, although not correlated with transcription levels nor with acetylation status. *P. falciparum* H2A.Z-H2B.Z promoter occupancy in *var* genes was found to be strongly associated with transcriptional activity, whereas silent or poised *var* genes would be depleted of double-variant nucleosome (see **Figure 3**) [65, 86]. The authors have speculated that it may function

as a similar physical switch to control gene expression in response to temperature change (for example, during fever or as *P. falciparum* is transmitted between its two hosts), as a thermosensory response that was seen in *Arabidopsis thaliana* and yeast [96]. This could be due to reduced DNA wrapping of H2A.Z containing nucleosomes at higher temperatures, resulting in a relaxed chromatin structure, although this variant histone has also been associated with a tighter relationship with DNA, especially in heterotypic H2A.Z/H2A nucleosomes [97]. While heterochromatic intergenic regions showed to contain low levels of histone variant H2B.Z [64], it is interesting that double-variant nucleosomes are depleted from silent *var* gene promoters but not from silent promoters of heterochromatic invasion gene families, which have similar patterns of variegated expression [65]. Besides, this correlation between double-variant nucleosome presence and expression was only seen in *var* genes, while this nucleosome was also found enriched in intergenic regions across the genome, associated with euchromatic histone modifications and not necessarily associated with transcription [64, 65, 86]. Moreover, long promoter containing intergenic regions that maintain higher variant histone levels as compared with 3'UTR containing regions, which are considerably shorter, presents higher AT content, so this correlation could simply be due to the minimal length of the AT-rich content in these short 3'UTR regions [64]. As it was previously observed for H3.3 variant histone (see Section 3), a correlation between nucleosome occupancy and GC/AT content in the genome was observed, although contrary to H3.3 that was correlated with rich GC regions [42], here both H2B.Z and H2A.Z histone variant occupancy displayed a clear positive correlation toward genomic AT content [64]. In **Figure 2A**, a schematic representation of *P. falciparum* and *T. gondii* nucleosome occupancy is proposed.

## 6. Heterochromatin, telomeres, and subtelomeres

The telomere-associated sequences (TAS), also named subtelomeres, are heterochromatic regions adjacent to the telomeric-end looking toward the centromere. The telomeres and the TAS regions are the final structures at the chromosomes and integrate with the centromere the constitutive heterochromatin in the genome. These TAS regions have been described in *Plasmodium* and *Toxoplasma* with a size of 20–40 and near 30 Kpb, respectively (**Figure 3**) [98–100]. In *T. gondii*, the structure contains three tandem repeated elements (TARE), separated by noncoding DNA and flanked at one end by the telomere and at the other, downstream TARE 3, by a *Toxoplasma*-specific gene family, the *tsf* gene, of unknown function [100]. In general, there is only one *tsf* gene per TAS. Interestingly, based on predicted amino acidic sequence, TSF proteins present a high degree of conservation in the N-tail and middle regions while being highly variable at the C-terminal end. Up to now, only few studies were performed on chromatin modulation of *T. gondii* TAS.

The TAS element in *Plasmodium*, instead, has been deeply studied because of the presence of different families of genes associated to virulence and pathogenicity with a clonal pattern of expression [101, 102]. Telomeres are spatially restricted to nuclear periphery, where they form clusters of three to seven heterologous chromosome ends [103–105]. *Plasmodium* TAS is composed of six different TAREs, and the coding part of the genome is localized directly



downstream TARE 6, and is characterized by members of multiple antigen gene families including *var*, *rif*, *stevor*, and *pfmc-2tm* genes [94, 95].

The telomeres and TAS regions are dynamic structures associated to a plethora of specific factors that not only give it structure, but also configures all the regions as constitutive heterochromatin that participates in an epigenetic way to regulate subtelomeric genes expression (**Figure 3**). This epigenetic mechanism is carried out by proteins that introduce, recognize, and implement a repressive state over the gene expression under normal environmental conditions. It has been reported that under nutritional or environmental stress, the repressed subtelomeric genes activate their expression in response to events promoting growth and survival [87, 106–109].

It is important to highlight that the *T. gondii* TAS regions show a nucleosome composition enriched in H2A.X and heterochromatin markers [100]. An *in silico* analysis using the *Plasmodium* and *Toxoplasma* databases reveals the presence of only some orthologs to the yeast and mammal's telomeric-subtelomeric proteins as TRF1-2, HP1, KU70/KU80, and Sir2 proteins (**Table 1**). But interestingly, the principal actor in this scenario would be the histone deacetylase type III -Sir2. This NAD<sup>+</sup> deacetylase-dependent has also been implicated in different signaling pathways. *P. falciparum* has two Sir2 paralogues, Sir2A and B; with overlapping but distinct roles that regulate different subsets of *var* genes [110], binding reversibly with the promoter regions of silent but not active subtelomeric *var* genes [111]. PfSir2A is implicated in telomere length regulation [112]. In *T. gondii*, two deacetylases containing the Sir2-domain were identified: TgSir2A and TgSir2B, but their function has not been characterized yet. Another protein that had been described in *Plasmodium* is PfOrc1 (origin recognition complex 1), which together with Sir2 promotes the epigenetic silencing in *P. falciparum* TAS [113]. PfOrc1 has a role in DNA replication but also cooperates with Sir2 to coordinate the spreading of heterochromatin and regulation of *var* gene expression [114]. In general, Sir2 proteins act by removing acetyl groups in cytosolic targets and at the nuclear level at H3K9, K14 and K56, but it also was described to act on the histone mark H4K16 promoting the deposition of methyl groups on H4K20, H4K20me3 being a chromatin mark associated with heterochromatin [115]. Thus, Sir2 seems to play a very important role in linking signaling processes to gene expression and chromosome architecture.

Additionally, a member of the Alba protein family (PfAlba3) was demonstrated via ChIP assays to bind to telomeric and subtelomeric regions co-localizing with Sir2A in the periphery of the nucleus. PfAlba3 inhibits transcription *in vitro* by binding to DNA. PfSir2A was shown to interact with PfAlba3 deacetylating the lysine residue of N-terminal peptide of PfAlba3 specific for DNA binding [116] (**Figure 3**). In archaea, this interaction had been reported, in which Sir2 regulates silencing through deacetylation of the major archaeal chromatin protein Alba, highlighting an ancestrally conserved mechanism of gene regulation [117].

As stated above, heterochromatin protein 1 (HP1) is a very important protein that has been described to recognize the trimethylation on H3K9, a critical mark for the establishment, maintenance and silencing of centromeric and telomeric heterochromatic regions in various model organisms. In *P. falciparum*, it has been identified as PfHP1 [118] and the H3K9me3 mark was mainly associated with *var* genes at TAS regions, as said before [119]. Moreover, high levels of H3K9me3 correlate with genes localized to the nuclear periphery, implying

chromosome loop formation. In addition, an association between PfSir2 and H3K9me3 was found, since the lack of the sirtuin deacetylases causes changes in H3K9me3 localization at the chromosome and generates disruption of the monoallelic transcription of *var* genes, suggesting the existence of perinuclear repressive centers associated with control of expression of malaria parasite genes involved in phenotypic variation and pathogenesis [119].

Flueck et al. [120] described the presence of an ApiAP2 family member in *P. falciparum*, designated as SIP2 that binds to TARE-2 and TARE-3 regions and the upstream regions of *var* upSB *in vivo*. Immunofluorescence and genome-wide high-resolution ChIP analyses demonstrated that *P. falciparum* SIP2 and HP1 proteins co-localize and associate with the same subtelomeric region, suggesting that both proteins participate in the assembly of telomeric heterochromatin. A recent report from Gupta et al. [121] has demonstrated that the protein CAF-1, a chaperone that loads the H3-H4 to the nucleosome assembly after DDR, co-localizes with PfHP1

Yeast	Mammals	<i>T. gondii</i> (ToxoDB number)	<i>P. falciparum</i> (PlasmoDB PF3D7 number)
Sir2 (P06700)	Sir2 (Q8IXJ6)	Sir2A (227020)	Sir2A (1328800)
		Sir2B (267360)	Sir2B (1451400)
Sir3 (P06701)	Sir3 (Q9NTG7)	ATPase, AAA family protein (283900)*	Orc1/Sir3 like activity (1203000)*
Sir4 (P11978)	Sir4 (Q9Y6E7)	NF	NF
RAP1 (P11938)	RAP1 (Q9NYB0)	NF	NF
RIF1 (P29539)	RIF1 (Q5UIP0)	NF	NF
RIF2 (Q06208)	NF	NF	NF
Ku70 (P32807)	XRCC6 (B1AHC9)	Ku70 (248160)	NF
Ku80 (Q04437)	XRCC5 (P13010)	Ku80 (312510)	NF
Taz1 (P79005)	TRF1 (P54274)	NF	NF
NF	TRF2 (Q15554)	NF	NF
NF	TIN2 (Q9BSI4)	NF	NF
NF	HP1 (Q13185)	Chromo1 (268280)	HP1 (1220900)
Stn1 (P38960)	NF	NF	NF
Ten1 (Q07921)	NF	NF	NF
Cdc13 (P32797)	NF	NF	NF
NF	TPP1 (Q96AP0)	NF	NF
NF	POT1 (Q9NUX5)	NF	NF
Pif1 (P07271)	Pif1 (Q9H611)	NF	NF

NF: Not found. \**P. falciparum* Orc1 complement yeast Sir3 activity [113]. *T. gondii* counterpart was detected by searching ToxoDB with PfOrc1 amino acid sequence by BlastP.

**Table 1.** Telomeric proteins.

at the same subtelomeric localization, in the nuclear periphery, and also demonstrated its binding to TARE1-3 and co-localization with H3K56ac, a signal of completion on chromatin reassembly after DDR [122]. Interestingly, immunoprecipitation with PfCAF1 followed by LC-MS/MS analysis demonstrated that this protein would be interacting not only with PfHP1 but also with PfAlba3 among others [121].

In *T. gondii*, an HP1 protein was identified as TgChromo1, linked to the sequestration of chromosomes at the nuclear periphery and the process of cell division of the parasite [22]. TgChromo1 has shown to localize at *T. gondii* telomeres but not subtelomeres. However, by that time, subtelomeric regions had not yet been described and, in some cases, the sequences in these regions were not correctly assembled. Also, the presence of H4K20me3 and H2A.X at some TARE sequences and a region near *tsf* gene, previously named TgIRE, was observed [62, 63, 100, 123].

## 7. Double-strand break repair: H2A.X and chromatin

Cells are exposed to DNA lesions produced by exogenous (e.g., chemicals, UV-irradiation, and ionization) or endogenous factors (e.g., DNA replication stress, meiotic recombination). One of the most deleterious forms of DNA damage is the double-strand break (DSB) [124]. DSBs activate the signal transduction pathway to induce DNA damage checkpoints that delay cell cycle progression, which allows the cell to activate DNA repair mechanism [125]. The phosphorylation of SQE/D $\Phi$  motif (where  $\Phi$  represents a hydrophobic residue) on histone H2A.X (referred to as  $\gamma$ H2A.X) is one of the earliest responses to DSB [126, 127]. H2A.X seems to be incorporated randomly in the genome of resting cells [128], whereas  $\gamma$ H2A.X is clearly observed forming foci, labeling the DSB and replication fork sites, spreading along the chromosome up to 2 Mb from the damaged site. In addition, chromatin is subjected to several changes at damage sites playing an important role in regulating DNA repair [129]. DSB can be produced by various events, either external as ionizing and UV radiations or internal such as collapse of replication forks and transcription-associated damage, among others [130, 131]. DSB can be repaired by two main mechanisms: nonhomologous end joining (NHEJ) and homologous recombination repair (HRR); the first is an error-prone mechanism available along the cell cycle, and the second is an error-free mechanism active at S/G2 phases of cell cycle because of the requirement of sister chromatid as template [131–134]. Both mechanisms were described in *T. gondii* [14, 135], but *Plasmodium* genus is thought to rely only on HRR [136–138].

Before the election of NHEJ or HRR mechanism, DSB triggers a cascade of events that starts with Mre11-RAD50-Nbs1/Xrs2 (MRN in mammals and MRX in yeast) complex binding to the damaged site, which recruits and activates ATM kinase (**Figure 4A**) [139]. ATM is able to phosphorylate H2A.X at SQE motif as well as other DSB repair enzymes allowing the spreading of  $\gamma$ H2A.X and a correct DNA damage response (DDR) at DSB site (**Figure 4A**). ATM kinase is present in *T. gondii* and *P. falciparum* [14]. In *T. gondii*, the MYST family lysine acetyltransferase TgMYST-B has shown to mediate DDR induced by methyl methanesulfonate (MMS) and to stimulate the ATM expression at gene level [140]. In addition to this


finding, histone acetyltransferases (HATs) have a predominant role in DDR on the basis of chromatin modulation. Chromatin responds to DSB first by increasing the compaction stage by replacing H2A/H2A.X with the H2A.Z variant and by methylating H3K9 by suv39h1 methyltransferase, which is recruited after spreading the DDR response at both sides of DSB sites (**Figure 4B**) [141, 142]. The arrival of H3K9me3 allows its interaction with the HAT Tip60 and the acetylation of H4 on K16 together with the acetylation of ATM kinase, an important PTM for the activation of autophosphorylation and subsequent activation of ATM (**Figure 4**) [143, 144]. The H3K9me3 and H4K16ac marks were identified in *T. gondii* and *P. falciparum* by mass spectrometry analysis [48–51]. However, in the case of *T. gondii*, an acetylated residue was also detected in H3K9 in a more frequent fashion than H3K9me1,2,3, suggesting that chromatin is preferentially in an open state and that this lysine PTM can be regulated [51]. As it was stated before, H3K9me2/3 is also enriched in centromeres in *T. gondii* [21]. In addition, *T. gondii* H4K16ac was one of the most abundant PTMs found in the mass spectrometry analysis [51]. In the case of *P. falciparum*, the treatment with MMS has increased the level of H4K8ac and H4K16ac and reduction of H3K9ac [145]. Both, *T. gondii* and *P. falciparum* present H3K9me1,2,3 and H3K9ac in normal conditions suggesting a conserved mechanism of chromatin modulation [51]. The role of these histone marks on Apicomplexan histones and the connections with DNA repair remain to be elucidated.

As mentioned above,  $\gamma$ H2A.X spreading is a crucial step to initiate a correct DDR at DSB sites. In *T. gondii*, this PTM mark is accompanied by other DDR marks such as H3K9me2,3 and H4K16 in normal conditions of growth, opening the question whether DSBs are being produced during parasite replication [51, 135]. The *T. gondii* tachyzoite replicates at high rates, in a range of 5–9 hours [146]. So, a putative collapse of replication fork could be occurring in this stage. However, *T. gondii* ATM kinase could not be detected in normal conditions by Western blot, but it was detected by tachyzoites overexpressing MYST-B HAT [140].

The chromatin compaction that occurs early during DDR includes the remodeling of chromatin at DSB sites in which the H2A-H2B dimer is replaced by H2A.Z-H2B [142, 147]. This event is transient, allowing the recruitment of repressive kap-1(TRIM28)/HP1/suv39h1 complex that can be important to inhibit transcription. The presence of H2A-H2B dimer in the nucleosomal core particle produces a unique negatively charged region on the surface of the nucleosome, called the “acidic patch,” which is extended in H2A.Z (**Figure 4B**) [148–150]. The acidic patch favors the binding of H4 N-tail, resulting in an increase in the interaction between nucleosomes and chromatin compaction [150]. Interestingly, this seems a necessary step to continue with a relaxed chromatin state, since this compaction and recruitment of kap-1(TRIM28)/HP1/suv39h1 complex lead to methylation of H3K9 and phosphorylation of KAP-1 by ATM kinase, which in turn promote H4K16 acetylation by Tip60 and release kap-1(TRIM28)/HP1/suv39h1 (**Figure 4B**) (see [142]). *T. gondii* and *P. falciparum* have the novel H2A.Z-H2B.Z double-variant nucleosome (see Section 5). However, *T. gondii* and *P. falciparum* H2A-H2B and variants conserve the acidic patch (**Figure 4B**). To note, *T. gondii* and *P. falciparum* do not appear to have KAP-1 protein at ToxoDB and PlasmoDB databases [151].

In higher eukaryotes, another important PTM mark associated to DDR is ubiquitination by E3 ubiquitin ligases RNF168 and RNF8 at DSB site after  $\gamma$ H2A.X and MDC1 protein foci





**Figure 4.** (A) Recognition of DSB and initial steps of DDR pathways. MRN complex and ATM kinase are recruited to a DSB. ATM phosphorylates several DDR proteins and checkpoint kinases. Phosphorylation by ATM allows the spreading of  $\gamma$ H2A.X and DDR foci. RNF8 E3 ubiquitin ligase complexes ubiquitinate H2A and H1 histones to compact and generate histone marks, which will be read by other DDR factors such as MDC1 and E3 ubiquitin ligase RNF168. (B) Model of chromatin modulation after DSB. During PARylation, the p400 ATPase (NuA4-Tip60 complex) replaces H2A-H2B dimers by H2A.Z containing nucleosomes. Since *T. gondii* and *P. falciparum* present a double-variant nucleosome, we speculate that H2A-H2B or H2A.X-H2Ba dimers may be replaced by H2A.Z-H2B.Z dimers. This exchange might increase the interaction of the acidic patch with the N-tail of H4 increasing the chromatin compaction (see the sequence alignment: in red letters are written the acidic residues involved in generating the acidic patch). PARylation also produces an increase in H3K9me2/3 and histone deacetylation repressive marks. After a short time, the H2A.Z containing nucleosome is replaced by H2A-H2B dimer and N-tail of H4 is acetylated leading to a relaxed chromatin and recruitment of different DDR factors, among them those related to the DDR pathway choice, such as 53BP1 or BRCA1.

spreading (**Figure 4A**). MDC1 is also phosphorylated by ATM kinase allowing the recruitment of RNF168 and RNF8 [152]. Ubiquitination on H1 and H2A recruits several BRCT domain containing proteins such as BRCA1 and 53BP1 [129]. In the case of 53BP1, its binding requires the H2AK13/15ub and H4K20me2 and addresses the DDR to NHEJ pathway (**Figure 4B**). By contrast, the presence of H4K16ac impairs the 53BP1 binding to the nucleosome allowing the recruitment of BRCA, which addresses the DDR to HRR (**Figure 4B**) (see [141, 142]). As stated

above, in *T. gondii* and *P. falciparum*, the mark H4K20me<sub>1,2,3</sub> was found [48, 51, 53]. However, *T. gondii* and *P. falciparum* H2As did not contain H2AK15ub and lysine 13 ubiquitylation was not detected either [51]. In addition, *T. gondii* and *P. falciparum* did not show the presence of orthologs of BRCA1 and/or 53BP1, though *T. gondii* presents three different BRCT domain containing proteins [14].

*T. gondii* and *P. falciparum* conserve several histone marks present in chromatin-associated DDR to DBS, as well as histone variants—in the case of *T. gondii*, the DDR, well studied H2A.X, is present, whereas *Plasmodium* has only canonical H2A [55], involved in the recruitment of several factors that spread and choose the DDR pathway in higher eukaryotes. Although, *T. gondii* and *P. falciparum* lack some key DDR regulators such as KAP-1, 53BP1, BRCA1, MDC1, RNF168 and RNF8 [14], both parasites present the HRR mechanism of DNA repair, whereas NHEJ is present only in *T. gondii*. So, the modulation of both DDR pathways is still an intriguing issue.

## 8. Concluding remarks

In protozoan parasites, the modulation of chromatin seems to be a key biological process to regulate gene expression, pathogenicity and DNA repair, the latter probably associated to DNA replication, ergo, the cell cycle. In Apicomplexa, highly evident in *Plasmodium* genus, the TAS or subtelomeric regions play an important role in the control of a group of genes essential in parasite pathogenicity. This fact suggests that subtelomeres have not a trivial impact in the evolution of these organisms, and their structure can influence the features of the cell. How this genomic domain has evolved within the Apicomplexa phylum remains to be elucidated. *T. gondii*, in which to date a scenario of variant antigenicity was not detected, has shown a somewhat conserved structure with the presence of tandem repeated boxes and a gene family of unknown function (*tsf*). Different from *Plasmodium*, which variant antigen-associated gene is represented by hundreds of members, *T. gondii* has only one gene per TAS. However, the predicted protein sequences show conserved N-tail and middle regions, with highly variable C-terminal ends. We believe that the elucidation of the localization, role, and antigenic potential of these gene family proteins will be of high impact in our knowledge of this parasite. Also, it could be interesting to know if the members of this gene family show a regulation of gene expression similar to *Plasmodium* variable antigen gene family.

In addition to the presence of PTM marks similar to other organisms but with currently less-well characterized readers and erasers, Apicomplexa chromatin presents a double-variant nucleosome based on the new histone variant H2B.Z. If considering the partitioned knowledge in these parasites, specially *P. falciparum*, where H3.3 variant has been found in the same regions as this double-variant nucleosome, but in different studies, it would be possible that a triple-variant nucleosome exists in Apicomplexa. Since the presence of H2B.Z arose early in Apicomplexa evolution, it is expected that the double-variant nucleosome could have been important in the expansion of the phylum, maybe modulating chromatin structure during the execution of different biological processes. Interestingly, the genome-wide analyses seem to indicate that *Plasmodium* and *Toxoplasma* double-variant nucleosomes do not have the same behavior. In *T. gondii*, it is enriched in active and poised genes, whereas in *P. falciparum*, it is

localized in active and silent promoters, excepting the *var* genes, in which the presence of the double-variant nucleosome is associated to active promoters. The analysis of this novel nucleosome in the different genera of the phylum can give more information to elucidate the reason of the presence of this H2B variant.

Chromatin is also important to the DDR and has an important role in determining the different pathways of DNA repair after DSB. *T. gondii* seems to have every histone variant and histone mark as well as important proteins associated to every DDR pathway to repair a DSB: NHEJ (e.g., Ku70/Ku80) and HRR (RAD51). Different from *T. gondii*, *Plasmodium* does not present the histone variant H2A.X, whose phosphorylation ( $\gamma$ H2A.X) is linked to the localization of DSB on DNA. Moreover, *T. gondii* has shown a basal level of  $\gamma$ H2A.X, even without damage. Not expected, the proteins associated to the DDR pathway choice (NHEJ or HRR), which read the chromatin, were not detected in *T. gondii* nor in *Plasmodium*. So, it is unknown if these marks are associated to other proteins (*T. gondii* has three BRCT domain containing proteins) with similar roles and/or chromatin modulates DDR in another way.

Taken all together, these differences are not only interesting at the light of evolution but also can be analyzed in the context of the identification of new parasite-specific drug targets. Gene regulation, DNA replication, pathogenicity, and DNA repair are crucial biological processes, and all of them may offer new targets to exploit as future treatments against Apicomplexan pathogens.

## Acknowledgements

SO Angel (Researcher), L. Vanagas (Researcher), and S.M. Contreras (Fellow) are members of National Research Council of Argentina (CONICET). SO Angel and L. Vanagas are professors of Universidad Nacional General San Martín (UNSAM). This chapter was supported by ANPCyT PICT 2015 1288 (S.O.A.) and National Institute of Health (NIH, NIAID 1R01AI083162) grants.

## Author details

Laura Vanagas, Susana Marisol Contreras and Sergio Oscar Angel\*

\*Address all correspondence to: sangel@intech.gov.ar

Instituto Tecnológico de Chascomús (INTECH), CONICET/UNSAM, Chascomús, Provincia de Buenos Aires, Argentina

## References

- [1] Black MW, Boothroyd JC. Lytic cycle of *Toxoplasma gondii*. Microbiology and Molecular Biology Reviews. 2000;64(3):607-623

- [2] Zuber JA, Takala-Harrison S. Multidrug-resistant malaria and the impact of mass drug administration. *Infection and Drug Resistance*. 2018;**11**:209-336
- [3] Tenter AM, Heckeroth AR, Weiss LM. *Toxoplasma gondii*: From animals to humans. *International Journal for Parasitology*. 2000;**30**(12-13):1217-1258
- [4] Fuller Torrey E, Bartko JJ, Yolken RH. *Toxoplasma gondii* and other risk factors for schizophrenia: An update. 2012;**38**(3):642-647
- [5] Fuglewicz AJ, Piotrowski P, Stodolak A. Relationship between toxoplasmosis and schizophrenia: A review. *Advances in Clinical and Experimental Medicine*. 2017;**26**(6):1031-1036
- [6] Flegr J, Horáček J. Toxoplasma-infected subjects report an obsessive-compulsive disorder diagnosis more often and score higher in obsessive-compulsive inventory. *European Psychiatry*. 2017;**40**:82-87
- [7] Sutherland AL et al. Beyond the association. *Toxoplasma gondii* in schizophrenia, bipolar disorder, and addiction: Systematic review and meta-analysis. *Acta Psychiatrica Scandinavica*. 2015;**132**(3):161-179
- [8] Yolken R, Fuller Torrey E, Dickerson F. Evidence of increased exposure to *Toxoplasma gondii* in individuals with recent onset psychosis but not with established schizophrenia. *PLoS Neglected Tropical Diseases*. 2017;**6**(11):e0006040
- [9] Kissinger JC, Gajria B, Li L, Paulsen IT, Roos DS. ToxoDB: Accessing the *Toxoplasma gondii* genome. *Nucleic Acids Research*. 2003;**31**(1):234-236
- [10] Gardner MJ et al., Genome sequence of the human malaria parasite *Plasmodium falciparum*. *Nature*. 2002;**419**(6906):498-511
- [11] De Silva EK et al. Specific DNA-binding by apicomplexan AP2 transcription factors. *Proceedings of the National Academy of Sciences of the United States of America*. 2008;**105**(16):8393-8398
- [12] Behnke MS et al. Coordinated progression through two subtranscriptomes underlies the tachyzoite cycle of *Toxoplasma gondii*. *PLoS One*. 2010;**5**(8):e12354
- [13] Dixon SE, Bhatti MM, Uversky VN, Dunker AK, Sullivan WJ. Regions of intrinsic disorder help identify a novel nuclear localization signal in *Toxoplasma gondii* histone acetyltransferase TgGCN5-B. *Molecular and Biochemical Parasitology*. 2011;**175**(2):192-195
- [14] Fenoy IM, Bogado SS, Contreras SM, Gottifredi V, Angel SO. The knowns unknowns: Exploring the homologous recombination repair pathway in *Toxoplasma gondii*. *Frontiers in Microbiology*. 2016;**7**(May):1-15
- [15] Batugedara G, Lu XM, Bunnik EM, Le Roch KG. The role of chromatin structure in gene regulation of the human malaria parasite. *Trends in Parasitology*. 2017;**33**:364-377
- [16] Striepen B, Jordan CN, Reiff S, van Dooren GG. Building the perfect parasite: Cell division in apicomplexa. *PLoS Pathogens*. 2007;**3**(6):e78
- [17] Olins AL, Ernst A, Zwerger M, Herrmann H, Olins DE. An in vitro model for Pelger-Huët anomaly. *Nucleus*. 2010;**1**(6):506-512



- [18] Vanagas L, Dalmaso MC, Dubremetz JF, Portiansky EL, Olins DE, Angel SO. Epichromatin is conserved in *Toxoplasma gondii* and labels the exterior parasite chromatin throughout the cell cycle. *Parasitology*. 2013;**140**(9):1-1110
- [19] Erenpreisa J, Krigerts J, Salmina K, Selga T, Sorokins H, Freivalds T. Differential staining of peripheral nuclear chromatin with Acridine orange implies an A-form epichromatin conformation of the DNA. *Nucleus*. 2018;**9**(1):171-181
- [20] Weiner A et al. 3D nuclear architecture reveals coupled cell cycle dynamics of chromatin and nuclear pores in the malaria parasite *Plasmodium falciparum*. *Cellular Microbiology*. 2011;**13**(7):967-977
- [21] Brooks CF, Francia ME, Gissot M, Croken MM, Kim K, Striepen B. *Toxoplasma gondii* sequesters centromeres to a specific nuclear region throughout the cell cycle. *Proceedings of the National Academy of Sciences of the United States of America*. 2011;**108**(9):3767-3772
- [22] Gissot M, Walker R, Delhay S, Huot L, Hot D, Tomavo S. *Toxoplasma gondii* chromo-domain protein 1 binds to heterochromatin and colocalises with centromeres and telomeres at the nuclear periphery. *PLoS One*. 2012;**7**(3):e32671
- [23] Hoeijmakers WAM et al. *Plasmodium falciparum* centromeres display a unique epigenetic makeup and cluster prior to and during schizogony. *Cellular Microbiology*. 2012;**14**(9):1391-1401
- [24] Ahmad K, Henikoff S. Histone H3 variants specify modes of chromatin assembly. *Proceedings of the National Academy of Sciences USA*. 2002;**99**(Suppl 4):16477-16484
- [25] Smith S, Stillman B. Purification and characterization of CAF-I, a human cell factor required for chromatin assembly during DNA replication in vitro. *Cell*. 1989;**58**(1):15-25
- [26] Verreault A, Kaufman PD, Kobayashi R, Stillman B. Nucleosome assembly by a complex of CAF-1 and acetylated histones H3/H4. *Cell*. 1996;**87**(1):95-104
- [27] Tagami H, Ray-Gallet D, Ve Almouzni G, Nakatani Y. Histone H3.1 and H3.3 complexes mediate nucleosome assembly pathways dependent or independent of DNA synthesis. *Cell*. 2004;**116**:51-61
- [28] Konev AY et al. The CHD1 motor protein is required for deposition of histone variant H3.3 into chromatin in vivo *Science*. 2007;**317**(5841):1087-1090
- [29] Drané P, Ouarrhni K, Depaux A, Shuaib M, Hamiche A. The death-associated protein DAXX is a novel histone chaperone involved in the replication-independent deposition of H3.3 *Genes and Development*. 2010;**24**(12):1253-1265
- [30] Lewis PW, Elsaesser SJ, Noh K-M, Stadler SC, Allis CD. Daxx is an H3.3-specific histone chaperone and cooperates with ATRX in replication-independent chromatin assembly at telomeres. *Proceedings of the National Academy of Sciences*. 2010;**107**(32):14075-14080
- [31] Wirbelauer C, Bell O, Schübeler D. Variant histone H3.3 is deposited at sites of nucleosomal displacement throughout transcribed genes while active histone modifications show a promoter-proximal bias. *Genes and Development*. 2005;**19**(15):1761-1766

- [32] Loyola A, Almouzni G. Marking histone H3 variants: How, when and why? Trends in Biochemical Sciences. 2007;**32**(9):425-433
- [33] Goldberg AD et al. Distinct factors control histone variant H3.3 localization at specific genomic regions. Cell. 2010;**140**(5):678-691
- [34] Szenker E, Ray-Gallet D, Almouzni G. The double face of the histone variant H3.3. Cell Research. 2011;**21**(3):421-434
- [35] Mckittrick E, Gafken PR, Ahmad K, Henikoff S. Histone H3.3 is enriched in covalent modifications associated with active chromatin. Proceedings of the National Academy of Sciences USA. 2004;**101**(6):1525-1530
- [36] Jin C et al. H3.3/H2A.Z double variant-containing nucleosomes mark 'nucleosome-free regions' of active promoters and other regulatory regions in the human genome. Nature Genetics. 2009;**41**(8):941-945
- [37] Jin C et al. H3.3/H2A.Z double variant-containing nucleosomes mark 'nucleosome-free regions' of active promoters and other regulatory regions. Nature Genetics. 2009;**41**(8):941-945
- [38] Chen PB et al. *Plasmodium falciparum* PfSET7: Enzymatic characterization and cellular localization of a novel protein methyltransferase in sporozoite, liver and erythrocytic stage parasites. Scientific Reports. 2016;**6**:21802
- [39] Jang C-W, Shibata Y, Starmer J, Yee D, Magnuson T. Histone H3.3 maintains genome integrity during mammalian development. Genes & Development. 2015;**29**(13):1377-1392
- [40] Sullivan WJ. Histone H3 and H3.3 variants in the protozoan pathogens *Plasmodium falciparum* and *Toxoplasma gondii*. DNA Sequence. 2003;**14**(3):227-231
- [41] Miao J, Fan Q, Cui L, Li J, Cui L. The malaria parasite *Plasmodium falciparum* histones: Organization, expression, and acetylation. Gene. 2006;**369**:53-65
- [42] Anne-Kristin Fraschka S, Wilhelmus Maria Henderson R, Bártfai R. H3.3 demarcates GC-rich coding and subtelomeric regions and serves as potential memory mark for virulence gene expression in *Plasmodium falciparum*. Scientific Reports. 2016;**6**:31965
- [43] Scherf A, Lopez-Rubio JJ, Riviere L. Antigenic Variation in *Plasmodium falciparum*. Annual Review of Microbiology. 2008;**62**(1):445-470
- [44] Baruch DI et al. Cloning the gene encoding PfEMP1, a malarial variant antigen and adherence receptor on the surface of parasitized human erythrocytes. Cell. 1995;**92**:77-97
- [45] Smith JD et al. Switches in expression of *Plasmodium falciparum* var genes correlate with changes in antigenic and cytoadherent phenotypes of infected erythrocytes. Cell. 1995;**82**(1):101-110
- [46] Su XZ et al. The large diverse gene family var encodes proteins involved in cytoadherence and antigenic variation of *Plasmodium falciparum*-infected erythrocytes. Cell. 1995;**82**(1):89-100

- [47] Jiang L et al. PfSETvs methylation of histone H3K36 represses virulence genes in *Plasmodium falciparum*. *Nature*. 2013;**499**(7457):223-227
- [48] Trelle MB, Salcedo-Amaya AM, Cohen AM, Stunnenberg HG, Jensen ON. Global histone analysis by mass spectrometry reveals a high content of acetylated lysine residues in the malaria parasite *Plasmodium falciparum*. *Journal of Proteome Research*. 2009;**8**(7):3439-3450
- [49] Salcedo-Amaya AM et al. Dynamic histone H3 epigenome marking during the intra-erythrocytic cycle of *Plasmodium falciparum*. *Proceedings of the National Academy of Sciences*. 2009;**106**(24):9655-9660
- [50] Cui L, Miao J. Chromatin-mediated epigenetic regulation in the malaria parasite *Plasmodium falciparum*. *Eukaryotic Cell*. 2010;**9**(8):1138-1149
- [51] Nardelli SC et al. The histone code of *Toxoplasma gondii* comprises conserved and unique posttranslational modifications. *MBio Journal*. 2013;**4**(6):e00922-13
- [52] Dastidar EG et al. Comprehensive histone phosphorylation analysis and identification of Pf14-3-3 protein as a histone H3 phosphorylation reader in malaria parasites. *PLoS One*. 2013;**8**(1):e53179
- [53] Croken MM, Nardelli SC, Kim K. Chromatin modifications, epigenetics, and how protozoan parasites regulate their lives. *Trends in Parasitology*. 2012;**28**(5):202-213
- [54] Gissot M, Kelly KA, Ajioka JW, Greally JM, Kim K. Epigenomic modifications predict active promoters and gene structure in *Toxoplasma gondii*. *PLoS Pathogens*. 2007;**3**(6):e77
- [55] Dalmaso MC, Sullivan WJ, Angel SO. Canonical and variant histones of protozoan parasites. *Frontiers in Bioscience (Landmark Ed.)*. 2011;**16**:2086-2105
- [56] Fischer T et al. Diverse roles of HP1 proteins in heterochromatin assembly and functions in fission yeast. *Proceedings of the National Academy of Sciences of the United States of America*. 2009;**106**(22):8998-9003
- [57] Verma G, Surolia N. *Plasmodium falciparum* CENH3 is able to functionally complement Cse4p and its C-terminus is essential for centromere function. *Molecular and Biochemical Parasitology*. 2013;**192**(1-2):21-29
- [58] Talbert PB, Henikoff S. Histone variants—Ancient wrap artists of the epigenome. *Nature Reviews. Molecular Cell Biology*. 2010;**11**(4):264-275
- [59] Henikoff S, Smith MM. Histone variants and epigenetics. *Cold Spring Harbor Perspectives in Biology*. 2015;**7**(1):a019364
- [60] Talbert PB et al. A unified phylogeny-based nomenclature for histone variants. *Epigenetics & Chromatin*. 2012;**5**:7
- [61] Malik HS, Henikoff S. Phylogenomics of the nucleosome. *Nature Structural Biology*. 2003;**10**(11):882-891

- [62] Dalmaso MC, Onyango DO, Naguleswaran A, Sullivan WJ, Angel SO. Toxoplasma H2A variants reveal novel insights into nucleosome composition and functions for this histone family. *Journal of Molecular Biology*. 2009;**392**:33-47
- [63] Bogado SS et al. Canonical histone H2Ba and H2A.X dimerize in an opposite genomic localization to H2A.Z/H2B.Z dimers in *Toxoplasma gondii*. *Molecular and Biochemical Parasitology*. 2014;**197**(1-2):36-42
- [64] Hoeijmakers WAM et al. H2A.Z/H2B.Z double-variant nucleosomes inhabit the AT-rich promoter regions of the *Plasmodium falciparum* genome. *Molecular Microbiology*. 2013;**87**(5):1061-1073
- [65] Petter M et al. H2A.Z and H2B.Z double-variant nucleosomes define intergenic regions and dynamically occupy *var* gene promoters in the malaria parasite *Plasmodium falciparum*. *Molecular Microbiology*. 2013;**87**(6):1167-1182
- [66] Talbert PB, Henikoff S. Environmental responses mediated by histone variants. *Trends in Cell Biology*. 2014;**24**(11):642-650
- [67] Millar CB, Xu F, Zhang K, Grunstein M. Acetylation of H2AZ Lys 14 is associated with genome-wide gene activity in yeast. *Genes & Development*. 2006;**20**(6):711-722
- [68] Bruce K et al. The replacement histone H2A.Z in a hyperacetylated form is a feature of active genes in the chicken. *Nucleic Acids Research*. 2005;**33**(17):5633-5639
- [69] Ren Q, Gorovsky MA. Histone H2A.Z acetylation modulates an essential charge patch. *Molecular Cell*. 2001;**7**(6):1329-1335
- [70] Hartley PD, Madhani HD. Mechanisms that specify promoter nucleosome location and identity. *Cell*. 2009;**137**(3):445-458
- [71] Saeed S et al. Chromatin accessibility, p300, and histone acetylation define PML-RAR and AML1-ETO binding sites in acute myeloid leukemia. *Blood*. 2012;**120**(15):3058-3068
- [72] Valdés-Mora F et al. Acetylation of H2A.Z is a key epigenetic modification associated with gene deregulation and epigenetic remodeling in cancer. *Genome Research*. 2012;**22**(2):307-321
- [73] Valdés-Mora F et al. Acetylated histone variant H2A.Z is involved in the activation of neo-enhancers in prostate cancer. *Nature Communications*. 2017;**8**(1):1346
- [74] Sarcinella E, Zuzarte PC, Lau PNI, Draker R, Cheung P. Monoubiquitylation of H2A.Z distinguishes its association with euchromatin or facultative heterochromatin. *Molecular and Cellular Biology*. 2007;**27**(18):6457-6468
- [75] Draker R, Sarcinella E, Cheung P. USP10 deubiquitylates the histone variant H2A.Z and both are required for androgen receptor-mediated gene activation. *Nucleic Acids Research*. 2011;**39**(9):3529-3542
- [76] Talbert PB, Henikoff S. Histone variants on the move: Substrates for chromatin dynamics. *Nature Reviews. Molecular Cell Biology*. 2017;**18**(2):115-126



- [77] Clapier CR, Cairns BR. The biology of chromatin remodeling complexes. *Annual Review of Biochemistry*. 2009;**78**(1):273-304
- [78] Van C, Williams JS, Kunkel TA, Peterson CL. Deposition of histone H2A.Z by the SWR-C remodeling enzyme prevents genome instability. *DNA Repair (Amst)*. 2015;**25**:9-14
- [79] Ruhl DD et al. Purification of a human SRCAP complex that remodels chromatin by incorporating the histone variant H2A.Z into nucleosomes. *Biochemistry*. 2006;**45**(17):5671-5677
- [80] Watanabe S, Radman-Livaja M, Rando OJ, Peterson CL. A histone acetylation switch regulates H2A.Z deposition by the SWR-C remodeling enzyme. *Science (80-)*. 2013;**340**(6129):195-199
- [81] Wang F, Ranjan A, Wei D, Wu C. Comment on "A histone acetylation switch regulates H2A.Z deposition by the SWR-C remodeling enzyme". *Science (80-)*. 2016;**353**(6297):358-358
- [82] Watanabe S, Peterson CL. Response to comment on "A histone acetylation switch regulates H2A.Z deposition by the SWR-C remodeling enzyme". *Science (80-)*. 2016;**353**(6297):358-358
- [83] Altaf M et al. NuA4-dependent acetylation of nucleosomal histones H4 and H2A directly stimulates incorporation of H2A.Z by the SWR1 complex. *The Journal of Biological Chemistry*. 2010;**285**(21):15966-15977
- [84] Perell GT, Mishra NK, Sudhamalla B, Ycas PD, Islam K, Pomerantz WCK. Specific acetylation patterns of H2A.Z form transient interactions with the BPTF bromodomain. *Biochemistry*. 2017;**56**(35):4607-4615
- [85] Draker R, Ng MK, Sarcinella E, Ignatchenko V, Kislinger T, Cheung P. A combination of H2A.Z and H4 acetylation recruits Brd2 to chromatin during transcriptional activation. *PLoS Genetics*. 2012;**8**(11):e1003047
- [86] Ren Q, Gorovsky MA. The nonessential H2A N-terminal tail can function as an essential charge patch on the H2A.Z variant N-terminal tail. *Molecular and Cellular Biology*. 2003;**23**(8):2778-2789
- [87] Jeffers V, Tampaki Z, Kim K, Sullivan WJ. A latent ability to persist: Differentiation in *Toxoplasma gondii*. *Cellular and Molecular Life Sciences*. 2018;**75**(13):2355-2373
- [88] Whittle CM et al. The genomic distribution and function of histone variant HTZ-1 during *C. elegans* embryogenesis. *PLoS Genetics*. 2008;**4**(9):e1000187
- [89] Updike DL, Mango SE. Temporal regulation of foregut development by HTZ-1/H2A.Z and PHA-4/FoxA. *PLoS Genetics*. 2006;**2**(9):e161
- [90] Ridgway P, Brown KD, Rangasamy D, Svensson U, Tremethick DJ. Unique residues on the H2A.Z containing nucleosome surface are important for *Xenopus laevis* development. *The Journal of Biological Chemistry*. 2004;**279**(42):43815-43820

- [91] Law C, Cheung P. Expression of non-acetylatable H2A.Z in myoblast cells blocks myoblast differentiation through disruption of MyoD expression. *The Journal of Biological Chemistry*. 2015;**290**(21):13234-13249
- [92] Jeffers V, Yang C, Huang S, Sullivan WJ. Bromodomains in protozoan parasites: Evolution, function, and opportunities for drug development. *Microbiology and Molecular Biology Reviews*. 2017;**81**(1):e00047-e00016
- [93] Bártfai R et al. H2A.Z demarcates intergenic regions of the *Plasmodium falciparum* epigenome that are dynamically marked by H3K9ac and H3K4me3. *PLoS Pathogens*. 2010;**6**(12):e1001223
- [94] Westenberger SJ, Cui L, Dharia N, Winzeler E, Cui L. Genome-wide nucleosome mapping of *Plasmodium falciparum* reveals histone-rich coding and histone-poor intergenic regions and chromatin remodeling of core and subtelomeric genes. *BMC Genomics*. 2009;**10**(1):610
- [95] Petter M et al. Expression of *P. falciparum* var genes involves exchange of the histone variant H2A.Z at the promoter. *PLoS Pathogens*. 2011;**7**(2):e1001292
- [96] Kumar SV, Wigge PA. H2A.Z-containing nucleosomes mediate the thermosensory response in arabidopsis. *Cell*. 2010;**140**(1):136-147
- [97] Horikoshi N, Arimura Y, Taguchi H, Kurumizaka H. Crystal structures of heterotypic nucleosomes containing histones H2A.Z and H2A. *Open Biology*. 2016;**6**(6):160127
- [98] Scherf A, Figueiredo LM, Freitas-junior LH. Plasmodium telomeres: A pathogen's perspective. *Current Opinion in Microbiology*. 2001;**4**(4):409-414
- [99] Figueiredo LM, Freitas-Junior LH, Bottius E, Olivo-Marin J-C, Scherf A. A central role for *Plasmodium falciparum* subtelomeric regions in spatial positioning and telomere length regulation. *The EMBO Journal*. 2002;**21**(4):815-824
- [100] Dalmasso MC, Carmona SJ, Angel SO, Agüero F. Characterization of *Toxoplasma gondii* subtelomeric-like regions: Identification of a long-range compositional bias that is also associated with gene-poor regions. *BMC Genomics*. 2014;**15**(1):21
- [101] Kaviratne M, Khan SM, Jarra W, Preiser PR. Small variant STEVOR antigen is uniquely located within Maurer's clefts in *Plasmodium falciparum*-infected red blood cells. *Eukaryotic Cell*. 2002;**1**(6):926-935
- [102] Petter M, Haeggström M, Khattab A, Fernandez V, Klinkert M-Q, Wahlgren M. Variant proteins of the *Plasmodium falciparum* RIFIN family show distinct subcellular localization and developmental expression patterns. *Molecular and Biochemical Parasitology*. 2007;**156**(1):51-61
- [103] Freitas-Junior LH et al. Frequent ectopic recombination of virulence factor genes in telomeric chromosome clusters of *P. falciparum*. *Nature*. 2000;**407**(6807):1018-1022
- [104] Vernick KD, McCutchan TF. A novel class of supercoil-independent nuclease hypersensitive site is comprised of alternative DNA structures that flank eukaryotic genes. *Journal of Molecular Biology*. 1998;**279**(4):737-751

- [105] Gardner MJ et al. Genome sequence of the human malaria parasite *Plasmodium falciparum*. Nature. 2002;**419**(6906):498-511
- [106] Ai W, Bertram PG, Tsang CK, Chan TF, Zheng XFS. Regulation of subtelomeric silencing during stress response. Molecular Cell. 2002;**10**(6):1295-1305
- [107] Buck MJ, Lieb JD. A chromatin-mediated mechanism for specification of conditional transcription factor targets. Nature Genetics. 2006;**38**(12):6-1451
- [108] Tomar RS, Zheng S, Brunke-Reese D, Wolcott HN, Reese JC. Yeast Rap1 contributes to genomic integrity by activating DNA damage repair genes. The EMBO Journal. 2008;**27**(11):1575-1584
- [109] Harari Y, Romano G-H, Ungar L, Kupiec M. Nature vs nurture: Interplay between the genetic control of telomere length and environmental factors. Cell Cycle. 2013;**12**(22):3465-3470
- [110] Tonkin CJ et al. Sir2 paralogues cooperate to regulate virulence genes and antigenic variation in *Plasmodium falciparum*. PLoS Biology. 2009;**7**(4):0771-0788
- [111] Freitas LH et al. Telomeric heterochromatin propagation and histone acetylation control mutually exclusive expression of antigenic variation genes in malaria parasites. Cell. 2005;**121**(1):25-36
- [112] Merrick CJ et al. Functional analysis of sirtuin genes in multiple *Plasmodium falciparum* strains. PLoS One. 2015;**10**(3):e0118865
- [113] Mancio-Silva L, Rojas-Meza AP, Vargas M, Scherf A, Hernandez-Rivas R. Differential association of Orc1 and Sir2 proteins to telomeric domains in *Plasmodium falciparum*. Journal of Cell Science. 2008;**121**:2046-2053
- [114] Deshmukh AS et al. The role of N-terminus of *Plasmodium falciparum* ORC1 in telomeric localization and var gene silencing. Nucleic Acids Research. 2012;**40**(12):5313-5331
- [115] Serrano L et al. The tumor suppressor SirT2 regulates cell cycle progression and genome stability by modulating the mitotic deposition of H4K20 methylation. Genes & Development. 2013;**27**(6):639-653
- [116] Goyal M et al. Identification and molecular characterization of an Alba-family protein from human malaria parasite *Plasmodium falciparum*. Nucleic Acids Research. 2012;**40**(3):1174-1190
- [117] Bell SD. The interaction of Alba, a conserved archaeal chromatin protein, with Sir2 and its regulation by acetylation. Science (80-). 2002;**296**(5565):148-151
- [118] Pérez-Toledo K et al. *Plasmodium falciparum* heterochromatin protein 1 binds to trimethylated histone 3 lysine 9 and is linked to mutually exclusive expression of var genes. Nucleic Acids Research. 2009;**37**(8):2596-2606
- [119] Lopez-Rubio JJ, Mancio-Silva L, Scherf A. Genome-wide analysis of heterochromatin associates clonally variant gene regulation with perinuclear repressive centers in malaria parasites. Cell Host & Microbe. 2009;**5**(2):179-190

- [120] Flueck C et al. A major role for the *Plasmodium falciparum* ApiAP2 protein PfSIP2 in chromosome end biology. PLoS Pathogens. 2010;**6**(2):e1000784
- [121] Gupta MK, Agarawal M, Banu K, Reddy KS, Gaur D, Dhar SK. Role of chromatin assembly factor 1 in DNA replication of *Plasmodium falciparum*. Biochemical and Biophysical Research Communications. 2018;**495**(1):1285-1291
- [122] Masumoto H, Hawke D, Kobayashi R, Verreault A. A role for cell-cycle-regulated histone H3 lysine 56 acetylation in the DNA damage response. Nature. 2005;**436**(7048):294-298
- [123] Echeverria PC, Rojas PA, Martin V, Guarnera EA, Pszeny V, Angel SO. Characterisation of a novel interspersed *Toxoplasma gondii* DNA repeat with potential uses for PCR diagnosis and PCR-RFLP analysis. FEMS Microbiology Letters. 2000;**184**(1):23-27
- [124] Fillingham J, Keogh M-C, Krogan NJ. GammaH2AX and its role in DNA double-strand break repair. Biochemistry and Cell Biology. 2006;**84**(4):568-577
- [125] Nyberg KA, Michelson RJ, Putnam CW, Weinert TA. Toward maintaining the genome: DNA damage and replication checkpoints. Annual Review of Genetics. 2002;**36**(1):617-656
- [126] Redon C, Pilch D, Rogakou E, Sedelnikova O, Newrock K, Bonner W. Histone H2A variants H2AX and H2AZ. Current Opinion in Genetics and Development. 2002;**12**(2):162-169
- [127] Turinetto V, Giachino C. Histone variants as emerging regulators of embryonic stem cell identity. Epigenetics. 2015;**10**(7):563-573
- [128] Seo J et al. Genome-wide profiles of H2AX and  $\gamma$ -H2AX differentiate endogenous and exogenous DNA damage hotspots in human cells. Nucleic Acids Research. 2012;**40**:5965-5974
- [129] Lukas J, Lukas C, Bartek J. More than just a focus: The chromatin response to DNA damage and its role in genome integrity maintenance. Nature Cell Biology. 2011;**13**(10):1161-1169
- [130] Ciccia A, Elledge SJ. The DNA damage response: Making it safe to play with knives. Molecular Cell. 2010;**40**(2):179-204
- [131] Jackson SP, Bartek J. The DNA-damage response in human biology and disease. Nature. 2009;**461**(7267):1071-1078
- [132] Bétermier M, Bertrand P, Lopez BS. Is non-homologous end-joining really an inherently error-prone process? PLoS Genetics. 2014;**10**(1):e1004086
- [133] Guirouilh-Barbat J, Lambert S, Bertrand P, Lopez BS. Is homologous recombination really an error-free process? Frontiers in Genetics. 2014;**5**:175
- [134] Sancar A, Lindsey-Boltz LA, Unsal-Kacmaz K, Linn S. Molecular mechanisms of mammalian DNA repair and the DNA damage checkpoints. Annual Review of Biochemistry. 2004;**73**(1):39-85 (0066-4154 (Print) LA-eng PT-Journal Article PT-Review RN-0 (Cross-Linking Reagents) RN-9007-49-2 (DNA) SB-IM)



- [135] Smolarz B, Wilczyński J, Nowakowska D. DNA repair mechanisms and *Toxoplasma gondii* infection. Archives of Microbiology. 2014;**196**(1):1-8
- [136] Kelso AA, Waldvogel SM, Luthman AJ, Sehorn MG. Homologous recombination in protozoan parasites and recombinase inhibitors. Frontiers in Microbiology. 2017;**8**:1716
- [137] Fidock DA, Lee AH, Symington LS, Fidock A. DNA repair mechanisms and their biological roles in the malaria parasite *Plasmodium falciparum*. Microbiology and Molecular Biology Reviews. 2014;**3**:78
- [138] Calhoun SF, Reed J, Alexander N, Mason CE, Deitsch KW, Kirkman LA. Chromosome end repair and genome stability in *Plasmodium falciparum*. MBio Journal. 2017;**8**(4): e00547-e00517
- [139] Paull TT. Mechanisms of ATM activation. Annual Review of Biochemistry. 2015;**84**(1): 711-738
- [140] Vonlaufen N, Naguleswaran A, Coppens I, Sullivan WJ. MYST family lysine acetyltransferase facilitates ataxia telangiectasia mutated (ATM) kinase-mediated DNA damage response in *Toxoplasma gondii*. The Journal of Biological Chemistry. 2010;**285**(15): 11154-11161
- [141] Agarwal P, Miller KM. The nucleosome: Orchestrating DNA damage signaling and repair within chromatin. Biochemistry and Cell Biology. 2016;**94**(5):381-395
- [142] Gursoy-Yuzugullu O, House N, Price BD. Patching broken DNA: Nucleosome dynamics and the repair of DNA breaks. Journal of Molecular Biology. 2016;**428**(9):1846-1860
- [143] Sun Y, Jiang X, Chen S, Fernandes N, Price BD. A role for the Tip60 histone acetyltransferase in the acetylation and activation of ATM. Proceedings of the National Academy of Sciences of the United States of America. 2005;**102**(37):13182-13187
- [144] Francia ME, Striepen B. Cell division in apicomplexan parasites. Nature Reviews Microbiology. 2014;**12**(2):125-136
- [145] Gupta DK, Patra AT, Zhu L, Gupta AP, Bozdech Z. DNA damage regulation and its role in drug-related phenotypes in the malaria parasites. Scientific Reports. 2016;**6**(November):1-15
- [146] Radke JR, Striepen B, Guerini MN, Jerome ME, Roos DS, White MW. Defining the cell cycle for the tachyzoite stage of *Toxoplasma gondii*. Molecular and Biochemical Parasitology. 2001;**115**(2):165-175
- [147] Xu Y, Ayrapetov MK, Xu C, Gursoy-Yuzugullu O, Hu Y, Price BD. Histone H2A.Z controls a critical chromatin remodeling step required for DNA double-strand break repair. Molecular Cell. 2012;**48**(5):723-733
- [148] Luger K, Rechsteiner TJ, Flauss AJ, Waye MM, Richmond TJ. Characterization of nucleosome core particles containing histone proteins made in bacteria. Journal of Molecular Biology. 1997;**272**(3):301-311

- [149] Luger K, Suto RK, Clarkson MJ, Tremethick DJ. Crystal structure of a nucleosome core particle containing the variant histone H2A.Z. *Nature Structural Biology*. 2000; **7**(12):1121-1124
- [150] Park Y-J, Dyer PN, Tremethick DJ, Luger K. A new fluorescence resonance energy transfer approach demonstrates that the histone variant H2AZ stabilizes the histone octamer within the nucleosome. *The Journal of Biological Chemistry*. 2004; **279**(23):24274-24282
- [151] Aurrecochea C et al. EuPathDB: The eukaryotic pathogen database. *Nucleic Acids Research*. 2013; **41**(D1):D684-D691
- [152] Jackson SP, Durocher D. Regulation of DNA damage responses by ubiquitin and SUMO. *Molecular Cell*. 2013; **49**(5):795-807

IntechOpen

