

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

186,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Hip Fracture: Anatomy, Causes, and Consequences

Masoud Nasiri Sarvi

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/intechopen.75946>

Abstract

Fall-induced hip fracture is a major worldwide health problem among the elderly population. Nowadays, hip replacement surgery represents a big part of the orthopedic surgeons' workload and has associated remarkable clinical and social cost implications. Hip fractures have several complications including medical and surgical treatment. A significant number of biomechanical models have been introduced to study hip fracture risk. The purpose of proposing the biomechanical models for predicting the hip fracture risk is to introduce prevention and protection activities that may reduce the number of hip fractures. For accurate prediction of hip fracture risk, the fracture procedure and the parameters that affect the risk of hip fracture should be well studied. The objective of this study is to investigate in-depth the hip fracture anatomy, causes, and consequences.

Keywords: hip fracture causes, hip anatomy, fall, hip impact force, hip fracture consequences

1. Introduction

Low-trauma hip fracture has become a common health problem among the elderly all over the world [1–21], mainly due to the population aging and the prevalence of osteoporosis. Of all osteoporotic fractures, hip fracture has the highest morbidity and mortality rate [22]. Approximately 50% of patients have permanent functional disability greater than that before fracture [23, 24]. The incidence of hip fracture appears to be increasing in many countries [10], and the total number of hip fractures is estimated to be more than five million by 2050 [25]. Socioeconomic impacts of hip fracture are twofold. On the one hand, hip fracture increases the morbidity and mortality in the elderly [26–28]; on the other hand, it is a substantial source of healthcare expenditure [29, 30]. Therefore, there is an urgent need to accurately assess hip fracture risk and then develop preventive and protective measures. In this chapter, hip

anatomy is first reviewed, and hip fractures are classified by anatomic location. Then, prevalence of hip fracture is presented, followed by a description of the significance of accurately assessing hip fracture risk.

2. Hip fracture anatomy

Hip fracture is a medical condition in which there is a break in the continuity of the femoral bone. Hip fracture is generally affected by hip anatomy [31], the applied forces to the hip [32], and bone mechanical properties [33]. In this section, hip anatomy is explained to show why the hip is likely to experience fracture in a fall.

The hip joint is one of the most important joints in the human body. It is also one of the most flexible joints allowing a great range of motions. To better understand hip fracture, it helps to know the anatomy of the hip joint. The hip is a joint formed by the ball-shaped head of the femur and the socket of the pelvis. The femurs are the longest and the strongest bones in the human body, extending from the hip to the knee. Important geometric features of femur bones include the head, neck, and greater and lesser trochanters, as shown in **Figure 1(a)**. A femur is composed of two types of bones, cortical and cancellous. The cortical bone forms the outer layer of the femur and withstands most of the forces and moments. Cancellous bone is mostly enclosed by the cortical bone and mainly absorbs the shock energy produced in walking and running [34]. The hip joint is a stable ball-and-socket joint, much more stable than the shoulder joint. The stability in the hip mainly attributes to the deep socket, i.e., the acetabulum. Additional stability is provided by the strong joint capsule and its surrounding muscles and ligaments. The high level of stability of the hip joint is required to support the upper body [34].




Figure 1. (a) Anatomic structure of the hip [35]. (b) Concentration of applied forces on the proximal femur in a lateral fall which increases the risk of fracture.

More than 90% of all hip fractures occur in falls [36] as the femur is subjected to a high-level impact force. As shown in **Figure 1(b)**, in a sideways fall, the greater trochanter and the femoral head are subjected to the impact and the joint force, respectively, from the ground and the acetabulum. The forces produce a moment at the intersection of the neck-shaft axes. Muscles that are attached to the femur also produce forces during the fall. As it is shown in **Figure 1(b)**, the applied forces in a fall are mainly on the proximal femur, and it may explain why the majority of fall-induced hip fractures occur at the proximal femur [37]. A hip fracture refers to any fracture of the proximal femur down to a level of approximately 5 cm below the lower border of the lesser trochanter [38]. The extent of the break depends on the forces that are involved.

3. Hip fracture causes

Hip fracture is usually caused by an applied force that exceeds the strength of the femur bone [39]. Therefore, any situation that either induces a high level of force on the femur bone or decreases the bone strength should be considered as a hip fracture cause.

The main cause of hip fracture is falling (90–92%) [36, 40–42], in particular falling in sideways direction (63–69% in fall-related fractures) [8, 43], as it induces a high level of force on the femur. Parameters that increase the risk of fall and apply a high level of force on the femur, especially in the elderly, are:

- Mental impairment and confusion
- Impaired vision
- Impaired muscle reactions
- Slow reflex response
- Inability to effectively use the arms to reduce the energy of the fall
- Impaired neuromuscular coordination and neurological diseases (e.g., hemiplegia, Parkinson's disease)
- Reduced soft tissue padding over the hip [44, 45]

In the elderly, most fractures occur after a low-trauma fall, which would not cause any severe injury to a healthy individual. Therefore, low bone strength is another main cause of hip fracture. Osteoporosis as a progressive bone disease, which is characterized by decreases in bone mass and density, has been identified as one of the main contributors of hip fracture [46, 47]. Osteoporosis advances when bone resorption exceeds bone formation, and therefore it is more common among the elderly [48]. Approximately three to four out of ten women over the age of 50, and one in eight men, suffer osteoporotic fracture in their lifetime [49].

Apart from osteoporosis, several other causes may reduce the strength of the bone such as bone cancer and medical side effects [38]. Other factors associated with reduction in bone strength include [38]:

- Genetic and family history
- Sedentary lifestyle
- Impaired nutrition
- Smoking
- Excess alcohol
- Medications (including tranquilizers, hypnotics, anticonvulsant drugs, and steroids)
- Osteomalacia from vitamin D deficiency, malabsorption, and liver or renal disease
- Cardiovascular disease and cardiac arrhythmias
- Underlying bone disease (e.g., Paget's disease, bone tumors, and secondary bone tumors)
- Endocrine abnormalities: hyperthyroidism, hyperparathyroidism, or hypercortisolism

In addition to the mentioned causes, high-trauma falls and accidents such as car and motorcycle accidents can lead to hip fracture [50]. But they are not studied in this dissertation. **Figure 2** shows how different factors contribute to the hip fracture [6, 38].




Figure 2. Conceptual model of the fall-induced hip fracture procedure and associated effective factors [15].

4. Hip fracture consequences

Hip fractures are associated with significant morbidity, mortality, loss of independence, and financial burden [3, 9, 25, 42, 51–53]. It has been reported that approximately 20% of hip fracture patients died within 1 year of the fracture [54]. Generally, the first year after hip fracture appears to be the most critical time. A recent meta-analysis revealed that women sustaining a hip fracture had a fivefold increase and men almost an eightfold increase in relative likelihood of death within the first 3 months compared with age- and sex-matched controls [29]. The relative death risk decreases substantially over the second year but still much higher than that of the controls [55]. Many lose their ability to walk mainly due to the pain caused by the hip fracture. In fact, only 40–79% of patients regain their previous ambulatory function a year after the fracture, and less than half return to their pre-fracture status of daily activities [56].

In addition to functional impairments, hip fracture can have a negative impact on self-esteem, body image, and mood [57], which may lead to psychological problems [58]. Individuals who suffer fractures may be immobilized by a fear of falling again and suffering more fractures. They may feel isolated and helpless. The National Osteoporosis Foundation conducted a survey [59] among 1000 women with osteoporotic fracture in the United States to investigate the psychological effects of the fracture on the patients. Eighty-nine percent of said they feared breaking another bone; 80% were afraid that they would be less able to perform their daily activities and lose their independence; 73% worried that they would have to reduce activities with family and friends; and 68% were concerned that another fracture would result in their having to enter a nursing home [59]. If not addressed, fear about the future and a sense of helplessness can produce significant anxiety and depression. These problems may be compounded by an inability to fulfill occupational, domestic, or social duties, thus leading to further social isolation.

The disability, reduced functional status, and poor mental health caused by hip fracture can have a profound impact on the quality of the individual's life. Survivors of hip fracture reported a 52% reduction in the quality of life in the first 12 months and a 21% reduction after 2 years [60].

Also, hip fracture is a major cause of the need for long-term nursing home care and a major contributor to healthcare costs [30, 61, 62]. There are approximately 23,000 cases of hip fracture every year in Canada with associated treatment costs of about \$1 billion [63]. In the United States, 310,000 hip fractures occurred in 2003, and the total Medicare cost was estimated between \$10.3 and \$15.2 billion, including acute medical care and nursing home services [53, 64, 65]. As the population of the elderly is still continuously increasing, the number of hip fractures is expected to rise dramatically, and it will put more burdens on the community healthcare system [2, 66].

5. Classification of hip fractures

In general, there are three types of hip fractures, depending on what region of the proximal femur is involved [67]:

1. Femoral neck fractures occur in the narrow section of the proximal femur that lies between the femoral head and the intertrochanteric cross section. Most femoral neck fractures occur within the capsule surrounding the hip joint and are, therefore, termed intracapsular fracture. The blood supply to the femoral head is carried by a number of arteries that pass through the femoral neck region. Therefore, femoral neck fractures may disrupt the blood supply to the femoral head, causing death of the femoral head bone tissues, called osteonecrosis or avascular necrosis. Femoral neck fractures are further grouped into nondisplaced and displaced fractures by the alignment of the fractured segments in relation to the original anatomic position of the femur [68].
2. Intertrochanteric fractures occur at a lower location than femoral neck fractures, in the area between the greater and lesser trochanters. The trochanters are bony projections where major hip muscles are attached. Intertrochanteric hip fractures occur outside of the joint capsule and are therefore also called extracapsular fracture in the literature. Intertrochanteric fractures are complicated by the pull of the hip muscles on the bony muscle attachments, which can exert competing forces against fractured bone segments and pull them out of alignment. Thus, healing of the fracture in a misaligned position is considered as a complication for intertrochanteric fractures. Intertrochanteric fractures may be further grouped into stable and unstable fractures, depending on the location, number, and size of the fractured bony segments [68].
3. Subtrochanteric fractures occur in the zone about 5 cm below the lesser trochanter of the proximal femur. The blood supply to the bone of the subtrochanteric region is not as good as the blood supply to the bone of the intertrochanteric region, and thus subtrochanteric fracture heals more slowly [68]. Similar to the intertrochanteric fractures, subtrochanteric fractures are likely to cause femur misalignment [68].

In more complicated cases, the fracture of the bone can involve more than one of these zones. **Figure 3** shows different types of proximal femur fracture.




Figure 3. Three main types of hip fractures: femoral neck fracture (subcapital and transcervical fractures), intertrochanteric fracture, and subtrochanteric fracture [69].

6. Demographic feature of hip fractures

A variety of studies have examined hip fracture rates in different regions of the world [10, 51, 52, 70]. Greater than tenfold differences have been found on the basis of studies undertaken at a regional or national level for different calendar years. The studies show that the main demographic risk factors for hip fractures include increased age and female gender [10, 25]. The geographic distribution by fracture risk is shown for men and women combined in **Figure 4**. Heterogeneity in hip fracture risk in countries can be seen in this figure. Based on statistical results [10], for women, the lowest annual incidences are found in Nigeria (2/100,000), South Africa (20), Tunisia (58), and Ecuador (73). The highest rates were observed in Denmark (574/100,000), Norway (563), Sweden (539), and Austria (501). The incidence of hip fracture in men is approximately half of that noted in women. The highest annual incidence in men has been found in Denmark (290/100,000) and the lowest in Ecuador (35/100,000) [10].

As it is shown in **Figure 4**, the high-risk countries are Iceland, the United Kingdom, Ireland, Denmark, Sweden, and Norway in Northwestern Europe; Belgium, Germany, Austria, Switzerland, and Italy in Central Europe; Greece, Hungary, Czech Republic, Slovakia, and Slovenia in southwestern Europe; Lebanon, Oman, Iran, Hong Kong, Singapore, Malta, and Taiwan in Asia; and Argentina in South America. Regions of moderate risk include North America, Oceania, the Russian Federation, and southern countries of Latin America. Low-risk regions include the northern regions of Latin America, Africa, Jordan, Saudi Arabia,




Figure 4. Hip fracture rates (men and women combined) in different countries of the world categorized by risk. Where estimates are available, countries are color-coded red (annual incidence >250/100,000), orange (150–250/100,000), or green (<150/100,000) [10] (reproduced with permission).




Figure 5. Estimated number of hip fractures by sex in the year 1990 and the number expected in 2025 and 2050 by region assuming no increase in age- and sex-specific rates, a 1% annual increase worldwide, or no increase in North America and northern Europe but an increase in age- and sex-specific incidence elsewhere of 2, 3, or 4%. (ROW is rest of world) [25] (reproduced with permission).

India, China, Indonesia, and the Philippines. It is notable that in Europe, the majority of countries are categorized as high or moderate risk. Low risk is identified only in Croatia and Romania [10].

Hip fracture incidence rates are known to increase exponentially with age in both men and women for the most regions of the world [71–74]. The increasing rate of hip fracture in the elderly is mainly associated with their slower reflex response and the inability to effectively use their arms to reduce the energy of the fall and low bone density of the proximal femur [44, 45].

Epidemiological studies show that the number of hip fractures will rise from 1.66 million in 1990 to 4.5–21.3 million by 2050 (**Figure 5**) depending on the underlying assumptions about age- and gender-specific incidence trends [9, 25, 51, 75].

7. Significance of accurately assessing hip fracture risk

The aim of accurately assessing hip fracture risk is to identify patients at high risk of hip fracture and to start intine prevention and protection measures to reduce the number of hip fractures. These measures are accepted by the patients only after they are accurately diagnosed with the high fracture risk. Also, accurate assessment of hip fracture risk is the prerequisite step before starting a therapy. For example, during the process of osteoporosis treatment, it is required to monitor the change of fracture risk and subsequently track the effectiveness of the therapy. By knowing the risk of fracture, people can improve their bone health and change their environment to reduce the likelihood of the fall.

Patients diagnosed with high fracture risk may consider the following prevention measurements:

- Individualized exercise programs:
 - Muscle-strengthening exercises [76]
 - Practicing balance exercises [77]
 - Increasing the lower extremity joint function [32]
- Management of visual impairment:
 - Obtaining maximum vision correction [6, 78]
- Examination of basic neurological function, including mental status, muscle strength, lower extremity peripheral nerves, and reflexes [79]
- Using mobility assisting devices (e.g., walking stick, frames)
- Implementing surveillance and observation strategies

Protection measurements must be provided to patients with high fracture risk, for example:

- Remembering that sideways falling is more likely to result in a hip fracture than falling in other directions [8]:
 - Trying to fall forward or backward not from sides
- Taking steps to reduce the potential energy and subsequently decrease the risk of fracture [80]
- Landing with the aid of hands or reachable objects around to break the fall [81]
- Using hip protectors [82–87]
- Environmental modification (e.g., flooring) [31]
- Medication and nutritional improvement:
 - Consuming a calcium-rich diet that provides about 1000 mg (milligrams) daily for men and women up to age 50 [88]. Women over age 50 and men over age 70 should increase their intake to 1200 mg daily from a combination of foods and supplements.
 - Obtaining 600 IU (international units = 0.025 µg) of vitamin D daily up to age 70 [88]. Men and women over age 70 should increase their uptake to 800 IU daily.
 - 5–15 min' exposure to sunlight 4–6 times per week [89].

8. Bone fracture criterion and hip fracture risk measurement

From biomechanics point of view, assessment of hip fracture under stance loading or lateral impact force has been performed using three criteria: factor of safety (FOS) [90], risk factor

(RF) [70], and fracture risk index (FRI) [91]. In this section, a review is performed on previously adopted bone fracture criteria in both 2D and 3D FE models.

Keyak et al. [90] assessed FOS under two loading conditions: one representing loading during the stance phase of gait and the other simulating the impact from a fall. Their study was based on a 3D FE model generated from CT data of the patient. They calculated FOS to compare the actual element strength with the applied von Mises stress.

Schileo et al. [92] applied maximum principle strain, von Mises stress, and maximum principle stress criteria to calculate risk factor and to predict fracture location of the femur. RF compares the applied stress/strain with the yield one to predict the bone fracture. Lotz et al. [93, 94] also used von Mises stress yield criterion for the cortical bone and crushing-cracking stress criterion for the trabecular bone. The performance of nine stress- and strain-based failure theories in assessment of hip fracture is investigated by Keyak and Rossi [95]. They evaluated the distortion energy (DE), maximum normal stress, maximum normal strain, maximum shear strain, maximum shear stress, Coulomb-Mohr, modified Mohr, Hoffman, and strain-based Hoffman failure theories, using CT-based FE models of the femur [95].

The abovementioned fracture risk measurements are all derived from CT images. The most recent DXA-based fracture risk criterion is proposed by Luo et al. [91]. They calculated the averaged FRI as a ratio between the effective stress (von Mises stress) by applied forces and the allowable stress (yield stress) of the bone over a region of interest (ROI). FRI is a local fracture risk measurement, while FOS and RF are global ones.

Acknowledgements

This chapter is concluded from a research that was supported by Dr. Yunhua Luo, and therefore, he is gratefully acknowledged.

Conflict of interest

Masoud Nasiri Sarvi declares that he has no conflict of interest.

Author details

Masoud Nasiri Sarvi^{1,2*}

*Address all correspondence to: nasirism@myumanitoba.ca

1 Faculty of Engineering, Department of Mechanical Engineering, University of Manitoba, Canada

2 AI Incorporated, Toronto, Canada

References

- [1] Shao CJ, Hsieh YH, Tsai CH, Lai KA. A nationwide seven-year trend of hip fractures in the elderly population of Taiwan. *Bone*. 2009;**44**:125-129
- [2] Green C, Molony D, Fitzpatrick C, ORourke K. Age-specific incidence of hip fracture in the elderly: A healthy decline. *The Surgeon*. 2010;**8**:310-313
- [3] Gronskag AB, Forsmo S, Romundstad P, Langhammer A, Schei B. Incidence and seasonal variation in hip fracture incidence among elderly women in Norway. The HUNT study. *Bone*. 2010;**46**:1294-1298
- [4] Alvarez-Nebreda ML, Jimenez AB, Rodriguez P, Serra JA. Epidemiology of hip fracture in the elderly in Spain. *Bone*. 2008;**42**:278-285
- [5] Wilson RT, Chase GA, Chrischilles EA, Wallace RB. Hip fracture risk among community-dwelling elderly people in the United States: A prospective study of physical, cognitive, and socioeconomic indicators. *American Journal of Public Health*. 2006;**96**:1210-1218
- [6] Marks R, Allegrante JP, Ronald MacKenzie C, Lane JM. Hip fractures among the elderly: Causes, consequences and control. *Ageing Research Reviews*. 2003;**2**:57-93
- [7] Testi D, Viceconti M, Baruffaldi F, Cappello A. Risk of fracture in elderly patients: A new predictive index based on bone mineral density and finite element analysis. *Computer Methods and Programs in Biomedicine*. 1999;**60**:23-33
- [8] Greenspan SL, Myers ER, Kiel DP, Parker RA, Hayes WC, Resnick NM. Fall direction, bone mineral density, and function: Risk factors for hip fracture in frail nursing home elderly. *The American Journal of Medicine*. 1998;**104**:539-545
- [9] Cooper C, Campion G, LJ MI. Hip fractures in the elderly: A world-wide projection. *Osteoporosis International*. 1992;**2**:285-259
- [10] Kanis JA, Oden A, McCloskey EV, Johansson H, Wahl DA, Cooper C. A systematic review of hip fracture incidence and probability of fracture worldwide. *Osteoporosis International*. 2012;**23**:2239-2256
- [11] Nasiri Sarvi M, Luo Y. Sideways fall-induced impact force and its effect on hip fracture risk: A review. *Osteoporosis International*. 2017;**28**:2759-2780
- [12] Nasiri M, Luo Y. Study of sex differences in the association between hip fracture risk and body parameters by DXA-based biomechanical modeling. *Bone*. 2016;**90**:90, 98
- [13] Nasiri Sarvi M, Luo Y. A two-level subject-specific biomechanical model for improving prediction of hip fracture risk. *Clinical biomechanics*. 2015;**30**:881-887
- [14] Luo Y, Nasiri Sarvi M. A subject-specific inverse-dynamics approach for estimating joint stiffness in sideways fall. *International Journal of Experimental and Computational Biomechanics*. 2015;**3**:137-160
- [15] Nasiri Sarvi M. Assessment of hip fracture risk by a two-level subject-specific biomechanical model. Ph.D. thesis. Canada: Mechanical Engineering, University of Manitoba; 2015. p. 164

- [16] Nasiri SM, Luo Y. Development of an image-based biomechanical model for assessment of hip fracture risk. In: ASME 2015 International Design Engineering Technical Conferences and Computers and Information in Engineering Conference, IDETC/CIE; Boston; 2015
- [17] Nasiri SM, Luo Y. A compound risk indicator for subject-specific prediction of hip fracture in sideways falls. In: ASME 2015 International Design Engineering Technical Conferences and Computers and Information in Engineering Conference, IDETC/CIE 2015; Boston, 2015
- [18] Luo Y, Nasiri Sarvi M, Sun P, Ouyang J. A subject specific dynamics model for predicting impact force in elderly lateral fall. *Applied Mechanics and Materials*. 2014;**446-447**:339-343
- [19] Luo Y, Nasiri Sarvi M, Sun P, Leslie WD, Ouyang J. Prediction of impact force in sideways fall by image-based subject-specific dynamics model. *International Biomechanics*. 2014;1-14
- [20] Nasiri Sarvi M, Luo Y, Sun P, Ouyang J. Experimental validation of subject-specific dynamics model for predicting impact force in sideways fall. *Journal of Biomedical Science and Engineering*. 2014;**7**:405-418
- [21] Nasiri Sarvi M, Luo Y. Estimation of body segment masses using whole-body DXA image. In: 24th CANCAM Conference, June 2-6, 2013; Saskatoon
- [22] WHO. Assessment of fracture risk and its application to screening for osteoporosis. WHO technical report series 843 1994; Geneva
- [23] Sernbo I, Johnell O. Consequences of a hip fracture: A prospective study over 1 year. *Osteoporosis International*. 1993;**3**:148-153
- [24] Chrischilles E, Butler C, Davis C, Wallace R. A model of lifetime osteoporosis impact. *Archives of Internal Medicine*. 1991;**151**:2026-2032
- [25] Gullberg B, Johnell O, Kanis JA. World-wide projections for hip fracture. *Osteoporosis International*. 1997;**7**:407-413
- [26] Roche JJW, Wenn RT, Sahota O, Moran CG. Effect of comorbidities and postoperative complications on mortality after hip fracture in elderly people: Prospective observational cohort study. *British Medical Journal*. 2005;**331**:1-5
- [27] Boonen S, Autier P, Barette M, Vanderschueren D, Lips P, Haentjens P. Functional outcome and quality of life following hip fracture in elderly women: A prospective controlled study. *Osteoporosis International*. 2004;**15**:87-94
- [28] Lieberman D, Friger M, Fried V, Grinshpun Y, Mytlis N, Tylis R, Galinsky D. Characterization of elderly patients in rehabilitation: Stroke versus hip fracture. *Disability and Rehabilitation*. 1999;**21**:542-547
- [29] Phy MP, Vanness DJ, Melton L, Long KH, et al. Effects of a hospitalist model on elderly patients with hip fracture. *Archives of Internal Medicine*. 2005;**165**:796-801
- [30] Huddleston JM, Whitford KJ. Medical care of elderly patients with hip fractures. *Mayo Clinic Proceedings*. 2001;**76**:295-298

- [31] Majumder S, Roychowdhury A, Pal S. Hip fracture and anthropometric variations: Dominance among trochanteric soft tissue thickness, body height and body weight during sideways fall. *Clinical biomechanics*. 2013;**28**:1034-1040
- [32] Van den Kroonenberg AJ, Hayes WC, McMahon TA. Hip impact velocities and body configurations for voluntary falls from standing height. *Journal of Biomechanics*. 1996;**29**:807-811
- [33] Roberts BJ, Thrall E, Muller JA, Bouxsein ML. Comparison of hip fracture risk prediction by femoral aBMD to experimentally measured factor of risk. *Bone*. 2010;**46**:742-746
- [34] Marieb EN, Mallatt J. *Human Anatomy and Physiology*. Pearson plc; 2002
- [35] Hirshorn K. <http://stpetehipandknee.com/anatomy-hip-joint/>. 2014
- [36] Cumming R, Klineberg R. Fall frequency and characteristics and the risk of hip fractures. *Journal of the American Geriatrics Society*. 1994;**42**:774-778
- [37] Ford CM, Keaveny TM, Hayes WC. The effect of impact direction on the structural capacity of the proximal femur during falls. *Journal of Bone and Mineral Research*. 1996;**11**:377-383
- [38] Parker MJ. *Fractures of the hip*. Surgery (Oxford). 2003;**21**:221-224
- [39] Zioupos P, Wang XT, Currey JD. Experimental and theoretical quantification of the development of damage in fatigue tests of bone and antler. *Journal of Biomechanics*. 1996;**29**:989-1002
- [40] Youm T, Koval KJ, Kummer FJ, Zuckerman JD. Do all hip fractures result from a fall? *American Journal of Orthopedics*. 1999;**28**:190-194
- [41] Teppo LNJ, Harri S, Karim MK, Ari H, Pekka K. Shifting the focus in fracture prevention from osteoporosis to falls. *BMJ*. 2008;**336**:124-126
- [42] Cummings SR, Melton LJ. Epidemiology and outcomes of osteoporotic fractures. *The Lancet*. 2002;**359**:1761-1767
- [43] Kannus P, Leiponen P, Parkkari J, Palvanen M, Jarvinen M. A sideways fall and hip fracture. *Bone*. 2006;**39**:383-384
- [44] Greenspan SL, Myers ER, Maitland LA, Resnick NM, Hayes WC. Fall severity and bone mineral density as risk factors for hip fracture in ambulatory elderly. *JAMA*. 1994;**271**:128-133
- [45] Sabick MB, Hay JG, Goel VK, Banks SA. Active responses decrease impact forces at the hip and shoulder in falls to the side. *Journal of Biomechanics*. 1999;**32**:993-998
- [46] Kanis JA. Diagnosis of osteoporosis and assessment of fracture risk. *The Lancet*. 2002;**359**:1929-1936
- [47] Carpintero P, Caeiro JR, Carpintero R, Morales A, Silva S, Mesa M. Complications of hip fractures: A review. *World Journal of Orthopedics*. 2014;**5**:402-411
- [48] Rizzoli R, Bruyere O, Cannata-Andia JB, Devogelaer JP, Lyritis G, Ringe JD, Vellas B, Reginster JY. Management of osteoporosis in the elderly. *Current Medical Research and Opinion*. 2009;**25**:2373-2387

- [49] Jee W. Integrated bone tissue physiology: Anatomy and physiology. In: Cowin Bone Mechanics Handbook. New York: Informa Healthcare; 2001
- [50] Cummings SR, Nevitt MC, Browner WS, Stone K, Fox KM, Ensrud KE, Cauley J, Black D, Vogt TM. Risk factors for hip fracture in white women. *New England Journal of Medicine*. 1995;**332**:767-774
- [51] Fisher AA, O'Brien ED, Davis MW. Trends in hip fracture epidemiology in Australia: Possible impact of bisphosphonates and hormone replacement therapy. *Bone*. 2009;**45**:246-253
- [52] Johnell O, Kanis JA. An estimate of the worldwide prevalence and disability associated with osteoporotic fractures. *Osteoporosis International*. 2006;**17**:1726-1733
- [53] Brauer CA, Coca-Perrillon M, Cutler DM, Rosen AB. Incidence and mortality of hip fractures in the United States. *The Journal of the American Medical Association*. 2009;**302**:1573-1579
- [54] Leibson CL, Tosteson ANA, Gabriel SE, Ransom JE, Melton LJ. Mortality, disability, and nursing home use for persons with and without hip fracture: A population-based study. *Journal of the American Geriatrics Society*. 2002;**50**:1644-1650
- [55] Haentjens P, Magaziner J, Colon-Emeric CS, Vanderschueren D, Milisen K, Velkeniers B, Boonen S. Meta-analysis: Excess mortality after hip fracture among older women and men. *Annals of Internal Medicine*. 2010;**152**:380-390
- [56] Greendale G, Barrett-Connor E. *Outcomes of Osteoporotic Fractures*. 2nd ed. San Diego: Academic Press; 2001
- [57] Ross PD. Clinical consequences of vertebral fractures. *The American Journal of Medicine*. 1997;**103**:S30-S43
- [58] Gold D, Lyles K, Shipp K, Drezner M. *Osteoporosis and its Nonskeletal Consequences: Their Impact on Treatment Decisions*. 2nd ed. San Diego: Academic Press; 2001
- [59] Gallup. *Osteoporosis Prevalence Figures; State-By-State Report*. Washington, D.C.: National Osteoporosis Foundation; 1997
- [60] Tosteson ANA, Gabriel SE, Grove MR, Moncur MM, Kneeland TS, Melton III LJ. Impact of hip and vertebral fractures on quality-adjusted life years. *Osteoporosis International*. 2001;**12**:1042-1049
- [61] Phillips S, Fox N, Jacobs J, Wright WE. The direct medical costs of osteoporosis for American women aged 45 and older. *Bone*. 1988;**9**:271-279
- [62] Lonnroos E, Kautiainen H, Karppi P, Huusko T, Hartikainen S, Kiviranta I, Sulkava R. Increased incidence of hip fractures. A population based-study in Finland. *Bone*. 2006;**39**:623-627
- [63] Wiktorowicz ME, Goeree R, Papaioannou A, Adachi JD, Papadimitropoulos E. Economic implications of hip fracture: Health service use, institutional care and cost in Canada. *Osteoporosis International*. 2001;**12**:271-278

- [64] Dy CJ, McCollister KE, Lubarsky DA, Lane JM. An economic evaluation of a systems-based strategy to expedite surgical treatment of hip fractures. *The Journal of Bone & Joint Surgery*. 2011;**93**:1326-1334
- [65] Hayes WC, Myers ER, Robinovitch SN, Van Den Kroonenberg A, Courtney AC, McMahon TA. Etiology and prevention of age-related hip fractures. *Bone*. 1996;**18**:S77-S86
- [66] Kannus P, Parkkari J, Sievänen H, Heinonen A, Vuori I, Järvinen M. Epidemiology of hip fractures. *Bone*. 1996;**18**:S57-S63
- [67] An YH, Draughn RA. *Mechanical Testing of Bone and the Bone-Implant Interface*. Florida: CRC Press; 2000. pp. 407-438
- [68] Butler M, Forte M, Kane R. Treatment of common hip fractures. Evidence Report/Technology Assessment. 2009;**184**:1-85
- [69] Peters M. <http://advancedortho.net/info/hipfractureinfo.php>. 2000
- [70] Dhanwal D, Dennison E, Harvey N, Cooper C. Epidemiology of hip fracture: Worldwide geographic variation. *Indian Journal of Orthopaedics*. 2011;**45**:15-22
- [71] Melton LJ. Hip fractures: A worldwide problem today and tomorrow. *Bone*. 1993;**14**:1-8
- [72] Jacobsen SJ, Goldberg J, Miles TP. Hip fracture among the old and very old: A population-based study of 745 435 cases. *The American Journal of Public Health*. 1990;**80**:871-873
- [73] Cummings SR, Black DM, Rubin SM. Lifetime risks of hip, colles, or vertebral fracture and coronary heart disease among white postmenopausal women. *Archives of Internal Medicine*. 1989;**149**:2445-2448
- [74] Melton JL. Perspectives: How many women have osteoporosis now? *Journal of Bone and Mineral Research*. 1995;**10**:175-177
- [75] Winner SJ, Morgan CA, Evans JG. Perimenopausal risk of falling and incidence of distal forearm fracture. *BMJ*. 1989;**298**:1486-1488
- [76] Iwamoto J, Takeda T, Sato Y. Effect of muscle strengthening exercises on the muscle strength in patients with osteoarthritis of the knee. *The Knee*. 2007;**14**:224-230
- [77] Wijnhuizen GJ, de Jong R, Hopman-Rock M. Older persons afraid of falling reduce physical activity to prevent outdoor falls. *Preventive Medicine*. 2007;**44**:260-264
- [78] Chang WW, Friedman S. Hip fracture management. *Hospital Medicine Clinics*. 2013;**2**:e399-e421
- [79] Kubota M, Uchida K, Kokubo Y, Shimada S, Matsuo H, Yayama T, Miyazaki T, Sugita D, Watanabe S, Baba H. Postoperative gait analysis and hip muscle strength in patients with pelvic ring fracture. *Gait & Posture*. 2013;**38**:385-390
- [80] Groen BE, Weerdesteyn V, Duysens J. Martial arts fall techniques decrease the impact forces at the hip during sideways falling. *Journal of Biomechanics*. 2007;**40**:458-462

- [81] Van der Zijden AM, Groen BE, Tanck E, Nienhuis B, Verdonschot N, Weerdesteyn V. Can martial arts techniques reduce fall severity? An in vivo study of femoral loading configurations in sideways falls. *Journal of Biomechanics*. 2012;**45**:1650-1655
- [82] Derler S, Spierings AB, Schmitt KU. Anatomical hip model for the mechanical testing of hip protectors. *Medical Engineering & Physics*. 2005;**27**:475-485
- [83] Holzer G, Holzer L. Hip protectors and prevention of hip fractures in older persons. *Geriatrics*. 2007;**62**:15-22
- [84] Holzer LA, von Skrbensky G, Holzer G. Mechanical testing of different hip protectors according to a European standard. *Injury*. 2009;**40**:1172-1175
- [85] Laing AC, Robinovitch SN. The force attenuation provided by hip protectors depends on impact velocity, pelvic size, and soft tissue stiffness. *Journal of Biomechanical Engineering*. 2008;**130**:1-9
- [86] Li N, Tsushima E, Tsushima H. Comparison of impact force attenuation by various combinations of hip protector and flooring material using a simplified fall-impact simulation device. *Journal of Biomechanics*. 2013;**46**:1140-1146
- [87] Robinovitch SN, Evans SL, Minns J, Laing AC, Kannus P, Crompton PA, Derler S, Birge SJ, Plant D, Cameron ID, Kiel DP, Howland J, Khan K, Lauritzen JB. Hip protectors: Recommendations for biomechanical testing-an international consensus statement (part I). *Osteoporosis International*. 2009;**20**:1977-1988
- [88] Lilliu H, Pamphile R, Chapuy M-C, Schulten J, Arlot M, Meunier PJ. Calcium-vitamin D3 supplementation is cost-effective in hip fractures prevention. *Maturitas*. 2003;**44**:299-305
- [89] Ascherio A, Munger KL, Simon KC. Vitamin D and multiple sclerosis. *The Lancet Neurology*. 2010;**9**:599-612
- [90] Keyak JH, Rossi SA, Jones KA, Skinner HB. Prediction of femoral fracture load using automated finite element modeling. *Journal of Biomechanics*. 1998;**31**:125-133
- [91] Luo Y, Ferdous Z, Leslie WD. Precision study of DXA-based patient-specific finite element modeling for assessing hip fracture risk. *International Journal for Numerical Methods in Biomedical Engineering*. 2013;**29**:615-629
- [92] Schileo E, Taddei F, Cristofolini L, Viceconti M. Subject-specific finite element models implementing a maximum principal strain criterion are able to estimate failure risk and fracture location on human femurs tested in vitro. *Journal of Biomechanics*. 2008;**41**:356-367
- [93] Lotz JC, Cheal EJ, Hayes WC. Fracture prediction for the proximal femur using finite element models: Part I-linear analysis. *Journal of Biomechanical Engineering*. 1991;**113**:353-360
- [94] Lotz JC, Cheal EJ, Hayes WC. Fracture prediction for the proximal femur using finite element models: Part II-nonlinear analysis. *Journal of Biomechanical Engineering*. 1991;**113**:361-365
- [95] Keyak JH, Rossi SA. Prediction of femoral fracture load using finite element models: An examination of stress- and strain-based failure theories. *Journal of Biomechanics*. 2000;**33**:209-214