

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

186,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.  
For more information visit [www.intechopen.com](http://www.intechopen.com)



# Dietary Antioxidants in Experimental Models of Liver Diseases

*Sila Appak-Baskoy, Mustafa Cengiz, Ozgun Teksoy  
and Adnan Ayhanci*

## Abstract

Oxidative stress is caused by the imbalance between the amount of reactive oxygen species (ROS) and antioxidant capacity in the body. A balanced diet involving the daily intake of antioxidant-rich foods makes improvements in the total antioxidant capacity of individuals and would therefore reduce the incidence of oxidation-related diseases. It may also regulate the degree of oxidative stress. In fact, dietary micronutrients are either direct antioxidants or components of antioxidant enzymes, which may contribute positively to certain indicators of hepatic function. Liver plays an important role in the regulation of various processes such as metabolism, secretion, storage, and the clearance of endogenous and exogenous substances. Once liver is damaged by pursuing a wrong diet and inflammation takes place, most of these physiological functions get altered. Apart from drugs that used to treat the ailments, it is also necessary to determine the pharmaceutical alternatives for the drugs that are used in the treatment of liver diseases. Therefore, this chapter aims to summarize all known information on the effects of dietary nutrients on oxidative stress in experimental liver models.

**Keywords:** dietary antioxidant, free radicals, oxidative stress, liver

## 1. Introduction


The liver is the major organ metabolizing xenobiotics and endogenous molecules in order to maintain metabolic homeostasis in the organism, which is why it is a target of many toxic substances that cause dysregulated hepatic homeostasis. One of the mostly found clinical liver diseases is nonalcoholic fatty liver disease (NAFLD) (**Figure 1**) [2, 3]. In NAFLD, hepatocytes get filled up with triglycerides, liver expands and its normal functions may get altered. Although it is a disease, the triglyceride accumulation can still be reversed and normal functions may be restored by proper nutrition and exercise. If these lifestyle changes are not pursued, the damage goes on and results in inflammation followed with fibrosis which is unfortunately irreversible.

The liver is made up of hepatocytes, Kupffer cells, liver sinusoidal endothelial cells, pit cells, and hepatic stellate cells (HSC) [4]. Activation of stellate cells by injury caused by many etiological factors would lead to cirrhosis, and it would mark the end stage of progressive fibrosis [5]. Oxidative stress has a vital part in establishing

fibrosis and consequently cirrhosis [6]. For this reason, using molecules with antioxidant properties has been proposed as a treatment for not only fibrosis but also oxidative stress-related cirrhosis. Liver diseases are considered a major medical problem worldwide. There are known to be a large number of liver diseases caused by different insults. Furthermore, the disease type depends on lifestyle factors. For example, the main causes of liver diseases are reported to be viral and parasitic infections in regions like Africa and Asia. For Europe and America, alcohol consumption is thought to be the most important cause of this disease. However, viral hepatitis has showed an increase in recent times in most of the countries [7]. Lifestyle and unhealthy diet is the leading cause of liver diseases in almost all western countries. Until today, no medication is approved for the treatment of this disease; however, improving diet habits and physical exercise works if the disease is not accompanied by inflammation. On the other hand, biologically active food compounds that regulate gene expressions in lipogenesis, fibrosis, and inflammation serve as good therapeutic means to ameliorate these pathological states observed in liver [1] (**Figure 1**).

1.1 Oxidative stress

Oxidative stress is recognized as a disproportion between the production of free radicals (FR) and the antioxidant defenses [8]. Increased levels of prooxidants result in damage to the cell in terms of lipid peroxidation as well as oxidative DNA damage




**Figure 1.** Molecular mechanisms explaining the hepatoprotective effect of food bioactives. Development of NAFLD/ NASH is induced by different risk factors, such as Western-type diet, physical inactivity, and genetic predisposition. In the presence of obesity and IR, there is an increased flux of FFAs to the liver. These FFAs are stored as TG in lipid droplets leading to hepatic fat accumulation or undergo  $\beta$ -oxidation increasing oxidative stress and the inflammatory pathway. The damaged hepatocyte leads to a further increase of inflammatory signaling (IL-1, TNF $\alpha$ , IL-6) and the recruitment of circulating and residual macrophages (KCs). All of these mechanisms can directly induce the activation of HSCs, the major cell type involved in extracellular matrix deposition and liver fibrosis. The bioactive compounds may exert beneficial effects on NAFLD development and progression by inhibiting lipogenesis,  $\beta$ -oxidation of FFAs, inflammation, and HSCs activation. In the cartoon, we have listed the food bioactives indicating the putative mechanisms by which they may improve liver damage in NAFLD [1].

and thus protein damage [9]. One or more unelectrified FR atoms or molecules may be present as radical cations or radical anions. They are usually unstable and highly reactive because they can react with molecules and abstract electrons. Oxygen can reduce and produce reactive oxygen species (ROS) with exciting electrons, secondary to the interaction of transition metals or by the addition of energy [9, 10]. Oxidative stress causes fibrogenesis by increasing transforming cytokines including transforming grown factor-beta-1 (TGFβ1), interleukin-6 (IL-6), and tumor necrosis factor-alpha (TNFα) [3]. Disruption of the liver metabolism arises from increased quantity of ROS to amplified electron transfer in mitochondrial B-oxidation and increased expression and activity of Cyp2e1 that is from CYP450 family [11]. Intense production of Cyp2e1 is present because of much more consumption of ethanol which is produced by virtue of a lot of direct and indirect mechanisms [12–14] (Figure 2).

## 1.2 Basic sources of free radicals

Cells produce FR as a result of metabolic events; however, this is not the only source that can cause oxidative stress in body. The pollutants in the environment such as toxic chemicals as well as radiation cause a significant increase in amount of FR, ROS, and reactive nitrogen species (RNS) [10]. In the body, variety of different cell types and chemical reactions produce ROS, the most important metabolism is the cytochrome P450 metabolism and mitochondria-catalyzed electron transport reactions. Most of the inflammatory conditions are also responsible from ROS production, and important cell types in these processes are neutrophils, eosinophils,



**Figure 2.**

*Mechanisms of enhanced ROS production during hepatocyte damage. Ethanol metabolism promotes strong ROS production in the ER by the inducible CYP. It impairs GSH import in the mitochondria, preventing ROS removal. It also impairs B-oxidation promoting lipid accumulation. ETOH induces lipid-raft clustering and increases iron uptake, promoting Fe<sup>2+</sup> leakage from lysosomes and increased Fe<sup>2+</sup> loads in mitochondria and ER, resulting in ROS production. Ethanol also reduced the autophagic removal of damaged cellular components. Viral infection challenges the ER protein folding process leading to ROS production and Ca<sup>2+</sup> leakage in the cytosol and mitochondria. Increased MAMs formation promotes Ca<sup>2+</sup> efflux from ER into mitochondria, increasing mitochondrial ROS production [15].*



and macrophages [16, 17]. The chief molecule responsible for the reduction of oxygen in mitochondria is ubiquinone. Mitochondria is such an important organelle to produce ROS and hydrogen peroxide ( $\text{H}_2\text{O}_2$ ) as it produces 2–3 nmol of superoxide/min per mg of protein, [17]. Different tissues of mammals and different species of mammals have an enzyme called xanthine oxidase, an enzyme belonging to molybdenum, iron-sulfur, flavin hydroxylases that play an important role in the hydroxylation of purines by the oxidation of hypoxanthine to xanthine. Resultant xanthine then oxidized to uric acid. Oxygen reduction takes place in both of these reactions and the first one produces  $\text{O}_2^-$ , while the second one produces  $\text{H}_2\text{O}_2$  [16]. Inflammation serves as another source of ROS generation. During inflammation, activated macrophages increase their oxygen uptake, and this process results in production of  $\text{O}_2^-$ , nitric oxide (NO), and  $\text{H}_2\text{O}_2$  [18]. Another mechanism of  $\text{O}_2^-$  production during inflammation is by neutrophils; the enzyme nicotinic adenine dinucleotide phosphate [NAD(P)H] oxidase generates  $\text{O}_2^-$  that is used to destroy bacteria and this nonphagocytic NAD(P)H oxidases produce  $\text{O}_2^-$  in a range of 1–10% [19]. Cytochrome P450 (CYP) enzymes are also important in the production of ROS by the breakdown and/or uncoupling of the P450 catalytic cycle. Hyperoxia would trigger 80% of the  $\text{H}_2\text{O}_2$  synthesis by microsomes, and under normoxic conditions, peroxisomes produce  $\text{H}_2\text{O}_2$  but not  $\text{O}_2^-$ , and most of the peroxisomal  $\text{H}_2\text{O}_2$  production takes place in liver [16]. Arginine is reduced to citrulline in a five-electron oxidative reaction by nitric oxide synthases (NOSs) and this reaction gives rise to NO. Immune cells can also produce NO in the oxidative burst during inflammation. NO can react with oxygen and water in an extracellular environment in order to form nitrate and nitrite anions. Also, the NO and  $\text{O}_2^-$  can react together and cause a more reactive FR called peroxynitrite anion ( $\text{ONOO}^-$ ) which can cause lipid peroxidation and fragmentation of DNA [20].

### 1.3 Antioxidants

Antioxidants are molecules that can help prevent or delay oxidation of an oxidizable substrate when in low concentrations and they have a high affinity to FR [21]. Antioxidants play an important role to maintain health of the organism by scavenging FR by donating electrons to it. This reduces the reactivity of FR and helps maintain prooxidant/antioxidant balance in cell. A lot of different molecules that have antioxidant activity have been identified. Different natural compounds have so far been studied extensively especially in liver diseases (Table 1).

### 1.4 Curcumin

Curcumin, diferuloylmethane or 1,7-bis (4-hydroxy-3-methoxyphenyl)-6-heptadieno-3,5-dione is obtained from the rhizomes of *Curcuma longa* (turmeric). Curcumin has many pharmacological properties as it is a strong antioxidant, antifibrogenic, anti-inflammatory, antimicrobial, and anticarcinogenic agent and it also aids in a wound healing [22]. The Food and Drug Administration (FDA) has classified turmeric as a safe substance and toxicity assays done on animals have shown curcumin to be safe even when used in high doses. On the other hand, prolonged high-dose intake of turmeric has been associated with incidences of hepatotoxicity in mice and rats [3]. Curcumin is known to have low bioavailability when administered orally. Arcaro et al. [23] used piperine (inhibitor of hepatic and intestinal absorption) together with curcumin. Even in the presence of piperine, antidiabetic and antioxidant activity of curcumin was not altered. But when higher dose of piperine (40 mg/kg) was used, the beneficial effects of curcumin vanished. On the other hand, Sehgal et al. [24] showed the effect of piperine on curcumin in benzo(a)pyrene toxicity in the liver. They found that pretreatment with 100 mg/kg of curcumin protects against

Antioxidant	Main clinical effects	Clinical relevance
Curcumin	Antioxidant	No studies available in human hepatic disorders
	Antifibrotic	
	Anti-inflammatory	
	Antimicrobial	
	Wound healing	
	Anticarcinogenic	
Resveratrol	Antioxidant	Current data are conflicting, so more clinical studies are needed
	Anti-inflammatory	
	Anticarcinogenic	
	Lipid modulation	
	Antifibrotic	
Coffee	Antinecrotic	Inverse relationship between coffee-cirrhosis has been demonstrated, but more basic research and prospective clinical trials are necessary
	Antifibrotic	
	Antioxidant	
	Anticholestatic	
	Chemoprotective	
Quercetin	Chelation of transition metal ions	No studies available in human hepatic disorders
	Anticarcinogenic	
	Cardioprotective	
	Bacteriostatic	
	Antioxidant	
	Antifibrotic	
	Anti-inflammatory	
	Antiapoptotic	
	Antiaggregatory	
	Vasodilating	
Silymarin	Antioxidant	Silymarin has been shown to be effective, but it is necessary to do more clinical trials focused on survival rates of patients with cirrhosis
	Antifibrotic	
	Anti-inflammatory	
	Anticarcinogenic	
	Immunomodulation	
Naringenin	Antioxidant	No studies available in human hepatic disorders
	Hypocholesterolemic	
	Anti-estrogenic	
	Hypolipidemic	
	Antihypertensive	
	Anti-inflammatory	
	Antifibrotic	
	Anticarcinogenic	
	Antiatherogenic	

Antioxidant	Main clinical effects	Clinical relevance
Green tea	Anti-inflammatory	More clinical studies are needed
	Antiarthritic	
	Antimicrobial	
	Antioxidant	
	Neuroprotective	
	Antidiabetic	
	Antiangiogenesis	
	Anticarcinogenic	
L-carnitine	Chelation of transition metal ions	More clinical studies are needed
	Antioxidant	
	Cardioprotective	
	Neuroprotective	
	Anti-inflammatory	
Lycopene	Antioxidant	More clinical studies are needed
	Anti-inflammatory	
	Chemoprotective	
	Anticarcinogenic	

**Table 1.**  
*Nutritional antioxidants.*

a single dose of benzo(a)pyrene; and at this dose, coadministration of piperine had a much better effect than did curcumin alone showing enhancer activity of piperine. In acute and chronic liver injury, curcumin has been shown to have hepatoprotective effects [25]. In 2007, Reyes-Gordillo et al. [26] showed that curcumin is able to inhibit the release of TNF- $\alpha$ , IL-1B, and IL-6. Additionally, curcumin reduces carbon tetra-chloride (CCl<sub>4</sub>)-mediated oxidative stress inactivating the nuclear factor-kB (NF-kB) pathway. Moreover, curcumin’s hepatoprotective effect takes place by its interactions with Fe<sup>3+</sup> and Cu<sup>2+</sup>. A study by Jiao et al. [27] suggested that curcumin could serve as an iron chelator since transferrin receptor 1 and iron regulatory proteins, indicators of iron depletion, showed an increase with curcumin administration. Charoensuk et al. [28] have indicated that curcumin increases antioxidant capacity of cells by increasing mRNA and protein levels of factors and enzymes such as nuclear factor erythroid 2-related factor 2 (Nrf2), heme oxygenase (HO-1), NAD(P)H quinone oxidoreduc-tase 1 (NQO1), glutamate cysteine ligase (GCL), transcription factor-3, peroxiredoxin 3 (Prdx3), and Prdx6. Curcumin also increases the activity of glutathione (GSH), superoxide dismutase (SOD), catalase (CAT), glutathione peroxidase (GPx), and glutathione-S-transferase (GST) activity [29, 30]. Curcumin also interacts with enzymes or genes that are important in liver cirrhosis. Hassan et al. [31] showed that curcumin modulates miRNA 199 and 200 which are associated with liver fibrosis in CCl<sub>4</sub>-induced experimental fibrosis model and that curcumin reduced these miRNAs levels close to their basal levels. Finally, in alcohol-induced liver damage, curcumin inhibits the activity of cytochrome P450 2E1 (Cyp2e1) and also its protein levels [32].

1.5 Resveratrol

The phytoalexin resveratrol (3,5,4'-trans-trihydroxystilbene) is a polyphen-ol mostly found in red grapes, red wine, peanuts, and berries [33]. Resveratrol has effects on lipid metabolism, and it also has antioxidant, anti-inflammatory,

anticarcinogenic, and antifibrogenic properties [34]. The rate of absorption of resveratrol is about 75% following an oral administration [35]. Resveratrol is metabolized to resveratrol sulfate, and in its low concentrations, it is converted into resveratrol glucuronide [36] by enzymes glucuronosyltransferase (UGT) or sulfotransferase (ST) [37]. In 2007, Chávez et al. [34] showed that under CCl<sub>4</sub>, resveratrol decreased cytokine TGF- $\beta$  levels and prevented hepatic fibrosis. It also inhibited NF- $\kappa$ B translocation to the nucleus. Resveratrol, as an antioxidant, has protective effects against ethanol-induced lipid peroxidation, toxicity by acetaminophen (APAP), and oxidative stress in animal models of cholestasis [38]. Important player in resveratrol's antioxidant activity is suggested to be run by the OH groups [7]. Blocking OH group methylation showed that resveratrol and trimethylated resveratrol provide some degree of protection, but the latter one has a better protective effect [39]. Another hepatoprotection mechanism of resveratrol comes from its ability to activate genes related to antioxidant system or from its ability to inhibit enzymes. A study by Cheng et al. [40] suggested that resveratrol could activate extracellular signal-regulated kinase (ERK) signaling pathway, which may, in turn, enhance the activation and translocation of Nrf2 to the nucleus, thus increasing the expression of HO-1 and glyoxalase. Price et al. [41] found that resveratrol activates AMP-activated protein kinase (AMPK) and increased nicotinamide adenine dinucleotide (NAD) levels in mice. Zhu et al. [42] have also shown that, in mice, administration of resveratrol increased the antioxidant system (SOD, GPx, and GSH) and also the levels of SIRT1 and p-AMPK were upregulated in liver. Resveratrol has also been shown to inhibit the activity of Cyp2e1 in microsomes of rat liver [43]. Resveratrol also inhibited the activity of P450 isoform APAP-induced liver injury model [44] and activity of Cyp2e1 was also inhibited in diethylnitrosamine (DEN)-induced hepatocarcinogenesis model [45]. The only clinical study that was performed to determine the resveratrol hepatoprotective effect demonstrated that a 500 mg resveratrol dose administered for 12 weeks caused a significant reduction in inflammatory cytokines, serum cytokeratin-18, NF- $\kappa$ B activation, liver alanine aminotransferase (ALT), and hepatic steatosis when compared to the placebo group in patients with nonalcoholic fatty liver disease (NAFLD) [46].

## 1.6 Coffee

Coffee is a mixture of several different molecules such as carbohydrates, vitamins, lipids, nitrogenous molecules, alkaloids, and phenolic compounds [47]. Caffeine, diterpene alcohols (cafestol and kahweol), and chlorogenic acid are the three major compounds found in coffee [48]. Coffee consumption has been linked to the reduction of several chronic diseases [49], probably due to the pharmacological properties that have antinecrotic, antifibrotic, anticholestatic, chemoprotective, and antioxidant functions [50]. Caffeine is the best-known active component of coffee, which is absorbed very rapidly once it has been taken orally (5 min), reaching its peak blood levels after 30 min. When consumed in high amounts, it may produce some side effects. Recommendations from Health Canada in 2013 demanded that the daily caffeine intake for children should not exceed 2.5 mg/kg of the body weight. What is more, tachycardia and arrhythmia typically arise when more than 200 mg of caffeine are ingested [51]. Smith et al. [52] reported in 2002 that the intake of 300 mg of caffeine resulted in a rise in anxiety and tension. Caffeine gets metabolized in the liver. The principal metabolite of caffeine is paraxanthine [53]. An important property of caffeine is that it can easily pass through the blood-brain barrier [54]. Coffee-cirrhosis relationship was shown by Klatsky et al. for the first time [55]. The study showed that the odds ratio for liver cirrhosis tend to decrease from 1.0 for people abstaining from coffee to 0.47, 0.23, 0.21, and 0.16 for 1, 2, 3, or 4 cups of coffee daily, respectively. Although coffee is generally beneficial to the liver, this study failed to show a causative



role of coffee in prevention of liver injury. Therefore, additional basic research and controlled prospective studies are needed in order to show exact effect of coffee on liver tissue. Arauz et al. [50] demonstrated that coffee has a protective effect upon liver injury caused by chronic administration of thioacetamide (TAA). Coffee ameliorated cholestasis and necrosis and this was seen by the measurement of  $\gamma$ -glutamyl transpeptidase ( $\gamma$ -GTP), alkaline phosphatase, and ALT levels. Arauz et al. [50] demonstrated in murine models that coffee prevents experimental liver cirrhosis. In these studies, coffee reduced the expression of the profibrogenic cytokine TGF- $\beta$ . Cavin et al. [56] reported coffee to be an inducer of GST, aldo-keto reductase, GSH, HO-1, GST-P1, which are enzymes involved in the detoxification process. Also, they suggested that a possible mechanism of chemoprotection of coffee by stimulating the Nrf2 pathway. In another study, coffee was able to elevate mRNA levels of NQO1 and glutathione-S transferase A1 in the liver and the small intestine [57].

### 1.7 Quercetin

Quercetin, 3,3,4,5,7-penta-hydroxyflavone, is a flavonol especially found in apples and onions [58]. Quercetin chelates heavy metals and has anticarcinogenic, cardioprotective, bacteriostatic, anti-inflammatory, and antioxidant properties [59]; it also functions as a hepatoprotective agent [60]. The normal daily intake of quercetin is less than 5–40 mg. However, if the peels of the foods that contain high amount of quercetin are also consumed, daily intake of quercetin increases to 200–500 mg [59]. In 2004, high-purity quercetin used in foods was grass that serves 10–125 mg of quercetin [59]. The functional groups responsible for quercetin's antioxidant activity were described by Bors et al. [61] in 1990, and they found that orthodihydroxy or catechol groups in the B-ring, a 2,3-double bond of the C-ring, and OH substitution on positions 3 and 5 of the C-ring and A-ring, respectively, are important players in antioxidant action of quercetin [61]. It can interact with both FR and metal ions like  $\text{Fe}^{3+}$  and  $\text{Cu}^{2+}$  for chelation. In a study by Mira et al. [62], it was reported that reduction of  $\text{Fe}^{3+}$  and  $\text{Cu}^{2+}$  takes place by quercetin's 2,3-double bond and the presence of catechol group in the B-ring. Following ingestion, quercetin is rapidly absorbed and its levels in blood peak at approximately in 30 min [63] before it is metabolized by glucuronidation and sulfation by the UGT and ST, respectively.

In experimental fibrosis model in rats, quercetin showed hepatoprotective properties under  $\text{CCl}_4$  treatment that lasted 8 weeks. The hepatoprotective effect of quercetin was found to be mediated by its ability to suppress the expression of profibrogenic expressions of TGF- $\beta$ , CTGF, and collagen-1 $\alpha$  (Col-1 $\alpha$ ). On the other hand, quercetin also activated enzymes such as metalloproteinases 2 and 9 (MMP2 and MMP9); it also improved the activity of SOD and CAT [64]. Pavanato et al. [65] extended  $\text{CCl}_4$  treatment for 16 weeks and also observed that quercetin improved the hepatic liver enzymes AST, ALT, and inducible NOS (iNOS) expressions; it also decreased collagen amount and reduced lipid peroxidation in liver. Granado-Serrano et al. [66] showed that quercetin modulated Nrf2 and p38 in HepG2 cells. Quercetin has also been shown to suppress the activity of Cyp2e1 in hepatocytes in the presence of ethanol [67]. In line with this finding, in a nonalcoholic steatohepatitis (NASH) model, quercetin was able to reduce Cyp2e1 activity [68].


### 1.8 Silymarin

Silymarin, milk thistle or Saint Mary's thistle, is a natural substance obtained from *Silybum marianum* [7]. Silymarin has not been associated with any side effects at acute consumption and the dose range used in literature ranges between 280 and 800 mg/kg of body weight per day. After oral administration, the silymarin peak

plasma concentration is achieved at approximately 6–8 h. The metabolites of silymarin get conjugated in the liver by UGT and ST (phase II reactions) [69]. This substance has many hepatoprotective effects (**Figure 3**). In fact, silybin, a major constituent of silymarin, has found to have iron chelating properties [71]. In a study performed by Najafzadeh et al. [72], hepatoprotective effect of silymarin in iron-overload-induced hepatotoxicity was attributed to its iron-chelator activity; however, no studies have proved the chelating properties per se of silymarin in liver diseases. Silymarin acts as hepatoprotector against several hepatotoxins including D-galactosamine [73]. Silymarin's ameliorating effects on oxidative stress, fibrosis, cirrhosis, and lipid peroxidation are modulated by its phosphatidylethanolamine amount [74]. This hepatoprotective effect is seen with the improvement of liver enzyme activities and levels of cholesterol/phospholipids and also sphingomyelin/phosphatidylcholine ratios in the membrane [75, 76]. Kim et al. [77] showed that silymarin increases nuclear translocation of Nrf2 in activated HSC. Also, silymarin increases the activity of antioxidant enzymes such as SOD, GPX [78], and CAT [79]. A clinical trial examining silymarin in a complex with phosphatidylcholine found reduced levels of the liver enzymes, ALT and  $\gamma$ -GGT, and serum bilirubin levels in a dose-dependent manner in patients suffering from hepatitis due to virus infection or alcohol abuse [80] (**Figure 3**).

### 1.9 Naringenin

Naringenin is also recognized as 5,7,4'-thihydroxyflavanone, and it is a flavanone found in citrus fruits and tomatoes [81]. In a recent study, Yang et al. [82] have



**Figure 3.**  
A schematic notation of the main pharmacological effects of silymarin in accordance with its hepatoprotective features: The effects of Sm upon cell membranes (upper left) and intracellular cascades are shown here. The metabolic paths are indicated by interrupted lines, while its signal effects are shown in full lines. LTB<sub>4</sub>, leukotriene B<sub>4</sub>; GSH, glutathione; NF- $\kappa$ B, nuclear factor kappa B; PG's, prostaglandins; Sm, silymarin; SOD, superoxide dismutase; ROS, reactive oxygen types=; TNF $\alpha$ , tumor necrosis factor  $\alpha$  [70].

reported that naringenin did not cause any harmful effects in beagle dogs, the maximum time of exposure being 180 days and with doses varying of 20, 100, or 500 mg/kg body weight per day. Naringenin has many pharmacological properties. It acts as a hypolipidemic, antihypertensive, anti-inflammatory, antioxidant, and antifibrotic agent [81]. The metabolism of naringenin takes place in small intestine where glycoside form of naringenin gets cleaved, resulting in sulfate and glucuronide metabolites in the small intestine wall; then, it gets absorbed [77]. Mira et al. [62] showed that naringenin can reduce  $\text{Fe}^{3+}$  and  $\text{Cu}^{2+}$  ions but it is less potent than quercetin. Chtourou et al. [83] found that naringenin averts depletion of SOD, CAT, GPx, and GSH. On the other hand, naringenin also prevents an increase in lipid peroxidation, and it also prevents increase of enzymes ALT and AST [78]. Yen et al. [84] also obtained similar results on liver enzymes and prevention of lipid peroxidation when they used naringenin alone and also naringenin-loaded nanoparticle system (NARN). In both treatments, naringenin exhibited antioxidant and hepatoprotective activities. In these experiments, treatment with naringenin also inhibited the activation of caspases 3 and 8. However, NARN was found to have better hepatoprotective and antioxidant effects than free naringenin, and it was also shown to inhibit caspase 9 during  $\text{CCl}_4$ -induced hepatotoxicity in rats. Han et al. [64] reported that a pre-treatment with naringenin-7-O-glucoside increases NQO1 and ERK phosphorylation and translocation of Nrf2 to the nucleus in H9c2 cardiomyocytes. It also upregulated the mRNA expression of GCLC and GCL modifier [64]. Similar findings have been reported by Esmaeili et al. [85] who showed that naringenin attenuates  $\text{CCl}_4$ -induced liver injury by downregulating TNF- $\alpha$ , INOS, and cyclooxygenase-2 and also by increasing Nrf2 and HO-1 expressions. Motawi et al. [86] showed that naringenin inhibits Cyp2e1 in liver microsomal assay done on rats [86].

### 1.10 Green tea

*Camellia sinensis* or green tea is a widely consumed beverage across the globe and it has antioxidant, anti-inflammatory, antiarthritic, and antiangiogenic effects. It is a mixture of polyphenols (the major class of active compounds) including catechins (also known as flavan-3-ols) which constitute about 30% (mass fraction) of green tea leaves. The major catechins in green tea are (+) -catechin, (–) -epicatechin, (–) -epigallocatechin, (–) -epicatechin-3-gallate, (–) -gallocatechin, (–) -gallocatechin gallate, and (–) -epigallocatechin-3-gallate (EGCG). EGCG is the most abundant catechin accounting for 50% of total polyphenols; thus, it is the main biological active compound of green tea [87]. However, polyphenols are not the only compounds that green tea exerts its antioxidant activity with through. The amino acid, L-theanine, in green tea accounts for 1–2% of the leaf dry weight that is synthesized in the roots of green tea and is concentrated in the leaves. Studies have reported that L-theanine protects the cell by maintaining its GSH levels in cancer and neurotoxic diseases [88]. The intake of green tea can be considered safe unless its consumption exceeds 1–2 cups a day. And higher consumption such as in attempts to lose weight resulted in hepatotoxicity [87]. At normal doses, Pérez-Vargas et al. [88] found that the main amino acid of green tea, L-theanine, reduced expression of NF-kB and downregulated IL-1 $\beta$  and IL-6 and the cytokines TGF- $\beta$  and CTGF. Haleboua-De Marzio et al. [89] tested a single high dose of green tea (400 mg), in patients with HCV-induced cirrhosis and found that it is well tolerated by patients and beneficial for treating cirrhosis.

### 1.11 L-Carnitine


L-Carnitine (LC), B-hydroxy-y-trimethylaminobutyric acid, is a water-soluble molecule important in mitochondrial oxidation of fatty acids in mammalian



metabolism (**Figure 4**). LC can exist in three different forms: as free LC, acetyl-L-carnitine (ALC), or other carnitine esters. About 25% of carnitine is obtained from methionine biosynthesis, but most LC is provided by the diet, especially through red meat and milk consumption [91]. LC acts as a carrier of fatty acids across the inner mitochondrial membrane for  $\beta$ -oxidation and ATP production. Apart from its role in the lipid metabolism, LC is also a potent antioxidant, and it protects tissues from oxidative damage. Reduced concentrations of LC in the body are mostly due to the accumulating toxic metabolites and also because of lack of protein in restricted diets. Thus, LC supplementation could be useful not only to supply the tissues in presence of but also in avoiding oxidative damage as a result of increased amounts of reactive species. Since LC can easily cross the blood-brain barrier, LC supplementation may also be beneficial in preventing oxidative injury-related neurological damage and further studies are needed in order to clearly establish LC's role in neurological diseases [92].

### 1.12 Lycopene

Lycopene (LYC) is an acyclic isomer of beta-carotene which has great antioxidant activities. It is synthesized by plants or autotrophic bacteria but not by animals. Red fruits and vegetables, such as tomatoes, watermelons, pink grapefruits, apricots, pink guavas, and papaya, contain LYC. Studies show that LYC consumption not only reduces the risk of cancer of many organs but also retards the growth of tumors. LYC has been shown to have protective effects on other pathologies such as cardiovascular diseases, osteoporosis, male infertility, and this action is mainly mediated by LC's ability to inhibit other toxic agents (**Figure 5**). Numerous *in vitro*



**Figure 4.** The mitochondrial carnitine system. Abbreviations: CPT I, carnitine-palmitoyl transferase I; CACT, carnitine acyl carnitine transferase; CPT II, carnitine-palmitoyl transferase II; CAT, carnitine acyl transferase; CoA, coenzyme A [90].




and *in vivo* studies showed that LYC could provide protection against ionizing radiation. Therefore, supplementation of LYC might be protective against damaging effects of radiotherapy in cancer treatments and it can also be protective against accidental radiation exposure [94].

1.13 Piperine

Piperine [1-[5-(1,3-benzodioxol-5-yl)-1-oxo-2,4, pentadienyl] piperidine] is the major pungent alkaloid present in the fruits of *Piper nigrum* L. [95]. Piperine at low concentrations acts as a hydroxyl radical scavenger, but at higher concentrations, it activates the Fenton reaction, resulting in increased generation of hydroxyl radicals. Piperine has hepatoprotective effects and it was shown to inhibit lipid peroxidation in the rat liver microsomes at a concentration of 600  $\mu$ M [96].

1.14 Capsaicin

Capsaicin (trans-8-methyl-N-vanillyl-6-nonenamide) is the major strong and irritating ingredient of red pepper. It may inhibit copper ion-induced lipid peroxidation of human LDL, which suggests that it is an effective antioxidant offering protection against oxidation of human LDL [97].



**Figure 5.** General mechanisms of action of lycopene. The proposed mechanisms of action of lycopene (oxidative and nonoxidative) that decreases the risk of oxidative stress-mediated diseases. Lycopene most likely acts via the oxidative mechanism of action to prevent oxidative stress and its detrimental effects on male infertility. ROS: reactive oxygen species [93].

### 1.15 Garlic and onion

Diallyl sulfides and diallyl disulfides, the active components of garlic, have anti-inflammatory and antimutagenic activities. Onion is a major source of flavonoids, especially the two quercetin glycosides, quercetin 4-*o*- $\beta$ -glucoside and quercetin 3,4-*o*- $\beta$ -diglucosides, which are recognized as bioactive substances. In order to show the antioxidant properties and protective effects of garlic and onion, a study was carried out on rats. Animals were treated with 0.6 mg nicotine/kg and also given 100 mg garlic or onion oils/kg for 21 days. Nicotine increased concentrations of thiobarbituric acid, conjugated dienes, and hydroperoxides in the tissues. Supplementation with both the garlic oil and onion oil increased resistance not only to lipid peroxidation but they also increased levels of antioxidant enzymes and glutathione. These conclusions state that oils of garlic and onion are effective antioxidants against nicotine-related oxidative stress and damage [98].

### 1.16 Vitamins C and E

Vitamin C, substrate for ascorbate peroxidase, is not only a highly effective antioxidant but also an essential component of a healthy diet. Vitamin E, the major antioxidant found in lipid composition of membranes, is a fat-soluble antioxidant. During fat oxidization, vitamin E helps to inhibit formation of ROS [99]. Several studies showed that vitamin E serum levels are significantly reduced in alcoholic liver disease [100]. It is also shown that vitamin E levels are inversely proportional to formation of oxidative stress products that correlate with the extent of liver damage [101]. For this reason, maintenance of normal concentrations of vitamin E appears to be necessary for preventing lipid peroxidation due to alcohol consumption. Works from several laboratories have so far indicated that mitochondrial damage may present a common early event in cell injury [102]. It is possible to prevent mitochondrial damage through vitamin E [103]. Vitamin E or C alone, or in combination, can ease scavenging free radicals that are generated in the liver tissue [104]. In the mouse model, vitamin E supplementation restores alcohol-induced redox status, reduces apoptosis, and prevents oxidative stress [105]. What is more, vitamin E is effective in doses of 600 mg daily when it comes to suppressing HBV replication and normalizing ALT in a significant proportion of chronically infected patients with CLD [106].

### 1.17 Trace minerals

Trace minerals act as a cofactor of antioxidant enzymes thus enabling the antioxidant activities to take place. These trace minerals include selenium (Se), zinc (Zn), manganese (Mn), iron (Fe), and copper (Cu) [103].  $O_2^-$  radicals are eliminated by the enzyme Cu-Zn-SOD and Cu and Zn are the co-factors for the enzyme. One of the enzymes responsible for  $H_2O_2$  clearance from the cells is CAT and Fe is the essential cofactor of this enzyme. Levels of ferritin may decline with exercise and increasing dietary or supplemental Fe can improve performance. It was shown that moderate-level supplementation of Fe to competitive swimmers increased their performance and helped to maintain normal ferritin levels [107]. Selenium (Se) is a cofactor for the antioxidant enzyme GPx, which is like the enzyme CAT, responsible for removing  $H_2O_2$  and other organic  $H_2O_2$  from the cell. A study by Akil et al. [108] showed in rats, that upon acute swimming exercise, lipid peroxidation in the brain was increased and Se supplementation to these rats increased antioxidant activity resulting in inhibition of the free radical production [108]. Manganese (Mn) is a cofactor for the enzyme Mn-SOD. It eliminates  $O_2^-$  radicals produced during oxidative phosphorylation [109].

## 2. Conclusion

Investigations done on antioxidants have shown that these compounds are candidates for the treatment and candidates to prevent oxidative stress-related diseases. This chapter focuses on antioxidants that can be investigated in experimental and clinical trials of many diseases but especially in diseases of liver. Main nutritional components involved in the production and/or removal of free radicals and the role of free radicals in the pathogenesis of several hepatic diseases and related comorbidities have been described in this chapter.

Among the antioxidants that were described, curcumin, naringenin, and quercetin have been found to be effective antioxidants in treatment of experimental liver injury. Green tea has been shown to protect against different kinds of cancer in clinical trials but not on hepatocellular carcinoma. Resveratrol has been extensively studied in experimental models of liver diseases and has been shown to have protective effects on fibrosis. So far, there are not much clinical trials on ameliorating and disease preventing effects of most potent antioxidants on liver and these antioxidants are good candidates for clinical trials not only because they show great disease preventing and ameliorating effects but also because they are derived from food sources and have a good metabolic tolerance.

### Author details

Sila Appak-Baskoy<sup>1</sup>, Mustafa Cengiz<sup>2\*</sup>, Ozgun Teksoy<sup>3</sup> and Adnan Ayhanci<sup>3</sup>

<sup>1</sup> Faculty of Science, Ryerson University, Toronto, ON, Canada

<sup>2</sup> Faculty of Education, Department of Elementary Education, Siirt University, Siirt, Turkey

<sup>3</sup> Faculty of Arts and Science, Department of Biology, Eskisehir Osmangazi University, Eskisehir, Turkey

\*Address all correspondence to: mustafacengizogu@gmail.com

### IntechOpen

© 2019 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. 

## References

- [1] Dongiovanni P, Lanti C, Riso P, Valenti L. Nutritional therapy for nonalcoholic fatty liver disease. *Journal of Nutritional Biochemistry*. 2016;**29**:1-11
- [2] Muriel P. Some experimental models of liver damage. In: Sahu SC, editor. *Hepatotoxicity: From Genomics to In Vitro and In Vivo Models*. West Sussex, England: Wiley; 2007. pp. 119-137
- [3] Muriel P. Cytokines in liver diseases. In: Sahu SC, editor. *Hepatotoxicity: From Genomics to In Vitro and In Vivo Models*. West Sussex, England: Wiley; 2007. pp. 371-389
- [4] Muriel P, Arauz J. Coffee and liver health. In: Chu Y, editor. *Coffee Emerging Health Effects and Disease Prevention*. West Sussex, UK: IFT Press/Wiley-Blackwell; 2012. pp. 123-139
- [5] Reyes-Gordillo K, Segovia J, Shibayama M, et al. Curcumin prevents and reverses cirrhosis induced by bile duct obstruction or CCl<sub>4</sub> in rats: Role of TGF-beta modulation and oxidative stress. *Fundamental & Clinical Pharmacology*. 2008;**22**:417-427. DOI: 10.1111/j.1472-8206.2008.00611.x
- [6] Muriel P. Peroxidation of lipids and liver damage. In: Baskin SI, Salem H, editors. *Oxidants, Antioxidants and Free Radicals*. Washington D.C., USA: Taylor & Francis; 1997. pp. 237-357
- [7] Muriel P, Rivera-Espinoza Y. Beneficial drugs for liver diseases. *Journal of Applied Toxicology*. 2008;**28**:93-103. DOI: 10.1002/jat.1310
- [8] Betteridge DJ. What is oxidative stress? *Metabolism*. 2000;**49**:3-8. DOI: 10.1016/S0026-0495(00)80077-3
- [9] Yoshikawa T, Naito Y. What is oxidative stress? *JMAJ*. 2002;**45**:271-276
- [10] Ternay JAL, Sorokin V. Redox, radicals, and antioxidants. In: Baskin SI, Salem H, editors. *Oxidants, Antioxidants, and Free Radicals*. Washington DC, USA: Taylor and Francis; 1997. p. 121
- [11] Ji LL, Gomez-Cabrera MC, Vina J. Exercise and hormesis: Activation of cellular antioxidant signaling pathway. *Annals of the New York Academy of Sciences*. 2006;**1067**(1):425-435
- [12] Ceni E, Mello T, Galli A. Pathogenesis of alcoholic liver disease: Role of oxidative metabolism. *World Journal of Gastroenterology*. 2014;**20**(47):17756-17772
- [13] Mello T, Ceni E, Surrenti C, Galli A. Alcohol induced hepatic fibrosis: Role of acetaldehyde. *Molecular Aspects of Medicine*. 2008;**29**(1-2):17-21
- [14] Zambo V, Simon-Szabo L, Szelenyi P, Kereszturi E, Banhegyi G, Csala M. Lipotoxicity in the liver. *World Journal of Hepatology*. 2013;**5**(10):550-557
- [15] Mello T, Zanieri F, Ceni E, Galli A. Oxidative stress in the healthy and wounded hepatocyte: A cellular organelles perspective. *Oxidative Medicine and Cellular Longevity*. 2016;**2016**:8327410
- [16] Cadenas E. Biochemistry of oxygen toxicity. *Annual Review of Biochemistry*. 1989;**58**:79-110. DOI: 10.1146/annurev.bi.58.070189.000455
- [17] Valko M, Rhodes CJ, Moncol J, Izakovic M, Mazur M. Free radicals, metals and antioxidants in oxidative stress-induced cancer. *Chemico-Biological Interactions*. 2006;**160**:1-40. DOI: 10.1016/j.cbi.2005.12.009
- [18] Inoue M, Sato EF, Nishikawa M, Park AM, Kira Y, Imada I, et al.



Mitochondrial generation of reactive oxygen species and its role in aerobic life. *Current Medicinal Chemistry*. 2003;**10**:2495-2505. DOI: 10.2174/0929867033456477

[19] Conner EM, Grisham MB. Inflammation, free radicals, and antioxidants. *Nutrition*. 1996;**12**:274-277. DOI: 10.1016/S0899-9007(96)00000-8

[20] Carr AC, McCall MR, Frei B. Oxidation of LDL by myeloperoxidase and reactive nitrogen species: Reaction pathways and antioxidant protection. *Arteriosclerosis, Thrombosis, and Vascular Biology*. 2000;**20**:1716-1723. DOI: 10.1016/01.atv.20.7.1716

[21] Halliwell B. How to characterize a biological antioxidant. *Free Radical Research Communications*. 1990;**9**:1-32. DOI: 10.3109/10715769009148569

[22] Maheshwari RK, Singh AK, Gaddipati J, Srimal RC. Multiple biological activities of curcumin: A short review. *Life Sciences*. 2006;**78**:2081-2087. DOI: 10.1016/j.lfs.2005.12.007

[23] Arcaro CA, Gutierrez VO, Assis RP, et al. Piperine, a natural bioenhancer, nullifies the antidiabetic and antioxidant activities of curcumin in streptozotocin-diabetic rats. *PLoS One*. 2014;**9**:e113993. DOI: 10.1371/journal.pone.0113993

[24] Sehgal A, Kumar M, Jain M, Dhawan DK. Piperine as an adjuvant increases the efficacy of curcumin in mitigating benzo(a)pyrene toxicity. *Human & Experimental Toxicology*. 2012;**31**:473-482. DOI: 10.1177/0960327111421943

[25] Matés JM, Segura JA, Alonso FJ, Márquez J. Natural antioxidants: Therapeutic prospects for cancer and neurological diseases. *Mini Reviews in*

*Medicinal Chemistry*. 2009;**9**:1202-1214. DOI: 10.2174/138955709789055180

[26] Reyes-Gordillo K, Segovia J, Shibayama M, Vergara P, Moreno MG, Muriel P. Curcumin protects against acute liver damage in the rat by inhibiting NF-kappaB, proinflammatory cytokines production and oxidative stress. *Biochimica et Biophysica Acta*. 2007;**1770**:989-996. DOI: 10.1016/j.bbagen.2007.02.004

[27] Jiao Y, Wilkinson J, Christine Pietsch E, et al. Iron chelation in the biological activity of curcumin. *Free Radical Biology & Medicine*. 2006;**40**:1152-1160. DOI: 10.1016/j.freeradbiomed.2005.11.003

[28] Charoensuk L, Pinlaor P, Prakobwong S, Hiraku Y, Laothong U, Ruangjirachuporn W, et al. Curcumin induces a nuclear factor-erythroid 2-related factor 2-driven response against oxidative and nitrative stress after praziquantel treatment in liver fluke-infected hamsters. *International Journal for Parasitology*. 2011;**41**:615-626. DOI: 10.1016/j.ijpara.2010.12.011

[29] Li B, Wang L, Lu Q, Da W. Liver injury attenuation by curcumin in a rat NASH model: An Nrf2 activation-mediated effect? *Irish Journal of Medical Science*. 2014;**185**:93-100. DOI: 10.1007/s11845-014-1226-9

[30] Reddy ACP, Lokesh BR. Alterations in lipid peroxides in rat liver by dietary n-3 fatty acids: Modulation of antioxidant enzymes by curcumin, eugenol, and vitamin E. *The Journal of Nutritional Biochemistry*. 1994;**5**:181-188. DOI: 10.1016/0955-2863(94)90070-1

[31] Hassan ZK, Al-Olayan EM. Curcumin reorganizes miRNA expression in a mouse model of liver fibrosis. *Asian Pacific Journal of Cancer Prevention*. 2012;**13**:5405-5408. DOI: 10.7314/apjcp.2012.13.11.5405

- [32] Lee HI, McGregor RA, Choi MS, et al. Low doses of curcumin protect alcohol-induced liver damage by modulation of the alcohol metabolic pathway, CYP2E1 and AMPK. *Life Sciences*. 2013;**93**:693-699. DOI: 10.1016/j.lfs.2013.09.014
- [33] Matés JM, Segura JA, Alonso FJ, Márquez J. Anticancer antioxidant regulatory functions of phytochemicals. *Current Medicinal Chemistry*. 2011;**18**:2315-2338. DOI: 10.2174/092986711795656036
- [34] Chávez E, Reyes-Gordillo K, Segovia J, et al. Resveratrol prevents fibrosis, NF-kappaB activation and TGF-beta increases induced by chronic CCl4 treatment in rats. *Journal of Applied Toxicology*. 2008;**28**:35-43. DOI: 10.1002/jat.1249
- [35] Walle T. Bioavailability of resveratrol. *Annals of the New York Academy of Sciences*. 2011;**1215**:9-15. DOI: 10.1111/j.1749-6632.2010.05842.x
- [36] Yu C, Shin YG, Chow A, et al. Human, rat, and mouse metabolism of resveratrol. *Pharmaceutical Research*. 2002;**19**:1907-1914
- [37] Wenzel E, Somoza V. Metabolism and bioavailability of transresveratrol. *Molecular Nutrition & Food Research*. 2005;**49**:472-481. DOI: 10.1002/mnfr.200500010
- [38] Sener G, Toklu HZ, Sehirli AO, Velioğlu-Oğünç A, Cetinel S, Gedik N. Protective effects of resveratrol against acetaminophen-induced toxicity in mice. *Hepatology Research*. 2006;**35**:62-68. DOI: 10.1016/j.hepres.2006.02.005
- [39] Rivera H, Shibayama M, Tsutsumi V, Perez-Alvarez V, Muriel P. Resveratrol and trimethylated resveratrol protect from acute liver damage induced by CCl4 in the rat. *Journal of Applied Toxicology*. 2008;**28**:147-155. DOI: 10.1002/jat.1260
- [40] Cheng AS, Cheng YH, Chiou CH, Chang TL. Resveratrol upregulates Nrf2 expression to attenuate methylglyoxal-induced insulin resistance in Hep G2 cells. *Journal of Agricultural and Food Chemistry*. 2012;**60**:9180-9187. DOI: 10.1021/jf302831d
- [41] Price NL, Gomes AP, Ling AJ, et al. SIRT1 is required for AMPK activation and the beneficial effects of resveratrol on mitochondrial function. *Cell Metabolism*. 2012;**15**:675-690. DOI: 10.1016/j.cmet.2012.04.003
- [42] Zhu W, Chen S, Li Z, et al. Effects and mechanisms of resveratrol on the amelioration of oxidative stress and hepatic steatosis in KKAY mice. *Nutrition and Metabolism*. 2014;**11**:35. DOI: 10.1186/1743-7075-11-35
- [43] Piver B, Berthou F, Dreano Y, Lucas D. Inhibition of CYP3A, CYP1A and CYP2E1 activities by resveratrol and other non volatile red wine components. *Toxicology Letters*. 2001;**125**:83-91. DOI: 10.1016/S0378-4274(01)00418-0
- [44] Wang Y, Jiang Y, Fan X, et al. Hepato-protective effect of resveratrol against acetaminophen-induced liver injury is associated with inhibition of CYP-mediated bioactivation and regulation of SIRT1-p53 signaling pathways. *Toxicology Letters*. 2015;**236**:82-89. DOI: 10.1016/j.toxlet.2015.05.001
- [45] Wu X, Li C, Xing G, Qi X, Ren J. Resveratrol Downregulates Cyp2e1 and Attenuates Chemically Induced Hepatocarcinogenesis in SD Rats. *Journal of Toxicologic Pathology*. 2013;**26**:385-392. DOI: 10.1293/tox.2013-0020
- [46] Faghihzadeh F, Adibi P, Rafiei R, Hekmatdoost A. Resveratrol supplementation improves inflammatory biomarkers in patients with nonalcoholic fatty liver disease.

Nutrition Research. 2014;**34**:837-843. DOI: 10.1016/j.nutres.2014.09.005

[47] Shin JW, Wang JH, Kang JK, Son CG. Experimental evidence for the protective effects of coffee against liver fibrosis in SD rats. *Journal of the Science of Food and Agriculture*. 2010;**90**:450-455. DOI: 10.1002/jsfa.3838

[48] McCusker RR, Goldberger BA, Cone EJ. Caffeine content of specialty coffees. *Journal of Analytical Toxicology*. 2003;**27**:520-522. DOI: 10.1093/jat/27.7.520

[49] Muriel P, Arauz J. Coffee and liver diseases. *Fitoterapia*. 2010;**81**:297-305. DOI: 10.1016/j.fitote.2009.10.003

[50] Arauz J, Moreno MG, Cortés-Reynosa P, Salazar EP, Muriel P. Coffee attenuates fibrosis by decreasing the expression of TGF- $\beta$  and CTGF in a murine model of liver damage. *Journal of Applied Toxicology*. 2013;**33**:970-979. DOI: 10.1002/jat.2788

[51] Ibrahim NK, Iftikhar R. Energy drinks: Getting wings but at what health cost? *Pakistan Journal of Medical Sciences*. 2014;**30**:1415-1419. DOI: 10.12669/pjms.306.5396

[52] Smith A. Effects of caffeine on human behavior. *Food and Chemical Toxicology*. 2002;**40**:1243-1255. DOI: 10.1016/S0278-6915(02)00096-0

[53] Gajewska M, Paini A, Sala Benito JV, et al. In vitro-to-in vivo correlation of the skin penetration, liver clearance and hepatotoxicity of caffeine. *Food and Chemical Toxicology*. 2015;**75**:39-49. DOI: 10.1016/j.fct.2014.10.017

[54] McCall AL, Millington WR, Wurtman RJ. Blood-brain barrier transport of caffeine: Dose-related restriction of adenine transport. *Life Sciences*. 1982;**31**:2709-2715. DOI: 10.1016/0024-3205(82)90715-9

[55] Klatsky AL, Armstrong MA. Alcohol, smoking, coffee, and cirrhosis. *American Journal of Epidemiology*. 1992;**136**:1248-1257

[56] Cavin C, Marin-Kuan M, Langouët S, et al. Induction of Nrf2-mediated cellular defenses and alteration of phase I activities as mechanisms of chemoprotective effects of coffee in the liver. *Food and Chemical Toxicology*. 2008;**46**:1239-1248. DOI: 10.1016/j.fct.2007.09.099

[57] Higgins LG, Cavin C, Itoh K, Yamamoto M, Hayes JD. Induction of cancer chemopreventive enzymes by coffee is mediated by transcription factor Nrf2. Evidence that the coffee-specific diterpenes cafestol and kahweol confer protection against acrolein. *Toxicology and Applied Pharmacology*. 2008;**226**:328-337. DOI: 10.1016/j.taap.2007.09.018

[58] Hertog MG, Hollman PC, Katan MB, Kromhout D. Intake of potentially anticarcinogenic flavonoids and their determinants in adults in The Netherlands. *Nutrition and Cancer*. 1993;**20**:21-29. DOI: 10.1080/01635589309514267

[59] Harwood M, Danielewska-Nikiel B, Borzelleca JF, Flamm GW, Williams GM, Lines TC. A critical review of the data related to the safety of quercetin and lack of evidence of in vivo toxicity, including lack of genotoxic/carcinogenic properties. *Food and Chemical Toxicology*. 2007;**45**:2179-2205. DOI: 10.1016/j.fct.2007.05.015

[60] Hernández-Ortega LD, Alcántar-Díaz BE, Ruiz-Corro LA, et al. Quercetin improves hepatic fibrosis reducing hepatic stellate cells and regulating pro-fibrogenic/anti-fibrogenic molecules balance. *Journal of Gastroenterology and Hepatology*. 2012;**27**:1865-1872. DOI: 10.1111/j.1440-1746.2012.07262.x



- [61] Bors W, Heller W, Michel C, Saran M. Radical chemistry of flavonoid antioxidants. In: Emerit I, Packer L, Auclair C, editors. *Antioxidants in Therapy and Preventive Medicine (Advances in Experimental Medicine and Biology)*. New York, USA: Plenum Press; 1990. pp. 165-170
- [62] Mira L, Fernandez MT, Santos M, Rocha R, Florêncio MH, Jennings KR. Interactions of flavonoids with iron and copper ions: A mechanism for their antioxidant activity. *Free Radical Research*. 2002;**36**:1199-1208. DOI: 10.1080/1071576021000016463
- [63] Murota K, Terao J. Quercetin appears in the lymph of unanesthetized rats as its phase II metabolites after administered into the stomach. *FEBS Letters*. 2005;**579**:5343-5346. DOI: 10.1016/j.febslet.2005.08.060
- [64] Han X, Pan J, Ren D, Cheng Y, Fan P, Lou H. Naringenin-7-O-glucoside protects against doxorubicin-induced toxicity in H9c2 cardiomyocytes by induction of endogenous antioxidant enzymes. *Food and Chemical Toxicology*. 2008;**46**:3140-3146. DOI: 10.1016/j.fct.2008.06.086
- [65] Pavanato A, Tuñón MJ, Sánchez-Campos S, et al. Effects of quercetin on liver damage in rats with carbon tetrachloride-induced cirrhosis. *Digestive Diseases and Sciences*. 2003;**48**:824-829. DOI: 10.1023/A:1022869716643
- [66] Granado-Serrano AB, Martín MA, Bravo L, Goya L, Ramos S. Quercetin modulates Nrf2 and glutathione-related defenses in HepG2 cells: Involvement of p38. *Chemico-Biological Interactions*. 2012;**195**:154-164. DOI: 10.1016/j.cbi.2011.12.005
- [67] Tang Y, Tian H, Shi Y, et al. Quercetin suppressed CYP2E1-dependent ethanol hepatotoxicity via depleting heme pool and releasing CO. *Phytomedicine*. 2013;**20**:699-704. DOI: 10.1016/j.phymed.2013.03.010
- [68] Surapaneni KM, Priya VV, Mallika J. Pioglitazone, quercetin and hydroxy citric acid effect on cytochrome P450 2E1 (CYP2E1) enzyme levels in experimentally induced non alcoholic steatohepatitis (NASH). *European Review for Medical and Pharmacological Sciences*. 2014;**18**:2736-2741
- [69] Dixit N, Baboota S, Kohli K, Ahmad S, Ali J. Silymarin: A review of pharmacological aspects and bioavailability enhancement approaches. *Indian Journal of Pharmacology*. 2007;**39**:172-179. DOI: 10.4103/0253-7613.36534
- [70] Saller R, Meier R, Brignoli R. The use of silymarin in the treatment of liver diseases. *Drugs*. 2001;**61**:2035-2020
- [71] Hutchinson C, Bomford A, Geissler CA. The iron-chelating potential of silybin in patients with hereditary haemochromatosis. *European Journal of Clinical Nutrition*. 2010;**64**:1239-1241. DOI: 10.1038/ejcn.2010.136
- [72] Najafzadeh H, Jalali MR, Morovvati H, Taravati F. Comparison of the prophylactic effect of silymarin and deferoxamine on iron overload-induced hepatotoxicity in rat. *Journal of Medical Toxicology*. 2010;**6**:22-26
- [73] Cengiz M, Kutlu HM, Burukoglu DD, Ayhancı A. A comparative study on the therapeutic effects of silymarin and silymarin-loaded solid lipid nanoparticles on d-GaIN/TNF- $\alpha$ -induced liver damage in balb/c mice. *Food and Chemical Toxicology*. 2015;**77**:93-100
- [74] Mata-Santos HA, Dutra FF, Rocha CC, et al. Silymarin reduces profibrogenic cytokines and reverses hepatic fibrosis in chronic murine schistosomiasis. *Antimicrobial Agents*



and Chemotherapy. 2014;**58**:2076-2083. DOI: 10.1128/AAC.01936-13

[75] Muriel P, Mourelle M. Prevention by silymarin of membrane alterations in acute CCl<sub>4</sub> liver damage. *Journal of Applied Toxicology*. 1990;**10**:275-279. DOI: 10.1002/jat.2550100408

[76] Muriel P, Moreno MG, Hernández Mdel C, Chávez E, Alcantar LK. Resolution of liver fibrosis in chronic CCl<sub>4</sub> administration in the rat after discontinuation of treatment: Effect of silymarin, silibinin, colchicine and trimethylcolchicinic acid. *Basic & Clinical Pharmacology & Toxicology*. 2005;**96**:375-380. DOI: 10.1111/j.1742-7843.2005.pto\_06.x

[77] Kim M, Yang SG, Kim JM, Lee JW, Kim YS, Lee JI. Silymarin suppresses hepatic stellate cell activation in a dietary rat model of non-alcoholic steatohepatitis: Analysis of isolated hepatic stellate cells. *International Journal of Molecular Medicine*. 2012;**30**:473-479. DOI: 10.3892/ijmm.2012.1029

[78] Tzeng JI, Chen MF, Chung HH, Cheng JT. Silymarin decreases connective tissue growth factor to improve liver fibrosis in rats treated with carbon tetrachloride. *Phytotherapy Research*. 2013;**27**:1023-1028. DOI: 10.1002/ptr.4829

[79] Kiruthiga PV, Pandian SK, Devi KP. Silymarin protects PBMC against B(a)P induced toxicity by replenishing redox status and modulating glutathione metabolizing enzymes—An in vitro study. *Toxicology and Applied Pharmacology*. 2010;**247**:116-128. DOI: 10.1016/j.taap.2010.06.004

[80] Vailati A, Aristia L, Sozze E, et al. Randomized open study of the dose-effect relationship of a short course of IdB 1016 (Silybin Phytosome) in patients with viral or alcoholic hepatitis. *Fitoterapia*. 1993;**94**:219-228

[81] Patel K, Singh GK, Patel DK. A review on pharmacological and analytical aspects of naringenin. *Chinese Journal of Integrative Medicine*. 2014. DOI: 1-13 DOI: 10.1007/s11655-014-1960-x

[82] Yang CP, Liu MH, Zou W, Guan XL, Lai L, Su WW. Toxicokinetics of naringin and its metabolite naringenin after 180-day repeated oral administration in beagle dogs assayed by a rapid resolution liquid chromatography/tandem mass spectrometric method. *Journal of Asian Natural Products Research*. 2012;**14**:68-75. DOI: 10.1080/10286020.2011.632369

[83] Chtourou Y, Fetoui H, Jemai R, Ben Slima A, Makni M, Gdoura R. Naringenin reduces cholesterol-induced hepatic inflammation in rats by modulating matrix metalloproteinases-2, 9 via inhibition of nuclear factor  $\kappa$ B pathway. *European Journal of Pharmacology*. 2015;**746**:96-105. DOI: 10.1016/j.ejphar.2014.10.027

[84] Yen FL, Wu TH, Lin LT, Cham TM, Lin CC. Naringenin-loaded nanoparticles improve the physicochemical properties and the hepatoprotective effects of naringenin in orally-administered rats with CCl<sub>4</sub>(4)-induced acute liver failure. *Pharmaceutical Research*. 2009;**26**:893-902. DOI: 10.1007/s11095-008-9791-0

[85] Esmaeili MA, Alilou M. Naringenin attenuates CCl<sub>4</sub> –induced hepatic inflammation by the activation of an Nrf2-mediated pathway in rats. *Clinical and Experimental Pharmacology & Physiology*. 2014;**41**:416-422. DOI: 10.1111/1440-1681.12230

[86] Motawi TK, Teleb ZA, El-Boghdady NA, Ibrahim SA. Effect of simvastatin and naringenin coadministration on rat liver DNA fragmentation and cytochrome P450 activity: An in vivo and in vitro study. *Journal of Physiology*

- and Biochemistry. 2014;**70**:225-237. DOI: 10.1007/s13105-103-0296-x
- [87] Mazzanti G, Sotto AD, Vitalone A. Hepatotoxicity of green tea: An update. *Archives of Toxicology*. 2015;**89**:1175-1191. DOI: 10.1007/s00204-015-1521-x
- [88] Pérez-Vargas JE, Zarco N, et al. l-Theanine prevents carbon tetrachloride-induced liver fibrosis via inhibition of nuclear factor  $\kappa$ B and downregulation of transforming growth factor  $\beta$  and connective tissue growth factor. *Human & Experimental Toxicology*. 2015;**35**:135-146. DOI: 10.1177/0960327115578864
- [89] Halegoua-De Marzio D, Kraft WK, Daskalakis C, Ying X, Hawke RL, Navarro VJ. Limited sampling estimates of epigallocatechin gallate exposures in cirrhotic and noncirrhotic patients with hepatitis C after single oral doses of green tea extract. *Clinical Therapeutics*. 2012;**34**:2279-2285.e1. DOI: 10.1016/j.clinthera.2012.10.009
- [90] Available from: <http://lpi.oregonstate.edu/mic/dietary-factors/L-carnitine>
- [91] Walter JH. L-Carnitine. *Archives of Disease in Childhood*. 1996;**74**:475-478
- [92] Ribas GS, Vargas CR, Wajner M. L-carnitine supplementation as a potential antioxidant therapy for inherited neurometabolic disorders. *Gene*. 2014;**533**:469-476
- [93] Durairajanayagam D, Agarwal A, Ong C, Prashast P. Lycopene and male infertility. *Asian Journal of Andrology*. 2014;**16**(3):420-425
- [94] Gajowik A, Dobrzyńska MM. Lycopene—antioxidant with radioprotective and anticancer properties. A review. *Roczniki Państwowego Zakładu Higieny*. 2014;**65**:263-271
- [95] Srinivasan K. Black pepper and its pungent principle-piperine: A review of diverse physiological effects: Critical Reviews in Food Science and Nutrition. *Critical Reviews in Food Science and Nutrition*. 2007;**47**:735-748. DOI: 10.1080/10408390601062054
- [96] Reddy AC, Lokesh BR. Studies on spice principles as antioxidants in the inhibition of lipid peroxidation of rat liver microsomes. *Molecular and Cellular Biochemistry*. 1992;**111**:117-124
- [97] Naidu KA, Thippeswamy NB. Inhibition of human low density lipoprotein oxidation by active principles from spices. *Molecular and Cellular Biochemistry*. 2002;**229**:19-23
- [98] Helen A, Rajasree CR, Krishnakumar K, Augusti KT, Vijayammal PL. Antioxidant role of oils isolated from garlic (*Allium sativum* Linn.) and onion (*Allium cepa* Linn.) on nicotine-induced lipid peroxidation. *Veterinary and Human Toxicology*. 1999;**41**:316-319
- [99] Herrera E, Barbas C. Vitamin E: Action, metabolism and perspectives. *Journal of Physiology and Biochemistry*. 2001;**57**:43-56
- [100] Bjørneboe GE, Johnsen J, Bjørneboe A, Bache-Wiig JE, Mørland J, Drevon CA. Diminished serum concentration of vitamin E in alcoholics. *Annals of Nutrition & Metabolism*. 1988;**32**:56-61
- [101] Masalkar PD, Abhang SA. Oxidative stress and antioxidant status in patients with alcoholic liver disease. *Clinica Chimica Acta*. 2005;**355**:61-65
- [102] Fulda S, Scaffidi C, Susin SA, Krammer PH, Kroemer G, Peter ME, et al. Activation of mitochondria and release of mitochondrial apoptogenic factors by betulinic acid. *The Journal of Biological Chemistry*. 1998;**273**:33942-33948

[103] Sakurai K, Cederbaum AI. Oxidative stress and cytotoxicity induced by ferric-nitrilotriacetate in HepG2 cells that Express cytochrome P450 2E1. *Molecular Pharmacology*. 1998;**54**:1024-1035

[104] Zaidi SM, Al-Qirim TM, Banu N. Effects of antioxidant vitamins on glutathione depletion and lipid peroxidation induced by restraint stress in the rat liver. *Drugs in R&D*. 2005;**6**:157-165

[105] Kaur J, Shalini S, Bansal MP. Influence of vitamin E on alcohol-induced changes in antioxidant defenses in mice liver. *Toxicology Mechanisms and Methods*. 2010;**20**:82-89

[106] Andreone P, Fiorino S, Cursaro C, et al. Vitamin E as treatment for chronic hepatitis B: Results of a randomized controlled pilot trial. *Antiviral Research*. 2001;**49**:75-81

[107] Powers SK, DeRuisseau KC, Quindry J, Hamilton KL. Dietary antioxidants and exercise. *Journal of Sports Sciences*. 2004;**22**:81-94

[108] Akil M, Bicer M, Menevse E, Baltaci AK, Mogulkoc R. Selenium supplementation prevents lipid peroxidation caused by arduous exercise in rat brain tissue. *Bratislavské Lekárske Listy*. 2011;**112**:314-317

[109] Zimmermann MB. Vitamin and mineral supplementation and exercise performance. *Schweizerische Zeitschrift für Medizin und Traumatologie*. 2003;**51**:53-57