

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

186,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.  
For more information visit [www.intechopen.com](http://www.intechopen.com)



# Excitability of Vascular Smooth Muscle

*Alexandra V. Ulyanova*

## Abstract

Regulation of pressure and local blood flow occurs at the level of resistance arteries and arterioles. Under physiological conditions, these small vessels exist in a state of partial constriction, termed myogenic tone. Myogenic tone is considered to be an intrinsic property of arteriolar smooth muscle cells, which membranes depolarize in response to increase in the intraluminal pressure. Oscillations of membrane potential in smooth muscles are mediated by the activity of voltage-gated L-type  $\text{Ca}^{2+}$  channels, which provide an influx of  $\text{Ca}^{2+}$  to activate various voltage-gated and  $\text{Ca}^{2+}$ -sensitive channels of smooth muscle cells and to initiate endothelial  $\text{Ca}^{2+}$  signaling needed for vasodilation. Although a relationship between change in membrane potential and myogenic response is considered to be universal throughout various smooth muscle tissues, it may be regulated differently based on autoregulatory responses and channels expression. Here we review electrophysiological signature of arteriolar smooth muscle in various tissues, with an emphasis and specific examples of the excitability of 4th order arterioles isolated from skeletal muscle.

**Keywords:** vasculature, arteriolar smooth muscle, excitability, electrophysiology, voltage-gated  $\text{Ca}^{2+}$  and  $\text{Na}^{+}$  channels

## 1. Introduction

Regulation of pressure and local blood flow occurs at the level of resistance arteries and arterioles. Once blood exits the heart, it first flows into large elastic arteries, followed by smaller distributing arteries, which branch further into small resistance arteries and, finally, arterioles. The branching and reduction in vessel diameter actually results in the increase in total cross-sectional area of circulation. Because of their small diameter, resistance arteries and arterioles are the place of the largest pressure drop.


## 2. Myogenic tone

Resistance arteries and arterioles typically exhibit a state of partial constriction termed myogenic tone [1]. Myogenic tone is related to the level of the intraluminal pressure and provides a level of tone that vasodilators can act upon [2]. When the intraluminal pressure increases, resistance arteries and arterioles first dilate due to their elastic properties and then constrict to the new steady-state level. Myogenic tone is an intrinsic property of arteriolar smooth muscle cells and does not require endothelium [3, 4]. Resistance to blood flow is actively controlled by contraction

or relaxation of arteriolar smooth muscle cells wrapped around the vessel so that their tone regulates the vessel diameter [2]. Arterial walls are made up of three layers: the tunica intima, tunica media and tunica adventitia. While the tunica intima contains endothelial cells and a thin layer of connective tissue, the tunica media supplies mechanical strength and contractile power. It is composed of several layers of spindle-shaped smooth muscle cells arranged helically in a matrix of elastin and collagen fibers. In some places, endothelial cells make contacts with smooth muscle cells to transmit signals between the intima and media. The tunica adventitia is mostly a connective tissue sheath with no distinct outer border. Its role is to tether the vessel loosely to the surrounding tissue [5]. Arterioles, the smallest resistance arteries placed right before capillaries, have a single layer of spindle-shaped smooth muscle cells [6]. Endothelial cells frame the lumen of arterioles. The shape and orientation within the vessels help to distinguish between these two distinct cells types (**Figure 1**).

Development of the tone is associated with depolarization of smooth muscle cells. While the mechanisms underlying depolarization are not completely understood, it is known that development of myogenic tone depends on extracellular  $\text{Ca}^{2+}$ . At 0  $\text{Ca}^{2+}$ , pressurized resistance arteries and arterioles are fully dilated. Adding  $\text{Ca}^{2+}$  up to  $\approx 2$  mM to the extracellular space causes maximal tone [7]. In addition to the development of basal tone, the myogenic mechanism is believed to underlie the response to acute changes in pressure and contribute to spontaneous vasomotor activity [7].

Myogenic tone is controlled by intrinsic as well as extrinsic mechanisms. Intrinsic mechanisms include constriction of arterioles in response to pressure increase (Bayliss effect), endothelial secretions (nitric oxide, EDHF, prostacyclin, endothelin), vasoactive metabolites (e.g. adenosine in exercising muscle), autacoids (local vasoactive paracrine secretions such as histamine), and temperature. Important physiological responses mediated entirely by intrinsic regulation include the autoregulation of flow, and functional and reactive hyperemia. Intrinsic regulation also contributes to pathological responses such as inflammation and arterial vasospasm. Extrinsic regulation is brought about by factors originating outside the organ, namely the vasomotor nerves (sympathetic, parasympathetic and others) and circulating hormones such as adrenaline, vasopressin and insulin (for review, see [8]).



**Figure 1.** 4th order skeletal muscle arteriole loaded with 10 mM Fluo-4. A single layer of spindle-like vascular smooth muscle cells (VSMC) runs perpendicular to the vessel's length (left panel). Long endothelial cells (EC) frame the lumen of skeletal muscle arteriole (right panel, adapted from [6]). Scale bar = 50  $\mu\text{m}$ .

### 3. Skeletal muscle vasculature

Skeletal muscle is the largest organ in the body that receives about 20% of cardiac output at rest and up to 80% during exercise. In skeletal muscle, the local blood flow is regulated over a 20× range to meet the demands of exercise. Therefore, vascular resistance of skeletal muscle is a critical determinant of total peripheral resistance and blood pressure. Arterioles of skeletal muscle have a high myogenic tone at rest as they are subject to major hyperemia (increased blood flow) during exercise [8]. In case of orthostatic hypotension, a potential contributor to cardiovascular adaptation to prolonged periods of bed rest or microgravity, skeletal muscle arterioles may develop functional or structural adaptations. Despite the physiological importance of skeletal muscle arterioles, little is known about ionic mechanisms underlying their excitability. Even within the same skeletal muscle, arterioles might respond differently to pressure changes. For instance, the first-order arterioles isolated from fast-twitch (e.g., white gastrocnemius) skeletal muscle fiber demonstrated both functional and structural changes such as reduced myogenic tone, decreased contractile responsiveness, and reduced wall thickness with no change in luminal diameter [9, 10]. In contrast, arterioles isolated from slow-twitch (e.g., soleus) fibers show no difference in myogenic tone or contractile responsiveness but rather a structural remodeling resulting in a decreased arteriole diameter [11].

### 4. Excitability of vascular smooth muscle

After development of isolated vessel techniques, the electrophysiology of smooth muscle cells was extensively studied in small arteries isolated from various vascular beds [12].

#### 4.1 Membrane potential of vascular smooth muscle

Both slow changes (myogenic tone) and fast spikes in the membrane potential have been observed in arteriolar smooth muscle cells. Arteriolar smooth muscle cells can generate rhythmical contractions (vasomotion) over an extended period of time. Vasomotion occurs spontaneously or in response to vasoactive stimulation with a frequency of 3–20 per minute. The exact physiological role of vasomotion is not clear. In cases of ischemia and hypertension, it serves as a protective mechanism. Vasomotion is not a consequence of heartbeat, respiration, or neuronal input. It is generated within arteriolar smooth muscle cells by an endogenous pacemaker mechanism driven by a cytosolic  $\text{Ca}^{2+}$  oscillator. The cytosolic  $\text{Ca}^{2+}$  oscillator depends on  $\text{Ca}^{2+}$  entry and is regulated by transmitters and hormones, which increase the formation of  $\text{InsP}_3$  and diacylglycerol (DAG) and promote oscillatory activity (for review, see [13]).


From the early studies, it has been appreciated that an increase in intraluminal pressure leads to depolarization and consequent contraction of vascular smooth muscle cell (electromechanical coupling). Mechanisms of depolarization are still not well understood. Depolarization increases  $\text{Ca}^{2+}$  concentration inside the cell, which leads to activation of myosin light chain kinase by  $\text{Ca}^{2+}$ /calmodulin and consequent contraction. Propagation of depolarization is achieved through gap junctions between neighboring smooth muscle cells. The signaling between endothelial and smooth muscle cells via gap junctions is essential for normal vascular function. Gap junctions in arteriolar smooth muscle cells and endothelial cells are formed by connexins Cx37, Cx40, and Cx43. Coupling between arteriolar smooth muscle cells appears to be essential for the maintenance of oscillations in membrane potential, the intracellular  $[\text{Ca}^{2+}]$ , and vasomotion.

The exact signaling mechanisms that underlie detection of the mechanical stimulus and membrane depolarization are not completely understood (for review, see [7, 14]). Depolarization of vascular smooth muscle's membrane activates various voltage-gated ion channels, pumps and exchangers, including voltage-gated  $\text{Ca}^{2+}$  and  $\text{K}^+$  channels,  $\text{Ca}^{2+}$ -activated  $\text{BK}_{\text{Ca}}$  and  $\text{Cl}_{\text{Ca}}$  channels,  $\text{Na}^+/\text{Ca}^{2+}$  exchanger,  $\text{Ca}^{2+}$ -ATP pump and  $\text{Na}^+/\text{K}^+$ -ATPase, ATP-sensitive P2X receptor, and transient receptor potential (TRP) ion channels. Voltage-gated L-type  $\text{Ca}^{2+}$  channels are activated upon depolarization and increase  $\text{Ca}^{2+}$  concentration inside the cell.  $\text{BK}_{\text{Ca}}$   $\text{K}^+$  channels hyperpolarize smooth muscle cell back to its resting potential [15, 16]. Some vascular myocytes, particularly in large vessels, contract via pathways that appear to be unrelated to significant changes in the membrane potential (pharmacomechanical coupling).

Membrane potential of arteriolar smooth muscle cells is difficult to measure because of the changes in the intraluminal pressure and active vasomotor responses. Resting membrane potentials range from approximately  $-60$  to  $-35$  mV for physiological pressures (for review, see [17, 18]). Resting membrane potentials between  $-80$  to  $-60$  mV were previously reported for un-pressurized mice mesenteric artery [19], submucosal arterioles [20], and cerebral arterioles [21, 22]. A range of membrane potentials was reported for un-pressurized mice skeletal muscle arterioles (**Figure 2**). The average resting potential was reported to be around  $-68$  mV (**Figure 2**, adapted from [6]). However, it's significantly more negative than  $-55$  mV previously reported for un-pressurized arterioles in rat cremaster muscle [23]. Resting membrane potential in vascular smooth muscle cells is determined to a large extent by  $\text{K}^+$  conductance [18]. Intracellular  $[\text{Cl}^-]$  in vascular smooth muscle cells is around 55 mM, which is unusually high and mediated probably through  $\text{Cl}^-/\text{HCO}_3^-$  exchanger. A small influx of  $\text{Ca}^{2+}$  and efflux of  $\text{Cl}^-$  ions at rest reduce the membrane potential from the Nernst equilibrium for  $\text{K}^+$  [24].


## 4.2 Excitability of vascular smooth muscle

Most of the arteriolar smooth muscle cells are quiescent. In some cases, a drop in the intraluminal pressure (during trauma to a vessel) that leads to hyperpolarization and dilation generates membrane action potentials. Physiological role of membrane action potentials in arteriolar smooth muscle is unclear, as relatively small changes in the membrane potential (between  $-55$  and  $-35$  mV) are sufficient to control  $\text{Ca}^{2+}$  entry and to initiate contraction in response to stimulation by mechanical stress of the blood flow [14]. Spontaneous action potentials could be associated either with rhythmic vasomotor activity [23]; follow  $\text{K}^+$ -induced hyperpolarization and dilation [25], or precede injury-induced constriction [26].



**Figure 2.** Resting membrane potential of un-pressurized 4th order arterioles isolated from mice skeletal muscle. Distribution of the resting membrane potential values was fitted by a single Gaussian function peaking at  $-77 \pm 2$  mV ( $n = 81$ , smooth line). The average resting potential was  $V_{\text{rest}} = -68 \pm 2$  mV ( $n = 81$ ). Adapted from [6].





**Figure 3.**  
*Spontaneous action potentials were observed in un-pressurized skeletal muscle arterioles (2 mM Ca bath solution). Smooth muscle cells were current-clamped using gramicidin-perforated configuration, with intracellular solution containing 150 KCl. Adapted from [6].*

Only a few recordings of action potentials in skeletal muscle arterioles were made so far [6, 23, 25, 27]. Spontaneous action potential spikes could be observed in pressurized small arteries [23, 25], as well as in un-pressurized arterioles as shown in **Figure 3** [6, 21].

### 4.3 Voltage-gated channels of arterial smooth muscle


Unlike in many other excitable tissues, action potentials in smooth muscle cells of arteries [28] and arterioles [21] are thought to be independent of voltage-gated  $\text{Na}^+$  channels, as they could not be blocked by the application of voltage-gated  $\text{Na}^+$  channel blocker tetrodotoxin (TTX). Depolarization of smooth muscle cells induced by the change in intraluminal pressure and/or tissue-stretch produces an increase in the intracellular  $[\text{Ca}^{2+}]$ , consequent myosin light chain phosphorylation and contraction (for review, see [14]). Since specific blockers of L-type voltage-gated  $\text{Ca}^{2+}$  channels suppress both the upstroke and the after-depolarizing components of action potentials, these channels are thought to be the main pathway for the depolarizing current. Nevertheless, several groups have demonstrated that voltage-gated  $\text{Na}^+$  channels are present in arterial beds and are activated upon membrane depolarization [6, 29].

#### 4.3.1 Voltage-gated $\text{Ca}^{2+}$ channels

Voltage-gated  $\text{Ca}^{2+}$  channels mediate influx of  $\text{Ca}^{2+}$  ions in response to membrane depolarization and regulate intracellular processes such as contraction, secretion, neurotransmission, and gene expression in many different cell types. Their activity is essential to couple electrical signals in the cell surface to physiological events in cells. Voltage-gated  $\text{Ca}^{2+}$  channels are comprised of the pore-forming  $\alpha_1$  subunit in complex with auxiliary subunits. A transmembrane disulfide-linked  $\alpha_2\delta$ , an intracellular  $\beta$ , and a  $\gamma$  subunit are components of most types of calcium channels. The  $\alpha_1$  subunit contains four repeats of a domain with six transmembrane segments, the fourth of which is the voltage-sensing S4 segment. The pore loop between transmembrane segments S5 and S6 in each domain determines ion conductance and selectivity (for review, see [30]). Ten distinct genes encode mammalian  $\alpha_1$  subunits of three subfamilies of voltage-gated  $\text{Ca}^{2+}$  channels. The amino acid sequences of these  $\alpha_1$  subunits are more than 70% identical within a subfamily but less than 40% identical among subfamilies. Voltage-gated  $\text{Ca}^{2+}$  channels are named using the chemical symbol of the principal permeating ion with the principal physiological regulator (voltage) indicated as a subscript according to the nomenclature developed for voltage-gated  $\text{K}^+$  channels [31]. The  $\text{Ca}_v1$  subfamily ( $\text{Ca}_v1.1$ – $\text{Ca}_v1.4$ ) represents high-voltage activated L-type  $\text{Ca}^{2+}$  channels, the  $\text{Ca}_v2$  subfamily ( $\text{Ca}_v2.1$ – $\text{Ca}_v2.3$ ) represents neuronal N-, P/Q-, and R-types  $\text{Ca}^{2+}$  channels, and  $\text{Ca}_v3$  subfamily ( $\text{Ca}_v3.1$ – $\text{Ca}_v3.3$ ) represents low-voltage activated T-type  $\text{Ca}^{2+}$  channels.

The pharmacology and biophysics of  $\text{Ca}^{2+}$  channels subfamilies are quite distinct. L-type  $\text{Ca}^{2+}$  calcium channels typically require a strong depolarization for activation and do not inactivate at positive potentials. They are the main pathway for  $\text{Ca}^{2+}$  currents recorded in muscle cells, where they initiate contraction and secretion. The  $\text{Ca}_v1$  subfamily is the molecular target of the organic  $\text{Ca}^{2+}$  channel blockers used widely in the therapy of cardiovascular diseases. L-type  $\text{Ca}^{2+}$  channels are blocked by the organic antagonists, including dihydropyridines (e.g., nifedipine), phenylalkylamines, and benzothiazepines. Dihydropyridines can be channel activators or inhibitors and therefore are thought to act allosterically to shift the channel toward the open or closed state rather than by occluding the pore. T-type  $\text{Ca}^{2+}$  channels are activated by weak depolarization and are transient at sustained depolarization. They are expressed in a wide variety of cell types, where they are involved in shaping the action potential and controlling patterns of repetitive firing. The  $\text{Ca}_v3$  subfamily of voltage-gated  $\text{Ca}^{2+}$  channels is insensitive to both the dihydropyridines that block  $\text{Ca}_v1$  channels and the spider and cone snail toxins that block the  $\text{Ca}_v2$  channels. There are no widely useful pharmacological agents that block T-type  $\text{Ca}^{2+}$  currents specifically. Mibefradil blocks both T-type  $\text{Ca}^{2+}$  channels and with less potency L-type  $\text{Ca}^{2+}$  channels. Currents through expressed  $\text{Ca}_v3.2$  channels could be selectively blocked by application of 40  $\mu\text{M}$  of  $\text{Ni}^{2+}$  [32].

Currents through dihydropyridine-sensitive  $\text{Ca}^{2+}$  channels were recorded in arteries of various vascular beds [21, 33–36], including from smooth muscle cells of skeletal muscle arterioles (**Figure 4**, left panel).  $\text{Ca}_v1.2$  considered to be the principal sub-type of voltage-gated  $\text{Ca}^{2+}$  channels involved in excitation-contraction coupling of vascular smooth muscle cells [37]. Since L-type  $\text{Ca}^{2+}$  blocker nifedipine did not eliminate basal tone in skeletal muscle arterioles, other  $\text{Ca}^{2+}$  entry mechanisms are believed to contribute to myogenic tone along with L-type  $\text{Ca}^{2+}$  channels [38]. In addition to L-type, vascular smooth muscle cells also express T-type  $\text{Ca}^{2+}$  channels (for review, see [30, 39]). Currents through T-type  $\text{Ca}^{2+}$  channels have been found in smooth muscle cells of arteries and arterioles of cerebral [21], mesenteric [40], renal [41], coronary [35], and skeletal [6] vascular beds as shown in **Figure 4** (right panel). Although the messenger RNAs for both,  $\text{Ca}_v3.1$  and  $\text{Ca}_v3.2$  T-type  $\text{Ca}^{2+}$  channels were found in rat cremaster arterioles [42], only L-type  $\text{Ca}^{2+}$  currents were recorded in single smooth muscle cells isolated from resistance arteries of hamster cremaster muscle [43]. The pressure/stretch stimulus initiates a depolarization that



**Figure 4.**


Voltage-gated T-type and L-type  $\text{Ca}^{2+}$  channels are present in arteriolar smooth muscle. Whole-cell  $\text{Ba}^{2+}$  currents were recorded in the presence of 1  $\mu\text{M}$  tetrodotoxin (TTX), voltage-gated  $\text{Na}^{+}$  channel blocker as described in more detail in [6]. Two kinetically different components were observed: while voltage steps from  $-30$  up to  $-10$  mV produced fast inactivating T-type  $\text{Ca}^{2+}$  current (left panel), further depolarization activated slowly inactivating L-type  $\text{Ca}^{2+}$  current (right panel). Adapted from [6].

causes  $\text{Ca}^{2+}$  influx through voltage-gated L-type  $\text{Ca}^{2+}$  channels and initiates  $\text{Ca}^{2+}$  sparklets [44–46]. Both,  $\text{Ca}^{2+}$  sparklets and  $\text{Ca}^{2+}$  sparks ( $\text{Ca}^{2+}$  release from intracellular stores) signaling events, activate negative feedback mechanisms via  $\text{Ca}^{2+}$ -dependent  $\text{K}^+$  currents thus preventing unrestrained depolarization [47–49]. While  $\text{Ca}^{2+}$  influx through L-type  $\text{Ca}^{2+}$  channels are believed to be involved in contraction of vascular smooth muscle, T-type  $\text{Ca}^{2+}$  channels, particularly  $\text{Ca}_v3.2$  appear to be mostly involved in relaxation of coronary arteries, acting through activation of  $\text{BK}_{\text{Ca}}$  channels [40–43, 50, 51].

#### 4.3.2 Voltage-gated $\text{Na}^+$ channels

In addition to voltage-gated  $\text{Ca}^{2+}$  channels, significant TTX-sensitive  $\text{Na}^+$  currents were found in smooth muscle cells from rabbit main pulmonary artery [52], human aorta [53], murine mesenteric arteries [29], and skeletal muscle arterioles [6] as shown in **Figure 5**. The above observations suggest that more complex mechanisms are likely to generate action potential in vascular smooth muscle cells. When voltage-gated  $\text{Ca}^{2+}$  and  $\text{Na}^+$  channels coexist in the same cell,  $\text{Na}^+$  conductance is thought to be responsible for generation and propagation of action potential in excitable cells such as neurons, striated muscle, and neuroendocrine cells. It has lower threshold and faster rise time than  $\text{Ca}^{2+}$  conductance. After  $\text{Na}^+$  channels rapidly inactivate, L-type voltage-gated  $\text{Ca}^{2+}$  channels open and further depolarize the cell, thus prolonging the plateau of an action potential. In contrast, T-type voltage-gated  $\text{Ca}^{2+}$  channels may open at resting potential and produce depolarizing current that brings the cell to threshold for  $\text{Na}^+$  spike [54, 55].

Voltage-gated  $\text{Na}^+$  channels consist of the subunit associated with auxiliary subunits. The pore-forming subunit is sufficient for functional expression, but the kinetics and voltage dependence of channel gating are modified by the subunits. These auxiliary subunits are involved in channel localization and interaction with cell adhesion molecules, extracellular matrix, and intracellular cytoskeleton. Similar to the voltage-gated  $\text{Ca}^{2+}$  channels, the subunits of  $\text{Na}^+$  channels are organized in four homologous domains, each composed of six transmembrane segments. S4 segment is a voltage sensor, and a pore loop located between the S5 and S6 segments determines ion conductance and selectivity (for review, see [56]).



**Figure 5.** TTX-sensitive voltage-gated  $\text{Na}^+$  channels are present in arteriolar smooth muscle. In arteriolar smooth muscle of skeletal muscle arterioles, voltage steps produced inward currents with at least two kinetically distinct components in 2 mM  $\text{Ca}$  solution (left panel). The fast component was through voltage-gated  $\text{Na}^+$  channels as it was blocked by application of 1  $\mu\text{M}$  TTX (right panel). Adapted from [6].



According to the nomenclature, the  $\text{Na}_v1$  superfamily ( $\text{Na}_v1.1$ – $\text{Na}_v1.9$ ) represents voltage-gated  $\text{Na}^+$  channels. Unlike the different classes of voltage-gated  $\text{Ca}^{2+}$  channels, the functional properties of  $\text{Na}^+$  channels are relatively similar. Amino acid sequences of the nine mammalian  $\text{Na}^+$  channel isoforms are more than 50% identical to each other, but separate families are difficult to define. By this criterion, nine isoforms are considered to be members of one  $\text{Na}_v1$  gene subfamily. Some voltage-gated  $\text{Na}^+$  channels ( $\text{Na}_v1.1$ ,  $\text{Na}_v1.2$ ,  $\text{Na}_v1.3$ ,  $\text{Na}_v1.4$ ,  $\text{Na}_v1.6$ , and  $\text{Na}_v1.7$ ) could be blocked by tetrodotoxin that binds to the extracellular side of the pore.

Several isoforms of TTX-sensitive voltage-gated channels have been found previously smooth muscle cells of in vasa recta ( $\text{Na}_v1.3$ ), portal vein ( $\text{Na}_v1.6$  and  $\text{Na}_v1.8$ ), and vas deference ( $\text{Na}_v1.6$ ) [57–59]. The activity of these voltage-gated  $\text{Na}^+$  channels may be tightly controlled by elevations of intracellular  $\text{Ca}^{2+}$ , potentially suppressing activity of  $\text{Na}^+$  channels [6]. For instance,  $\text{Ca}^{2+}$ /CaM binding was shown to down-regulates skeletal muscle isoform  $\text{Na}_v1.4$  by shifting its steady-state inactivation curve in the hyperpolarizing direction [60].  $\text{Na}_v1.3$  channels in descending vasa recta are suppressed by calmodulin inhibitors, while elevation of the intracellular  $[\text{Ca}^{2+}]$  shifts the voltage-dependence of their activation to more positive voltages [57]. In addition,  $\text{Ca}^{2+}$ -dependent down-regulation of  $\text{Na}^+$  channels can also occur via the PKC pathway since activation of PKC decreases peak sodium currents through brain  $\text{Na}_v1.2$  and skeletal muscle  $\text{Na}_v1.4$  channels by up to 80% [61, 62].

## 5. Conclusion

L-type voltage-gated  $\text{Ca}^{2+}$  channels are involved in excitation-contraction coupling of vascular smooth muscle cells. In addition, T-type  $\text{Ca}^{2+}$  channels were detected in arteriolar smooth muscle cells. These two types of voltage-gated  $\text{Ca}^{2+}$  channels together play an important role in the constriction/relaxation of smooth muscle cells by regulating  $\text{Ca}^{2+}$  signaling during myogenic tone. However, there are indications that other type of voltage-gated channels, specifically  $\text{Na}^+$  channels are present in various vascular beds. While the role of voltage-gated  $\text{Ca}^{2+}$  channels is well established, contribution of voltage-gated  $\text{Na}^+$  channel remains to be determined.

## Acknowledgements

I would like to thank Dr. Roman E. Shirokov for invaluable discussion related to this work.

## Conflict of interest

The author declares no conflict of interest.

## Abbreviations

DAG	diacylglycerol
EC	endothelial cell
EDHF	endothelium-derived hyperpolarizing factor
$\text{InsP}_3$	inositol 1,4,5-triphosphate
TTX	tetrodotoxin
VSMC	vascular smooth muscle cell

IntechOpen

IntechOpen

### **Author details**

Alexandra V. Ulyanova

Center for Brain Injury and Repair, Department of Neurosurgery, Perelman School of Medicine, University of Pennsylvania, Philadelphia, USA

\*Address all correspondence to: [ulyanova@pennmedicine.upenn.edu](mailto:ulyanova@pennmedicine.upenn.edu).

### **IntechOpen**

© 2019 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. 

## References

- [1] Bayliss N. On the local reactions of the arterial wall to changes of internal pressure. *The Journal of Physiology*. 1902;**28**:220-231
- [2] Mellander S, Johansson B. Control of resistance, exchange, and capacitance functions in the peripheral circulation. *Pharmacological Reviews*. 1968;**20**(3):117-196
- [3] Kuo L, Chilian WM, Davis MJ. Coronary arteriolar myogenic response is independent of endothelium. *Circulation Research*. 1990;**66**(3):860-866
- [4] Falcone JC, Davis MJ, Meininger GA. Endothelial independence of myogenic response in isolated skeletal muscle arterioles. *American Journal of Physiology. Heart and Circulatory Physiology*. 1991;**260**(1):H130-H135
- [5] Greensmith JE, Duling BR. Morphology of the constricted arteriolar wall: Physiological implications. *American Journal of Physiology. Heart and Circulatory Physiology*. 1984;**247**(5):H687-HH98
- [6] Ulyanova AV, Shirokov RE. Voltage-dependent inward currents in smooth muscle cells of skeletal muscle arterioles. *PLoS One*. 2018;**13**(4):e0194980
- [7] Davis MJ, Hill MA. Signaling mechanisms underlying the vascular myogenic response. *Physiological Reviews*. 1999;**79**(2):387-423
- [8] Herring N, Paterson DJ. *Levick's Introduction to Cardiovascular Physiology*. 6th Ed. Boca Raton, FL: CRC Press; 2018
- [9] Delp MD, Collieran PN, Wilkerson MK, McCurdy MR, Muller-Delp J. Structural and functional remodeling of skeletal muscle microvasculature is induced by simulated microgravity. *American Journal of Physiology*. *Heart and Circulatory Physiology*. 2000;**278**(6):H1866-H1873
- [10] Delp MD. Myogenic and vasoconstrictor responsiveness of skeletal muscle arterioles is diminished by hindlimb unloading. *Journal of Applied Physiology*. 1999;**86**(4):1178-1184
- [11] Heaps CL, Bowles DK. Nonuniform changes in arteriolar myogenic tone within skeletal muscle following hindlimb unweighting. *Journal of Applied Physiology*. 2002;**92**(3):1145-1151
- [12] Duling BR, Gore RW, Dacey RG Jr, Damon DN. Methods for isolation, cannulation, and in vitro study of single microvessels. *American Journal of Physiology. Heart and Circulatory Physiology*. 1981;**241**(1):H108-H116
- [13] Berridge MJ. Smooth muscle cell calcium activation mechanisms. *The Journal of Physiology*. 2008;**586**(21):5047-5061
- [14] Hill MA, Davis MJ, Meininger GA, Potocnik SJ, Murphy TV. Arteriolar myogenic signalling mechanisms: Implications for local vascular function. *Clinical Hemorheology and Microcirculation*. 2006;**34**(1):67-79
- [15] Brayden J, Nelson M. Regulation of arterial tone by activation of calcium-dependent potassium channels. *Science*. 1992;**256**(5056):532-535
- [16] Jaggar JH, Porter VA, Lederer WJ, Nelson MT. Calcium sparks in smooth muscle. *American Journal of Physiology. Cell Physiology*. 2000;**278**(2):C235-C256
- [17] Hirst GD, Edwards FR. Sympathetic neuroeffector transmission in arteries and arterioles. *Physiological Reviews*. 1989;**69**(2):546-604

- [18] Nelson MT, Patlak JB, Worley JF, Standen NB. Calcium channels, potassium channels, and voltage dependence of arterial smooth muscle tone. *American Journal of Physiology. Cell Physiology*. 1990;**259**(1):C3-C18
- [19] Kuriyama H, Suzuki H. Adrenergic transmissions in the guinea-pig mesenteric artery and their cholinergic modulations. *The Journal of Physiology*. 1981;**317**:383-396
- [20] Hirst GD, van Helden DF. Ionic basis of the resting potential of submucosal arterioles in the ileum of the guinea-pig. *The Journal of Physiology*. 1982;**333**:53-67
- [21] Hirst GD, Silverberg GD, van Helden DF. The action potential and underlying ionic currents in proximal rat middle cerebral arterioles. *The Journal of Physiology*. 1986;**371**:289-304
- [22] Knot HJ, Nelson MT. Regulation of arterial diameter and wall  $[Ca^{2+}]$  in cerebral arteries of rat by membrane potential and intravascular pressure. *The Journal of Physiology*. 1998;**508**(Pt 1):199-209
- [23] Kotecha N, Hill MA. Myogenic contraction in rat skeletal muscle arterioles: Smooth muscle membrane potential and  $Ca^{2+}$  signaling. *American Journal of Physiology. Heart and Circulatory Physiology*. 2005;**289**(4):H1326
- [24] Gerstheimer FP, Mühleisen M, Nehring D, Kreye VA. A chloride-bicarbonate exchanging anion carrier in vascular smooth muscle of the rabbit. *Pflügers Archiv*. 1987;**409**(1-2):60-66
- [25] Burns WR, Cohen KD, Jackson WF.  $K^+$ -induced dilation of hamster cremasteric arterioles involves both the  $Na^+/K^+$ -ATPase and inward-rectifier  $K^+$  channels. *Microcirculation*. 2004;**11**(3):279-293
- [26] Graham JM, Keatinge WR. Responses of inner and outer muscle of the sheep carotid artery to injury. *The Journal of Physiology*. 1975;**247**(2):473-482
- [27] Bartlett IS, Crane GJ, Neild TO, Segal SS. Electrophysiological basis of arteriolar vasomotion in vivo. *Journal of Vascular Research*. 2000;**37**(6):568-575
- [28] Keatinge WR. Sodium flux and electrical activity of arterial smooth muscle. *The Journal of Physiology*. 1968;**194**(1):183-200
- [29] Berra-Romani R, Blaustein MP, Matteson DR. TTX-sensitive voltage-gated  $Na^+$  channels are expressed in mesenteric artery smooth muscle cells. *American Journal of Physiology. Heart and Circulatory Physiology*. 2005;**289**(1):H137-H145
- [30] Catterall WA, Perez-Reyes E, Snutch TP, Striessnig J. International union of pharmacology. XLVIII. Nomenclature and structure-function relationships of voltage-gated calcium channels. *Pharmacological Reviews*. 2005;**57**(4):411-425
- [31] Chandy KG, Gutman GA. Nomenclature for mammalian potassium channel genes. *Trends in Pharmacological Sciences*. 1993;**14**(12):434
- [32] Lee J-H, Gomora JC, Cribbs LL, Perez-Reyes E. Nickel block of three cloned T-type calcium channels: Low concentrations selectively block  $\alpha_1H$ . *Biophysical Journal*. 1999;**77**(6):3034-3042
- [33] Aarons PI, Bolton TB, Lang RJ, MacKenzie I. Calcium currents in single isolated smooth muscle cells from the rabbit ear artery in normal-calcium and high-barium solutions. *The Journal of Physiology*. 1988;**405**(1):57-75
- [34] Bean B, Sturek M, Puga A, Hermsmeyer K. Calcium channels in muscle cells isolated from rat mesenteric



- arteries: Modulation by dihydropyridine drugs. *Circulation Research*. 1986;**59**(2):229-235
- [35] Ganitkevich VY, Isenberg G. Contribution of two types of calcium channels to membrane conductance of single myocytes from guinea-pig coronary artery. *The Journal of Physiology*. 1990;**426**(1):19-42
- [36] Smirnov SV, Aaronson PI.  $\text{Ca}^{2+}$  currents in single myocytes from human mesenteric arteries: Evidence for a physiological role of L-type channels. *The Journal of Physiology*. 1992;**457**(1):455-475
- [37] Koch W, Ellinor P, Schwartz A. cDNA cloning of a dihydropyridine-sensitive calcium channel from rat aorta. Evidence for the existence of alternatively spliced forms. *The Journal of Biological Chemistry*. 1990;**265**(29):17786-17791
- [38] Hill MA, Meininger GA. Calcium entry and myogenic phenomena in skeletal muscle arterioles. *American Journal of Physiology. Heart and Circulatory Physiology*. 1994;**267**(3):H1085-H1092
- [39] Perez-Reyes E. Molecular physiology of low voltage-activated T-type calcium channels. *Physiological Reviews*. 2003;**83**(1):117-161
- [40] Morita H, Cousins H, Onoue H, Ito Y, Inoue R. Predominant distribution of nifedipine-insensitive, high voltage-activated  $\text{Ca}^{2+}$  channels in the terminal mesenteric artery of guinea pig. *Circulation Research*. 1999;**85**(7):596-605
- [41] Gordienko DV, Clausen C, Goligorsky MS. Ionic currents and endothelin signaling in smooth muscle cells from rat renal resistance arteries. *American Journal of Physiology. Renal Physiology*. 1994;**266**(2):F325-F341
- [42] VanBavel E, Sorop O, Andreasen D, Pfaffendorf M, Jensen BL. Role of T-type calcium channels in myogenic tone of skeletal muscle resistance arteries. *American Journal of Physiology. Heart and Circulatory Physiology*. 2002;**283**(6):H2239-H2243
- [43] Cohen KD, Jackson WF. Hypoxia inhibits contraction but not calcium channel currents or changes in intracellular calcium in arteriolar muscle cells. *Microcirculation*. 2003;**10**(2):133-141
- [44] Amberg GC, Navedo MF, Nieves-Cintrón M, Molkentin JD, Santana LF. Calcium sparklets regulate local and global calcium in murine arterial smooth muscle. *Journal of Physiology (London)*. 2007;**579**(1):187-201
- [45] Navedo MF, Amberg GC, Nieves M, Molkentin JD, Santana LF. Mechanisms underlying heterogeneous  $\text{Ca}^{2+}$  sparklet activity in arterial smooth muscle. *The Journal of General Physiology*. 2006;**127**(6):611-622
- [46] Navedo MF, Amberg GC, Votaw VS, Santana LF. Constitutively active L-type  $\text{Ca}^{2+}$  channels. *PNAS*. 2005;**102**(31):11112-11117
- [47] Jaggar JH, Wellman GC, Heppner TJ, Porter VA, Perez GJ, Gollasch M, et al.  $\text{Ca}^{2+}$  channels, ryanodine receptors and  $\text{Ca}^{2+}$ -activated  $\text{K}^{+}$  channels: A functional unit for regulating arterial tone. *Acta Physiologica Scandinavica*. 2008;**164**(4):577-587
- [48] Ledoux J, Werner ME, Brayden JE, Nelson MT. Calcium-activated potassium channels and the regulation of vascular tone. *Physiology*. 2006;**21**(1):69-78
- [49] Nelson MT, Cheng H, Rubart M, Santana LF, Bonev AD, Knot HJ, et al. Relaxation of arterial smooth muscle by calcium sparks. *Science*. 1995;**270**(5236):633
- [50] Chen CC, Lamping KG, Nuno DW, Barresi R, Prouty SJ, Lavoie JL, et al.

- Abnormal coronary function in mice deficient in  $\alpha_1H$  T-type  $Ca^{2+}$  channels. *Science*. 2003;**302**(5649):1416-1418
- [51] Morita H, Shi J, Ito Y, Inoue R. T-channel-like pharmacological properties of high voltage-activated, nifedipine-insensitive  $Ca^{2+}$  currents in the rat terminal mesenteric artery. *British Journal of Pharmacology*. 2002;**137**(4):467-476
- [52] Okabe K, Kitamura K, Kuriyama H. The existence of a highly tetrodotoxin sensitive Na channel in freshly dispersed smooth muscle cells of the rabbit main pulmonary artery. *Pflügers Archiv—European Journal of Physiology*. 1988;**411**(4):423-428
- [53] Cox RH, Zhou Z, Tulenko TN. Voltage-gated sodium channels in human aortic smooth muscle cells. *Journal of Vascular Research*. 1998;**35**(5):310-317
- [54] Llinás R, Yarom Y. Electrophysiology of mammalian inferior olivary neurones in vitro. Different types of voltage-dependent ionic conductances. *The Journal of Physiology*. 1981;**315**:549-567
- [55] Llinás R, Yarom Y. Properties and distribution of ionic conductances generating electroresponsiveness of mammalian inferior olivary neurones in vitro. *The Journal of Physiology*. 1981;**315**:569-584
- [56] Catterall WA, Goldin AL, Waxman SG. International union of pharmacology. XLVII. Nomenclature and structure-function relationships of voltage-gated sodium channels. *Pharmacological Reviews*. 2005;**57**(4):397
- [57] Lee-Kwon W, Goo JH, Zhang Z, Silldorff EP, Pallone TL. Vasa recta voltage-gated  $Na^+$  channel  $Na_v1.3$  is regulated by calmodulin. *American Journal of Physiology. Renal Physiology*. 2007;**292**(1):F404-F414
- [58] Saleh S, Yeung SYM, Prestwich S, Pucovsky V, Greenwood I. Electrophysiological and molecular identification of voltage-gated sodium channels in murine vascular myocytes. *The Journal of Physiology*. 2005;**568**(1):155-169
- [59] Zhu H-L, Shibata A, Inai T, Nomura M, Shibata Y, Brock JA, et al. Characterization of  $Na_v1.6$ -mediated  $Na^+$  currents in smooth muscle cells isolated from mouse vas deferens. *Journal of Cellular Physiology*. 2010;**223**(1):234-243
- [60] Biswas S, Deschenes I, DiSilvestre D, Tian Y, Halperin VL, Tomaselli GF. Calmodulin regulation of  $Na_v1.4$  current: Role of binding to the carboxyl terminus. *The Journal of General Physiology*. 2008;**131**(3):197-209
- [61] Numann R, Catterall WA, Scheuer T. Functional modulation of brain sodium channels by protein kinase C phosphorylation. *Science*. 1991;**254**(5028):115-118
- [62] Numann R, Hauschka S, Catterall W, Scheuer T. Modulation of skeletal muscle sodium channels in a satellite cell line by protein kinase C. *The Journal of Neuroscience*. 1994;**14**(7):4226-4236