

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

185,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Primary Prevention of Alzheimer's Disease (AD)

Ettore Bergamini and Gabriella Cavallini

Abstract

Alzheimer dementia (AD) is a complex, aging-associated disease whose effects on the brain (an organ made up by nonreplaceable cells) are devastating. Disease is not curable, but progress in pathobiology shows that intervention on aging can make primary prevention of AD feasible. According to the amyloid-cascade hypothesis, mechanisms of AD include: an age-related alteration of free radical metabolism in membranes, leading to a higher yield in the toxic A β 1-42 peptide and an overwhelming impact on the weaker repair mechanisms of the aging cells. The proposed intervention on aging with anti-AD effects includes a daily assumption of antioxidants (red wine polyphenols enriched with resveratrol), a reinforcement of membrane antioxidant defenses by the assumption of polyunsaturated fatty acids at the first meal after fasting, and an enhancement of cell repair function (at the proteasome and autophagy level by an intermittent feeding regimen and physical exercise plus the assumption of antilipolytic agents during time of fasting). The beneficial effects of diet and physical activity on the endogenous production of protective nerve growth factors are magnified by an enriched environment. Treatment has already been started on healthy individuals at a higher risk of AD in the city of Volterra.

Keywords: Alzheimer's disease, proteasome, autophagy, antioxidants, PUFAs, APP, cholesterol, antilipolytic drugs, calorie restriction, physical exercise, brain plasticity, nerve growth factors

1. Introduction

Like many other degenerative diseases, AD is an irreversible and progressive cerebropathy, the cause of one of the most common types of dementia affecting the elderly. It is a brain disorder characterized by the accumulation of two main protein aggregates, senile plaques and neurofibrillary tangles, leading to a progressive neuronal degeneration. It causes death after years of disability, progressive loss of memory, inability to perform normal daily activities, and, finally, dementia. The senile plaques are generated by the deposition in human brain of fibrils of the β -amyloid peptide (A β), a fragment derived from the proteolytic processing of the amyloid precursor protein (APP). There is evidence that oxidative stress might be the main factor that turns APP into a proteolytically processable substrate. The neurofibrillary tangles (NFTs) are seen as a compact filamentous network formed by paired helical filaments (PHFs), whose major component is a hyperphosphorylated Tau protein. Two main protein kinases appear to be involved in the anomalous tau phosphorylation: the cyclin-dependent kinase Cdk5 and glycogen synthase kinase GSK3. Dysfunction of the ubiquitin-proteasome system may be the cause [1]

possibly together with a secondary failure of the engulfed lysosomal degradation resulting in apoptosis. Pathology is a consequence of the extensive neuronal death, and when it manifests clinically, it is already incurable. Human, social, and health costs of this incurable disease are immense. It is highly desirable indeed an effective primary prevention to postpone the appearance of the debilitating manifestations, hopefully to the time of death.

2. Biological aging: Is it a major risk or the basic causative factor for neurodegeneration?

Age is the main factor in major debilitating and life-threatening conditions, including cancer, cardiovascular disease, diabetes, and neurodegeneration, all of which are therefore increasing in prevalence [2]. Nowadays almost all can make it to the old and nobody can say anymore that reaching old age is a fortune, and between those who should solve social and health problems of older people prevails the old slogan “old age is not a disease.” For many scholars, however, modern scientific discoveries confirm the deductions of ancient philosophers: all living beings, without exception, would be affected by an innate chronic degenerative disease, we call it aging, characterized by having an incubation period as long as be compatible with the reproductive success of the species. If aging is a disease and not a simple disability and disease risk factor, it is not surprising that the diseases associated with it (cancer, neurodegeneration, atherosclerosis, diabetes, etc.) can be regarded as signs or easily preventable complications, all together, by fighting the underlying disease. The prescription is simple but hard to follow a sober physically active lifestyle; to eat fruits, vegetables, fish, and little red meat; to take food after having been hungry for a few hours; to take supplements rich in polyunsaturated fatty acids (PUFAs) after dinner regularly; and to make good use of all functions of all organs of our body, including brain. This is exactly what cardiologists, diabetologists, oncologists, and neurologists all recommend for primary prevention. Understanding exactly the causes of aging and of all age-associated diseases may help to tackle the growing problem of neurodegeneration. Free radicals are the root of all evil. They may be generated either by endogenous (metabolic) causes or by environmental factors, including pollution (living less than 50 m from a major traffic road may increase hazards ratio of incident dementia by a 10% [3] and even oral hygiene and chronic inflammation [4]). Perhaps, we should remind here the oxygen paradox: without oxygen we die in minutes; with oxygen we grow old and die [5]. Oxygen is actually slowly poisonous, and it just takes 75–100 years to kill us, difference depending on how much we use and how we deal with it and repair the endogenous oxidative damage to protein, lipids, and (most important) DNA responsible for intrinsic aging, as well as the additional free radical-mediated damage from the inflammatory responses and environmental factors (e.g., ionizing radiation) [6]. It was computed that the oxygen consumption of human brain may be higher than 3 mL (i.e., about 1020 oxygen molecules)/g/min. In humans, over 99% of these molecules do generate water safely, but 10^{18} per min will produce free radicals, approximately 10^6 free radicals per cell per min. It was estimated that the number of oxidative hits to DNA in the human cell per day is about 10,000 and that DNA-repair enzymes efficiently remove 99.9% of the lesions formed so that only one oxidative lesion accumulate in the DNA of any cell every day. This is not nothing: it makes over 30,000 lesions in a long life [7]. Since there are about 20,000 genes in human cells, by the age of 100 years, all neurons may carry about two mutations a gene on the average. Why be surprised if over time cell functions are reduced?

3. Pathology of AD

With AD patients, the brain is smaller than normal and of reduced weight. A reduction of the thickness of the convolutions is evident. Atrophy is more evident in the temporal lobe, particularly in the parahippocampus, but also in the frontal and parietal regions. The occipital lobe and the motor cortex may be spared. Histologically, several major changes are recognized in AD. Amyloid, consisting of accumulations of A β peptide, is deposited in the cerebral cortex in the form of spherical deposits called senile plaques. Intraneuronal inclusions are formed in the cortical neurons, constituted by abnormal, often flame-shaped, bundles of filaments called neurofibrillary tangles, which occupy a large part of the neuronal cytoplasm, and are made up of a protein that binds to the microtubules, called tau protein. The processes of the cortical nerve cells diverge, twist, and dilate due to the accumulation of filaments in the form of tangles [8]. Changes are due to oxidative stress damage and to the relative failure of repair mechanisms at the molecular and subcellular (autophagy) level and result in the disruption of the neural network (**Table 1**). It is customary to distinguish two forms of AD. There is indeed a precocious rare form that occurs between 30 and 60 years, with a peak in the fifties, with a formation of the amyloidogenic peptide A β (1-42) from APP genetically favored by particular isoforms of presenilin 1 and 2 or of APP. The other form, sporadic, more frequent, late-onset (observed after age 65 with a frequency that increases with age) is due to the progressive increase in oxidative stress and decline with increasing age in mitochondrial and peroxisomal maintenance and in the efficiency of the mechanisms that neutralize free radicals and repair damage. This latter form might be postponed successfully to time of death by anti-aging interventions (nutrition, physical activity, damage repair, nerve growth factors).

4. The primary cause of AD and the roles of cholesterol and unsaturated fatty acids

It is obvious that native APP cannot be the ultimate substrate for γ -secretase trimming. Hence, both with the earlier and the later form of AD, the rate of A β

Mechanism of repair	Effects of the age-related decline
Molecular level	
DNA repair	Accumulation of DNA lesion
Proteasome	Accumulation of altered proteins
Phospholipid repair	Changes in polyunsaturated fatty acids
Subcellular Level	
Autophagy and lysosomal function	Accumulation of altered mitochondria Accumulation of protein aggregates Changes in membrane proteins and lipids
Cellular and tissue level	
Apoptosis	Accumulation of damaged, misfunctional cells in all tissues
<i>There is evidence that the above-mentioned repair mechanisms are responsible for cleaning cells from any produced waste and for getting tissues rid of irreversibly altered cells. In younger persons, functions are redundant but progressively decline with increasing age, and may gradually fail in older persons resulting in the accumulation of "waste" in cells and tissues. On a healthy life, "waste" is the limiting factor for cell and tissue "cleaning" activities, but waste recognition-acuity co-varies with cell and body request for nutrients and repair. Function of all repair mechanisms are inducible (or suppressible) depending on life style. Benefits from healthy life style, diet restriction, and physical activity depend at least in part from the induction of repair mechanisms.</i>	

Table 1.
Effects of failure of repair mechanisms at the molecular, subcellular, and cellular level.

production from APP should depend on a higher production of the ultimate substrate, possibly by a posttranslational modification of the APP molecules, likely to be secondary to oxidative stress. Age-related changes in the machinery that protects membrane proteins from free radical-mediated injury might help to account for age dependency [9]. With regard to the higher production of A β (1-42) in AD patients, a displacement of the free radical-mediated attack from the wanted site in the APP molecule might account for the higher involvement of the β and γ secretase pathways and the higher yield in A β 1-42. The effect may be favored either by genetic factors and/or aging.

4.1 Role of cholesterol

Elevated cholesterol levels may be associated with a higher risk of AD [10]. Evidence was found suggesting an intimate connection between APP processing and lipid rafts [11]. An age-related increase in cholesterol and oxidized cholesterol products (namely 24-hydroxycholesterol and 27-hydroxycholesterol) was shown indeed to be increasingly associated with AD progression (brain levels are higher in AD patients, and levels of 24-hydroxycholesterol, 27-hydroxycholesterol and cholesterol in the cerebrospinal fluid appear to be useful biomarkers for the evaluation of mild cognitive impairment (MCI) and AD, together with A β 42, total tau, and phospho-tau) [12]. It has been shown recently that higher total cholesterol levels in the blood are observed long before the clinical manifestation of MCI and AD in patients without psychiatric or somatic comorbidities and are independent of APOE genotype [13]. However, evidence was produced that changes in cholesterol metabolism in AD may not be the primary cause of the disease (see below): they may simply be a tightly associated sign with A β production by sharing a common cause (a higher intramembrane oxidative stress) (see below).

4.2 Role of polyunsaturated fatty acids (PUFAs)

PUFAs and their oxylipins may affect the onset of AD [14]. The administration of omega-3 fatty acids may cause a dose-dependent reduction of triglyceridemia and cholesterolemia and exert an antiatherogenic effect [15]. With regard to mechanisms, in view of the proposed protective role of unsaturated fatty acids in phospholipids against the free radical-mediated injury of membrane proteins [9], it should be mentioned that the distribution of unsaturation (the trap for unpaired electrons) across the membrane leaflets is not uniform, and minima (the least protected areas from oxidative stress) were observed close to the C-6 site (i.e., very close to the membrane exterior, to the phosphorylatable site of HMGCoAR, and to the vulnerable site of APP) and at the C-15 and C-17 levels (closer to the free radical conductor dolichol) [16]. Quite interestingly, signal might help focus free radical-mediated injury on the right target; more interestingly, in ad-libitum-fed (shorter-lived) rats, is its recognizability that may fade on aging: the abundance of double bonds near the C-6 site (but not at the C-15 and C-17) indeed appears to increase up to a doubling by age 24 months. Furthermore, this age-related change is prevented in part by nutritional anti-aging intervention [17]. Perhaps age-related changes in the production of A β 1-42, in the activity of HMGCoA reductase, and in the PUFA content of membrane phospholipids are all somehow bound together and involved all in the risk of AD. As an additional comment, beneficial intervention on A β 1-42 production may require the administration of antioxidants (e.g., polyphenols and resveratrol) to curb oxidative stress and of omega-3 fatty acids at a high dosage at the first good meal after fast (on the anabolic phase of metabolism) to counteract age-related changes in phospholipid unsaturated fatty acids and increase membrane

resistance to oxidative stress. An enhancement of the membrane turnover rate may be useful (e.g., by dietary restriction and/or pharmacological stimulation of autophagy) [18].

5. The pathogenesis of AD

The hypothesis of the amyloid cascade attributes to beta-amyloid, the responsibility of all cases of AD, and considers the tau pathology and other degenerative changes secondary to the A β pathology. Indeed, the extracellular accumulation of A β , a hallmark of AD, produces ROS, including hydrogen peroxide (H₂O₂) in the presence of Fe³⁺ or Cu²⁺ [19, 20]. The amyloid β -peptide responsible for AD is generated within the transmembrane domain (TMD) of a C-terminal fragment of the amyloid β protein-precursor (APP CTF β) by the proteolytic action of the γ -secretase complex [21]. Very interestingly, it was shown that some mutation(s) that promote destabilization of TMD helix might affect the length of the accumulated A β species and the ratio of the toxic A β 42 to the safer A β 40 peptide and may result in a young-onset AD [22]. Hence, the speculation may be invited that a free radical-induced modification in (some) amino acids of APP close to the C-6 carbon of phospholipid fatty acids might have a similar effect and enhance amyloid deposition. With regard to disease progression, many different active factors in sequence would determine timing of neuronal damage and symptom development, such as: (1) an overproduction of the toxic A β peptide (1-42) for genetic reasons and/or decrease in the quality of the antioxidant device of the membranes, (2) an ensuing increase in A β excretion in the oligomeric form may concentrate the redox-active copper at neuronal membranes before stacking in amyloid fibrils to form an ROS generating complex, (3) oxidative stress may be enhanced with an ensuing increase in lipid peroxidation and di-tyrosine formation, and (4) inflammation and activation of the microglia, which may further increase free radical generation, spread lytic enzymes, and cause cytotoxicity and anticipation of apoptotic neuronal death [23].

6. AD-associated changes in cholesterol metabolism

Abnormal cholesterol metabolism is an established feature of AD: levels of cholesterol are increased in MCI subjects and levels of 24-hydroxycholesterol and 27-hydroxycholesterol are elevated compared to controls both in AD and MCI subjects [24]. It appears that the rate of cholesterol production may be boosted by a free radical attack via a constitutive activation of HMGCoA-reductase, the rate-limiting step in sterol biosynthesis. Mechanism was clarified by Bergamini group in Pisa in cooperation with Trentalance group in Rome: in the rat, a free radical attack may prevent AMP-dependent protein kinase from phosphorylating a serine residue close to the C-terminus of HMGCoA reductase, Ser 872 with human enzyme [25, 26]. It has been proposed by several authors that higher cholesterol may disturb the lipid raft domains in various membrane organelles and affect the functioning of α , β , and γ secretases as well as APP itself and the production of A β 42; and that by converse, the toxicity of A β may be produced, in part, by disturbing the composition of the lipid raft domains in which they reside [27, 28]. Both with in vitro experiments and with animal models, statins strongly lowered blood cholesterol and reduced the levels of A β peptides, A β 42 and A β 40 [29, 30]. However, a statistically significant correlation between two events does not necessarily imply the existence of a causal link: an alternative explanation is that the events share a common cause. This might

be the case here: aging is known to lower antioxidant defenses, and any increase in oxidative stress either in vitro (e.g., by UV radiation of isolated rat liver cells) or in vivo (by chronic deprivation either of vitamin E or PUFA) may cause the constitutive increase in HMGCoA reductase and deregulate cholesterol synthesis [26, 31].

7. Alzheimer's disease: treatment or prevention strategies?

The pharmacological treatments for AD can be divided into two categories: symptomatic treatments such as acetylcholinesterase inhibitors and N-methyl-D-aspartate (NMDA) receptor antagonists and etiology-based treatments such as secretase inhibitors, amyloid binders, and tau therapies [32].

Despite significant investments in therapeutic drug discovery programs, no drugs to alter the course of disease have been found so far. Only four drugs with cholinergic (donepezil, galantamine, rivastigmine) or glutamatergic activity (Memantine) are currently approved and marketed for the treatment of AD-associated dementia, and their utility is very limited; several trials on inhibitors of γ -secretase and β -secretase have been discontinued; the available drug treatments of AD are merely symptomatic and unsatisfactory, and only minor benefits are obtained even if therapy is started at a very early time [33]. By the way, the physiological function of the amyloidogenic peptide has not been clarified yet, though it may be synchronized with life history [34].

In conclusion, no effective therapy is available to cure AD so far, and attention had to be shifted to the primary prevention of the disease. It was realized indeed that there is an extremely long, symptom-free prodromal phase in the path toward dementia in which deficits in synaptic density and plasticity are the principal alteration [35]. As an additional comment, better diagnostic tools and earlier diagnosis are needed (earlier timing is an important factor for the success rate of intervention), and novel strategies toward primary intervention may be wanted [36].

Problems in primary prevention were tackled recently. Qiu et al. [37] stressed the potential risk roles of vascular risk factors and disorders (e.g., cigarette smoking, midlife high blood pressure and obesity, diabetes, and cerebrovascular lesions) and the possible beneficial roles of psychosocial factors (e.g., high education, active social engagement, physical exercise, and mentally stimulating activity) in the pathogenetic process and clinical manifestation of the dementing disorders. Paillard-Borg et al. [38] showed that the participation in activities (mental, physical, or social activity) can retard the onset of dementia significantly (a 17 months' delay was seen in mean age at dementia onset between an inactive group and the most active group).

It may be worthwhile to remind here that the primary risk factor for AD is old age and that the prevalence of AD and other age-related dementias increases with increasing age [2]. It is very surprising, indeed, that little attention has been given so far to benefits from the most effective antiaging interventions (dietary restriction and physical activity) on the age of onset of neurodegeneration [39].

Antiaging diet restriction is known to be the most effective intervention that retards aging and extends lifespan and health span. Effects are known to involve the activation of macroautophagy, a cell repair mechanism [40] that can be intensified by the administration of antilipolytic agents during fasting to safely improve cell housekeeping and boost the benefits of caloric restriction [41]. Physical exercise, which is available at low cost and largely free of adverse effects, is another powerful antiaging strategy that can influence, at least partly, most of the hallmarks of biological aging [42]. It is known that greater levels of physical activity are associated with decreased risk of a future diagnosis of MCI or AD [43], extend longevity, and

reduce the risk of physical disability and may be an important adjunct to pharmacological treatment of AD [44].

Looking at mechanisms, it appears that dietary restriction and physical exercise share common neuroprotective mechanisms and should be included both in primary prevention of AD to increase the quality of nerve cells and oppose neurodegeneration and apoptosis (the third item in the **Table 1**, a repair mechanism harmful to brain) and give synergic support in a “train body and brain” program aimed to enhance neurotrophic antiapoptotic signals and defer or suppress neuron misfunctioning and death [45].

Antiaging diet restriction improves metabolism and promotes rejuvenation (by stimulating autophagy) of visceral organs that talk to brain via the vagus nerve and spinal afferent nerves [46]. Physical exercise improves metabolism and promotes rejuvenation of the lean body mass by the process of autophagy, which is very active in skeletal muscle and more intense when strenuous exercise is performed in the fasted state [47] and helps communication of the exercising muscle with the brain via increase in the discharge frequency of thinly myelinated (Group III) and unmyelinated (Group IV) nerve fibers [48]. Under these conditions, physical exercise is good both for physical health and mental health and abilities, and constitutes a practical neuroprotective strategy that provides a remarkable protection against brain insults of different etiology and anatomy [49].

Quite interesting, it has been clarified that dietary restriction and physical exercise share also common neuroprotective metabolic mechanism: the increased availability to brain of 3-hydroxybutyrate and an ensuing endogenous production of the brain-derived-neurotrophic-factor BDNF [50], and thus, both shelter the aging brain from memory loss and neurodegeneration, ameliorate mitochondrial function, and reduce the expression of apoptotic and inflammatory mediators [51]. As an additional evidence: both treatments safely modulate the endogenous production of BDNF, a neurotrophin that is vital to the survival, growth, and maintenance of neurons in key brain circuits involved in emotional and cognitive function [52, 53]. As an additional benefit, sustained levels of physical exercise together with dietary intervention may increase brain uptake of physiologically relevant neuroprotective trophic factors, such as IGF-I [49].

In conclusion, both (dietary and physical) interventions should be included in primary prevention of AD to increase quality of nerve cells and oppose neurodegeneration and apoptosis to implement reported programs aimed to enhance neurotrophic antiapoptotic signals and defer or suppress neuron misfunctioning and death.

8. The “train body and brain” protocol for a primary intervention on AD in the city of Volterra (Tuscany, Italy)

In view of the continuing increase in the prevalence of dementia to a magnitude growing to emergency [54, 55], a free of charge program to train people at higher risk of disease has been started in the Italian city of Volterra. Since AD is an aging-associated multifaceted disease that is hard to treat by single-modal treatment, a corresponding multifaceted preventive approach was included in a teaching program on how to counteract in practice the effects of biological and pathological aging on human body and brain.

Program includes activation of the global antioxidant defense system in order to attenuate the AD-causative oxidative stress, improvement of the function of the free radical conducting mechanism responsible for membrane resistance to oxidative stress, reinforcement of cell repair mechanisms at the molecular and subcellular

level by dietary and physical intervention, and targeted high-intensity training of cognitive brain functions to boost the induction of neurotrophic factors by the physiological mechanism. Here, a few details on protocol are given.

- a. Strengthening antioxidant defenses by nutritional intervention. It is said that the efficiency of antioxidant defenses may be boosted by eating the colors in the peel of fruits and vegetables that are known to contain complex mixtures of polyphenols and other phytochemicals. (e.g., polyphenols in blueberries and red grapes and wine [56]; epigallocatechin-3-gallate in green tea [57]; total phenolic, flavonoids, and flavonols in pomegranate [58, 59]). Red wine is particularly rich in specific polyphenolic compounds that appear to affect the biological processes of AD and Parkinson's disease, such as quercetin, myricetin, catechins, tannins, anthocyanidins, resveratrol, and ferulic acid. Indeed, there is now a consistent body of in vitro and in vivo data on the neuroprotective activity of red wine polyphenols [60], and it is known that effects may be boosted by adding more resveratrol [61]. Obviously, to enjoy optimum benefits, participants are taught to assume phytochemicals in a way to ensure protection throughout the day.
- b. Increasing the resistance of cell membranes to oxidative stress. In order to get a better resistance to ROS-induced damage to brain and prevent from neurodegeneration and neuronal apoptosis, participants are instructed to a timed assumption of omega-3-rich fish oil. PUFAs are essential to control oxidative damage to neurons [62] when they are incorporated in the phospholipid molecules as a part in the proposed antioxidant machinery that protects membrane proteins from peroxidation (**Figure 1**) (On the contrary, free PUFAs are very sensitive to a free radical attack and ready to peroxidation.) To get full benefit from supplement, participants are requested to take pills at the first good meal after the long time of fasting, when chances to be incorporated in membrane phospholipids are paramount [63].




Figure 1. Model of the membrane antioxidant machinery based on the proposed locations of dolichol (disc, green), ubiquinone (hexagon, red), and vitamin E (rhomb, pink) (see Cathcart et al. [69]; Sharma et al. [70]).

c. Promoting cell cleaning and rejuvenation. It is done by teaching participants in practice how to eat a healthy calorie-restricted diet and to make aerobic physical exercise during fasting. It is known indeed that calorie restriction is the most robust antiaging intervention known so far [64], and that benefit comes from the alternation of a long time of fasting and good meal (twice a week, participants spend a great part of their time in a state of fasting) [65] in order to get cells free of altered ROS-hypergenerating organelles in older cells [66, 67]. Good maintenance will be finalized later, thanks to a good meal. If needed, benefit might be magnified by taking a pill of an antilipolytic drug on the time of fasting [41]. Benefits of protocol might be monitored noninvasively by the assay of the urinary excretion of 8-hydroxy-2'-deoxyguanosine, a recently recommended biomarker for monitoring oxidative status over time [68].

d. A targeted high-intensity training of cognitive brain functions. In addition to the previously described interventions on diet and physical activity, the mental and the social activity protocols tested by Paillard-Borg et al. [38] are being practiced. In view of the results obtained by Dahlgren et al. [51], high intensity interval training (HIIT) and memory training using the PEAK brain training app are included in the protocol.

9. Conclusion

A dynamic antiaging nutritional and physical intervention protocol including enriched living conditions is described useful to prevent the appearance of aging-associated AD. Treatment is already granted for free to the citizens of the city of Volterra known to be at higher risk of AD (relatives of AD patients and persons with mild cognitive impairment, likely to progress to clinically probable AD at a considerably accelerated rate compared with healthy age-matched individuals) to teach them how to counteract and retard the disease. Treatment was designed to delay biological aging and empower adult brain plasticity in order to retard the progress of brain aging and associated diseases. It includes a daily assumption of antioxidants (red wine polyphenols enriched with resveratrol), a reinforcement of membrane antioxidant defenses by the timed assumption of polyunsaturated fatty acids, and an enhancement of cell repair function at the molecular and subcellular level by an intermittent feeding regimen and physical exercise, whose efficiency is empowered by the pharmacological intensification of autophagy by an antilipolytic drug (Acipimox) taken at a very low dosage while fasting. The beneficial effects on neurodegeneration are magnified by living an enriched environment. The effectiveness of the treatment will be detected by comparison with the expected frequencies of disease appearance.

Acknowledgements

Research is a part in the “train body and brain” project organized and supported by Volterra Research Foundation, Rotary District 2071, and Alberto Sordi Association, Rome, under the patronage of the city of Volterra. The project was born to give school-age citizens the most modern information on how to achieve an effective primary prevention of all chronic age-associated diseases, including neurodegenerative diseases, and to offer adult people at higher risk of Alzheimer's

disease (relatives of patients and patients with MCI) a free service of practical education on how to prevent and slow down the progression of the disease. The first goal is achieved by donating to high school students a pdf entitled “How to keep in shape, preventing diseases and defending good health” (available on the site www.fondazionevolterraricerche.org). The “train body and brain” project was approved by the Istituto Superiore di Sanità and has patronage of the Italian Ministry of Education.

Conflict of interest

The authors disclose no conflicts.

Author details

Ettore Bergamini^{1,2*} and Gabriella Cavallini³

¹ University Campus Biomedico, Roma, Italy

² Volterra Ricerche Onlus Foundation, Volterra, Italy

³ Interdepartmental Research Centre on Biology and Pathology of Aging, University of Pisa, Italy

*Address all correspondence to: ettore.bergamini@gmail.com

IntechOpen

© 2019 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. 

References

- [1] Tai HC, Serrano-Pozo A, Hashimoto T, Frosch MP, Spires-Jones TL, Hyman BT. The synaptic accumulation of hyperphosphorylated tau oligomers in Alzheimer disease is associated with dysfunction of the ubiquitin-proteasome system. *The American Journal of Pathology*. 2012;**181**:1426-1435. DOI: 10.1016/j.ajpath.2012.06.033
- [2] Niccoli T, Partridge L. Ageing as a risk factor for disease. *Current Biology*. 2012;**22**:R741-R752. DOI: 10.1016/j.cub.2012.07.024
- [3] Chen H, Kwong JC, Copes R, Tu K, Villeneuve PJ, van Donkelaar A, et al. Living near major roads and the incidence of dementia, Parkinson's disease, and multiple sclerosis: A population-based cohort study. *Lancet*. 2017;**389**:718-726. DOI: 10.1016/S0140-6736(16)32399-6
- [4] Singhrao SK, Harding A, Poole S, Kesavalu L, Crean S. *Porphyromonas gingivalis* periodontal infection and its putative links with Alzheimer's disease. *Mediators of Inflammation*. 2015;**2015**:137357. DOI: 10.1155/2015/137357
- [5] Davies KJ. The oxygen paradox, oxidative stress, and ageing. *Archives of Biochemistry and Biophysics*. 2016;**595**:28-32. DOI: 10.1016/j.abb.2015.11.015
- [6] Davies JMS, Cillard J, Friguet B, Cadenas E, Cadet J, Cayce R, et al. The oxygen paradox, the French paradox, and age-related diseases. *Geroscience*. 2017;**39**:499-550. DOI: 10.1007/s11357-017-0002-y
- [7] Ames BN, Shigenaga MK. Oxidants are a major contributor to aging. *Annals of the New York Academy of Sciences*. 1992;**663**:85-96. DOI: 10.1111/j.1749-6632.1992.tb38652.x
- [8] Stevens A, Lowe J, Scott I. *Core Pathology*. 3rd ed. Edinburgh: Mosby Elsevier; 2009
- [9] Bergamini E, Bizzarri R, Cavallini G, Cerbai B, Chiellini E, Donati A, et al. Ageing and oxidative stress: A role for dolichol in the antioxidant machinery of cell membranes? *Journal of Alzheimer's Disease*. 2004;**6**:129-135. DOI: 10.3233/JAD-2004-6204
- [10] Xue-Shan Z, Juan P, Qi W, Zhong R, Li-Hong P, Zhi-Han T, et al. Imbalanced cholesterol metabolism in Alzheimer's disease. *Clinica Chimica Acta*. 2016;**456**:107-114. DOI: 10.1016/j.cca.2016.02.024
- [11] Cheng H, Vetrivel KS, Gong P, Meckler X, Parent A, Thinakaran G. Mechanisms of disease: New therapeutic strategies for Alzheimer's disease-targeting APP processing in lipid rafts. *Nature Clinical Practice. Neurology*. 2007;**3**:374-382
- [12] Wang HL, Wang YY, Liu XG, Kuo SH, Liu N, Song QY, et al. Cholesterol, 24-hydroxycholesterol, and 27-hydroxycholesterol as surrogate biomarkers in cerebrospinal fluid in mild cognitive impairment and Alzheimer's disease: A meta-analysis. *Journal of Alzheimer's Disease*. 2016;**51**:45-55. DOI: 10.3233/JAD-150734
- [13] Toro P, Degen C, Pierer M, Gustafson D, Schröder J, Schönknecht P. Cholesterol in mild cognitive impairment and Alzheimer's disease in a birth cohort over 14 years. *European Archives of Psychiatry and Clinical Neuroscience*. 2014;**264**:485-492. DOI: 10.1007/s00406-013-0468-2
- [14] Devassy JG, Leng S, Gabbs M, Monirujjaman M, Aukema HM. Omega-3 polyunsaturated fatty acids and oxylipins in neuroinflammation and management of Alzheimer disease.

Advances in Nutrition. 2016;7:905-916. DOI: 10.3945/an.116.012187

[15] Gladine C, Newman JW, Durand T, Pedersen TL, Galano JM, Demougeot C, et al. Lipid profiling following intake of the omega 3 fatty acid DHA identifies the peroxidized metabolites F4-neuroprostanes as the best predictors of atherosclerosis prevention. PLoS One. 2014;9:e89393. DOI: 10.1371/journal.pone.0089393

[16] Tamburini I, Quartacci MF, Izzo R, Bergamini E. Effects of dietary restriction on age-related changes in the phospholipid fatty acid composition of various rat tissues. Aging Clinical and Experimental Research. 2004;16:425-431. DOI: 10.1007/BF03327396

[17] Tamburini I. Invecchiamento e stress ossidativo: Effetti dell'invecchiamento, di trattamenti anti-invecchiamento e di modulazioni acute e croniche dello stress ossidativo su molecole perossidabili lipofile di membrane [PhD thesis]. Pisa: University of Pisa; 2006

[18] Bergamini E. Targets for antiageing drugs. Expert Opinion on Therapeutic Targets. 2005;9:77-82

[19] Huang X, Cuajungco MP, Atwood CS, Hartshorn MA, Tyndall JD, Hanson GR, et al. Cu(II) potentiation of alzheimer abeta neurotoxicity. Correlation with cell-free hydrogen peroxide production and metal reduction. The Journal of Biological Chemistry. 1999;274:37111-37116. DOI: 10.1074/jbc.274.52.37111

[20] Nadal RC, Rigby SE, Viles JH. Amyloid beta-Cu²⁺ complexes in both monomeric and fibrillar forms do not generate H₂O₂ catalytically but quench hydroxyl radicals. Biochemistry. 2008;47:11653-11664. DOI: 10.1021/bi8011093

[21] Fernandez MA, Biette KM, Dolios G, Seth D, Wang R, Wolfe MS. Transmembrane substrate determinants for γ -secretase processing of APP CTF β . Biochemistry. 2016;55:5675-5688. DOI: 10.1021/acs.biochem.6b00718

[22] Liu CY, Ohki Y, Tomita T, Osawa S, Reed BR, Jagust W, et al. Two novel mutations in the first transmembrane domain of presenilin1 cause young-onset Alzheimer's disease. Journal of Alzheimer's Disease. 2017;58:1035-1041. DOI: 10.3233/JAD-161203

[23] Bergamini E. Autophagy: A cell repair mechanism that retards ageing and age-associated diseases and can be intensified pharmacologically. Molecular Aspects of Medicine. 2006;27:403-410. DOI: 10.1016/j.mam.2006.08.001

[24] Chang T-Y, Yamauchi Y, Hasan MT, Chang C. Cellular cholesterol homeostasis and Alzheimer's disease. Journal of Lipid Research. 2017;58:2239-2254. DOI: 10.1194/jlr.R075630

[25] Clarke PR, Hardie DG. Regulation of HMG-CoA reductase: Identification of the site phosphorylated by the AMP-activated protein kinase in vitro and in intact rat liver. The EMBO Journal. 1990;9:2439-2446

[26] Pallottini V, Martini C, Pascolini A, Cavallini G, Gori Z, Bergamini E, et al. 3-Hydroxy-3-methylglutaryl coenzyme A reductase deregulation and age-related hypercholesterolemia: A new role for ROS. Mechanisms of Ageing and Development. 2005;126:845-851. DOI: 10.1016/j.mad.2005.02.009

[27] Vetrivel KS, Thinakaran G. Membrane rafts in Alzheimer's disease beta-amyloid production. Biochimica et Biophysica Acta. 2010;1801:860-867. DOI: 10.1016/j.bbalip.2010.03.007

- [28] Barrett PJ, Song Y, Van Horn WD, Hustedt EJ, Schafer JM, Hadziselimovic A, et al. The amyloid precursor protein has a flexible transmembrane domain and binds cholesterol. *Science*. 2012;**336**:1168-1171. DOI: 10.1126/science.1219988
- [29] Fassbender K, Simons M, Bergmann C, Stroick M, Lutjohann D, Keller P, et al. Simvastatin strongly reduces levels of Alzheimer's disease beta-amyloid peptides Abeta 42 and Abeta 40 in vitro and in vivo. *Proceedings of the National Academy of Sciences of the United States of America*. 2001;**98**:5856-5861. DOI: 10.1073/pnas.081620098
- [30] Shinohara M, Sato N, Kurinami H, Takeuchi D, Takeda S, Shimamura M, et al. Reduction of brain beta-amyloid (Abeta) by fluvastatin, a hydroxymethylglutaryl-CoA reductase inhibitor, through increase in degradation of amyloid precursor protein C-terminal fragments (APP-CTFs) and Abeta clearance. *The Journal of Biological Chemistry*. 2010;**285**:22091-22102. DOI: 10.1074/jbc.M110.102277
- [31] Pallottini V, Martini C, Cavallini G, Bergamini E, Mustard KJ, Hardie DG, et al. Age-related HMG-CoA reductase deregulation depends on ROS-induced p38 activation. *Mechanisms of Ageing and Development*. 2007;**128**:688-695. DOI: 10.1016/j.mad.2007.10.001
- [32] Mendiola-Precoma J, Berumen LC, Padilla K, Garcia-Alcocer G. Therapies for prevention and treatment of Alzheimer's disease. *BioMed Research International*. 2016;**2016**:2589276. DOI: 10.1155/2016/2589276
- [33] Graham WV, Bonito-Oliva A, Sakmar TP. Update on Alzheimer's disease therapy and prevention strategies. *Annual Review of Medicine*. 2017;**68**:413-430. DOI: 10.1146/annurev-med-042915-103753
- [34] Maldonado TA, Jones RE, Norris DO. Distribution of beta-amyloid and amyloid precursor protein in the brain of spawning (senescent) salmon: A natural, brain-aging model. *Brain Research*. 2000;**858**:237-251. DOI: 10.1016/S0006-8993(99)02328-8
- [35] Dubois B, Hampel H, Feldman HH, Scheltens P, Aisen P, Andrieu S, et al. Proceedings of the meeting of the international working group (IWG) and the American Alzheimer's Association on "the preclinical state of AD"; July 23, 2015; Washington DC, USA. *Preclinical Alzheimer's disease: Definition, natural history, and diagnostic criteria. Alzheimer's & Dementia*. 2016;**12**:292-323. DOI: 10.1016/j.jalz.2016.02.002
- [36] Tan CC, Yu JT, Tan L. Biomarkers for preclinical Alzheimer's disease. *Journal of Alzheimer's Disease*. 2014;**42**:1051-1069. DOI: 10.3233/JAD-140843
- [37] Qiu C, Kivipelto M, von Strauss E. Epidemiology of Alzheimer's disease: Occurrence, determinants, and strategies toward intervention. *Dialogues in Clinical Neuroscience*. 2009;**11**:111-128
- [38] Paillard-Borg S, Fratiglioni L, Xu W, Winblad B, Wang HX. An active lifestyle postpones dementia onset by more than one year in very old adults. *Journal of Alzheimer's Disease*. 2012;**31**:835-842. DOI: 10.3233/JAD-2012-120724
- [39] Blagosklonny MV. Prospective treatment of age-related diseases by slowing down aging. *The American Journal of Pathology*. 2012;**181**:1142-1146. DOI: 10.1016/j.ajpath.2012.06.024
- [40] Bergamini E, Cavallini G, Donati A, Gori Z. The role of macroautophagy in the ageing process, anti-ageing intervention and age-associated diseases. *The International Journal*

of Biochemistry & Cell Biology. 2004;**36**:2392-2404. DOI: 10.1016/j.biocel.2004.05.007

[41] Bergamini E. Searching for the fountain of autophagy-dependent youth: Ettore Bergamini. Interview by Daniel J. Klionsky. *Autophagy*. 2012;**8**:1169-1174. DOI: 10.4161/auto.20788

[42] Fiuza-Luces C, Garatachea N, Berger NA, Lucia A. Exercise is the real polypill. *Physiology (Bethesda)*. 2013;**28**:330-358. DOI: 10.1152/physiol.00019.2013

[43] Schlosser Covell GE, Hoffman-Snyder CR, Wellik KE, Woodruff BK, Geda YE, Caselli RJ, et al. Physical activity level and future risk of mild cognitive impairment or dementia: A critically appraised topic. *The Neurologist*. 2015;**19**:89-91. DOI: 10.1097/NRL.0000000000000013

[44] Aguirre LE, Villareal DT. Physical exercise as therapy for frailty. Nestlé Nutrition Institute Workshop Series. 2015;**83**:83-92. DOI: 10.1159/000382065

[45] Sale A, Berardi N, Maffei L. Environment and brain plasticity: Towards an endogenous pharmacotherapy. *Physiological Reviews*. 2014;**94**:189-234. DOI: 10.1152/physrev.00036.2012

[46] Spencer NJ, Zagorodnyuk V, Brookes SJ, Hibberd T. Spinal afferent nerve endings in visceral organs: Recent advances. *American Journal of Physiology Gastrointestinal and Liver Physiology*. 2016;**311**:G1056-G1063. DOI: 10.1152/ajpgi.00319.2016

[47] Martin-Rincon M, Morales-Alamo D, Calbet JAL. Exercise-mediated modulation of autophagy in skeletal muscle. *Scandinavian Journal of Medicine & Science in Sports*. 2018;**28**:772-781. DOI: 10.1111/sms.12945

[48] Amann M. Significance of group III and IV muscle afferents for the

endurance exercising human. *Clinical and Experimental Pharmacology & Physiology*. 2012;**39**:831-835. DOI: 10.1111/j.1440-1681.2012.05681.x

[49] Carro E, Trejo JL, Busiguina S, Torres-Aleman I. Circulating insulin-like growth factor I mediates the protective effects of physical exercise against brain insults of different etiology and anatomy. *The Journal of Neuroscience*. 2001;**21**:5678-5684. DOI: 10.1523/JNEUROSCI.21-15-05678.2001

[50] Marosi K, Kim SW, Moehl K, Scheibye-Knudsen M, Cheng A, Cutler R, et al. 3-Hydroxybutyrate regulates energy metabolism and induces BDNF expression in cerebral cortical neurons. *Journal of Neurochemistry*. 2016;**139**:769-781. DOI: 10.1111/jnc.13868

[51] Dahlgren K, Gibas KJ. Ketogenic diet, high intensity interval training (HIIT) and memory training in the treatment of mild cognitive impairment: A case study. *Diabetes and Metabolic Syndrome: Clinical Research and Reviews*. 2018;**12**:819-822. DOI: 10.1016/j.dsx.2018.04.031

[52] Cherif A, Roelands B, Meeusen R, Chamari K. Effects of intermittent fasting, caloric restriction, and Ramadan intermittent fasting on cognitive performance at rest and during exercise in adults. *Sports Medicine*. 2016;**46**:35-47. DOI: 10.1007/s40279-015-0408-6

[53] Huang T, Larsen KT, Ried-Larsen M, Møller NC, Andersen LB. The effects of physical activity and exercise on brain-derived neurotrophic factor in healthy humans: A review. *Scandinavian Journal of Medicine & Science in Sports*. 2014;**24**:1-10. DOI: 10.1111/sms.12069

[54] von Strauss E, Viitanen M, De Ronchi D, Winblad B, Fratiglioni L. Aging and the occurrence of dementia: Findings from a

population-based cohort with a large sample of nonagenarians. *Archives of Neurology*. 1999;**56**:587-592. DOI: 10.1001/archneur.56.5.587

[55] Corrada MM, Brookmeyer R, Berlau D, Paganini-Hill A, Kawas CH. Prevalence of dementia after age 90: Results from the 90+ study. *Neurology*. 2008;**71**:337-343. DOI: 10.1212/01.wnl.0000310773.65918.cd

[56] Li YR, Li S, Lin CC. Effect of resveratrol and pterostilbene on ageing and longevity. *BioFactors*. 2018;**44**: 69-82. DOI: 10.1002/biof.1400

[57] Cascella M, Bimonte S, Muzio MR, Schiavone V, Cuomo A. The efficacy of Epigallocatechin-3-gallate (green tea) in the treatment of Alzheimer's disease: An overview of pre-clinical studies and translational perspectives in clinical practice. *Infectious Agents and Cancer*. 2017;**12**:36. DOI: 10.1186/s13027-017-0145-6

[58] Derakhshan Z, Ferrante M, Tadi M, Ansari F, Heydari A, Hosseini MS, et al. Antioxidant activity and total phenolic content of ethanolic extract of pomegranate peels, juice and seeds. *Food and Chemical Toxicology*. 2018;**114**: 108-111. DOI: 10.1016/j.fct.2018.02.023

[59] AlMatar M, Islam MR, Albarri O, Var I, Koksai F. Pomegranate as a possible treatment in reducing risk of developing wound healing, obesity, neurodegenerative disorders, and diabetes mellitus. *Mini Reviews in Medicinal Chemistry*. 2018;**18**:507-526. DOI: 10.2174/1389557517666170419114722

[60] Caruana M, Cauchi R, Vassallo N. Putative role of red wine polyphenols against brain pathology in Alzheimer's and Parkinson's disease. *Frontiers in Nutrition*. 2016;**12**(3):31. DOI: 10.3389/fnut.2016.00031

[61] Cavallini G, Straniero S, Donati A, Bergamini E. Resveratrol requires

red wine polyphenols for optimum antioxidant activity. *The Journal of Nutrition, Health & Aging*. 2016;**20**:540-545. DOI: 10.1007/s12603-015-0611-z

[62] Kim SJ, Zhang Z, Saha A, Sarkar C, Zhao Z, Xu Y, et al. Omega-3 and omega-6 fatty acids suppress ER- and oxidative stress in cultured neurons and neuronal progenitor cells from mice lacking PPT1. *Neuroscience Letters*. 2010;**479**:292-296. DOI: 10.1016/j.neulet.2010.05.083

[63] Cavallini G, Sgarbossa A, Parentini I, Bizzarri R, Donati A, Lenci F, et al. Dolichol: A component of the cellular antioxidant machinery. *Lipids*. 2016;**51**:477-486. DOI: 10.1007/s11745-016-4137-x

[64] Masoro EJ. *Caloric Restriction: A Key to Understanding and Modulating Aging*. 1st ed. Vol. 192. Amsterdam, The Netherlands: Elsevier Science; 2002

[65] Bergamini E, Cavallini G, Donati A, Gori Z. The role of autophagy in ageing: Its essential part in the anti-ageing mechanism of caloric restriction. *Annals of the New York Academy of Sciences*. 2007;**1114**:69-78. DOI: 10.1196/annals.1396.020

[66] Donati A, Taddei M, Cavallini G, Bergamini E. Stimulation of macroautophagy can rescue older cells from 8-OHdG mtDNA accumulation: A safe and easy way to meet goals in the SENS agenda. *Rejuvenation Research*. 2006;**9**:408-412. DOI: 10.1089/rej.2006.9.408

[67] Cavallini G, Donati A, Taddei M, Bergamini E. Peroxisomes proliferation and pharmacological stimulation of autophagy in rat liver: Evidence to support that autophagy may remove the "older" peroxisomes. *Molecular and Cellular Biochemistry*. 2017;**431**:97-102. DOI: 10.1007/s11010-017-2979-z

[68] Cavallini G, Donati A, Capasso B, Fella M, Leone V, Pezzella G, et al. Effects of stimulation of autophagy on the urinary excretion of biomarkers of the oxidative status. *Aging Clinical and Experimental Research*. 2014;**26**:13-18. DOI: 10.1007/s40520-013-0116-8

[69] Cathcart R, Schwieters E, Ames BN. Detection of picomole levels of hydroperoxides using a fluorescent dichlorofluorescein assay. *Analytical Biochemistry*. 1983;**134**:111-116. DOI: 10.1016/0003-2697(83)90270-1

[70] Sharma MK, Buettner GR. Interaction of vitamin C and vitamin E during free radical stress in plasma: An ESR study. *Free Radical Biology & Medicine*. 1993;**14**:649-653. DOI: 10.1016/0891-5849(93)90146-L