We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists



186,000

200M



Our authors are among the

TOP 1% most cited scientists





WEB OF SCIENCE

Selection of our books indexed in the Book Citation Index in Web of Science™ Core Collection (BKCI)

Interested in publishing with us? Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected. For more information visit www.intechopen.com



Chapter

Stability of Vanadium Chalcone Complexes

Vikas Vilas Borge and Raju M. Patil

Abstract

The vanadyl(IV) complexes of substituted chalcones were prepared by refluxing vanadyl sulphate with different substituted chalcones in ethanoic medium. The chalcones were prepared with different aromatic aldehydes like benzaldehyde, hydroxy benzaldehyde, nitro benzaldehyde and chloro benzaldehyde. The synthesized Vanadium complexes were characterized by different spectral techniques. The IR spectral studies revealed that the chalcone derivatives are bidentate ligand. Magnetic studies, electron spin resonance and UV studies suggest that the complexes are in square pyramidal geometry. Conductance measurements suggest that all complexes are non-electrolyte in DMF. The thermal study explained the stability of complex and its decomposition. The synthesized ligand and metal complexes were screened for their antibacterial activity against *E. coli* and *Staphylococcus aureus* bacterial strains and for antifungal activity against *P. notatum*.

Keywords: vanadyl sulphate, chalcone ligands, complexes, spectral studies, TGA, magnetic study, antifungal and antibacterial study

1. Introduction

Chemically chalcones consist of open chain flavonoids in which the two aromatic ring joined by a three carbon. Chalcones are α , β -unsaturated ketones and contain a reactive keto-ethylenic group 2-hydroxychalcones and their heterocyclic analogues are also reported to form coordination complexes.

Chalcones are widely distributed in fruits, vegetables, tea, spices and also have applications in biological activities. These are abundant in edible plants and are considered to be the precursors of flavonoids and isoflavonoids [1]. Some chalcones has been employed for the gravimetric estimation of divalent ions of palladium, copper and nickel. Calcium determination by complexometric titration by using chalcone-metanil as indicator also has been reported. Chalcones possesses several biological activities with wide ranging. Chalcones use for the treatment of chronic diseases which involve inflammatory processes, for example, diabetes mellitus [2]. The study of chalcone derivatives has become of much interest in recent years on account of their antibacterial, antiviral activities [3]. Many chalcones based have been recognized in a number of screening assays for modulating important pathways or molecular targets in cancer [4, 5].

The coordination chemistry of oxovanadium(IV) (i.e. VO²⁺ or vanadyl ion) is more interesting and rather more important because of two main reasons. Firstly, the vanadyl complexes are finding more and more importance in biological systems. Secondly, the coordination number and geometry of this metal is highly ligand dependent. Moreover vanadyl ion is less toxic. In last year's research has been directed towards the synthesis of efficient bioactive compounds with low toxicity, in order to achieve this goal the type and the position of substituent into ligand were varied [6].

A wide variety of oxovanaduim(IV) complexes have been prepared and characterized by various physicochemical methods. Oxovanadium(IV) complexes have a square pyramidal geometry with apical oxygen and the vanadium atom lying above the plane defined by the donor atoms of the equatorial ligands. These square pyramidal complexes generally exhibit strong tendency to remain five coordinate [7, 8]. The coordination chemistry of vanadium(V) compounds is dominated by oxo complexes, containing the VO³⁺ or the VO²⁺ moiety [9]. The absorption band due to V=O stretching vibration of oxovanadium(IV) complexes is usually observed at a higher wavenumber compared to those of vanadate(V) complexes. However, the V=O stretching vibration is susceptible to a number of influences including electron donation from basal plane ligand atoms, solid-state effects, and coordination of additional molecules. Therefore, there has been considerable work done to assign the V=O stretching frequencies in oxovanadium(IV) compounds [10].

An increasing research interest of vanadium in coordination chemistry is not only due to its exhibiting a range of oxidation dates from +5 to -1 but also complexities exhibited by vanadium complexes and their industrial [11, 12], biological [13, 14] and medicinal [15, 16] applications. Vanadium complexes have been reported to have interesting antibacterial activities [17–20]. It has been reported that V(IV) complexes high effect on anticancer activity [21].

2. Preparation of substituted chalcone

The substituted chalcones prepared by stirring the equimolar concentration mixture of 2-hydroxy-4,5-dimethyl acetophenone (0.01 mol) and substituted aromatic benzaldehyde (0.01 mol) in 20 ml ethanol for 1 h in presence of 50% NaOH. The mixture stirred till completion of the reaction (progress of reaction checked by TLC). The crude mixture poured into ice water then acidified the product with 10% hydrochloric acid. The coloured compound formed was filtered, washed with water and dried. The compounds recrystallized from absolute ethanol (**Figure 1**).



Figure 1. *Reaction scheme for chalcones (R* = *H*, 4-OH, 4-NO₂ *and* 3-*chloro).*

3. Preparation of complexes

To a hot suspension of ligand (0.02 M) in ethanol, ethanolic solution (0.01 M) of the metal salt vanadyl sulphate (VOSO₄) was added drop wise with constant stirring and refluxed for 2–3 h. The resulting reaction mixture was cooled to room temperature and maintained up to pH 8.0 by adding ammonia then refluxed further for 30 min. The resultant product was filtered through Whatman filter paper no. 1 and repeatedly washed with ethanol until the washing were free from the excess of ligand. These complexes were finally dried under vacuum in desiccator over fused CaCl₂.

4. Instrumental methods

The IR spectra of complexes were recorded on a Perkin-Elmer instrument in KBr pallets in the range of 4000–400 cm⁻¹. TGA analysis of metal complexes were carried out in nitrogen atmosphere in the range 25–900°C on Rigaku Thermo Plus-8120 TG-DTA instrument with a heating rate 10°C min⁻¹ using Alumina as a standard. UV-Visible spectra were recorded using DMF as solvent on Shimadzu UV-VIS spectrophotometer in the range 250–950 nm. Electron spin resonance spectra complexes were recorded on E-112 ESR Spectrometer as 'g' marker (g = 2.00277) at room temperature. The conductance was measured in DMF solvent on Equiptronics Conductivity meter (EQ-664A).

5. Results and discussion

All the vanadium chalcone complexes are insoluble in water and most common organic solvents but sparingly soluble in DMF. The complexes are stable at room temperature. The elemental analysis shown in **Table 1** indicates that, all metal complexes have 1:2 (metal: ligand) stoichiometry for all the complexes (**Figure 2**).

$$VL_1:R_1 = H \& R_2 = H; VL_2:R_1 = H \& R_2 = OH$$

$$VL_3:R_1 = H \& R_2 = NO_2 and VL_4:R_1 = Cl \& R_2 = H$$

All synthesized complexes are green in colour. All complexes are decomposed above 300°C. The molar conductance values obtained for these complexes at the concentration are in the range of 15–27 mhos.cm² mol⁻¹ suggesting their non-electrolytic nature [22].

5.1 IR spectral studies

Some structurally significant IR bands for uncoordinated chalcones and complexes have been scanned in the $4000-400 \text{ cm}^{-1}$ region and presented in **Table 2**. The complexes shows a band in the region 1162–1205 cm⁻¹ which corresponds to

Complex	Empirical	Mol	Yield	% Elemental analysis found (calculated)						Molar
	formula	wt.	%	М	С	Н	N	Cl	0	conductance mhos.cm ² mol ⁻¹
V L ₁	$C_{34}H_{30}O_5V$	569	62	9.10	71.52	5.15	_	_	14.13	18.21
			_	(8.96)	(71.70)	(5.27)			(14.06)	
V L ₂	$C_{34}H_{30}O_7V$	601	59	8.55	67.94	4.91		_	18.58	23.46
			-	(8.48)	(67.88)	(4.99)			(18.63)	
VL ₃	$C_{34}H_{28}N_2O_9V$	659	65	7.68	61.93	4.22	4.28	_	21.88	15.62
			_	(7.74)	(61.91)	(4.25)	(4.25)		(21.85)	
$V L_4$	C34H28O5Cl	638	56	8.05	63.88	4.35	_	11.13	12.58	26.33
	V		-	(7.99)	(63.95)	(4.39)		(11.1)	(12.54)	

 Table 1.

 Physical and analytical data of metal complexes.

strong band in the region 1185–1221 in free ligand of chalcones due to ν (C—O). The shifts of this band were observed to move to lower wave numbers indicates bonding of phenolic oxygen atom to metal center via deprotonation. The band in the region 552–573 cm⁻¹ can be assigned to the stretching modes of the metal to ligand bonds, ν (V—O) [23]. In addition the compound exhibit a strong band in the 952–983 cm⁻¹ range due to the terminal V=O stretching and this value is close to the majority of oxovanadium(IV) complexes [24]. In the IR spectra of all the ligands an intense band appearing around 1640 cm⁻¹ is attributed [25] to (C=O) group. In the complexes this band is observed in the 1618–1626 cm⁻¹ range. The shifting of band to lower wave number indicates coordination through oxygen of (C=O) group. The band appears around 740 cm⁻¹ in chloro complex indicating presence of C—Cl bond. The occurrence of band around 1500 cm⁻¹ in VL₃ complex is due to ν (C—NO₂) mode.



Figure 2. Structure of vanadium complex.

Compound	OH group	C=0	C==C	C=0	C=CH(Aro)	V—0	-NO ₂	C—Cl	V=0
L	2921	1640	1503	1221	2858	_			_
VL1	3029	1626	_	1198	2964	552	—	—	963
L ₂	3242	1639	1588	1221	2987	—			—
VL ₂	3174	1621	1404	1205	3018	561	—	—	952
L ₃	3390	1635	1513	1190	2921	—	1520		—
VL ₃	3353	1623	1406	1168	3018	556	1490	_	983
L ₄	3527	1635	1567	1185	2920	_		738	_
VL_4	3190	1618	1583	1162	3020	573	_	742	976

Table 2.Important IR spectral bands (cm^{-1}) of the complexes.

5.2 Electronic spectra

Electronic absorption spectral study is an important tool used [26] to determine the geometry around a central metal atom. The electronic spectra (**Figure 3**) of the metal complexes in DMF (10^{-3} M) were recorded on Shimadzu UV-VIS spectrophotometer in the range 200–950 nm. The Vanadium chalcone complexes show absorption band in 380–770 nm region (**Table 5**). Most of the VO(II) complexes are green in colour. The electronic absorption spectra of VO(II) complexes showed three bands at 13,015–13,305, 17,455–17,632 and 25,755–25,930 cm⁻¹, with 17,455–17,632 cm⁻¹ representing 10 Dq. These bands are assigned to the transitions ${}^{2}B_{2} \rightarrow [2]E(d_{xy} \rightarrow d_{xz}, d_{yz}), {}^{2}B_{2} \rightarrow {}^{2}B_{1}(d_{xy} \rightarrow dx^{2}-y [2]), and {}^{2}B_{2} \rightarrow [2]A_{1}(d_{xy} \rightarrow d_{z}^{2})$ respectively. From these electronic absorption bands and paramagnetic nature, square pyramidal geometry suggested for VO(II) chalcone complexes. As these oxovanadium(IV) belong to the 3d¹ system, there is no interelectronic repulsion in the metal ion so B₀ value of V⁴⁺ is not available. Therefore calculation of Racah parameter is not possible for these vanadium complexes (**Table 3**).



Figure 3. Electronic spectra of V(IV) complexes.

Complexes	(v1)	<i>(v2)</i>	(v3)	Dq (cm ⁻¹)	CFSE (KJ/mol)
VL ₁	13,275	17,455	25,810	1745	-83.50
VL ₂	13,305	17,585	25,930	1758	-84.12
VL ₃	13,015	17,632	25,755	1763	-84.36
VL ₄	13,135	17,490	25,890	1749	-83.69

Table 3.

Electronic spectral data for vanadium metal complexes.

Complex	Magnetic moment (B.M.)
VL ₁	1.73
VL ₂	1.71
VL ₃	1.76
VL ₄	1.72

Table 4.

Magnetic moment values of the vanadium (IV) chalcone complexes.

5.3 Magnetic susceptibility measurement

The magnetic susceptibility for all the Vanadium complexes at room temperature were recorded by the Gouy's method using Hg[Co(SCN)₄] as a calibrant and reported in **Table 4**. The effective magnetic moments were calculated after applying diamagnetic corrections for the ligand components using Pascal's constants. The room temperature μ_{eff} values for the vanadium complexes were found in the range 1.71–1.76 B.M. The magnetic susceptibilities of the complexes are consistent with square pyramidal geometry around the central metal ion [27, 28]. The magnetic moments of the compounds investigated are in agreement with the findings of electronic absorption spectra.

5.4 Electron spin resonance study

ESR spectrum of VO(II) complexes was recorded at room temperature as polycrystalline sample, on *x*-band at 9.1 GHz, under the magnetic field strength 3000 G using TCNE as marker. The value of g_{\parallel} and g_{\perp} is shown in **Table 5**. Paramagnetism of vanadyl ion arises from the single unpaired electron, its paramagnetism is due to spin angular momentum and VO²⁺ is most stable diatomic cation. In V(IV) complexes value of g is below value of free electron. The trend 2.0023 > g_{\perp} > g_{\parallel} , observed for the complexes. The EPR signal are split into eight lines suggested that V(IV) complexes are mononuclear [29]. The oxovanadium(IV) ion belong to the 3d¹ system. Because of strong multiple covalent vanadium oxo-bond, a tetragonal compression occurs and therefore square pyramidal geometry proposed for VO(II) complex. The d orbital split into different level, dx^2-y^2 orbital is least stable as it is face to face with all four equatorial ligands. d_{xy} being most stabilized orbital therefore d¹ electron lies in d_{xy} orbital. As G values of present complexes are lower than four indicates, the ligands involve in the complex formation are the strong field ligands.

5.5 Thermal measurements

A plot of change in weight vs. temperature or time is known as Thermo gravimetric curve. It records the loss in weight as a function of temperature for transition that involves dehydration or decomposition. The TGA studies of the complexes was recorded in nitrogen atmosphere on Rigaku Thermo Plus-8120 instrument by increasing the temperature from room temperature upto 900°C at the heating rate of 10°C min⁻¹. The decomposition pattern obtained for VO(II) complexes of chalcone ligands follow two major stages (**Figure 4**). The thermal dehydration of this complex in the first stage was attributed to the loss of water leaving behind the nonhydrated complex takes place between 80 and 120°C. In the second stage the complex starts decomposing gradually which occurs in the temperature range corresponds to 220–310°C, this range correspond to the decomposition of part of chalcone ligand and the observed mass loss was recorded as 29.81% for VL₄, 32.41% for VL₂, 38.78%

Complex	g∥	g⊥	g _{av}	G axial symmetry parameter
VL ₁	1.981	1.986	1.984	1.31
VL ₂	1.958	1.983	1.975	2.29
VL ₃	1.973	1.987	1.982	1.91
VL ₄	1.949	1.987	1.974	3.48

 Table 5.

 ESR spectral data of vanadium(IV) metal complexes.

Stability of Vanadium Chalcone Complexes DOI: http://dx.doi.org/10.5772/intechopen.88072

for VL₃ and 36.90% for VL₄ complex, which shows that at least unstable intermediate are also formed during this degradation stage. The third stage was decomposition of complete chalcone ligand in temperature range of 550–700°C. The overall total mass loss up to 700°C is around 88–90% which shows finally formation of VO. The residue left after 700°C is around 9–11% which resemble the theoretical values. Hence from thermal analysis it is clear that the complexes under study contains two water molecules for VL₁, VL₃ and VL₄ complex and three water molecules for VL₂ complex which are present in the ionic sphere of the complex (**Table 6**).



Figure 4.				
Thermogravimetry	spectra	of	V(IV)	complexes.

Complex	plex Temperature range (°C) % Weight loss		Decomposition product	
		Found	Calculated	
VL ₁	80–110	5.90	5.95	2H ₂ O
-	235–290	29.81	29.75	C ₁₄ H ₁₂
	550–680	53.26	53.20	$C_{20}H_{18}O_4$
6	700–780	11.03	11.07	VO
VL ₂	85–110	8.30	8.24	3H ₂ O
	230–300	32.41	32.36	C ₁₄ H ₁₂ O ₂
	570–660	49.09	49.16	$C_{20}H_{18}O_4$
	700–750	10.16	10.22	VO
VL ₃	80–115	5.12	5.17	2H ₂ O
	240–310	38.78	38.85	$C_{14}H_{10}O_4N_2$
-	580–700	46.41	46.33	$C_{20}H_{18}O_4$
	720–780	9.70	9.64	VO
VL ₄	90–120	5.27	5.34	2H ₂ O
	220–290	36.90	36.94	$C_{14}H_{10}Cl_2$
-	550–660	47.85	47.77	$C_{20}H_{18}O_4$
	710–760	9.95	9.94	VO

Table 6.

Thermal data for vanadium metal complexes.

	Antibacterial activity		Antifungal activity
Compound	E. coli	S. aureus	P. notatum
	$1.0\mathrm{mgml}^{-1}$	$1.0\mathrm{mgml^{-1}}$	250 μg disc ⁻¹
L ₁	6	5	8
VL ₁	9	8	9
L ₂	8	9	11
VL ₂	11	13	13
L ₃	5	4	6
VL ₃	7	6	7
L ₄		8	10
VL ₄	9	11	12
Neomycin	26	27	_

Table 7.

Antibacterial and antifungal activities (inhibition zone of bacterial growth, mm) of the ligand and metal complexes.

6. Antibacterial activity

It is known that killing of bacteria by chelated ligand is more powerful than non-chelated ligand, therefore chelated ligand acts as potent bactericidal agents. It was observed that due to delocalization of the π electrons over whole ligand and partial sharing of its positive charge with donor groups reduce the polarity of a metal ion in a complex. Thus, chelation increases the lipophilic character in complexes and results in an enhancement of activity [30, 31]. The antibacterial results, **Table 7**, showed that the complexes shows moderate activity against the bacteria *E. coli* and *S. aureus*.

7. Antifungal activity

Table 7 showed that metal complexes exhibited significant antifungal activity than chalcones at the same concentration against the fungi *P. notatum*. The order of inhibition with respect to metal complexes was $VL_2 > VL_4 > VL_4 > VL_3$.

It was concluded from above data that electron withdrawing substituents like nitro, cyno at para position in ring A decreases the antibacterial and antifungal activity. Electron donating group like hydroxyl, methoxy at para position increases the activity. Presence of halogen group at meta position in ring A shows good activity than the unsubstituted ring. The increased activity of the chelates can be explained based on the overtone concept and the Tweedy chelation theory [32]. According to the overtone concept of cell permeability, the lipid membrane that surrounds the cell favors the passage of only lipid-soluble material.

8. Conclusion

Based on the results the following conclusion maybe drawn. The higher decomposition temperature and electrical conductance studies show the presence of strong metal-ligand bonding and non-electrolytic nature of the complexes, respectively. Room temperature magnetic studies are indicative of paramagnetic nature Stability of Vanadium Chalcone Complexes DOI: http://dx.doi.org/10.5772/intechopen.88072

and square pyramidal geometry of the VO(II) complexes which is supported by the electronic spectra. The IR spectra shows bonding of the metal through O-donor atoms of the ligands. In this study VO(II) complexes shows the moderate activity against *Escherichia coli*, *Staph aureus* bacteria and against *P. notatum* fungi.



Intechopen

Author details

Vikas Vilas Borge^{*} and Raju M. Patil Department of Chemistry, The Institute of Science, Mumbai, India

*Address all correspondence to: vikasborge@gmail.com

IntechOpen

© 2020 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/3.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

References

[1] Ahmad MR, Sastry VG, Bano N, Anwar S. Synthesis of novel chalcone derivatives by conventional and microwave irradiation methods and their pharmacological activities. Arabian Journal of Chemistry. 2016;**9**:931-935

[2] Najafian M, Ebrahim-Habibi
A, Yaghmaei P, Parivar K, Larijani
B. Core structure of flavonoids
precursor as an antihyperglycemic and antihyperlipidemic agent: As in vivo
study in rats. Acta biochimica Polonica.
2010;57:553

[3] Khan SA, Asiri AM. Arabian Journal of Chemistry. 2017;**10**:2890-2895

[4] Evangelista FCG, Bandeira MO, Silva GD, Silva MG, Andrade SN, Marques DR, et al. Anticancer activity of ruthenocenyl chalcones and their molecular docking studies. Medicinal Chemistry Research. 2017;**26**:27-43

[5] Khanapure S, Jagadale M, Bansode P, Prafulla C, Rashinkar G. Synthesis, spectroscopic and antimicrobial studies of oxovanadium (IV) complexes incorporating tridentate ONO donor hydrazones. Journal of Molecular Structure. 2018;**1173**:142-147

[6] Patil SK, Naik VM, Mallur NB.
Comprehensive Coordination
Chemistry. The synthesis, reactions,
properties and applications of
coordination compounds. Journal of
Chemical and Pharmaceutical Research.
2012;4(4):2029-2036

[7] Wilkinson G, editor. Comprehensive Coordination Chemistry. Oxford: Pergamon Press; 1987

[8] Page EM, Wass SA. Vanadium 1995.Coordination Chemistry Reviews.1997;164:203

[9] Nicholls D. In vitro study of the insulin-like action of vanadyl-pyrone

and pyridinone complexes with a VO(O4) coordination mode. Coordination Chemistry Reviews. 1966;**1**:379-414

[10] Rangel M, Tamura A, Fukushima C, Sakurai H. Oxovanadium(V) Tetrathiacalix[4]arene complexes and their activity as oxidation catalysts. Journal of Biological Inorganic Chemistry. 2001;**6**:128

[11] Hoppe E, Limberg C. Vanadium catalysed asymmetric epoxidation of homoallylic alcohols. Chemistry: A European Journal. 2007;**13**:7006-7016

[12] Zhang W, Yamamoto H. Journal of the American Chemical Society.2007;**129**:286-287

[13] Aureliano M, Gandara RMC. The role of vanadium in the chemical defense of the solitary tunicate, phallusia nigra. Journal of Inorganic Biochemistry. 2005;**99**:979-985

[14] Shobu O, Joseph P, Chem J. Insulin-Enhancing vanadium(III) complexes. Ecology. 2007;**33**:643-654

[15] Melchior M, Rettig SJ, Liboiron BD, Thompson KH, Yuen VG, McNeill JH, et al. Solid state and solution studies of vanadium(III)-L-cysteine compound and demonstration of its antimetastatic antioxidant and inhibition of neutral endopeptidase activities. Inorganic Chemistry. 2001;**40**:4686-4690

[16] Papaioannou A, Manos M, Karkabounas S, Liasko R, Evangelou AM, Correia I, et al. Synthesis and reactivity in inorganic, metal-Organic, and nano-metal chemistry microwave assisted synthesis and antimicrobial study of schiff base vanadium(IV) complexes of phenyl esters of amino acids. Journal of Inorganic Biochemistry. 2004;**98**:959-968 Stability of Vanadium Chalcone Complexes DOI: http://dx.doi.org/10.5772/intechopen.88072

[17] Wazalwar SS, Bhave NS, Dikundwar AG, Ali P. Synthesis and Reactivity in Inorganic Metal-Organic and Nano-Metal Chemistry. 2011;**41**:459

[18] Liu J-L, Sun M-H, Ma J-J. Metal based biologically active compounds: Design, synthesis and antibacterial/antifungal/ cytotoxic properties of triazole-derived schiffs bases and their oxovanadium(IV) complexes. Synthesis and Reactivity in Inorganic Metal-Organic and Nano-Metal Chemistry. 2015;**45**:117

[19] Chohan ZH, Sumrra SH, Youssoufi MH, Hadda TB. Synthesis, crystal structures and antibacterial studies of oxidovanadium(IV) complexes of salen-type schiff base ligands derived from meso-1,2-diphenyl-1,2ethylenediamine. European Journal of Medicinal Chemistry. 2010;45:2739

[20] Taheri O, Behzad M, Ghaffari A, Kubicki M, Dutkiewicz G, Bezaatpour A, et al. New ni(II) complexes involving symmetrical bidentate N,O-donor Schiff base ligands: Synthesis at ambient temperature, crystal structures, electrochemical study, antioxidant and cytotoxic activities. Transition Metal Chemistry. 2014;**39**:253

[21] Abbasi Z, Salehi M, Kubicki M, Khaleghian A. Journal of Coordination Chemistry. 2017;**70**:2074

[22] Geasy MT. Spectroscopic studies of oxovanadium coordination compounds. Coordination Chemistry Reviews. 1971;7:81

[23] Baran EJ. Journal of Coordination Chemistry. 2001;**54**:215-238

[24] Syamal A, Kale KS. Inorganic Chemistry. 1979;**18**:992-995

[25] Dhar DN, Gupta VP. Indian Journal of Chemistry. 1971;**9**:818

[26] Cotton FA, Wilkinson G. Advanced Inorganic Chemistry. 5th ed. New York: John-Wiley & Sons; 1988 [27] Selwood PW. Magnetochemistry.2nd ed. New York: Interscience; 1956

[28] Ballhausen CJ. Introduction to Ligand Field Theory. New York: McGraw-Hill; 1962

[29] Mustafi D, Makinen MW. Inorganic Chemistry. 2005;44:5580-5590

[30] Shrivastava RS. Metal-based antibacterial and antifungal amino acid derived Schiff bases: Their synthesis, characterization and in vitrobiological activity. Inorganica Chimica Acta. 1981;**56**:165

[31] Chohan ZH, Arif M, Sarfaz M. Synthesis, structural characterization and electrochemical and antibacterial studies of schiff base copper complexes. Applied Organometallic Chemistry. 2007;**21**:294

[32] Raman N, Kulandaisamy A, Tungaraja C, Manishankar P, Viswanthan S, Vedhi S. Transition Metal Chemistry. 2004;**29**:129

