

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

185,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.  
For more information visit [www.intechopen.com](http://www.intechopen.com)



# Functional Value of Amaranth as Applied to Sports Nutrition

*Torregrosa-García Antonio and López-Román F. Javier*

## Abstract

Amaranth can be beneficial to health and sports, with multiple applications owing to varied concentration of phytochemical; the concentration of these compounds depends on the part or by-product employed. For example, as a food supplement, amaranth oil (rich in squalene) can exert cardiovascular properties, while amaranth extract (rich in nitrites) can increase nitric oxide production (therefore improving endurance performance). On the other hand, as a functional ingredient, its gluten-free flours (containing fibre) can act as probiotics, whereas its proteins (with a peculiar amino acid profile) support muscle recovery. The few clinical results with athletic population suggest it can contribute to improved aerobic metabolism, but there is insufficient clinical data to draw any conclusion. Further research warrants elucidation of amaranth phytochemicals as promising ergogenic aids in sports by high-quality human clinical trials with both standardised products and ingredients in athletic population.

**Keywords:** amaranth, sports nutrition, clinical trial, functional food, athletic performance, recovery, ergogenic aid

## 1. Introduction

The field of sports nutrition studies how food and other dietary supplements can improve performance and general health status of physically active subjects, or athletes (both recreational and professional). Usually, recreational physical activity aims to obtain the healthy benefits which can include improved mood and movement in daily physical activities, aesthetic reasons and maintenance of health status. On the other hand, competitive athletes (especially elite athletes) pursue to peak performance.

For this reason, the first category of subjects would prefer including functional foods and change their overall diet by healthier choices (whatever definition is in their minds), while competitive subjects might prefer a combination of nutritional strategies with promising dietary supplements. Surveys vary much to this respect, but statistics estimated that between 40 and 100% of athletes typically uses supplements such as sports foods, medical supplements, ergogenic supplements and functional foods (super foods) [1, 2]. Amaranth is a pseudo-cereal which has many edible parts like seeds and leaves and by-products (like amaranth oil, flour and powder) which has been attributed to many health benefits.

The area of sports nutrition studying all these aspects is an emerging field which shows some evidence showing efficacy for active compounds, while others still need further quality research to draw reliable initial conclusions. This chapter

reviews the existing literature of amaranth applied to the sports field and discusses the potential application of amaranth and its derivate products to sports and their functional value supported by the existing clinical and empirical data.

## 2. Existing literature of amaranth in the sports field

The number of published research articles of amaranth with physically active human population is still scarce. The few studies found (through Google Scholar and PubMed search engines) at the time of writing this article are summarised in **Table 1**. Two of the three articles found employed amaranth oil, while the other employed a different product (amaranth drink instead of amaranth oil). To our best knowledge, only one of the studies assessed amaranth supplementation in an exercise test, while the other two studies used athletic population to measure aerobic metabolism parameters and cardiovascular parameters (heart rate variability). Furthermore, the only trial with an exercise test administered an amaranth drink, while most of the clinical literature addresses the functional properties of amaranth oil—without a single clinical trial with exercise tests in athletic population to our best knowledge. Due to the small amount of exercise test-related data, it is not feasible to obtain strong conclusions of the contribution of amaranth products to sports, according to various guidelines to assess the level of clinical evidence [3, 4].

## 3. Prospective applications of functional value of amaranth to exercise

There is a good *rationale* to employ amaranth in the sports field. The following chapter discusses the potential application of the functional value of amaranth and its by-products supported by the existing clinical and empirical data of previous studies.

### 3.1 Amaranth sports performance and metabolism

#### 3.1.1 Amaranth impact on aerobic metabolism

Among amaranth products, the most studied by its functional properties is perhaps amaranth oil. Amaranth oil possesses large quantities of polyunsaturated fatty acids (PUFAs) especially linoleic acid (LA) (38.2%) (the most important FA in the omega-6 family) and oleic acid (33.3%) [5]. Action of PUFAs on heart rate variability was previously linked to omega-3 fatty acids (FAs) [6] translated in heart rate recovery after exercise partly attributed to incorporation in myocardial cells, affecting cardiovascular performance [7–9]. However, amaranth oil has a small quantity of alpha-linolenic acid (ALA) (compared to other seeds like walnuts and flaxseed oil [10] which are the one addressed to improve performance in sports [11]).

In spite of this fact, amaranth oil was studied in athletes showing activation of aerobic metabolism (important in long distances and low- to medium-intensity sports), concomitant with clinical biomarkers showing an improved utilisation of oxidation waste products (lipid peroxidation products), showing improved heart rate variability (HRV) [12], which is related with a better physical performance condition [13, 14]. This happened also in another study also for the group patients with diabetes mellitus 2 [15] later confirmed by studies in unhealthy subjects (duodenal peptic ulcer patients) [16].

Title	N	Trial design	Test	Supplementation protocol	Outcome/s
The influence of an amaranth-based beverage on cycling performance: a pilot study [28]	6 trained cyclists (5 men and 1 woman)	Randomised, counterbalanced, cross-over trial	2 time trials (TT): 10 minutes of warm-up, 32.20 km, 10 minutes of recovery, and 5 km	Periodic intake of 150 ml of an amaranth based (52.48 kcal/100 ml) or commercial drink (24 kcal/100 ml) rich in carbohydrates, every 10 minutes through the first TT and recovery time	↓ Time to complete 1st TT Trend to decreased time to complete 2nd TT (p = 0.06) No significant changes in rate or perceived exertion (RPE), muscular power during the TT or haematocrit
Activation of aerobic metabolism by amaranth oil improves heart rate variability both in athletes and patients with type 2 diabetes mellitus [15]	36 patients with diabetes mellitus type 2 (DM) (20 male and 16 female) 24 national level athletes (16 male and 8 female)	Comparative study	Heart rate variability (HRV) (standing and orthostatic test) Aerobic metabolism (by lipid peroxidation/ antioxidant activity and blood haemoproteins)	1 ml of concentrated amaranth oil ( <i>Amaranthus cruentus</i> ) obtained by vacuum CO2 cold extraction, at least 2 hours before meal, for 28 ± 2 days in the DM group and 22 ± 2 days for athletes	In the athlete group: ↑ HRV Haemoglobin ligands were redistributed ↑ levels of oxidative destruction products Moderate strength correlations of HRV with aerobic metabolism found
Study of aerobic metabolism parameters and heart rate variability and their correlations in elite athletes: a modulatory effect of amaranth oil [12]	68 (36 competitive male athletes and 32 healthy males)	Comparative study	HRV and autonomic profile Oxidative stress (catalase activity, superoxide dismutase, oxidatively modified proteins) Middle mass molecules Haemoglobin (Hb) and its ligands (oxyHb-, carboxyHb, sulfHb and metHb)	1 ml of concentrated amaranth oil ( <i>Amaranthus cruentus</i> ) obtained by vacuum CO2 cold extraction, at least 2 hours before meal, for 21 consecutive days	↑ HRV in both groups ↓ HbO <sub>2</sub> percentage ↑ HbCO, HbS and MetHb Moderate correlation of HRV with some aerobic metabolism and oxidative damage markers was found

**Table 1.**  
Summary of studies that used Amaranth in physically active population. Search was conducted through Google Scholar and PubMed search engines.

Another mechanism in which amaranth can contribute to aerobic performance is through participation in the nitric oxide pathway, thanks to its nitrate and nitrite composition. Dietary nitrate is firstly converted to nitrite by the oral microflora [17, 18] and then can enter into the system. A study with healthy human subjects supplemented with 2 g of amaranth extract showed a significant increase in plasma nitrate and nitrite, lasting up to 8 hours after administration [19].

Nitric oxide (NO) is important in physiological processes that may enhance exercise performance [20–22]. Some studies showed amaranth extract capacity to increase nitrite and nitrate in plasma, which is expected to increase the production of nitric oxide [19] (previously seen in cell cultures [23] and animal models [24]). Nitric oxide has been linked to improved endurance performance by reducing the oxygen cost during exercise, increasing the efficiency of energy production [25] (through an increase in mitochondrial efficiency [26] and ATP turnover functions [27]). Nitric oxide dietary supplements are widely consumed among athletes, and therefore future employment of amaranth extracts in new powder sports products (like pre-workouts) could be enticing if its efficacy is clinically validated.

### *3.1.2 Amaranth and carbohydrate metabolism*

Amaranth components and ingredients can be used both as an ergogenic aid or a healthy food depending on the glycaemic index (GI) (the speed at which glucose appears in the bloodstream). High-glycaemic index foods have been attributed to higher conversion in triglycerides (fat) but may be of interest when, after a workout, our aim is to provide a fast source of carbohydrates to quickly restore muscle glucose stores (glycogen). Previous work stated that after exercise, a transitory physiological phenomenon called “metabolic window” (also referred as “anabolic window”) occurs, so that utterly every carbohydrate (glucose) is synthesised to muscle glycogen and therefore stored the “healthy way” (not converted into fat).

A study with women showed that a snack with extruded amaranth presented a significantly higher GI (107) and insulinemic response than white bread, which can be recommended to athletes with this purpose as a post-workout snack [29]. amaranth starch has low viscosity and high solubility, greater than wheat and corn starches, with little content in resistant starch which provides a high glycaemic ingredient that can be dissolved in high concentrations in water, with high digestibility [30] which can be used efficiently in post-workout drinks to quickly restore glycogen stores. However, outside this metabolic window, the interest could be to avoid high GI foods, so they are not concomitantly synthesised as lipids, which can increase our fat stores (mainly as fat tissue under the skin).

Replacement of some traditional products with amaranth-based products can represent a healthier option, when they present a low GI, as in amaranth-based breakfast cereals. However, the format employed is the most determinant factor for its GI, showing higher values for a popped amaranth (105.2) than a regular cereal (74.1) but significantly lower for amaranth cereals (45.9) [31]. This is in line with a study with diabetic patients in which a diet with different amaranth proportions (grain, popped amaranth and amaranth-wheat flour) showed that its employment can represent a low or high GI [32]. This should be taken in consideration by athletes to decide when to use each product.

Apart from amaranth nutritional properties as an ingredient, amaranth oil can also improve carbohydrate metabolism as a food supplement. Previous studies with animal models showed that amaranth grain and oil can exert a hypoglycaemic effect in diabetic [33, 34] and in postmenopausal women (n = 90) when administered with amaranth leaf powder [35], showing a decreased fasting blood glucose level.



However, further research is needed to clarify its role in athletic population and if these clinical markers are translated in improvement of physical and/or biochemical markers and health condition.

### **3.2 Recovery and health maintenance of the athlete**

#### *3.2.1 Cardiovascular system health*

The main benefit associated with amaranth in cardiovascular health is through amaranth oil (very likely its high squalene content). Squalene is an unsaturated hydrocarbon ranging from 2.4 to 8.0% [36, 37], which has a similar structure to beta-carotene, and is an intermediate metabolite in the synthesis of cholesterol [38].

Animal models showed a decrease in total level of cholesterol by amaranth grain [39, 40] and also of triglycerides, LDL cholesterol [41] and VLDL [33] in addition to the fatty acid composition of the membranes of erythrocytes by amaranth oil [42] later confirmed in human patients after taking 600 mg of squalene [43]. This was also found in a study with a larger number of patients (n = 125) with amaranth oil intake from 3 to 18 ml daily (3 ml provides 100 mg of squalene) for 3 weeks, showing a dose-dependent relation with also reduced systolic and diastolic blood pressure [44].

In addition to its oil, other bioactive peptides have been found in amaranth with interesting applications to cardiovascular health. For example, globulin 11S [45] showed a potential to modulate the angiotensin system [through inhibition of angiotensin-I-converting enzyme (ACE)] tested to be resistant to gastrointestinal hydrolysis in a simulated digestive environment [46], suggesting resistance to peptic hydrolysis by human digestion. Seed protein hydrolysates also showed antihypertensive effect in vivo [47], which was also the case for amaranth oil in pulmonary arterial hypertension, although a lot of clarification is required to this respect [48]. Other in vitro studies showed that the pure peptides of amaranth inhibit cellular markers associated with the development of atherosclerosis [49] (a disease strongly linked to coronary artery disease [50]).

Apart from its regular edible parts, amaranth leaf powder also showed a different action to cardiovascular markers by increasing haemoglobin levels after administration of 9 g daily (for 3 months) in postmenopausal women [35], in line with the effect of amaranth oil, which showed a shift in haemoglobin ligands in athletes [15]. A combination of amaranth edible parts (seeds, oils and flours) as part of the diet, combined with supplements of its bioactive compounds (as per its leaves and peptides), seems to be linked to the cardiovascular physiology to some extent.

The cardiovascular system is also comprised of the blood vessels, whose inner tissue (endothelium) health status is a determinant and plays an important factor in the energetic chain, due to its role in the interchange of nutrients and oxygen. To this respect, NO is an important signalling molecule that can improve endothelial function and, as previously discussed, can be enhanced by the dietary intake of nitrite present in amaranth extracts [19]. It should be noted that NO is usually produced during exercise in absence of dietary supplementation [51] and that improvements in endothelial function are not only dependent on the cells of the arteries but also by the bone marrow whose main stimulator is also the NO produced by exercising [52]. However, the growth factor which is important during the first stages of training adaptation becomes significantly reduced after significant angiogenesis has occurred [53], and therefore, external sources (like in amaranth extracts) that can improve the endothelial function are of high value to preserve its health integrity.

### 3.2.2 Amaranth fibre as a probiotic for health and athletic performance

Probiotics and gut microbiota health are becoming popular among athletes and general population. Intense and strenuous exercise can impact gastrointestinal health status [54], while on the other hand, recent publications suggest that probiotics and gut microbiota are related to improved exercise performance [55]. Probiotics are an early field of study in sports nutrition, which can directly interact with gut microbiota and interact with the mucosal immune system and immune signalling to a variety of organs and systems [56].

Dietary fibre can serve as food for the microbiota, and to this respect, amaranth is a good source. Fibre from amaranth seeds is 78% insoluble (mainly composed of pectic substances: galacturonic acid, arabinose, galactose, xylose and glucose) and 22% soluble (mainly xyloglucans and arabinose-rich pectic polysaccharides) [57].

While clinical studies in humans are still missing to prove the effects of amaranth fibre to human microbiota, an in vitro study (emulating human intestinal ecosystems) suggests it has a probiotic effect, as it exerts some properties to specific culture of probiotic cultures [58], and a prebiotic effect (by increasing *Bifidobacterium* spp. and *Lactobacillus/Enterococcus*) as confirmed in human faeces [59] (and in vitro [60]). Apart from possible influence of amaranth fibre to microbiota, the fibre content can help to relieve common gut problems derived from the disturbance of mucosal surfaces because of prolonged or intensive exercise [61], which can be a beneficial side effect.

In addition, amaranth protein is surprisingly a source of dietary fibre with prospective applications to sports nutrition, since protein concentrate powders are widely used among athletes in protein shakes, being the main product consumed by muscle builders. The soluble dietary fibre content of amaranth protein concentrate powder is notably higher [12.90 g/100 g on dry weight (dw)] than in its flour (4.29 g/100 g dw) also with higher content of insoluble fibre (20.69 g/100 g dw against 5.54 g/100 g) [62]. This kind of amaranth product would provide large amounts of fibre for the development of gut microbiota, being also a source of vegetable protein. Clinical studies with protein and probiotics (whey + probiotics) showed improved recovery (diminished exercise-induced muscle damage) in recreationally trained males [63], as well as an improved strength restoration [64], compared to only protein (whey: probably the most widely used protein powder). Further research should be carried to study if this behaviour is mimicked when vegetal proteins (like amaranth's) are used, compared to that of milk (whey).

### 3.2.3 Antioxidant effect linked to sports

Polyphenols (the most abundant antioxidants in plants) have shown to improve athletic performance from large improvement to moderate impairment [65], while a meta-analysis showed an average improvement of 1.90% (95% CI 0.40–3.39) when at least supplemented by 7 days. They also have been blamed to impair training adaptations due to its antioxidative effects by chronic supplementation [66–68] which may hamper the development of antioxidant endogenous enzymes, while a recent review states that acute administration just before or during exercise can have beneficial effects [69]. Different parts of amaranth contain fractions of a variety of antioxidants which could be employed.

Amaranth sprouts are rich in red-coloured betacyanins—a subclass pigment belonging to the betalain family (and therefore a polyphenol)—with known antioxidant [68] and anti-inflammatory properties [70] in vitro [35], which at the same time can be applied to improve performance in sports. Betacyanins are also present

in other red-coloured plants like beet [71, 72] still requiring studies in humans to clarify its bioavailability, which suggests being low [73].

In addition, amaranth seeds and sprouts showed antioxidant flavonoids (such as rutin) and phenolics (like gallic acid, *p*-hydroxybenzoic acid and vanillic acid), while sprouting makes the appearance of *p*-coumaric and syringic acid (related with light exposure) [74].

Clinical studies showed that supplementation with amaranth leaf powder in postmenopausal women showed a significant increase in the endogenous anti-oxidant enzymes superoxide dismutase and glutathione peroxidase with decrease in the marker of oxidative stress malondialdehyde [35]. These are developed as a consequence of oxidation provoked by training in physical exercise, but more research is required to assess any possible contribution of the described anti-oxidants of amaranth ingredients to trained athletes (which presumably have developed this endogenous antioxidant system to its highest) and its link to sports performance outcomes.

#### *3.2.4 Amaranth protein, muscle recovery and improved body composition*

Early studies suggested that amino acid composition of amaranth grain protein had leucine as the limiting amino acid [75] (later confirmed in post-prandial analysis of children [76]), while it is pretty rich in lysine [62] and tryptophan [76] being suitable to combine with other cereals and provide a more balanced amino acid profile [77]. On the other hand, latter studies suggest that they are especially rich in the essential amino acids threonine and tryptophan [78] and that leucine and lysine are not the limiting amino acids in pseudo-cereals [79] (contrary to cereals) which are not true cereals from a botanical view and have nutritionally been considered as a mix of rice and beans. Protein in amaranth seeds is mainly in the embryo (instead of the endosperm) [80] showing also variation in their composition between species [81, 82]. Proteins have been appointed as the macronutrient with the most satiating effect [83] which can cause an improvement in body composition thanks to an increased satiety [84].

Protein powders which are popular among athletes can also include diverse forms like concentrates (the most common and cheap) or isolates (with virtually no carbohydrate fraction) and additionally be hydrolysed (with its proteins partly broken into smaller peptides and even amino acids). Protein hydrolysates of whey have shown to increase the insulin concentration due to a mechanism independent of gastric emptying, which can improve muscle repair by its accentuated anabolic response. Amaranth protein hydrolysate releases biopeptides which were investigated for its antihypertensive properties [47]. Studies in humans warrant further research in this field to study the same response in insulin of blood pressure in healthy and physically active population.

#### *3.2.5 Employment of amaranth for gluten-free diets and vegan diets*

Nutritional strategies through special diets are a common resource used by athletes to reduce fat mass, increase muscle mass and improve health [85]. Among these diets we find raw food diets, gluten-free diets and vegan diets [86].

To this respect, amaranth is a gluten-free alternative source of carbohydrates which can be used in gluten-free diets [87]. Amaranth can be cooked in water, extruded (appearance is of pellet form, which could be used as breakfast cereals (for a picture see [88])), toasted, incorporated into flakes [89, 90] or pastas and baked into bread, biscuits and cookies, resulting in gluten-free bakery food-stuffs [91]. Some work has been done to try to develop grain amaranth-based



nutrient-rich snack bars [92]. In Mexico, *Amaranth cruentus* cultivars are used to manufacture cookies called “Alegría” [93] (in which amaranth is used in the form of popcorn) and a commercial drink called “atole” made from milled and roasted amaranth seed, which contains large amounts of sugar. In Asia, the Indian diet finds amaranth grain as an alternative to wheat, easy to incorporate into the traditional cuisine [94].

On the other hand, amaranth protein is suitable for vegan diets of athletes, whose requirement in protein can be markedly high [86, 95] and whose nutritional value was previously discussed (see 3.2.4 *Amaranth protein, muscle recovery and improved body composition*).

## **4. Further research in the sports field of amaranth**

### **4.1 Clinical trials with amaranth and athletic population require more research**

Nutraceuticals are an area of recent research in which substances not traditionally associated with nutrients or drugs show physiological effect on the body [96]. The nutraceutical era is an emerging opportunity to find the inherent nutritional value in biology (like the functional value of amaranth discussed in this book) and requires an additional effort in its research to conduct robust and controlled clinical trials to throw light on the knowledge field. Compared to the research conducted in pharmacy, nutrition, nutraceuticals and dietary supplements in the sports nutrition field require clinical studies with good trial design and standardised products and ingredients. Uncontrolled factors, as the regular diet of subjects participating in a clinical trial, can contain one or different functional molecules that can affect the results of the trial, so its study as stand-alone substances is not so easy (and in fact is better controlled in clinical trials with patients staying in a hospital room). The following section focuses on some of the aspects that should be taken in consideration to conduct further high-quality research with amaranth.

### **4.2 Fluctuation in the nutritional value of amaranth plants should be considered**

Natural ingredients and by-products from plants are subject to a natural fluctuation in their composition and nutritional value (which can affect their functional effect) due to external factors or conversions until they are manufactured a final ingredient or product. These are some of the empirical data found in the literature when assessing the functional value of amaranth products used in clinical trials:

#### **4.2.1 External factors during crop growth**

External factors during the growth of the crop, its manufacturing and conversion until a final product is created can affect the nutritional value of amaranth:

- The lipidic content (which includes squalene) shows great variations depending on the species and genotype [37, 97] as well as the different parts of the seed [98].
- Stress and other external crop factors can influence their nutritional value [99] and in consequence its clinical outcomes.

- Enrichment with selenium of *Amaranthus cruentus* improved its anti-inflammatory effect in rat models, very likely due to its increased content in betacyanins and selenium which significantly decreased inflammatory interleukin 6 production [100].
- Sunlight exposure of *Amaranthus tricolor* showed more development of red-fleshed cultivars with greater exposure to direct sunlight, with an increased quantity of antioxidants like polyphenols [101].
- Stress like climate changes and harsh conditions can affect seed's morphology and nutritional value, which showed globulins paralogs and precursors in wild species, which could be a genetic source for improving the nutritional quality of amaranth seed [102].
- Amaranth sprouting showed increased vitamin content (significantly in biotin, folic acid and especially riboflavin (vitamin B2) and ascorbic acid (vitamin C) [103]) and decreased lipid and phytic acid content, which also is specie dependent [104].
- Amaranth contains varying amounts of vitamin C depending on the species and part used, with around 69–288 mg/100 g in the grains and 62–209 mg/100 g [105], which is suggested to have an important role in the defence of exercise-induced oxidative stress [106].

#### 4.2.2 Manufacturing and storage can also affect its nutritional and functional values

Manufacturing of amaranth to final product is also an important stage in the consecution of a dietary supplement for administration in clinical trials. Amaranth oil is extracted by squeezing amaranth grain, which can be extracted by cold extraction (considered as a gold standard in the manufacture of virgin oils)—which should be a well-controlled process to assure integrity maintenance of the oil—or solvent extraction (which requires further purification). On the other hand, amaranth flours can be converted to breakfast cereals by puffing or extruded, both processes that can affect its integrity, which is also affected by heat if, for example, used in bakery.

These are some of the facts found in literature which found how extraction technologies can affect amaranth bioactives and their functionality:

- Processing of amaranth may result in losses in protein content [107] as later confirmed by a study which abolished the sensitization potential of albumins with hypoallergenic properties in rats [108].
- Heating can affect antioxidant capacity [109], and phytic acid content has shown to be significantly decreased after undergoing a low-cost extrusion process.
- Storage conditions, as shown in a study which stored amaranth leaves, showed a reduction in beta-carotene of up to 85.0% [110] and a decrease in lysine of 4.8 and 9.6% in cracker and biscuits, respectively, after 4 months of storage [111].
- Extrusion improved anti-inflammatory effect of bioactive peptide hydrolysates studied afterwards in rat models [112], while other studies state

that antioxidant capacity (phenolic content) decreased on the favour of an improved digestibility (tested in vivo) [113].

- Puffing to convert amaranth seeds into popped amaranth (breakfast cereal-like product)—probably the most popular breakfast amaranth product [114]—can influence its nutritive value by decreasing the unsaturation of PUFAs, mainly linoleic acid (from 46.8 to 27.0%) with increased squalenes (by 15.5%) [115].
- In children, post-prandial amino acid analysis after the intake of toasted, popped or flaked amaranth consumption caused significant falls in leucine and threonine, suggesting that these were first- and second-limiting essential amino acids [76].

### 4.3 Standardised amaranth ingredients

Natural fluctuation is difficult to control, but there is an alternative solution to control the active ingredients in a dietary ingredient or supplement: Standardisation

Standardisation consists with describing a set of technical standards to guarantee constant qualitative and quantitative parameters and therefore its safety, quality and efficacy [116]. According to the same authors, these are some of the problems associated with nutraceuticals (not happening in synthetic drugs):

1. They are a mixture of many constituents.
2. The active ingredient/s is/are in some cases unknown.
3. Selective analytical methods or reference compounds (standard samples) may not be commercially available.
4. Plant materials are chemically and naturally variable.
5. Chemo-varieties and cultivars exist.
6. The source and quality of the raw material are variable.

Therefore, further research should focus on proper analysis of the composition of the product used for the trials, and pursue to guarantee, that the target compound is found in enough amounts, and not other compounds which might affect the outcomes of its experiments. Application of manufacturing processes to isolate active compounds like squalenes (by fractionation and distillation), as shown in a study [117], may allow the isolation of them for its further clinical study.

## 5. Conclusions

1. Amaranth still requires further investigation in athletes and physically active healthy subjects to study its functional effect on sports performance. The few results suggest it can activate the aerobic metabolism.
2. Clinical results with human and animal population and in vitro assays suggest physiological mechanisms that can contribute to aerobic performance through

activation of NO pathways, as a source of high and low glycaemic index carbohydrate, possible activation of gut flora (probiotic) which can be translated to athletic performance, antioxidant effect and employment of its protein and/or biopeptides.

3. Maintenance of athlete's health can be obtained through amaranth properties on cardiovascular health, as a probiotic for gut microbiota, as a source of dietary antioxidants and as a suitable ingredient for special diets like gluten-free diets (of its flours) and vegan diets (as a vegetal protein).
4. Further research should consider amaranth natural fluctuation of its nutritive value by external factors and control its composition by thorough analysis or employment of standardised products and supplements.

IntechOpen

## Author details

Torregrosa-García Antonio\* and López-Román F. Javier  
San Antonio Catholic University of Murcia (UCAM), Murcia, Spain

\*Address all correspondence to: [atorregrosa@ucam.edu](mailto:atorregrosa@ucam.edu)

## IntechOpen

© 2019 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. 



## References

- [1] Garthe I, Maughan RJ. Athletes and Supplements: Prevalence and Perspectives. *International Journal of Sport Nutrition and Exercise Metabolism*. 2018;**28**(2):126-138
- [2] Jovanov P, Đorđić V, Obradović B, Barak O, Pezo L, Marić A, et al. Prevalence, knowledge and attitudes towards using sports supplements among young athletes. *Journal of the International Society of Sports Nutrition*. 2019;**16**(1):27
- [3] Kerksick CM, Wilborn CD, Roberts MD, Smith-Ryan A, Kleiner SM, Jäger R, et al. ISSN exercise & sports nutrition review update: Research & recommendations. *Journal of the International Society of Sports Nutrition*. 2018;**15**(1):38
- [4] Kreider RB, Wilborn CD, Taylor L, Campbell B, Almada AL, Collins R, et al. ISSN exercise & sport nutrition review: research & recommendations. *Journal of the international society of sports nutrition*. Dec 2010;**7**(1):7
- [5] León-Camacho M, García-González DL, Aparicio R. A detailed and comprehensive study of amaranth (*Amaranthus cruentus* L.) oil fatty profile. *European Food Research and Technology*. 2001;**213**(4-5):349-355
- [6] Christensen JH, Christensen MS, Dyerberg J, Schmidt EB. Heart rate variability and fatty acid content of blood cell membranes: A dose-response study with n-3 fatty acids. *The American Journal of Clinical Nutrition*. 1999;**70**(3):331-337
- [7] Macartney MJ, Hingley L, Brown MA, Peoples GE, McLennan PL. Intrinsic heart rate recovery after dynamic exercise is improved with an increased omega-3 index in healthy males. *The British Journal of Nutrition*. 2014;**112**(12):1984-1992
- [8] O'Keefe JH, Abuissa H, Sastre A, Steinhaus DM, Harris WS. Effects of omega-3 fatty acids on resting heart rate, heart rate recovery after exercise, and heart rate variability in men with healed myocardial infarctions and depressed ejection fractions. *The American Journal of Cardiology*. 2006;**97**(8):1127-1130
- [9] Salvatore P, McLennan Peter L. Cardiac membrane fatty acid composition modulates myocardial oxygen consumption and postischemic recovery of contractile function. *Circulation*. 2002;**105**(19):2303-2308
- [10] DeFilippis AP, Sperling LS. Understanding omega-3's. *American Heart Journal*. 2006;**151**(3):564-570
- [11] Gammone MA, Riccioni G, Parrinello G, D'Orazio N. Omega-3 polyunsaturated fatty acids: Benefits and endpoints in sport. *Nutrients*. 2019;**11**(1):46
- [12] Yelisyeyeva O, Cherkas A, Semen K, Kaminsky D, Lutsyk A. Study of aerobic metabolism parameters and heart rate variability and their correlations in elite athletes: A modulatory effect of amaranth oil. *Clinical and Experimental Medical Journal*. 2009;**3**(2):293-307
- [13] Aubert AE, Seps B, Beckers F. Heart rate variability in athletes. *Sports Medicine*. 2003;**33**(12):889-919
- [14] De Meersman RE. Heart rate variability and aerobic fitness. *American Heart Journal*. 1993;**125**(3):726-731
- [15] Yelisyeyeva O, Semen K, Zarkovic N, Kaminsky D, Lutsyk O, Rybalchenko V. Activation of aerobic metabolism by Amaranth oil improves heart rate variability both in athletes and patients with type 2 diabetes mellitus. *Archives of Physiology and Biochemistry*. 2012;**118**(2):47-57

- [16] Cherkas A, Zarkovic K, Cipak Gasparovic A, Jaganjac M, Milkovic L, Abrahamovych O, et al. Amaranth oil reduces accumulation of 4-hydroxynonenal-histidine adducts in gastric mucosa and improves heart rate variability in duodenal peptic ulcer patients undergoing *Helicobacter pylori* eradication. *Free Radical Research*. 2018;**52**(2):135-149
- [17] Spiegelhalder B, Eisenbrand G, Preussmann R. Influence of dietary nitrate on nitrite content of human saliva: Possible relevance to in vivo formation of N-nitroso compounds. *Food and Cosmetics Toxicology*. 1976;**14**(6):545-548
- [18] Tannenbaum SR, Weisman M, Fett D. The effect of nitrate intake on nitrite formation in human saliva. *Food and Cosmetics Toxicology*. 1976;**14**(6):549-552
- [19] Subramanian D, Gupta S. Pharmacokinetic study of amaranth extract in healthy humans: A randomized trial. *Nutrition*. 2016;**32**(7):748-753
- [20] Jones AM. Dietary nitrate supplementation and exercise performance. *Sports Medicine*. 2014;**44**(1):35-45
- [21] Habib S, Ali A. Role of nitric oxide in sports nutrition. In *Nutrition and Enhanced Sports Performance*. Academic Press; 1 Jan 2019. pp. 317-325. Available from: <http://www.sciencedirect.com/science/article/pii/B9780128139226000278>
- [22] Bloomer RJ. Nitric oxide supplements for sports. *Strength & Conditioning Journal*. 2010;**32**(2):14
- [23] Barba de la Rosa AP, Barba Montoya A, Martínez-Cuevas P, Hernández-Ledesma B, León-Galván MF, De León-Rodríguez A, et al. Tryptic amaranth glutelin digests induce endothelial nitric oxide production through inhibition of ACE: Antihypertensive role of amaranth peptides. *Nitric Oxide*. 2010;**23**(2):106-111
- [24] Caselato-Sousa VM, Ozaki MR, de Almeida EA, Amaya-Farfan J. Intake of heat-expanded amaranth grain reverses endothelial dysfunction in hypercholesterolemic rabbits. *Food & Function*. 2014;**5**(12):3281-3286
- [25] Besco R, Sureda A, Tur JA, Pons A. The effect of nitric-oxide-related supplements on human performance. *Sports Medicine*. 2012;**42**(2):99-117
- [26] Larsen FJ, Schiffer TA, Borniquel S, Sahlin K, Ekblom B, Lundberg JO, et al. Dietary inorganic nitrate improves mitochondrial efficiency in humans. *Cell Metabolism*. 2011;**13**(2):149-159
- [27] Bailey SJ, Fulford J, Vanhatalo A, Winyard PG, Blackwell JR, DiMenna FJ, et al. Dietary nitrate supplementation enhances muscle contractile efficiency during knee-extensor exercise in humans. *Journal of Applied Physiology*. 2010;**109**(1):135-148
- [28] Espino-González E, Muñoz-Daw MJ, Rivera-Sosa JM, la Torre-Díaz MLD, Cano-Olivas GE, Lara-Gallegos JCD, et al. THE influence of an amaranth-based beverage on cycling performance: A pilot study. *Biotechnia*. 2018;**20**(2):31-36
- [29] Guerra-Matias AC, Arêas JAG. Glycemic and insulinemic responses in women consuming extruded amaranth (*Amaranthus cruentus* L). *Nutrition Research*. 2005;**25**(9):815-822
- [30] Capriles VD, Coelho KD, Guerra-Matias AC, Arêas JAG. Effects of processing methods on amaranth starch digestibility and predicted glycemic index. *Journal of Food Science*. 2008;**73**(7):H160-H164
- [31] Schneider I, Heinemann M, Hahn A. Comparison of glycemic index and

satiety of cereals containing amaranth. *Journal of Human Nutrition & Food Science*. 2015;**3**(5):1074

[32] Chaturvedi A, Sarojini G, Nirmala G, Nirmalamma N, Satyanarayana D. Glycemic index of grain amaranth, wheat and rice in NIDDM subjects. *Plant Food for Human Nutrition*. 1997;**50**(2):171-178

[33] Kim HK, Kim M-J, Shin D-H. Improvement of lipid profile by amaranth (*Amaranthus esculantus*) supplementation in streptozotocin-induced diabetic rats. *Annals of Nutrition and Metabolism*. 2006;**50**(3):277-281

[34] Kim HK, Kim MJ, Cho HY, Kim E-K, Shin DH. Antioxidative and anti-diabetic effects of amaranth (*Amaranthus esculantus*) in streptozotocin-induced diabetic rats. *Cell Biochemistry and Function*. 2006;**24**(3):195-199

[35] Kushwaha S, Chawla P, Kochhar A. Effect of supplementation of drumstick (*Moringa oleifera*) and amaranth (*Amaranthus tricolor*) leaves powder on antioxidant profile and oxidative status among postmenopausal women. *Journal of Food Science and Technology*. 2014;**51**(11):3464-3469

[36] Bruni R, Medici A, Guerrini A, Scalia S, Poli F, Muzzoli M, et al. Wild *Amaranthus caudatus* seed oil, a nutraceutical resource from Ecuadorian flora. *Journal of Agricultural and Food Chemistry*. 2001;**49**(11):5455-5460

[37] Rodas B, Bressani R. The oil, fatty acid and squalene content of varieties of raw and processed amaranth grain. *Archivos Latinoamericanos de Nutrición*. 2009;**59**(1):82-87

[38] Kelly GS. Squalene and its potential clinical uses. *Alternative Medicine Review*. 1999;**4**(1):29-36

[39] Shin DH, Heo HJ, Lee YJ, Kim HK. Amaranth squalene reduces serum and liver lipid levels in rats fed a cholesterol diet. *British Journal of Biomedical Science*. 2004;**61**(1):11-14

[40] Berger A, Gremaud G, Baumgartner M, Rein D, Monnard I, Kratky E, et al. Cholesterol-lowering properties of amaranth grain and oil in hamsters. *International Journal for Vitamin and Nutrition Research*. 2003;**73**(1):39-47

[41] Qureshi AA, Lehmann JW, Peterson DM. Amaranth and its oil inhibit cholesterol biosynthesis in 6-week-old female chickens. *Journal of Nutrition*. 1996;**126**(8):1972-1978

[42] Kulakova SN, Pozdniakov AL, Korf II, Karagodina ZV, Medvedev FA, Viktorova EV, et al. Amaranths oil: Peculiarities of chemical composition and influence on lipid metabolism by rats. *Voprosy Pitaniia*. 2006;**75**(3):36-42

[43] Gonor KV, Pogozeva AV, Kulakova SN, Medvedev FA, Miroshnichenko LA. The influence of diet with including amaranth oil on lipid metabolism in patients with ischemic heart disease and hyperlipoproteidemia. *Voprosy Pitaniia*. 2006;**75**(3):17-21

[44] Martirosyan DM, Miroshnichenko LA, Kulakova SN, Pogojeva AV, Zoloedov VI. Amaranth oil application for coronary heart disease and hypertension. *Lipids in Health and Disease*. 2007;**6**:1

[45] Silva-Sánchez C, de la Rosa APB, León-Galván MF, de Lumen BO, de León-Rodríguez A, de Mejía EG. Bioactive peptides in amaranth (*Amaranthus hypochondriacus*) seed. *Journal of Agricultural and Food Chemistry*. 2008;**56**(4):1233-1240

[46] Tiengo A, Faria M, Netto FM. Characterization and ACE-inhibitory



- activity of amaranth proteins. Journal of Food Science. 2009;**74**(5):H121-H126
- [47] Fritz M, Vecchi B, Rinaldi G, Añón MC. Amaranth seed protein hydrolysates have in vivo and in vitro antihypertensive activity. Food Chemistry. 2011;**126**(3):878-884
- [48] Yelisyyeva O, Semen K, Bielawska K, Biernacki M, Kaminsky D, Yavorskyi O, et al. Amaranth oil in prevalent pulmonary arterial hypertension: Changes in fatty acid panel and products of lipid peroxidation. Free Radical Biology & Medicine. 2018;**124**:576-577
- [49] Montoya-Rodríguez A, de Mejía EG. Pure peptides from amaranth (*Amaranthus hypochondriacus*) proteins inhibit LOX-1 receptor and cellular markers associated with atherosclerosis development in vitro. Foodservice Research International. 2015;**77**:204-214
- [50] Hansson GK. Inflammation, atherosclerosis, and coronary artery disease. The New England Journal of Medicine. 2005;**352**(16):1685-1695
- [51] Maiorana A, O'Driscoll G, Taylor R, Green D. Exercise and the nitric oxide vasodilator system. Sports Medicine. 2003;**33**(14):1013-1035
- [52] Haram PM. Adaptation of endothelium to exercise training: Insights from experimental studies. Frontiers in Bioscience. 2008;**13**(13):336
- [53] Richardson RS, Wagner H, Mudaliar SRD, Saucedo E, Henry R, Wagner PD. Exercise adaptation attenuates VEGF gene expression in human skeletal muscle. American Journal of Physiology. Heart and Circulatory Physiology. 2000;**279**(2):H772-H778
- [54] Peters HPF, Vries WRD, Vanberge-Henegouwen GP, Akkermans LMA. Potential benefits and hazards of physical activity and exercise on the gastrointestinal tract. Gut. 2001;**48**(3):435-439
- [55] Pane M, Amoruso A, Deidda F, Graziano T, Allesina S, Mogna L. Gut microbiota, probiotics, and sport: From clinical evidence to agonistic performance. Journal of Clinical Gastroenterology. 2018;**52**:S46
- [56] Pyne DB, West NP, Cox AJ, Cripps AW. Probiotics supplementation for athletes—Clinical and physiological effects. European Journal of Sport Science. 2015;**15**(1):63-72
- [57] Lamothe LM, Srichuwong S, Reuhs BL, Hamaker BR. Quinoa (*Chenopodium quinoa* W.) and amaranth (*Amaranthus caudatus* L.) provide dietary fibres high in pectic substances and xyloglucans. Food Chemistry. 2015;**167**:490-496
- [58] de Albuquerque MAC, Bedani R, Vieira ADS, LeBlanc JG, Saad SMI. Supplementation with fruit and okara soybean by-products and amaranth flour increases the folate production by starter and probiotic cultures. International Journal of Food Microbiology. 2016;**236**:26-32
- [59] Gullón B, Gullón P, Tavaría FK, Yáñez R. Assessment of the prebiotic effect of quinoa and amaranth in the human intestinal ecosystem. Food & Function. 2016;**7**(9):3782-3788
- [60] Vieira ADS, Bedani R, Albuquerque MAC, Biscola V, Saad SMI. The impact of fruit and soybean by-products and amaranth on the growth of probiotic and starter microorganisms. Foodservice Research International. 2017;**97**:356-363
- [61] de Oliveira EP, Burini RC. Food-dependent, exercise-induced gastrointestinal distress. Journal of the International Society of Sports Nutrition. 2011;**8**(1):12



- [62] Escudero NL, de Arellano ML, Luco JM, Giménez MS, Mucciarelli SI. Comparison of the chemical composition and nutritional value of *Amaranthus cruentus* flour and its protein concentrate. *Plant Foods for Human Nutrition*. 2004;**59**(1):15-21
- [63] Jäger R, Shields KA, Lowery RP, Souza EOD, Partl JM, Hollmer C, et al. Probiotic *Bacillus coagulans* GBI-30, 6086 reduces exercise-induced muscle damage and increases recovery. *PeerJ*. 2016;**4**:e2276
- [64] Jäger R, Purpura M, Stone JD, Turner SM, Anzalone AJ, Eimerbrink MJ, et al. Probiotic *Streptococcus thermophilus* FP4 and *Bifidobacterium breve* BR03 supplementation attenuates performance and range-of-motion decrements following muscle damaging exercise. *Nutrients*. 2016;**8**(10):642
- [65] Braakhuis AJ, Hopkins WG. Impact of dietary antioxidants on sport performance: A review. *Sports Medicine*. 2015;**45**(7):939-955
- [66] Draeger C, Naves A, Marques N, Baptistella A, Carnauba R, Paschoal V, et al. Controversies of antioxidant vitamins supplementation in exercise: Ergogenic or ergolytic effects in humans? *Journal of the International Society of Sports Nutrition*. 2014;**11**(1):4
- [67] Gomez-Cabrera MC, Ristow M, Viña J. Antioxidant supplements in exercise: Worse than useless? *American Journal of Physiology. Endocrinology and Metabolism*. 2012;**302**(4):E476-E477
- [68] Gomez-Cabrera MC, Salvador-Pascual A, Cabo H, Ferrando B, Viña J. Redox modulation of mitochondriogenesis in exercise. Does antioxidant supplementation blunt the benefits of exercise training? *Free Radical Biology & Medicine*. 2015;**86**:37-46
- [69] Pastor R, Tur JA. Antioxidant supplementation and adaptive response to training: A systematic review. *Current Pharmaceutical Design*. 2019;**25**(16):1889-1912. ISSN: 1381-6128/1873-4286. Available from: <http://europepmc.org/abstract/med/31267859>
- [70] Kanner J, Harel S, Granit R. Betalains A new class of dietary cationized antioxidants. *Journal of Agricultural and Food Chemistry*. 2001;**49**(11):5178-5185
- [71] Escribano J, Pedreño MA, García-Carmona F, Muñoz R. Characterization of the antiradical activity of betalains from *Beta vulgaris* L. roots. *Phytochemical Analysis*. 1998;**9**(3):124-127
- [72] Pedreño MA, Escribano J. Studying the oxidation and the antiradical activity of betalain from beetroot. *Journal of Biological Education*. 2000;**35**(1):49-51
- [73] Frank T, Stintzing FC, Carle R, Bitsch I, Quaas D, Straß G, et al. Urinary pharmacokinetics of betalains following consumption of red beet juice in healthy humans. *Pharmacological Research*. 2005;**52**(4):290-297
- [74] Paśko P, Sajewicz M, Gorinstein S, Zachwieja Z. Analysis of selected phenolic acids and flavonoids in *Amaranthus cruentus* and *Chenopodium quinoa* seeds and sprouts by HPLC. *Acta Chromatographica*. 2008;**20**(4):661-672
- [75] Becker R, Wheeler EL, Lorenz K, Stafford AE, Grosjean OK, Betschart AA, et al. A compositional study of amaranth grain. *Journal of Food Science*. 1981;**46**(4):1175-1180
- [76] Graham GG, Lembcke J, Morales E. Post-prandial plasma aminograms in the assessment of protein quality for young children: Maize and grain amaranth, alone and combined. *European Journal of Clinical Nutrition*. 1990;**44**(1):35-43

- [77] Pedersen B, Kalinowski LS, Eggum BO. The nutritive value of amaranth grain (*Amaranthus caudatus*). Plant Food for Human Nutrition. 1987;**36**(4):309-324
- [78] Nimbalkar MS, Pai SR, Pawar NV, Oulkar D, Dixit GB. Free amino acid profiling in grain amaranth using LC–MS/MS. Food Chemistry. 2012;**134**(4):2565-2569
- [79] Mota C, Santos M, Mauro R, Samman N, Matos AS, Torres D, et al. Protein content and amino acids profile of pseudocereals. Food Chemistry. 2016;**193**:55-61
- [80] Bressani R. The proteins of grain amaranth. Food Reviews International. 1989;**5**(1):13-38
- [81] Juan R, Pastor J, Alaiz M, Megías C, Vioque J. Seed protein characterisation of eleven species of *Amaranthus*. Grasas y Aceites. 2007;**58**(1):49-55
- [82] Pisarikova B, Kracmar S, Herzig I. Amino acid contents and biological value of protein in various amaranth species. Czech Journal of Animal Science. 2005;**50**:169-174. Available from: <http://agris.fao.org/agris-search/search.do?recordID=CZ2005000610>
- [83] Morell P, Fiszman S. Revisiting the role of protein-induced satiation and satiety. Food Hydrocolloids. 2017;**68**:199-210
- [84] Weigle DS, Breen PA, Matthys CC, Callahan HS, Meeuws KE, Burden VR, et al. A high-protein diet induces sustained reductions in appetite, ad libitum caloric intake, and body weight despite compensatory changes in diurnal plasma leptin and ghrelin concentrations. The American Journal of Clinical Nutrition. 2005;**82**(1):41-48
- [85] Rosenbloom C. Popular diets and athletes: Premises, promises, pros, and pitfalls of diets and what athletes should know about diets and sports performance. Nutrition Today. 2014;**49**(5):244
- [86] Rogerson D. Vegan diets: Practical advice for athletes and exercisers. Journal of the International Society of Sports Nutrition. 2017;**14**(1):36. DOI: 10.1186/s12970-017-0192-9
- [87] Saturni L, Ferretti G, Bacchetti T. The gluten-free diet: Safety and nutritional quality. Nutrients. 2010;**2**(1):16-34
- [88] Gearhart CM, Rosentrater KA. Extrusion processing of amaranth and quinoa. Montreal, Quebec Canada: 13. Jul 2014;**13**:1
- [89] Bressani R, Kalinowski LS, Ortiz MA, Elías LG. Nutritional evaluation of roasted, flaked and popped *A. caudatus*. Archivos Latinoamericanos de Nutrición. 1987;**37**(3):525-531
- [90] Bressani R, Martell ECMD, Godínez CMD. Protein quality evaluation of amaranth in adult humans. Plant Food for Human Nutrition. 1993;**43**(2):123-143
- [91] de la Barca AMC, Rojas-Martínez ME, Islas-Rubio AR, Cabrera-Chávez F. Gluten-free breads and cookies of raw and popped amaranth flours with attractive technological and nutritional qualities. Plant Foods for Human Nutrition. 2010;**65**(3):241-246
- [92] Sharanya Rani D. Development of grain amaranth based nutrient rich snack bars [thesis]. Acharya Ng Ranga Agricultural University; 2011
- [93] Singhal RS, Kulkarni PR. Amaranths—an underutilized resource. International Journal of Food Science and Technology. 1988;**23**(2):125-139
- [94] Dixit AA, Azar KM, Gardner CD, Palaniappan LP. Incorporation of whole, ancient grains into a modern

Asian Indian diet to reduce the burden of chronic disease. *Nutrition Reviews*. 2011;**69**(8):479-488

[95] Schoenfeld ML. Nutritional considerations for the female vegan athlete. *Strength & Conditioning Journal*. 3 Jun 2019

[96] Goncharov N, Maevsky E, Voitenko N, Novozhilov A, Kubasov I, Jenkins R, et al. Nutraceuticals in sports activities and fatigue. In *Nutraceuticals*. Academic Press; 1 Jan 2016. pp. 177-188. Available from: <http://linkinghub.elsevier.com/retrieve/pii/S09780128021477000140>

[97] He H-P, Cai Y, Sun M, Corke H. Extraction and purification of squalene from amaranthus grain. *Journal of Agricultural and Food Chemistry*. 2002;**50**(2):368-372

[98] Betschart AA, Irving DW, Shepherd AD, Saunders RM. *Amaranthus Cruentus*: Milling characteristics, distribution of nutrients within seed components, and the effects of temperature on nutritional quality. *Journal of Food Science*. 1981;**46**(4):1181-1187

[99] Espitia Rangel E. Amaranto: Ciencia y Tecnología. Celaya, Guanajuato: Instituto Nacional de Investigaciones Forestales, Agrícolas y Pecuarias, Mejoramiento Genético de Cereales (Amaranto, Avena y Trigo): Sistema Nacional de Recursos Fitogenéticos para la Alimentación y la Agricultura; 2012

[100] Tyszka-Czochara M, Pasko P, Zagrodzki P, Gajdzik E, Wietecha-Posluszny R, Gorinstein S. Selenium supplementation of amaranth sprouts influences betacyanin content and improves anti-inflammatory properties via NFκB in Murine RAW 264.7 macrophages. *Biological Trace Element Research*. 2016;**169**(2):320-330

[101] Khandaker L, Ali MB, Oba S. Total polyphenol and antioxidant activity of red amaranth (*Amaranthus tricolor* L.) as affected by different sunlight level. *Journal of the Japanese Society for Horticultural Science*. 2008;**77**(4):395-401

[102] Bojórquez-Velázquez E, Barrera-Pacheco A, Espitia-Rangel E, Herrera-Estrella A, Barba de la Rosa AP. Protein analysis reveals differential accumulation of late embryogenesis abundant and storage proteins in seeds of wild and cultivated amaranth species. *BMC Plant Biology*. 2019;**19**(1):59

[103] De Ruiz AC, Bressani R. Effect of germination on the chemical composition and nutritive value of amaranth grain. *Cereal Chemistry*. 1990;**67**(6):519-522

[104] Cornejo F, Novillo G, Villacrés E, Rosell CM. Evaluation of the physicochemical and nutritional changes in two amaranth species (*Amaranthus quitensis* and *Amaranthus caudatus*) after germination. *Food Research International*. 2019;**121**:933-939

[105] Prakash D, Joshi BD, Pal M. Vitamin C in leaves and seed oil composition of the *Amaranthus* species. *International Journal of Food Sciences and Nutrition*. 1995;**46**(1):47-51

[106] Tauler P, Aguiló A, Gimeno I, Fuentespina E, Tur JA, Pons A. Influence of vitamin C diet supplementation on endogenous antioxidant defences during exhaustive exercise. *Pflügers Archiv - European Journal of Physiology*. 2003;**446**(6):658-664

[107] Marcílio R, Amaya-Farfan J, Silva M, Spehar C. Avaliação da farinha de amaranto na elaboração de biscoito sem glúten do tipo cookie.

Brazilian Journal of Food Technology.  
2005;8(2):175-181

[108] Cárdenas-Torres FI, Reyes-Moreno C, de Jesús Vergara-Jiménez M, Cuevas-Rodríguez EO, Milán-Carrillo J, Gutiérrez-Dorado R, et al. Assessing the sensitizing and allergenic potential of the albumin and globulin fractions from amaranth (*Amaranthus hypochondriacus*) grains before and after an extrusion process. *Medicina*. 2019;55(3):E72

[109] Queiroz YS, Manólio RS, Capriles VD, Torres EA, Areas JA. Effect of processing on the antioxidant activity of amaranth grain. *Archivos Latinoamericanos de Nutrición*. 2009;59(4):419-424

[110] Negi PS, Roy SK. Changes in  $\beta$ -carotene and ascorbic acid content of fresh amaranth and fenugreek leaves during storage by low cost technique. *Plant Foods for Human Nutrition*. 2003;58(3):225-230

[111] Hozová B, Buchtová V, Dodok L, Zemanovic J. Microbiological, nutritional and sensory aspects of stored amaranth biscuits and amaranth crackers. *Die Nahrung*. 1997;41(3):155-158

[112] Montoya-Rodríguez A, de Mejía EG, Dia VP, Reyes-Moreno C, Milán-Carrillo J. Extrusion improved the anti-inflammatory effect of amaranth (*Amaranthus hypochondriacus*) hydrolysates in LPS-induced human THP-1 macrophage-like and mouse RAW 264.7 macrophages by preventing activation of NF- $\kappa$ B signaling. *Molecular Nutrition & Food Research*. 2014;58(5):1028-1041

[113] Repo-Carrasco-Valencia R, Peña J, Kallio H, Salminen S. Dietary fiber and other functional components in two varieties of crude and extruded kiwicha (*Amaranthus caudatus*). *Journal of Cereal Science*. 2009;49(2):219-224

[114] Caselato-Sousa VM, Amaya-Farfán J. State of knowledge on amaranth grain: A comprehensive review. *Journal of Food Science*. 2012;77(4):R93-R104

[115] Singhal RS, Kulkarni PR. Effect of puffing on oil characteristics of amaranth (Rajgeera) seeds. *Journal of the American Oil Chemists' Society*. 1990;67(12):952-954

[116] Kunle. Standardization of herbal medicines—A review. *International Journal of Biodiversity and Conservation*. 2012;4(3):101-112. Available from: <http://www.academicjournals.org/ijbc/abstracts/abstracts/abstracts2012/March/Kunle%20et%20al.htm>

[117] Sun H, Wiesenborn D, Tostenson K, Gillespie J, Rayas-Duarte P. Fractionation of squalene from amaranth seed oil. *Journal of the American Oil Chemists' Society*. 1997;74(4):413-418