

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

186,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Determinantal Representations of the Core Inverse and Its Generalizations

Ivan I. Kyrchei

Abstract

Generalized inverse matrices are important objects in matrix theory. In particular, they are useful tools in solving matrix equations. The most famous generalized inverses are the Moore-Penrose inverse and the Drazin inverse. Recently, it was introduced new generalized inverse matrix, namely the core inverse, which was later extended to the core-EP inverse, the BT, DMP, and CMP inverses. In contrast to the inverse matrix that has a definitely determinantal representation in terms of cofactors, even for basic generalized inverses, there exist different determinantal representations as a result of the search of their more applicable explicit expressions. In this chapter, we give new and exclusive determinantal representations of the core inverse and its generalizations by using determinantal representations of the Moore-Penrose and Drazin inverses previously obtained by the author.

Keywords: Moore-Penrose inverse, Drazin inverse, core inverse, core-EP inverse, 2000 AMS subject classifications: 15A15, 16W10

1. Introduction

In the whole chapter, the notations \mathbb{R} and \mathbb{C} are reserved for fields of the real and complex numbers, respectively. $\mathbb{C}^{m \times n}$ stands for the set of all $m \times n$ matrices over \mathbb{C} . $\mathbb{C}_r^{m \times n}$ determines its subset of matrices with a rank r . For $\mathbf{A} \in \mathbb{C}^{m \times n}$, the symbols \mathbf{A}^* and $\text{rk}(\mathbf{A})$ specify the conjugate transpose and the rank of \mathbf{A} , respectively, $|\mathbf{A}|$ or $\det \mathbf{A}$ stands for its determinant. A matrix $\mathbf{A} \in \mathbb{C}^{n \times n}$ is Hermitian if $\mathbf{A}^* = \mathbf{A}$.

\mathbf{A}^\dagger means the Moore-Penrose inverse of $\mathbf{A} \in \mathbb{C}^{n \times m}$, i.e., the exclusive matrix \mathbf{X} satisfying the following four equations:

$$\mathbf{AXA} = \mathbf{A} \quad (1)$$

$$\mathbf{XAX} = \mathbf{X} \quad (2)$$

$$(\mathbf{AX})^* = \mathbf{AX} \quad (3)$$

$$(\mathbf{XA})^* = \mathbf{XA} \quad (4)$$

For $\mathbf{A} \in \mathbb{C}^{n \times n}$ with index $\text{Ind } \mathbf{A} = k$, i.e., the smallest positive number such that $\text{rk}(\mathbf{A}^{k+1}) = \text{rk}(\mathbf{A}^k)$, the Drazin inverse of \mathbf{A} , denoted by \mathbf{A}^d , is called the unique matrix \mathbf{X} that satisfies Eq. (2) and the following equations,

$$\mathbf{A}\mathbf{X} = \mathbf{X}\mathbf{A}; \quad (5)$$

$$\mathbf{X}\mathbf{A}^{k+1} = \mathbf{A}^k \quad (6)$$

$$\mathbf{A}^{k+1}\mathbf{X} = \mathbf{A}^k. \quad (7)$$

In particular, if $\text{Ind } \mathbf{A} = 1$, then the matrix \mathbf{X} is called *the group inverse*, and it is denoted by $\mathbf{X} = \mathbf{A}^\#$. If $\text{Ind } \mathbf{A} = 0$, then \mathbf{A} is nonsingular and $\mathbf{A}^d = \mathbf{A}^\dagger = \mathbf{A}^{-1}$.

It is evident that if the condition (5) is fulfilled, then (6) and (7) are equivalent. We put both these conditions because they will be used below independently of each other and without the obligatory fulfillment of (5).

A matrix \mathbf{A} satisfying the conditions (i), (j), ... is called an $\{i, j, \dots\}$ -inverse of \mathbf{A} , and is denoted by $\mathbf{A}^{(i,j,\dots)}$. The set of matrices $\mathbf{A}^{(i,j,\dots)}$ is denoted $\mathbf{A}\{i, j, \dots\}$. In particular, $\mathbf{A}^{(1)}$ is called the inner inverse, $\mathbf{A}^{(2)}$ is called the outer inverse, $\mathbf{A}^{(1,2)}$ is called the reflexive inverse, $\mathbf{A}^{(1,2,3,4)}$ is the Moore-Penrose inverse, etc.

For an arbitrary matrix $\mathbf{A} \in \mathbb{C}^{m \times n}$, we denote by

- $\mathcal{N}(\mathbf{A}) = \{\mathbf{x} \in \mathbb{H}^{n \times 1} : \mathbf{A}\mathbf{x} = 0\}$, the kernel (or the null space) of \mathbf{A} ;
- $\mathcal{C}(\mathbf{A}) = \{\mathbf{y} \in \mathbb{H}^{m \times 1} : \mathbf{y} = \mathbf{A}\mathbf{x}, \mathbf{x} \in \mathbb{H}^{n \times 1}\}$, the column space (or the range space) of \mathbf{A} ; and
- $\mathcal{R}(\mathbf{A}) = \{\mathbf{y} \in \mathbb{H}^{1 \times n} : \mathbf{y} = \mathbf{x}\mathbf{A}, \mathbf{x} \in \mathbb{H}^{1 \times m}\}$, the row space of \mathbf{A} .

$\mathbf{P}_A := \mathbf{A}\mathbf{A}^\dagger$ and $\mathbf{Q}_A := \mathbf{A}^\dagger\mathbf{A}$ are the orthogonal projectors onto the range of \mathbf{A} and the range of \mathbf{A}^* , respectively.

The core inverse was introduced by Baksalary and Trenkler in [1]. Later, it was investigated by S. Malik in [2] and S.Z. Xu et al. in [3], among others.

Definition 1.1. [1] A matrix $\mathbf{X} \in \mathbb{C}^{n \times n}$ is called the core inverse of $\mathbf{A} \in \mathbb{C}^{n \times n}$ if it satisfies the conditions

$$\mathbf{A}\mathbf{X} = \mathbf{P}_A, \text{ and } \mathcal{C}(\mathbf{X}) = \mathcal{C}(\mathbf{A}).$$

When such matrix \mathbf{X} exists, it is denoted as \mathbf{A}^\oplus .

In 2014, the core inverse was extended to the core-EP inverse defined by K. Manjunatha Prasad and K.S. Mohana [4]. Other generalizations of the core inverse were recently introduced for $n \times n$ complex matrices, namely BT inverses [5], DMP inverses [2], CMP inverses [6], etc. The characterizations, computing methods, and some applications of the core inverse and its generalizations were recently investigated in complex matrices and rings (see, e.g., [7–18]).

In contrast to the inverse matrix that has a determinantal representation in terms of cofactors, for generalized inverse matrices, there exist different determinantal representations as a result of the search of their more applicable explicit expressions (see, e.g. [19–25]). In this chapter, we get new determinantal representations of the core inverse and its generalizations using recently obtained by the author determinantal representations of the Moore-Penrose inverse and the Drazin inverse over the quaternion skew field, and over the field of complex numbers as a special case [26–34]. Note that a determinantal representation of the core-EP generalized inverse in complex matrices has been derived in [4], based on the determinantal representation of an reflexive inverse obtained in [19, 20].

The chapter is organized as follows: in Section 2, we start with preliminary introduction of determinantal representations of the Moore-Penrose inverse and the

Drazin inverse. In Section 3, we give determinantal representations of the core inverse and its generalizations, namely the right and left core inverses are established in Section 3.1, the core-EP inverses in Section 3.2, the core DMP inverse and its dual in Section 3.3, and finally the CMP inverse in Section 3.4. A numerical example to illustrate the main results is considered in Section 4. Finally, in Section 5, the conclusions are drawn.

2. Preliminaries

Let $\alpha := \{\alpha_1, \dots, \alpha_k\} \subseteq \{1, \dots, m\}$ and $\beta := \{\beta_1, \dots, \beta_k\} \subseteq \{1, \dots, n\}$ be subsets with $1 \leq k \leq \min\{m, n\}$. By \mathbf{A}_β^α , we denote a submatrix of $\mathbf{A} \in \mathbb{H}^{m \times n}$ with rows and columns indexed by α and β , respectively. Then, \mathbf{A}_α^α is a principal submatrix of \mathbf{A} with rows and columns indexed by α , and $|\mathbf{A}|_\alpha^\alpha$ is the corresponding principal minor of the determinant $|\mathbf{A}|$. Suppose that

$$L_{k,n} := \{\alpha : \alpha = (\alpha_1, \dots, \alpha_k), 1 \leq \alpha_1 < \dots < \alpha_k \leq n\}$$

stands for the collection of strictly increasing sequences of $1 \leq k \leq n$ integers chosen from $\{1, \dots, n\}$. For fixed $i \in \alpha$ and $j \in \beta$, put $I_{r,m}\{i\} := \{\alpha : \alpha \in L_{r,m}, i \in \alpha\}$ and $J_{r,n}\{j\} := \{\beta : \beta \in L_{r,n}, j \in \beta\}$.

The j th columns and the i th rows of \mathbf{A} and \mathbf{A}^* denote $\mathbf{a}_{.j}$ and $\mathbf{a}_{.j}^*$ and $\mathbf{a}_{i.}$ and $\mathbf{a}_{i.}^*$, respectively. By $\mathbf{A}_{i.}(\mathbf{b})$ and $\mathbf{A}_{.j}(\mathbf{c})$, we denote the matrices obtained from \mathbf{A} by replacing its i th row with the row \mathbf{b} , and its j th column with the column \mathbf{c} .

Theorem 2.1. [28] If $\mathbf{A} \in \mathbb{H}_r^{m \times n}$, then the Moore-Penrose inverse $\mathbf{A}^\dagger = (a_{ij}^\dagger) \in \mathbb{C}^{n \times m}$ possesses the determinantal representations

$$a_{ij}^\dagger = \frac{\sum_{\beta \in J_{r,n}\{j\}} |(\mathbf{A}^* \mathbf{A})_{.i}(\mathbf{a}_{.j}^*)|_\beta^\beta}{\sum_{\beta \in J_{r,n}} |\mathbf{A}^* \mathbf{A}|_\beta^\beta} = \quad (8)$$

$$= \frac{\sum_{\alpha \in I_{r,m}\{i\}} |(\mathbf{A} \mathbf{A}^*)_{.j}(\mathbf{a}_{i.}^*)|_\alpha^\alpha}{\sum_{\alpha \in I_{r,m}} |\mathbf{A} \mathbf{A}^*|_\alpha^\alpha}. \quad (9)$$

Remark 2.2. For an arbitrary full-rank matrix $\mathbf{A} \in \mathbb{C}_r^{m \times n}$, a row vector $\mathbf{b} \in \mathbb{H}^{1 \times m}$, and a column-vector $\mathbf{c} \in \mathbb{H}^{n \times 1}$, we put, respectively,

$$|(\mathbf{A} \mathbf{A}^*)_{i.}(\mathbf{b})| = \sum_{\alpha \in I_{m,m}\{i\}} |(\mathbf{A} \mathbf{A}^*)_{i.}(\mathbf{b})|_\alpha^\alpha, \quad i = 1, \dots, m,$$

$$|\mathbf{A} \mathbf{A}^*| = \sum_{\alpha \in I_{m,m}} |\mathbf{A} \mathbf{A}^*|_\alpha^\alpha, \quad \text{when } r = m;$$

$$|(\mathbf{A}^* \mathbf{A})_{.j}(\mathbf{c})| = \sum_{\beta \in J_{n,n}\{j\}} |(\mathbf{A}^* \mathbf{A})_{.j}(\mathbf{c})|_\beta^\beta, \quad j = 1, \dots, n,$$

$$|\mathbf{A}^* \mathbf{A}| = \sum_{\beta \in J_{n,n}} |\mathbf{A}^* \mathbf{A}|_\beta^\beta, \quad \text{when } r = n.$$

Corollary 2.3. [21] Let $\mathbf{A} \in \mathbb{C}_r^{m \times n}$. Then, the following determinantal representations can be obtained

i. for the projector $\mathbf{Q}_A = (q_{ij})_{n \times n}$,

$$q_{ij} = \frac{\sum_{\beta \in J_{r,n}\{i\}} |(\mathbf{A}^* \mathbf{A})_{.i}(\dot{\mathbf{a}}_{.j})|_{\beta}^{\beta}}{\sum_{\beta \in J_{r,n}} |\mathbf{A}^* \mathbf{A}|_{\beta}^{\beta}} = \frac{\sum_{\alpha \in I_{r,n}\{j\}} |(\mathbf{A}^* \mathbf{A})_j(\dot{\mathbf{a}}_{i.})|_{\alpha}^{\alpha}}{\sum_{\alpha \in I_{r,n}} |\mathbf{A}^* \mathbf{A}|_{\alpha}^{\alpha}}, \quad (10)$$

where $\dot{\mathbf{a}}_{.j}$ is the j th column and $\dot{\mathbf{a}}_{i.}$ is the i th row of $\mathbf{A}^* \mathbf{A}$; and

ii. for the projector $\mathbf{P}_A = (p_{ij})_{m \times m}$,

$$p_{ij} = \frac{\sum_{\alpha \in I_{r,m}\{j\}} |(\mathbf{A} \mathbf{A}^*)_{.j}(\ddot{\mathbf{a}}_{i.})|_{\alpha}^{\alpha}}{\sum_{\alpha \in I_{r,m}} |\mathbf{A} \mathbf{A}^*|_{\alpha}^{\alpha}} = \frac{\sum_{\beta \in J_{r,m}\{i\}} |(\mathbf{A} \mathbf{A}^*)_i(\ddot{\mathbf{a}}_{.j})|_{\beta}^{\beta}}{\sum_{\beta \in J_{r,m}} |\mathbf{A} \mathbf{A}^*|_{\beta}^{\beta}}, \quad (11)$$

where $\ddot{\mathbf{a}}_{i.}$ is the i th row and $\ddot{\mathbf{a}}_{.j}$ is the j th column of $\mathbf{A} \mathbf{A}^*$.

The following lemma gives determinantal representations of the Drazin inverse in complex matrices.

Lemma 2.4. [21] *Let $\mathbf{A} \in \mathbb{C}^{n \times n}$ with $\text{Ind } \mathbf{A} = k$ and $\text{rk } \mathbf{A}^{k+1} = \text{rk } \mathbf{A}^k = r$. Then, the determinantal representations of the Drazin inverse $\mathbf{A}^d = (a_{ij}^d) \in \mathbb{C}^{n \times n}$ are*

$$a_{ij}^d = \frac{\sum_{\beta \in J_{r,n}\{i\}} |(\mathbf{A}^{k+1})_{.i}(\mathbf{a}_{.j}^{(k)})|_{\beta}^{\beta}}{\sum_{\beta \in J_{r,n}} |\mathbf{A}^{k+1}|_{\beta}^{\beta}} = \quad (12)$$

$$= \frac{\sum_{\alpha \in I_{r,n}\{j\}} |(\mathbf{A}^{k+1})_j(\mathbf{a}_{i.}^{(k)})|_{\alpha}^{\alpha}}{\sum_{\alpha \in I_{r,n}} |\mathbf{A}^{k+1}|_{\alpha}^{\alpha}}, \quad (13)$$

where $\mathbf{a}_{i.}^{(k)}$ is the i th row and $\mathbf{a}_{.j}^{(k)}$ is the j th column of \mathbf{A}^k .

Corollary 2.5. [21] *Let $\mathbf{A} \in \mathbb{C}^{n \times n}$ with $\text{Ind } \mathbf{A} = 1$ and $\text{rk } \mathbf{A}^2 = \text{rk } \mathbf{A} = r$. Then, the determinantal representations of the group inverse $\mathbf{A}^{\#} = (a_{ij}^{\#}) \in \mathbb{C}^{n \times n}$ are*

$$a_{ij}^{\#} = \frac{\sum_{\beta \in J_{r,n}\{i\}} |(\mathbf{A}^2)_{.i}(\mathbf{a}_{.j})|_{\beta}^{\beta}}{\sum_{\beta \in J_{r,n}} |\mathbf{A}^2|_{\beta}^{\beta}} = \frac{\sum_{\alpha \in I_{r,n}\{j\}} |(\mathbf{A}^2)_j(\mathbf{a}_{i.})|_{\alpha}^{\alpha}}{\sum_{\alpha \in I_{r,n}} |\mathbf{A}^2|_{\alpha}^{\alpha}}. \quad (14)$$

3. Determinantal representations of the core inverse and its generalizations

3.1 Determinantal representations of the core inverses

Together with the core inverse in [35], the dual core inverse was to be introduced. Since the both these core inverses are equipollent and they are different only in the position relative to the inducing matrix \mathbf{A} , we propose called them as the right and left core inverses regarding to their positions. So, from [1], we have the following definition that is equivalent to Definition 1.1.

Definition 3.1. A matrix $\mathbf{X} \in \mathbb{C}^{n \times n}$ is said to be the right core inverse of $\mathbf{A} \in \mathbb{C}^{n \times n}$ if it satisfies the conditions

$$\mathbf{A}\mathbf{X} = \mathbf{P}_A, \text{ and } \mathcal{C}(\mathbf{X}) = \mathcal{C}(\mathbf{A}).$$

When such matrix \mathbf{X} exists, it is denoted as \mathbf{A}^\oplus .

The following definition of the left core inverse can be given that is equivalent to the introduced dual core inverse [35].

Definition 3.2 A matrix $\mathbf{X} \in \mathbb{C}^{n \times n}$ is said to be the left core inverse of $\mathbf{A} \in \mathbb{C}^{n \times n}$ if it satisfies the conditions

$$\mathbf{X}\mathbf{A} = \mathbf{Q}_A, \text{ and } \mathcal{R}(\mathbf{X}) = \mathcal{R}(\mathbf{A}). \quad (15)$$

When such matrix \mathbf{X} exists, it is denoted as \mathbf{A}_\oplus .

Remark 3.3. In [35], the conditions of the dual core inverse are given as follows:

$$\mathbf{A}_\oplus \mathbf{A} = \mathbf{P}_{A^*}, \text{ and } \mathcal{C}(\mathbf{A}_\oplus) \subseteq \mathcal{C}(\mathbf{A}^*).$$

Since $\mathbf{P}_{A^*} = \mathbf{A}^* (\mathbf{A}^*)^\dagger = (\mathbf{A}^\dagger \mathbf{A})^* = \mathbf{A}^\dagger \mathbf{A} = \mathbf{Q}_A$, and $\mathcal{R}(\mathbf{A}) = \mathcal{C}(\mathbf{A}^*)$, then these conditions and (15) are analogous.

Due to [1], we introduce the following sets of quaternion matrices

$$\mathbb{C}_n^{\text{CM}} = \{\mathbf{A} \in \mathbb{C}^{n \times n} : \text{rk } \mathbf{A}^2 = \text{rk } \mathbf{A}\},$$

$$\mathbb{C}_n^{\text{EP}} = \{\mathbf{A} \in \mathbb{C}^{n \times n} : \mathbf{A}^\dagger \mathbf{A} = \mathbf{A} \mathbf{A}^\dagger\} = \{\mathcal{C}(\mathbf{A}) = \mathcal{C}(\mathbf{A}^*)\}.$$

The matrices from \mathbb{C}_n^{CM} are called group matrices or core matrices. If $\mathbf{A} \in \mathbb{C}_n^{\text{EP}}$, then clearly $\mathbf{A}^\dagger = \mathbf{A}^\#$. It is known that the core inverses of $\mathbf{A} \in \mathbb{C}^{n \times n}$ exist if and only if $\mathbf{A} \in \mathbb{C}_n^{\text{CM}}$ or $\text{Ind } \mathbf{A} = 1$. Moreover, if \mathbf{A} is nonsingular, $\text{Ind } \mathbf{A} = 0$, then its core inverses are the usual inverse. Due to [1], we have the following representations of the right and left core inverses.

Lemma 3.4. [1] Let $\mathbf{A} \in \mathbb{C}_n^{\text{CM}}$. Then,

$$\mathbf{A}^\oplus = \mathbf{A}^\# \mathbf{A} \mathbf{A}^\dagger, \quad (16)$$

$$\mathbf{A}_\oplus = \mathbf{A}^\dagger \mathbf{A} \mathbf{A}^\# \quad (17)$$

Remark 3.5. In Theorems 3.6 and 3.7, we will suppose that $\mathbf{A} \in \mathbb{C}_n^{\text{CM}}$ but $\mathbf{A} \notin \mathbb{C}_n^{\text{EP}}$. Because, if $\mathbf{A} \in \mathbb{C}_n^{\text{CM}}$ and $\mathbf{A} \in \mathbb{C}_n^{\text{EP}}$ (in particular, \mathbf{A} is Hermitian), then from Lemma 3.4 and the definitions of the Moore-Penrose and group inverses, it follows that $\mathbf{A}^\oplus = \mathbf{A}_\oplus = \mathbf{A}^\# = \mathbf{A}^\dagger$.

Theorem 3.6. Let $\mathbf{A} \in \mathbb{C}_n^{\text{CM}}$ and $\text{rk } \mathbf{A}^2 = \text{rk } \mathbf{A} = s$. Then, its right core inverse has the following determinantal representations

$$a_{ij}^{\oplus, r} = \frac{\sum_{\alpha \in I_{s,n}\{j\}} |(\mathbf{A} \mathbf{A}^*)_{j \cdot} (\mathbf{u}_i^{(1)})|_\alpha^\alpha}{\sum_{\beta \in J_{s,n}} |\mathbf{A}^2|_\beta^\beta \sum_{\alpha \in I_{s,n}} |\mathbf{A} \mathbf{A}^*|_\alpha^\alpha} = \quad (18)$$

$$= \frac{\sum_{\beta \in J_{s,n}\{i\}} |(\mathbf{A}^2)_{i \cdot} (\mathbf{u}_j^{(2)})|_\beta^\beta}{\sum_{\beta \in J_{s,n}} |\mathbf{A}^2|_\beta^\beta \sum_{\alpha \in I_{s,n}} |\mathbf{A} \mathbf{A}^*|_\alpha^\alpha}, \quad (19)$$

where

$$\mathbf{u}_i^{(1)} = \left[\sum_{\beta \in J_{s,n}\{i\}} |(\mathbf{A}^2)_{.i}(\tilde{\mathbf{a}}_f)|_{\beta}^{\beta} \right] \in \mathbb{C}^{1 \times n}, \quad f = 1, \dots, n$$

$$\mathbf{u}_j^{(2)} = \left[\sum_{\alpha \in I_{s,n}\{j\}} |(\mathbf{A}\mathbf{A}^*)_{j.}(\tilde{\mathbf{a}}_l)|_{\alpha}^{\alpha} \right] \in \mathbb{C}^{n \times 1}, \quad l = 1, \dots, n.$$

are the row and column vectors, respectively. Here $\tilde{\mathbf{a}}_f$ and $\tilde{\mathbf{a}}_l$ are the f th column and l th row of $\tilde{\mathbf{A}} := \mathbf{A}^2 \mathbf{A}^*$.

Proof. Taking into account (16), we have for $\#A$,

$$a_{ij}^{\#,r} = \sum_{l=1}^n \sum_{f=1}^n a_{il}^{\#} a_{lf} a_{jf}^{\dagger}. \quad (20)$$

By substituting (14) and (15) in (20), we obtain

$$a_{ij}^{\#,r} = \sum_{l=1}^n \frac{\sum_{f=1}^n \sum_{\beta \in J_{s,n}\{i\}} |(\mathbf{A}^2)_{.i}(\mathbf{a}_f)|_{\beta}^{\beta} a_{fl}}{\sum_{\beta \in J_{s,n}} |\mathbf{A}^2|_{\beta}^{\beta}} \frac{\sum_{\alpha \in I_{s,n}\{j\}} |(\mathbf{A}\mathbf{A}^*)_{j.}(\mathbf{a}_l^*)|_{\alpha}^{\alpha}}{\sum_{\alpha \in I_{s,n}} |\mathbf{A}\mathbf{A}^*|_{\alpha}^{\alpha}} =$$

$$\frac{\sum_{f=1}^n \sum_{l=1}^n \sum_{\beta \in J_{s,n}\{j\}} |(\mathbf{A}^2)_{.j}(\mathbf{e}_f)|_{\beta}^{\beta} \tilde{a}_{fl} \sum_{\alpha \in I_{s,n}\{j\}} |(\mathbf{A}\mathbf{A}^*)_{j.}(\mathbf{e}_l)|_{\alpha}^{\alpha}}{\sum_{\beta \in J_{s,n}} |\mathbf{A}^2|_{\beta}^{\beta} \sum_{\alpha \in I_{s,n}} |\mathbf{A}\mathbf{A}^*|_{\alpha}^{\alpha}},$$

where \mathbf{e}_l and \mathbf{e}_l are the unit column and row vectors, respectively, such that all their components are 0, except the l th components which are 1; \tilde{a}_{lf} is the (lf) th element of the matrix $\tilde{\mathbf{A}} := \mathbf{A}^2 \mathbf{A}^*$.

Let

$$u_{il}^{(1)} := \sum_{f=1}^n \sum_{\beta \in J_{s,n}\{i\}} |(\mathbf{A}^2)_{.i}(\mathbf{e}_f)|_{\beta}^{\beta} \tilde{a}_{fl} = \sum_{\beta \in J_{s,n}\{i\}} |(\mathbf{A}^2)_{.i}(\tilde{\mathbf{a}}_l)|_{\beta}^{\beta}, \quad i, l = 1, \dots, n.$$

Construct the matrix $\mathbf{U}_1 = (u_{il}^{(1)}) \in \mathbb{H}^{n \times n}$. It follows that

$$\sum_l u_{il}^{(1)} \sum_{\alpha \in I_{s,n}\{j\}} |(\mathbf{A}\mathbf{A}^*)_{j.}(\mathbf{e}_l)|_{\alpha}^{\alpha} = \sum_{\alpha \in I_{s,n}\{j\}} |(\mathbf{A}\mathbf{A}^*)_{j.}(\mathbf{u}_i^{(1)})|_{\alpha}^{\alpha},$$

where $\mathbf{u}_i^{(1)}$ is the i th row of \mathbf{U}_1 . So, we get (18). If we first consider

$$u_{if}^{(2)} := \sum_l \tilde{a}_{fl} \sum_{\alpha \in I_{s,n}\{j\}} |(\mathbf{A}\mathbf{A}^*)_{j.}(\mathbf{e}_l)|_{\alpha}^{\alpha} = \sum_{\alpha \in I_{s,n}\{j\}} |(\mathbf{A}\mathbf{A}^*)_{j.}(\tilde{\mathbf{a}}_f)|_{\alpha}^{\alpha}, \quad f, j = 1, \dots, n.$$

and construct the matrix $\mathbf{U}_2 = (u_{if}^{(2)}) \in \mathbb{H}^{n \times n}$, then from

$$\sum_{f=1}^n \sum_{\beta \in J_{s,n}\{i\}} |(\mathbf{A}^2)_{.i}(\mathbf{e}_f)|_{\beta}^{\beta} u_{if}^{(2)} = \sum_{\beta \in J_{s,n}\{i\}} |(\mathbf{A}^2)_{.i}(\mathbf{u}_f^{(2)})|_{\beta}^{\beta},$$

it follows (19). □

Taking into account (17), the following theorem on the determinantal representation of the left core inverse can be proved similarly.

Theorem 3.7. Let $\mathbf{A} \in \mathbb{C}_n^{\text{CM}}$ and $\text{rk } \mathbf{A}^2 = \text{rk } \mathbf{A} = s$. Then for its left core inverse $(\# \mathbf{A}) = (a_{ij}^{\#,l})$, we have

$$a_{ij}^{\#,l} = \frac{\sum_{\alpha \in I_{s,n}\{j\}} |(\mathbf{A}^2)_j(\mathbf{v}_i^{(1)})|_\alpha^\alpha}{\sum_{\beta \in J_{s,n}} |\mathbf{A}^* \mathbf{A}|_\beta^\beta \sum_{\alpha \in I_{s,n}} |\mathbf{A}^2|_\alpha^\alpha} = \frac{\sum_{\beta \in J_{s,n}\{i\}} |(\mathbf{A}^* \mathbf{A})_i(\mathbf{v}_j^{(2)})|_\beta^\beta}{\sum_{\beta \in J_{s,n}} |\mathbf{A}^* \mathbf{A}|_\beta^\beta \sum_{\alpha \in I_{s,n}} |\mathbf{A}^2|_\alpha^\alpha},$$

where

$$\mathbf{v}_i^{(1)} = \left[\sum_{\beta \in J_{s,n}\{i\}} |(\mathbf{A}^* \mathbf{A})_i(\bar{\mathbf{a}}_f)|_\beta^\beta \right] \in \mathbb{C}^{1 \times n}, \quad f = 1, \dots, n$$

$$\mathbf{v}_j^{(2)} = \left[\sum_{\alpha \in I_{s,n}\{j\}} |(\mathbf{A}^2)_j(\bar{\mathbf{a}}_l)|_\alpha^\alpha \right] \in \mathbb{C}^{n \times 1}, \quad l = 1, \dots, n.$$

Here $\bar{\mathbf{a}}_f$ and $\bar{\mathbf{a}}_l$ are the f th column and l th row of $\bar{\mathbf{A}} := \mathbf{A}^* \mathbf{A}^2$.

3.2 Determinantal representations of the core-EP inverses

Similar as in [4], we introduce two core-EP inverses.

Definition 3.8. A matrix $\mathbf{X} \in \mathbb{C}^{n \times n}$ is said to be the right core-EP inverse of $\mathbf{A} \in \mathbb{C}^{n \times n}$ if it satisfies the conditions

$$\mathbf{XAX} = \mathbf{A}, \text{ and } \mathcal{C}(\mathbf{X}) = \mathcal{C}(\mathbf{X}^*) = \mathcal{C}(\mathbf{A}^d).$$

It is denoted as \mathbf{A}^\oplus .

Definition 3.9. A matrix $\mathbf{X} \in \mathbb{C}^{n \times n}$ is said to be the left core-EP inverse of $\mathbf{A} \in \mathbb{C}^{n \times n}$ if it satisfies the conditions

$$\mathbf{XAX} = \mathbf{A}, \text{ and } \mathcal{R}(\mathbf{X}) = \mathcal{R}(\mathbf{X}^*) = \mathcal{R}(\mathbf{A}^d).$$

It is denoted as \mathbf{A}_\oplus .

Remark 3.10. Since $\mathcal{C}((\mathbf{A}^*)^d) = \mathcal{R}(\mathbf{A}^d)$, then the left core inverse \mathbf{A}_\oplus of $\mathbf{A} \in \mathbb{C}^{n \times n}$ is similar to the $*$ core inverse introduced in [4], and the dual core-EP inverse introduced in [35].

Due to [4], we have the following representations the core-EP inverses of $\mathbf{A} \in \mathbb{C}^{n \times n}$,

$$\mathbf{A}^\oplus = \mathbf{A}^{\{2,3,6a\}} \quad \text{and} \quad \mathcal{C}(\mathbf{A}^\oplus) \subseteq \mathcal{C}(\mathbf{A}^k),$$

$$\mathbf{A}_\oplus = \mathbf{A}^{\{2,4,6b\}} \quad \text{and} \quad \mathcal{R}(\mathbf{A}_\oplus) \subseteq \mathcal{R}(\mathbf{A}^k).$$

Thanks to [35], the following representations of the core-EP inverses will be used for their determinantal representations.

Lemma 3.11. Let $\mathbf{A} \in \mathbb{C}^{n \times n}$ and $\text{Ind } \mathbf{A} = k$. Then

$$\mathbf{A}^\oplus = \mathbf{A}^k \left(\mathbf{A}^{k+1} \right)^\dagger, \quad (21)$$

$$\mathbf{A}_\oplus = \left(\mathbf{A}^{k+1} \right)^\dagger \mathbf{A}^k. \quad (22)$$

Moreover, if $\text{Ind } \mathbf{A} = 1$, then we have the following representations of the right and left core inverses

$$\mathbf{A}^\oplus = \mathbf{A} (\mathbf{A}^2)^\dagger, \quad (23)$$

$$\mathbf{A}_\oplus = (\mathbf{A}^2)^\dagger \mathbf{A}. \quad (24)$$

Theorem 3.12. Suppose $\mathbf{A} \in \mathbb{C}^{n \times n}$, $\text{Ind } \mathbf{A} = k$, $\text{rk } \mathbf{A}^k = s$, and there exist \mathbf{A}^\oplus and \mathbf{A}_\oplus . Then $\mathbf{A}^\oplus = (a_{ij}^{\oplus, r})$ and $\mathbf{A}_\oplus = (a_{ij}^{\oplus, l})$ possess the determinantal representations, respectively,

$$a_{ij}^{\oplus, r} = \frac{\sum_{\alpha \in I_{s,n}\{j\}} \left| \left(\mathbf{A}^{k+1} \left(\mathbf{A}^{k+1} \right)^* \right)_j (\hat{\mathbf{a}}_i) \right|_\alpha^\alpha}{\sum_{\alpha \in I_{s,n}} \left| \mathbf{A}^{k+1} \left(\mathbf{A}^{k+1} \right)^* \right|_\alpha^\alpha}, \quad (25)$$

$$a_{ij}^{\oplus, l} = \frac{\sum_{\beta \in J_{s,n}\{i\}} \left| \left(\left(\mathbf{A}^{k+1} \right)^* \mathbf{A}^{k+1} \right)_i (\check{\mathbf{a}}_j) \right|_\beta^\beta}{\sum_{\beta \in J_{s,n}} \left| \left(\mathbf{A}^{k+1} \right)^* \mathbf{A}^{k+1} \right|_\beta^\beta}, \quad (26)$$

where $\hat{\mathbf{a}}_i$ is the i th row of $\hat{\mathbf{A}} = \mathbf{A}^k \left(\mathbf{A}^{k+1} \right)^*$ and $\check{\mathbf{a}}_j$ is the j th column of $\check{\mathbf{A}} = \left(\mathbf{A}^{k+1} \right)^* \mathbf{A}^k$.

Proof. Consider $\left(\mathbf{A}^{k+1} \right)^\dagger = (a_{ij}^{(k+1, \dagger)})$ and $\mathbf{A}^k = (a_{ij}^{(k)})$. By (21),

$$a_{ij}^{\oplus, r} = \sum_{t=1}^n a_{it}^{(k)} a_{tj}^{(k+1, \dagger)}.$$

Taking into account (9) for the determinantal representation of $\left(\mathbf{A}^{k+1} \right)^\dagger$, we get

$$a_{ij}^{\oplus, r} = \sum_{t=1}^n a_{it}^{(k)} \frac{\sum_{\alpha \in I_{s,n}\{j\}} \left| \left(\mathbf{A}^{k+1} \left(\mathbf{A}^{k+1} \right)^* \right)_j (\mathbf{a}_t^{(k+1, *)}) \right|_\alpha^\alpha}{\sum_{\alpha \in I_{r,n}} \left| \mathbf{A}^{k+1} \left(\mathbf{A}^{k+1} \right)^* \right|_\alpha^\alpha},$$

where $\mathbf{a}_t^{(k+1, *)}$ is the t th row of $\left(\mathbf{A}^{k+1} \right)^*$. Since $\sum_{t=1}^n a_{it}^{(k)} \mathbf{a}_t^{(k+1, *)} = \hat{\mathbf{a}}_i$, then it follows (25).

The determinantal representation (26) can be obtained similarly by integrating (8) for the determinantal representation of $\left(\mathbf{A}^{k+1} \right)^\dagger$ in (22). \square

Taking into account the representations (23)-(24), we obtain the determinantal representations of the right and left core inverses that have more simpler expressions than they are obtained in Theorems 3.6 and 3.7.

Corollary 3.13. Let $\mathbf{A} \in \mathbb{C}_s^{n \times n}$, $\text{Ind } \mathbf{A} = 1$, and there exist \mathbf{A}^\oplus and \mathbf{A}_\oplus . Then $\mathbf{A}^\oplus = (a_{ij}^{\oplus, r})$ and $\mathbf{A}_\oplus = (a_{ij}^{\oplus, l})$ can be expressed as follows

$$a_{ij}^{\oplus, r} = \frac{\sum_{\alpha \in I_{s,n}\{j\}} \left| \left(\mathbf{A}^2 (\mathbf{A}^2)^* \right)_{j \cdot} (\hat{\mathbf{a}}_i) \right|_\alpha^\alpha}{\sum_{\alpha \in I_{s,n}} \left| \mathbf{A}^2 (\mathbf{A}^2)^* \right|_\alpha^\alpha},$$

$$a_{ij}^{\oplus, l} = \frac{\sum_{\beta \in J_{s,n}\{i\}} \left| \left((\mathbf{A}^2)^* \mathbf{A}^2 \right)_i (\check{\mathbf{a}}_j) \right|_\beta^\beta}{\sum_{\beta \in J_{s,n}} \left| (\mathbf{A}^2)^* \mathbf{A}^2 \right|_\beta^\beta},$$

where $\hat{\mathbf{a}}_i$ is the i th row of $\hat{\mathbf{A}} = \mathbf{A}(\mathbf{A}^2)^*$ and $\check{\mathbf{a}}_j$ is the j th column of $\check{\mathbf{A}} = (\mathbf{A}^2)^* \mathbf{A}$.

3.3 Determinantal representations of the DMP and MPD inverses

The concept of the DMP inverse in complex matrices was introduced in [2] by S. Malik and N. Thome.

Definition 3.14. [2] Suppose $\mathbf{A} \in \mathbb{C}^{n \times n}$ and $\text{Ind } \mathbf{A} = k$. A matrix $\mathbf{X} \in \mathbb{C}^{n \times n}$ is said to be the DMP inverse of \mathbf{A} if it satisfies the conditions

$$\mathbf{XAX} = \mathbf{X}, \mathbf{XA} = \mathbf{A}^d \mathbf{A}, \text{ and } \mathbf{A}^k \mathbf{X} = \mathbf{A}^k \mathbf{A}^\dagger. \quad (27)$$

It is denoted as $\mathbf{A}^{d, \dagger}$.

Due to [2], if an arbitrary matrix satisfies the system of Eq. (27), then it is unique and has the following representation

$$\mathbf{A}^{d, \dagger} = \mathbf{A}^d \mathbf{A} \mathbf{A}^\dagger. \quad (28)$$

Theorem 3.15. Let $\mathbf{A} \in \mathbb{C}_s^{n \times n}$, $\text{Ind } \mathbf{A} = k$, and $\text{rk}(\mathbf{A}^k) = s_1$. Then, its DMP inverse $\mathbf{A}^{d, \dagger} = (a_{ij}^{d, \dagger})$ has the following determinantal representations.

$$a_{ij}^{d, \dagger} = \frac{\sum_{\alpha \in I_{s,n}\{j\}} \left| (\mathbf{A} \mathbf{A}^*)_{j \cdot} (\mathbf{u}_i^{(1)}) \right|_\alpha^\alpha}{\sum_{\beta \in J_{s_1,n}} \left| \mathbf{A}^{k+1} \right|_\beta^\beta \sum_{\alpha \in I_{s,n}} \left| \mathbf{A} \mathbf{A}^* \right|_\alpha^\alpha} = \quad (29)$$

$$= \frac{\sum_{\beta \in J_{s_1,n}\{i\}} \left| (\mathbf{A}^{k+1})_i (\mathbf{u}_j^{(2)}) \right|_\beta^\beta}{\sum_{\beta \in J_{s_1,n}} \left| \mathbf{A}^{k+1} \right|_\beta^\beta \sum_{\beta \in J_{s,n}} \left| \mathbf{A} \mathbf{A}^* \right|_\beta^\beta}, \quad (30)$$

where

$$\mathbf{u}_i^{(1)} = \left[\sum_{\beta \in J_{s_1,n}\{i\}} \left| (\mathbf{A}^{k+1})_i (\tilde{\mathbf{a}}_f) \right|_\beta^\beta \right] \in \mathbb{C}^{1 \times n}, \quad f = 1, \dots, n,$$

$$\mathbf{u}_j^{(2)} = \left[\sum_{\alpha \in I_{s,n}\{j\}} \left| (\mathbf{A} \mathbf{A}^*)_{j \cdot} (\hat{\mathbf{a}}_l) \right|_\alpha^\alpha \right] \in \mathbb{C}^{n \times 1}, \quad l = 1, \dots, n.$$

Here, $\tilde{\mathbf{a}}_f$ and $\hat{\mathbf{a}}_l$ are the f th column and the l th row of $\tilde{\mathbf{A}} := \mathbf{A}^{k+1} \mathbf{A}^*$.

Proof. Taking into account (28) for $\mathbf{A}^{d,\dagger}$, we get

$$a_{ij}^{d,\dagger} = \sum_{l=1}^n \sum_{f=1}^n a_{il}^d a_{lf} a_{fj}^\dagger. \quad (31)$$

By substituting (12) and (9) for the determinantal representations of \mathbf{A}^d and \mathbf{A}^\dagger in (31), we get

$$\begin{aligned} a_{ij}^{d,\dagger} = & \sum_{l=1}^n \sum_{f=1}^n \frac{\sum_{\beta \in J_{s_1,n}\{i\}} \left| \left(\mathbf{A}^{k+1} \right)_{.i} \left(\mathbf{a}_l^{(k)} \right) \right|_\beta^\beta}{\sum_{\beta \in J_{s_1,n}} \left| \mathbf{A}^{k+1} \right|_\beta^\beta} a_{lf} \frac{\sum_{\alpha \in I_{s,n}\{j\}} \left| \left(\mathbf{A} \mathbf{A}^* \right)_j \left(\mathbf{a}_f^* \right) \right|_\alpha^\alpha}{\sum_{\alpha \in I_{s,n}} \left| \mathbf{A} \mathbf{A}^* \right|_\alpha^\alpha} = \\ & \sum_{l=1}^n \sum_{f=1}^n \frac{\sum_{\beta \in J_{s_1,n}\{i\}} \left| \left(\mathbf{A}^{k+1} \right)_{.i} \left(\mathbf{e}_l \right) \right|_\beta^\beta}{\sum_{\beta \in J_{s_1,n}} \left| \mathbf{A}^{k+1} \right|_\beta^\beta} \tilde{a}_{lf} \frac{\sum_{\alpha \in I_{s,n}\{j\}} \left| \left(\mathbf{A} \mathbf{A}^* \right)_j \left(\mathbf{e}_f \right) \right|_\alpha^\alpha}{\sum_{\alpha \in I_{s,n}} \left| \mathbf{A} \mathbf{A}^* \right|_\alpha^\alpha}, \end{aligned} \quad (32)$$

where \mathbf{e}_l and \mathbf{e}_f are the l th and f th unit column and row vectors, and \tilde{a}_{lf} is the (lf) th element of the matrix $\tilde{\mathbf{A}} = \mathbf{A}^{k+1} \mathbf{A}^*$. If we put

$$u_{if}^{(1)} := \sum_{l=1}^n \sum_{\beta \in J_{s_1,n}\{i\}} \left| \left(\mathbf{A}^{k+1} \right)_{.i} \left(\mathbf{e}_l \right) \right|_\beta^\beta \tilde{a}_{lf} = \sum_{\beta \in J_{s_1,n}\{i\}} \left| \left(\mathbf{A}^{k+1} \right)_{.i} \left(\tilde{\mathbf{a}}_f \right) \right|_\beta^\beta,$$

as the f th component of the row vector $\mathbf{u}_i^{(1)} = [u_{i1}^{(1)}, \dots, u_{in}^{(1)}]$, then from

$$\sum_{f=1}^n u_{if}^{(1)} \sum_{\alpha \in I_{s,n}\{j\}} \left| \left(\mathbf{A} \mathbf{A}^* \right)_j \left(\mathbf{e}_f \right) \right|_\alpha^\alpha = \sum_{\alpha \in I_{s,n}\{j\}} \left| \left(\mathbf{A} \mathbf{A}^* \right)_j \left(\mathbf{u}_i^{(1)} \right) \right|_\alpha^\alpha,$$

it follows (29). If we initially obtain

$$u_{lj}^{(2)} := \sum_{f=1}^n \tilde{a}_{lf} \sum_{\alpha \in I_{s,n}\{j\}} \left| \left(\mathbf{A} \mathbf{A}^* \right)_j \left(\mathbf{e}_f \right) \right|_\alpha^\alpha = \sum_{\alpha \in I_{s,n}\{j\}} \left| \left(\mathbf{A} \mathbf{A}^* \right)_j \left(\tilde{\mathbf{a}}_l \right) \right|_\alpha^\alpha,$$

as the l th component of the column vector $\mathbf{u}_j^{(2)} = [u_{1j}^{(2)}, \dots, u_{nj}^{(2)}]$, then from

$$\sum_{l=1}^n \sum_{\beta \in J_{s_1,n}\{i\}} \left| \left(\mathbf{A}^{k+1} \right)_{.i} \left(\mathbf{e}_l \right) \right|_\beta^\beta u_{lj}^{(2)} = \sum_{\beta \in J_{s_1,n}\{i\}} \left| \left(\mathbf{A}^{k+1} \right)_{.i} \left(\mathbf{u}_j^{(2)} \right) \right|_\beta^\beta,$$

it follows (30). \square

The name of the DMP inverse is in accordance with the order of using the Drazin inverse (D) and the Moore-Penrose (MP) inverse. In that connection, it would be logical to consider the following definition.

Definition 3.16. Suppose $\mathbf{A} \in \mathbb{C}^{n \times n}$ and $\text{Ind } \mathbf{A} = k$. A matrix $\mathbf{X} \in \mathbb{C}^{n \times n}$ is said to be the MPD inverse of \mathbf{A} if it satisfies the conditions

$$\mathbf{X} \mathbf{A} \mathbf{X} = \mathbf{X}, \mathbf{A} \mathbf{X} = \mathbf{A} \mathbf{A}^d, \text{ and } \mathbf{X} \mathbf{A}^k = \mathbf{A}^\dagger \mathbf{A}^k.$$

It is denoted as $\mathbf{A}^{\dagger,d}$.

The matrix $\mathbf{A}^{\dagger,d}$ is unique, and it can be represented as

$$\mathbf{A}^{\dagger,d} = \mathbf{A}^{\dagger} \mathbf{A} \mathbf{A}^d. \quad (33)$$

Theorem 3.17. Let $\mathbf{A} \in \mathbb{C}_s^{n \times n}$, $\text{Ind } \mathbf{A} = k$, and $\text{rk } \mathbf{A}^k = s_1$. Then, its MPD inverse $\mathbf{A}^{\dagger,d} = (a_{ij}^{\dagger,d})$ has the following determinantal representations

$$a_{ij}^{\dagger,d} = \frac{\sum_{\beta \in J_{s,n}\{i\}} |(\mathbf{A}^* \mathbf{A})_{.i} (\mathbf{v}_{.j}^{(1)})|_{\beta}^{\beta}}{\sum_{\beta \in J_{s,n}} |\mathbf{A}^* \mathbf{A}|_{\beta}^{\beta} \sum_{\beta \in I_{s_1,n}} |\mathbf{A}^{k+1}|_{\alpha}^{\alpha}} = \frac{\sum_{\alpha \in I_{s_1,n}\{j\}} |(\mathbf{A}^{k+1})_{j.} (\mathbf{v}_{i.}^{(2)})|_{\alpha}^{\alpha}}{\sum_{\alpha \in I_{s_1,n}} |\mathbf{A}^* \mathbf{A}|_{\beta}^{\beta} \sum_{\alpha \in I_{s,n}} |\mathbf{A}^{k+1}|_{\alpha}^{\alpha}},$$

where

$$\mathbf{v}_{.j}^{(1)} = \left[\sum_{\alpha \in I_{s_1,n}\{j\}} |(\mathbf{A}^{k+1})_{j.} (\hat{\mathbf{a}}_{l.})|_{\alpha}^{\alpha} \right] \in \mathbb{C}^{n \times 1}, \quad l = 1, \dots, n$$

$$\mathbf{v}_{i.}^{(2)} = \left[\sum_{\beta \in J_{s,n}\{i\}} |(\mathbf{A}^* \mathbf{A})_{.i} (\hat{\mathbf{a}}_{f.})|_{\beta}^{\beta} \right] \in \mathbb{C}^{1 \times n}, \quad l = 1, \dots, n.$$

Here, $\hat{\mathbf{a}}_{l.}$ and $\hat{\mathbf{a}}_{f.}$ are the l th row and the f th column of $\hat{\mathbf{A}} := \mathbf{A}^* \mathbf{A}^{k+1}$.

Proof. The proof is similar to the proof of Theorem 3.15. □

3.4 Determinantal representations of the CMP inverse

Definition 3.18. [6] Suppose $\mathbf{A} \in \mathbb{C}^{n \times n}$ has the core-nilpotent decomposition $\mathbf{A} = \mathbf{A}_1 + \mathbf{A}_2$, where $\text{Ind } \mathbf{A}_1 = \text{Ind } \mathbf{A}$, \mathbf{A}_2 is nilpotent, and $\mathbf{A}_1 \mathbf{A}_2 = \mathbf{A}_2 \mathbf{A}_1 = 0$. The CMP inverse of \mathbf{A} is called the matrix $\mathbf{A}^{c,\dagger} := \mathbf{A}^{\dagger} \mathbf{A}_1 \mathbf{A}^{\dagger}$.

Lemma 3.19. [6] Let $\mathbf{A} \in \mathbb{C}^{n \times n}$. The matrix $\mathbf{X} = \mathbf{A}^{c,\dagger}$ is the unique matrix that satisfies the following system of equations:

$$\mathbf{X} \mathbf{A} \mathbf{X} = \mathbf{X}, \mathbf{A} \mathbf{X} \mathbf{A} = \mathbf{A}_1, \mathbf{A} \mathbf{X} = \mathbf{A}_1 \mathbf{A}^{\dagger}, \text{ and } \mathbf{X} \mathbf{A} = \mathbf{A}^{\dagger} \mathbf{A}_1.$$

Moreover,

$$\mathbf{A}^{c,\dagger} = \mathbf{Q}_A \mathbf{A}^d \mathbf{P}_A. \quad (34)$$

Taking into account (34), it follows the next theorem about determinantal representations of the quaternion CMP inverse.

Theorem 3.20. Let $\mathbf{A} \in \mathbb{C}_s^{n \times n}$, $\text{Ind } \mathbf{A} = m$, and $\text{rk}(\mathbf{A}^m) = s_1$. Then, the determinantal representations of its CMP inverse $\mathbf{A}^{c,\dagger} = (a_{ij}^{c,\dagger})$ can be expressed as

$$a_{ij}^{c,\dagger} = \frac{\sum_{\beta \in J_{s,n}\{i\}} |(\mathbf{A}^* \mathbf{A})_{.i} (\mathbf{v}_{.j}^{(l)})|_{\beta}^{\beta}}{\left(\sum_{\beta \in J_{s,n}} |\mathbf{A}^* \mathbf{A}|_{\beta}^{\beta} \right)^2 \sum_{\beta \in J_{s_1,n}} |\mathbf{A}^{m+1}|_{\beta}^{\beta}} \quad (35)$$

$$a_{ij}^{c,\dagger} = \frac{\sum_{\alpha \in I_{s,n}\{j\}} |(\mathbf{A}\mathbf{A}^*)_{j.}(\mathbf{w}_{i.}^{(l)})|_{\alpha}^{\alpha}}{\left(\sum_{\alpha \in I_{s,n}} |\mathbf{A}\mathbf{A}^*|_{\alpha}^{\alpha}\right)^2 \sum_{\beta \in J_{s_1,n}} |\mathbf{A}^{m+1}|_{\beta}^{\beta}} \quad (36)$$

for all $l = 1, 2$, where

$$\mathbf{v}_{.j}^{(1)} = \left[\sum_{\alpha \in I_{s,n}\{j\}} |(\mathbf{A}\mathbf{A}^*)_{j.}(\hat{\mathbf{u}}_{t.})|_{\alpha}^{\alpha} \right] \in \mathbb{C}^{n \times 1}, t = 1, \dots, n, \quad (37)$$

$$\mathbf{w}_{i.}^{(1)} = \left[\sum_{\beta \in J_{s,n}\{i\}} |(\mathbf{A}^* \mathbf{A})_{.i}(\hat{\mathbf{u}}_{.k})|_{\beta}^{\beta} \right] \in \mathbb{C}^{1 \times n}, k = 1, \dots, n, \quad (38)$$

$$\mathbf{v}_{.j}^{(2)} = \left[\sum_{\alpha \in I_{s,n}\{j\}} |(\mathbf{A}^* \mathbf{A})_{j.}(\tilde{\mathbf{g}}_{t.})|_{\alpha}^{\alpha} \right] \in \mathbb{C}^{n \times 1}, t = 1, \dots, n, \quad (39)$$

$$\mathbf{w}_{i.}^{(2)} = \left[\sum_{\beta \in J_{s,n}\{i\}} |(\mathbf{A}^* \mathbf{A})_{.i}(\tilde{\mathbf{g}}_{.k})|_{\beta}^{\beta} \right] \in \mathbb{C}^{1 \times n}, k = 1, \dots, n. \quad (40)$$

Here, $\hat{\mathbf{u}}_{t.}$ is the t th row and $\hat{\mathbf{u}}_{.k}$ is the k th column of $\hat{\mathbf{U}} := \mathbf{U}\mathbf{A}\mathbf{A}^*$, $\tilde{\mathbf{g}}_{t.}$ is the t th row and $\tilde{\mathbf{g}}_{.k}$ is the k th column of $\tilde{\mathbf{G}} := \mathbf{A}^* \mathbf{A}\mathbf{G}$, and the matrices $\mathbf{U} = (u_{ij}) \in \mathbb{H}^{n \times n}$ and $\mathbf{G} = (g_{ij}) \in \mathbb{H}^{n \times n}$ are such that

$$u_{ij} = \sum_{\alpha \in I_{s_1,n}\{j\}} |(\mathbf{A}^{m+1})_{j.}(\hat{\mathbf{a}}_{i.})|_{\alpha}^{\alpha}, \quad g_{ij} = \sum_{\beta \in J_{s_1,n}\{i\}} |(\mathbf{A}^{m+1})_{.i}(\tilde{\mathbf{a}}_{.j})|_{\beta}^{\beta},$$

where $\hat{\mathbf{a}}_{i.}$ is the i th row of $\hat{\mathbf{A}} := \mathbf{A}^* \mathbf{A}^{m+1}$ and $\tilde{\mathbf{a}}_{.j}$ is the j th column of $\tilde{\mathbf{A}} := \mathbf{A}^{m+1} \mathbf{A}^*$. *Proof.* Taking into account (34), we get

$$a_{ij}^{c,\dagger} = \sum_{l=1}^n \sum_{k=1}^n q_{il}^A a_{lk}^d p_{kj}^A, \quad (41)$$

where $\mathbf{Q}_A = (q_{il}^A)$, $\mathbf{A}^d = (a_{lk}^d)$, and $\mathbf{P}_A = (p_{kj}^A)$.

a. Taking into account the expressions (13), (10), and (11) for the determinantal representations of \mathbf{A}^d , \mathbf{Q}_A , and \mathbf{P}_A , respectively, we have

$$a_{ij}^{c,\dagger} = \sum_l \sum_t \frac{\sum_{\beta \in J_{s,n}\{i\}} |(\mathbf{A}^* \mathbf{A})_{.i}(\hat{\mathbf{a}}_{t.})|_{\beta}^{\beta}}{\sum_{\beta \in J_{s,n}} |\mathbf{A}^* \mathbf{A}|_{\beta}^{\beta}} \frac{\sum_{\alpha \in I_{s_1,n}\{l\}} |(\mathbf{A}^{m+1})_{l.}(\mathbf{a}_{t.}^{(m)})|_{\alpha}^{\alpha}}{\sum_{\alpha \in I_{s_1,n}} |\mathbf{A}^{m+1}|_{\alpha}^{\alpha}} \frac{\sum_{\alpha \in I_{s,n}\{j\}} |(\mathbf{A}\mathbf{A}^*)_{j.}(\hat{\mathbf{a}}_{l.})|_{\alpha}^{\alpha}}{\sum_{\alpha \in I_{s,n}} |\mathbf{A}\mathbf{A}^*|_{\alpha}^{\alpha}},$$

where $\hat{\mathbf{a}}_{t.}$ is the t th column of $\mathbf{A}^* \mathbf{A}$, $\hat{\mathbf{a}}_{l.}$ is the l th row of $\mathbf{A}\mathbf{A}^*$, and $\mathbf{a}_{t.}^{(m)}$ is the t th row of \mathbf{A}^m . So, it is clear that

$$a_{ij}^{c,\dagger} = \sum_l \sum_t \sum_k \frac{\sum_{\beta \in J_{s,n}\{i\}} |(\mathbf{A}^* \mathbf{A})_{.i}(\mathbf{e}_{t.})|_{\beta}^{\beta} \hat{a}_{tk} \sum_{\alpha \in I_{s_1,n}\{l\}} |(\mathbf{A}^{m+1})_{l.}(\mathbf{e}_{k.})|_{\alpha}^{\alpha} \sum_{\alpha \in I_{s,n}\{j\}} |(\mathbf{A}\mathbf{A}^*)_{j.}(\hat{\mathbf{a}}_{l.})|_{\alpha}^{\alpha}}{\sum_{\beta \in J_{s,n}} |\mathbf{A}^* \mathbf{A}|_{\beta}^{\beta} \sum_{\alpha \in I_{s_1,n}} |\mathbf{A}^{m+1}|_{\alpha}^{\alpha} \sum_{\alpha \in I_{s,n}} |\mathbf{A}\mathbf{A}^*|_{\alpha}^{\alpha}},$$

where \mathbf{e}_t is the t th unit column vector, \mathbf{e}_k is the k th row vector, and \hat{a}_{tk} is the (tk) th element of $\hat{\mathbf{A}} = \mathbf{A}^* \mathbf{A}^{m+1}$.

Denote

$$u_{tl} := \sum_k \hat{a}_{tk} \sum_{\alpha \in I_{s_1,n}\{j\}} |(\mathbf{A}^{m+1})_{l.}(\mathbf{e}_k.)|_{\alpha}^{\alpha} = \sum_{\alpha \in I_{s_1,n}\{j\}} |(\mathbf{A}^{m+1})_{l.}(\hat{\mathbf{a}}_t.)|_{\alpha}^{\alpha} \quad (42)$$

as the t th component of a column vector $\mathbf{u}_l = [u_{1l}, \dots, u_{nl}]$. Then from

$$\sum_t \sum_{\beta \in J_{s,n}\{i\}} |(\mathbf{A}^* \mathbf{A})_{.i}(\mathbf{e}_t)|_{\beta}^{\beta} u_{tl} = \sum_{\beta \in J_{s,n}\{i\}} |(\mathbf{A}^* \mathbf{A})_{.i}(\mathbf{u}_l)|_{\beta}^{\beta},$$

we have

$$a_{ij}^{c,\dagger} = \sum_l \frac{\sum_{\beta \in J_{s,n}\{i\}} |(\mathbf{A}^* \mathbf{A})_{.i}(\mathbf{u}_l)|_{\beta}^{\beta} \sum_{\alpha \in I_{s,n}\{j\}} |(\mathbf{A} \mathbf{A}^*)_{j.}(\hat{\mathbf{a}}_l.)|_{\alpha}^{\alpha}}{\sum_{\beta \in J_{r,n}} |\mathbf{A}^* \mathbf{A}|_{\beta}^{\beta} \sum_{\alpha \in I_{s_1,n}} |\mathbf{A}^{m+1}|_{\alpha}^{\alpha} \sum_{\alpha \in I_{s,n}} |\mathbf{A} \mathbf{A}^*|_{\alpha}^{\alpha}}.$$

Construct the matrix $\mathbf{U} = (u_{tl}) \in \mathbb{H}^{n \times n}$, where u_{tl} is given by (42), and denote $\hat{\mathbf{U}} := \mathbf{U} \mathbf{A} \mathbf{A}^*$. Then, taking into account that $|\mathbf{A}^* \mathbf{A}|_{\beta}^{\beta} = |\mathbf{A} \mathbf{A}^*|_{\alpha}^{\alpha}$, we have

$$a_{ij}^{c,\dagger} = \frac{\sum_t \sum_k \sum_{\beta \in J_{s,n}\{i\}} |(\mathbf{A}^* \mathbf{A})_{.i}(\mathbf{e}_t)|_{\beta}^{\beta} \hat{u}_{tk} \sum_{\alpha \in I_{s,n}\{j\}} |(\mathbf{A} \mathbf{A}^*)_{j.}(\mathbf{e}_k.)|_{\alpha}^{\alpha}}{\left(\sum_{\beta \in J_{s,n}} |\mathbf{A}^* \mathbf{A}|_{\beta}^{\beta} \right)^2 \sum_{\alpha \in I_{s_1,n}} |\mathbf{A}^{m+1}|_{\alpha}^{\alpha}}.$$

If we put that

$$v_{tj}^{(1)} := \sum_k \hat{u}_{tk} \sum_{\alpha \in I_{s,n}\{j\}} |(\mathbf{A} \mathbf{A}^*)_{j.}(\mathbf{e}_k.)|_{\alpha}^{\alpha} = \sum_{\alpha \in I_{s,n}\{j\}} |(\mathbf{A} \mathbf{A}^*)_{j.}(\hat{\mathbf{u}}_t.)|_{\alpha}^{\alpha}$$

is the t th component of a column vector $\mathbf{v}_j^{(1)} = [v_{1j}^{(1)}, \dots, v_{nj}^{(1)}]$, then from

$$\sum_t \sum_{\beta \in J_{s,n}\{i\}} |(\mathbf{A}^* \mathbf{A})_{.i}(\mathbf{e}_t)|_{\beta}^{\beta} v_{tj}^{(1)} = \sum_{\beta \in J_{s,n}\{i\}} |(\mathbf{A}^* \mathbf{A})_{.i}(\mathbf{v}_j^{(1)})|_{\beta}^{\beta},$$

it follows (35) with $\mathbf{v}_j^{(1)}$ given by (37). If we initially put

$$w_{ik}^{(1)} := \sum_t \sum_{\beta \in J_{s,n}\{i\}} |(\mathbf{A}^* \mathbf{A})_{.i}(\mathbf{e}_t)|_{\beta}^{\beta} \hat{u}_{tk} = \sum_{\beta \in J_{s,n}\{i\}} |(\mathbf{A}^* \mathbf{A})_{.i}(\hat{\mathbf{u}}_k.)|_{\beta}^{\beta}$$

as the k th component of the row vector $\mathbf{w}_i^{(1)} = [w_{i1}^{(1)}, \dots, w_{in}^{(1)}]$, then from

$$\sum_k w_{ik}^{(1)} \sum_{\alpha \in I_{s,n}\{j\}} |(\mathbf{A}^2)_{j.}(\mathbf{e}_k.)|_{\alpha}^{\alpha} = \sum_{\alpha \in I_{s,n}\{j\}} |(\mathbf{A}^2)_{j.}(\mathbf{w}_i^{(1)})|_{\alpha}^{\alpha},$$

it follows (36) with $\mathbf{w}_i^{(1)}$ given by (38).

b. By using the determinantal representation (12) for \mathbf{A}^d in (41), we have

$$a_{ij}^{c,\dagger} = \sum_k \sum_t \frac{\sum_{\beta \in J_{s,n}\{i\}} |(\mathbf{A}^* \mathbf{A})_{.i}(\dot{\mathbf{a}}_t)|_\beta^\beta \sum_{\beta \in J_{s_1,n}\{t\}} |(\mathbf{A}^{m+1})_{.t}(\mathbf{a}_{.k}^{(m)})|_\beta^\beta \sum_{\alpha \in I_{s,n}\{j\}} |(\mathbf{A}\mathbf{A}^*)_{.j}(\dot{\mathbf{a}}_{k.})|_\alpha^\alpha}{\sum_{\beta \in J_{s,n}} |\mathbf{A}^* \mathbf{A}|_\beta^\beta \sum_{\beta \in J_{s,n}} |\mathbf{A}^{m+1}|_\beta^\beta \sum_{\alpha \in I_{s,n}} |\mathbf{A}\mathbf{A}^*|_\alpha^\alpha}.$$

Therefore,

$$a_{ij}^{c,\dagger} = \sum_l \sum_k \sum_t \frac{\sum_{\beta \in J_{s,n}\{i\}} ((\mathbf{A}^* \mathbf{A})_{.i}(\dot{\mathbf{a}}_t))_\beta^\beta}{\sum_{\beta \in J_{s,n}} |\mathbf{A}^* \mathbf{A}|_\beta^\beta} \times \frac{\sum_{\beta \in J_{s_1,n}\{t\}} |(\mathbf{A}^{m+1})_{.t}(\mathbf{e}_{.k})|_\beta^\beta}{\sum_{\beta \in J_{s_1,n}} |\mathbf{A}^{m+1}|_\beta^\beta} \tilde{a}_{kl} \frac{\sum_{\alpha \in I_{s,n}\{j\}} |(\mathbf{A}\mathbf{A}^*)_{.j}(\mathbf{e}_{l.})|_\alpha^\alpha}{\sum_{\alpha \in I_{s,n}} |\mathbf{A}\mathbf{A}^*|_\alpha^\alpha}.$$

where \mathbf{e}_k is the k th unit column vector, \mathbf{e}_l is the l th unit row vector, and \tilde{a}_{kl} is the (kl) th element of $\tilde{\mathbf{A}} = \mathbf{A}^{m+1} \mathbf{A}^*$.

If we denote

$$g_{tl} := \sum_l \sum_{\beta \in J_{s_1,n}\{t\}} |(\mathbf{A}^{m+1})_{.t}(\mathbf{e}_{.k})|_\beta^\beta \tilde{a}_{kl} = \sum_{\beta \in J_{s_1,n}\{t\}} |(\mathbf{A}^{m+1})_{.t}(\tilde{\mathbf{a}}_{.l})|_\beta^\beta \quad (43)$$

as the l th component of a row vector $\mathbf{g}_t = [g_{t1}, \dots, g_{tn}]$, then

$$\sum_l g_{tl} \sum_{\alpha \in I_{s,n}\{j\}} |(\mathbf{A}\mathbf{A}^*)_{.j}(\mathbf{e}_{l.})|_\alpha^\alpha = \sum_{\alpha \in I_{s,n}\{j\}} |(\mathbf{A}\mathbf{A}^*)_{.j}(\mathbf{g}_{t.})|_\alpha^\alpha.$$

From this, it follows that

$$a_{ij}^{c,\dagger} = \sum_t \frac{\sum_{\beta \in J_{s,n}\{i\}} |(\mathbf{A}^* \mathbf{A})_{.i}(\dot{\mathbf{a}}_t)|_\beta^\beta \sum_{\alpha \in I_{s,n}\{j\}} |(\mathbf{A}\mathbf{A}^*)_{.j}(\mathbf{g}_{t.})|_\alpha^\alpha}{\sum_{\beta \in J_{s,n}} |\mathbf{A}^* \mathbf{A}|_\beta^\beta \sum_{\alpha \in I_{s_1,n}} |\mathbf{A}^{m+1}|_\alpha^\alpha \sum_{\alpha \in I_{s,n}} |\mathbf{A}\mathbf{A}^*|_\alpha^\alpha}.$$

Construct the matrix $\mathbf{G} = (g_{tl}) \in \mathbb{H}^{n \times n}$, where g_{tl} is given by (43). Denote $\tilde{\mathbf{G}} := \mathbf{A}^* \mathbf{A} \mathbf{G}$. Then,

$$a_{ij}^{c,\dagger} = \frac{\sum_t \sum_k \sum_{\beta \in J_{s,n}\{i\}} |(\mathbf{A}^* \mathbf{A})_{.i}(\mathbf{e}_{.t})|_\beta^\beta \sum_{\alpha \in I_{s,n}\{j\}} |(\mathbf{A}\mathbf{A}^*)_{.j}(\mathbf{e}_{k.})|_\alpha^\alpha}{\sum_{\beta \in J_{s,n}} (|\mathbf{A}^* \mathbf{A}|_\beta^\beta)^2 \sum_{\alpha \in I_{s_1,n}} |\mathbf{A}^{m+1}|_\alpha^\alpha}.$$

If we denote

$$v_{tj}^{(2)} := \sum_k \tilde{g}_{tk} \sum_{\alpha \in I_{s,n}\{j\}} |(\mathbf{A}\mathbf{A}^*)_{.j}(\mathbf{e}_{k.})|_\alpha^\alpha = \sum_{\alpha \in I_{s,n}\{j\}} |(\mathbf{A}\mathbf{A}^*)_{.j}(\tilde{\mathbf{g}}_{t.})|_\alpha^\alpha$$

as the t th component of a column vector $\mathbf{v}_{\cdot j}^{(2)} = [v_{1j}^{(2)}, \dots, v_{nj}^{(2)}]$, then

$$\sum_t \sum_{\beta \in J_{s,n}\{i\}} |(\mathbf{A}^* \mathbf{A})_{\cdot i}(\mathbf{e}_t)|_{\beta}^{\beta} v_{tj}^{(2)} = \sum_{\beta \in J_{s,n}\{i\}} |(\mathbf{A}^* \mathbf{A})_{\cdot i}(\mathbf{v}_{\cdot j}^{(2)})|_{\beta}^{\beta}.$$

Thus, we have (35) with $\mathbf{v}_{\cdot j}^{(2)}$ given by (39).

If, now, we denote

$$w_{ik}^{(2)} := \sum_t \sum_{\beta \in J_{s,n}\{i\}} |(\mathbf{A}^* \mathbf{A})_{\cdot i}(\mathbf{e}_t)|_{\beta}^{\beta} \tilde{g}_{tk} = \sum_{\beta \in J_{s,n}\{i\}} |(\mathbf{A}^* \mathbf{A})_{\cdot i}(\tilde{\mathbf{g}}_{\cdot k})|_{\beta}^{\beta}$$

as the k th component of a row vector $\mathbf{w}_{i\cdot}^{(2)} = [w_{i1}^{(2)}, \dots, w_{in}^{(2)}]$, then

$$\sum_k w_{ik}^{(2)} \sum_{\alpha \in I_{s,n}\{j\}} |(\mathbf{A} \mathbf{A}^*)_{j\cdot}(\mathbf{e}_k)|_{\alpha}^{\alpha} = \sum_{\alpha \in I_{s,n}\{j\}} |(\mathbf{A} \mathbf{A}^*)_{j\cdot}(\mathbf{w}_{i\cdot}^{(2)})|_{\alpha}^{\alpha}.$$

So, finally, we have (36) with $\mathbf{w}_{i\cdot}^{(2)}$ given by (40).

4. An example

Given the matrix

$$\mathbf{A} = \begin{bmatrix} 2 & 0 & 0 \\ -i & i & i \\ -i & -i & -i \end{bmatrix}.$$

Since

$$\mathbf{A} \mathbf{A}^* = \begin{bmatrix} 4 & 2i & 2i \\ -2i & 3 & -1 \\ -2i & -1 & 3 \end{bmatrix}, \quad \mathbf{A}^2 = \begin{bmatrix} 4 & 0 & 0 \\ 2-2i & 0 & 0 \\ -2-2i & 0 & 0 \end{bmatrix}, \quad \mathbf{A}^3 = \begin{bmatrix} 8 & 0 & 0 \\ 4-4i & 0 & 0 \\ -4-4i & 0 & 0 \end{bmatrix},$$

then $\text{rk } \mathbf{A} = 2$ and $\text{rk } \mathbf{A}^2 = \text{rk } \mathbf{A}^3 = 1$, and $k = \text{Ind } \mathbf{A} = 2$ and $r_1 = 1$. So, we shall find \mathbf{A}^{\oplus} and \mathbf{A}_{\oplus} by (25) and (26), respectively.

Since

$$\hat{\mathbf{A}} = \mathbf{A}^2 (\mathbf{A}^3)^* = 16 \begin{bmatrix} 2 & 1+i & -1+i \\ 1-i & 1 & i \\ -1-i & i & 1 \end{bmatrix},$$

then by (25),

$$a_{11}^{\oplus, r} = \frac{\sum_{\alpha \in I_{1,3}\{1\}} |(\mathbf{A}^3 (\mathbf{A}^3)^*)_{\cdot 1}(\hat{\mathbf{a}}_{\cdot 1})|_{\alpha}^{\alpha}}{\sum_{\alpha \in I_{1,3}} |\mathbf{A}^3 (\mathbf{A}^3)^*|_{\alpha}^{\alpha}} = \frac{1}{4}.$$

By similarly continuing, we get

$$\mathbf{A}^{\oplus} = \frac{1}{8} \begin{bmatrix} 2 & 1+\mathbf{i} & -1+\mathbf{i} \\ 1-\mathbf{i} & 1 & \mathbf{i} \\ -1-\mathbf{i} & \mathbf{i} & 1 \end{bmatrix}.$$

By analogy, due to (26), we have

$$\mathbf{A}_{\oplus} = \frac{1}{2} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}.$$

The DMP inverse $\mathbf{A}^{d,\dagger}$ can be found by Theorem 3.15. Since

$$\tilde{\mathbf{A}} = \mathbf{A}^3 \mathbf{A}^* = 4 \begin{bmatrix} 4 & 2\mathbf{i} & 2\mathbf{i} \\ 2-2\mathbf{i} & 1+\mathbf{i} & 1+\mathbf{i} \\ -2-2\mathbf{i} & 1-\mathbf{i} & 1-\mathbf{i} \end{bmatrix}.$$

and $\text{rk}(\mathbf{A}^3) = 1$, then

$$\mathbf{u}_1^{(1)} = \tilde{\mathbf{a}}_1, \quad \mathbf{u}_2^{(1)} = \tilde{\mathbf{a}}_2, \quad \mathbf{u}_3^{(1)} = \tilde{\mathbf{a}}_3.$$

Furthermore, by (29),

$$a_{11}^{d,\dagger} = \frac{\sum_{\alpha \in I_{2,3}\{1\}} |(\mathbf{A}\mathbf{A}^*)_{1.}(\mathbf{u}_1^{(1)})|_{\alpha}^{\alpha}}{\sum_{\beta \in J_{1,3}} |\mathbf{A}^3|_{\beta}^{\beta} \sum_{\alpha \in I_{2,3}} |\mathbf{A}\mathbf{A}^*|_{\alpha}^{\alpha}} = \frac{1}{192} \left(\det \begin{bmatrix} 16 & 8\mathbf{i} \\ -2\mathbf{i} & 3 \end{bmatrix} + \det \begin{bmatrix} 16 & 8\mathbf{i} \\ -2\mathbf{i} & 3 \end{bmatrix} \right) = \frac{1}{3}.$$

By similarly continuing, we get

$$\mathbf{A}^{d,\dagger} = \frac{1}{12} \begin{bmatrix} 4 & 2\mathbf{i} & 2\mathbf{i} \\ 2-2\mathbf{i} & 1+\mathbf{i} & 1+\mathbf{i} \\ -2-2\mathbf{i} & 1-\mathbf{i} & 1-\mathbf{i} \end{bmatrix}.$$

Similarly by Theorem 3.17, we get

$$\mathbf{A}^{\dagger,d} = \frac{1}{4} \begin{bmatrix} 2 & 0 & 0 \\ -\mathbf{i} & 0 & 0 \\ -\mathbf{i} & 0 & 0 \end{bmatrix}.$$

Finally, by theorem, we find the CMP inverse $\mathbf{A}^{c,\dagger} = (a_{ij}^{c,\dagger})$. Since $\text{rk} \mathbf{A}^3 = 1$, then $\mathbf{G} = \tilde{\mathbf{A}}$ and

$$\tilde{\mathbf{G}} = \mathbf{A}^* \mathbf{A} \tilde{\mathbf{A}} = 16 \begin{bmatrix} 6 & 3\mathbf{i} & 3\mathbf{i} \\ -2\mathbf{i} & 1 & 1 \\ -2\mathbf{i} & 1 & 1 \end{bmatrix}.$$

Furthermore, by (40),

$$w_{11}^{(2)} = \sum_{\beta \in J_{2,3}\{1\}} |(\mathbf{A}^* \mathbf{A})_{.1}(\tilde{\mathbf{g}}_{.1})|_{\beta}^{\beta} = \left(\det \begin{bmatrix} 6 & 0 \\ -2\mathbf{i} & 2 \end{bmatrix} + \det \begin{bmatrix} 6 & 0 \\ -2\mathbf{i} & 2 \end{bmatrix} \right) = 24.$$

By similar calculations, we get

$$\mathbf{w}_1^{(2)} = [384, 96\mathbf{i}, 96\mathbf{i}], \quad \mathbf{w}_2^{(2)} = [-192\mathbf{i}, 96, 96], \quad \mathbf{w}_3^{(2)} = [-192\mathbf{i}, 96, 06].$$

So, by (36), we get

$$\begin{aligned} a_{11}^{c,\dagger} &= \frac{\sum_{\alpha \in I_{2,3}\{1\}} |(\mathbf{A} \mathbf{A}^*)_{.1}(\mathbf{w}_1^{(2)})|_{\alpha}^{\alpha}}{\left(\sum_{\alpha \in I_{2,3}} |\mathbf{A} \mathbf{A}^*|_{\alpha}^{\alpha} \right)^2 \sum_{\beta \in J_{1,3}} |\mathbf{A}^3|_{\beta}^{\beta}} \\ &= \frac{1}{4608} \left(\det \begin{bmatrix} 384 & 192\mathbf{i} \\ -2\mathbf{i} & 3 \end{bmatrix} + \det \begin{bmatrix} 384 & 192\mathbf{i} \\ -2\mathbf{i} & 3 \end{bmatrix} \right) = \frac{1}{3}. \end{aligned}$$

By similarly continuing, we derive

$$\mathbf{A}^{c,\dagger} = \frac{1}{12} \begin{bmatrix} 4 & 2\mathbf{i} & 2\mathbf{i} \\ -2\mathbf{i} & 1 & 1 \\ -2\mathbf{i} & 1 & 1 \end{bmatrix}.$$

5. Conclusions

In this chapter, we get the direct method to find the core inverse and its generalizations that are based on their determinantal representations. New determinantal representations of the right and left core inverses, the right and left core-EP inverses, the DMP, MPD, and CMP inverses are derived.

IntechOpen


IntechOpen

Author details

Ivan I. Kyrchei
Pidstryhach Institute for Applied Problems of Mechanics and Mathematics,
NAS of Ukraine, Lviv, Ukraine

*Address all correspondence to: kyrchei@online.ua

IntechOpen

© 2019 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. 

References

- [1] Baksalary OM, Trenkler G. Core inverse of matrices. *Linear and Multilinear Algebra*. 2010;**58**:681-697
- [2] Malik S, Thome N. On a new generalized inverse for matrices of an arbitrary index. *Applied Mathematics and Computation*. 2014;**226**:575-580
- [3] Xu SZ, Chen JL, Zhang XX. New characterizations for core inverses in rings with involution. *Frontiers in Mathematics China*. 2017;**12**:231-246
- [4] Prasad KM, Mohana KS. Core EP inverse. *Linear and Multilinear Algebra*. 2014;**62**(3):792-802
- [5] Baksalary OM, Trenkler G. On a generalized core inverse. *Applied Mathematics and Computation*. 2014;**236**:450-457
- [6] Mehdipour M, Salemi A. On a new generalized inverse of matrices. *Linear and Multilinear Algebra*. 2018;**66**(5): 1046-1053
- [7] Chen JL, Zhu HH, Patrício P, Zhang YL. Characterizations and representations of core and dual core inverses. *Canadian Mathematical Bulletin*. 2017;**60**:269-282
- [8] Gao YF, Chen JL. Pseudo core inverses in rings with involution. *Communications in Algebra*. 2018;**46**: 38-50
- [9] Guterman A, Herrero A, Thome N. New matrix partial order based on spectrally orthogonal matrix decomposition. *Linear and Multilinear Algebra*. 2016;**64**(3):362-374
- [10] Ferreyra DE, Levis FE, Thome N. Maximal classes of matrices determining generalized inverses. *Applied Mathematics and Computation*. 2018;**333**:42-52
- [11] Ferreyra DE, Levis FE, Thome N. Revisiting the core EP inverse and its extension to rectangular matrices. *Quaestiones Mathematicae*. 2018;**41**(2): 265-281
- [12] Liu X, Cai N. High-order iterative methods for the DMP inverse. *Journal of Mathematics*. 2018;**8175935**:6
- [13] Ma H, Stanimirović PS. Characterizations, approximation and perturbations of the core-EP inverse. *Applied Mathematics and Computation*. 2019;**359**:404-417
- [14] Mielniczuk J. Note on the core matrix partial ordering. *Discussiones Mathematicae Probability and Statistics* 2011;**31**:71-75
- [15] Mosić D, Deng C, Ma H. On a weighted core inverse in a ring with involution. *Communications in Algebra*. 2018;**46**(6):2332-2345
- [16] Prasad KM, Raj MD. Bordering method to compute core-EP inverse. *Special Matrices*. 2018;**6**:193-200
- [17] Rakić DS, Dinčić ČN, Djordjević DS. Group, Moore-Penrose, core and dual core inverse in rings with involution. *Linear Algebra and its Applications*. 2014;**463**:115-133
- [18] Wang HX. Core-EP decomposition and its applications. *Linear Algebra and its Applications*. 2016;**508**:289-300
- [19] Bapat RB, Bhaskara Rao KPS, Prasad KM. Generalized inverses over integral domains. *Linear Algebra and its Applications*. 1990;**140**:181-196
- [20] Bhaskara Rao KPS. Generalized inverses of matrices over integral domains. *Linear Algebra and its Applications*. 1983;**49**:179-189

- [21] Kyrchei I. Analogs of the adjoint matrix for generalized inverses and corresponding Cramer rules. *Linear and Multilinear Algebra*. 2008;**56**(4): 453-469
- [22] Kyrchei I. Explicit formulas for determinantal representations of the Drazin inverse solutions of some matrix and differential matrix equations. *Applied Mathematics and Computation*. 2013;**219**:7632-7644
- [23] Kyrchei I. Cramer's rule for generalized inverse solutions. In: Kyrchei I, editor. *Advances in Linear Algebra Research*. New York: Nova Science Publ; 2015. pp. 79-132
- [24] Stanimirović PS. General determinantal representation of pseudoinverses of matrices. *Matematicki Vesnik*. 1996;**48**:1-9
- [25] Stanimirović PS, Djordjevic DS. Full-rank and determinantal representation of the Drazin inverse. *Linear Algebra and its Applications*. 2000;**311**:131-151
- [26] Kyrchei I. Determinantal representations of the Moore-Penrose inverse over the quaternion skew field. *Journal of Mathematical Sciences*. 2012; **180**(1):23-33
- [27] Kyrchei I. Determinantal representations of the Moore-Penrose inverse over the quaternion skew field and corresponding Cramer's rules. *Linear and Multilinear Algebra*. 2011; **59**(4):413-431
- [28] Kyrchei I. Determinantal representations of the Drazin inverse over the quaternion skew field with applications to some matrix equations. *Applied Mathematics and Computation*. 2014;**238**:193-207
- [29] Kyrchei I. Determinantal representations of the W-weighted Drazin inverse over the quaternion skew field. *Applied Mathematics and Computation*. 2015;**264**:453-465
- [30] Kyrchei I. Explicit determinantal representation formulas of W-weighted Drazin inverse solutions of some matrix equations over the quaternion skew field. *Mathematical Problems in Engineering*. 2016;**8673809**:13
- [31] Kyrchei I. Explicit determinantal representation formulas for the solution of the two-sided restricted quaternionic matrix equation. *Journal of Applied Mathematics and Computing*. 2018;**58** (1-2):335-365
- [32] Kyrchei I. Determinantal representations of the Drazin and W-weighted Drazin inverses over the quaternion skew field with applications. In: Griffin S, editor. *Quaternions: Theory and Applications*. New York: Nova Sci. Publ.; 2017. pp. 201-275
- [33] Kyrchei I. Weighted singular value decomposition and determinantal representations of the quaternion weighted Moore-Penrose inverse. *Applied Mathematics and Computation*. 2017;**309**:1-16
- [34] Kyrchei I. Determinantal representations of the quaternion weighted Moore-Penrose inverse and its applications. In: Baswell AR, editor. *Advances in Mathematics Research 23*. New York: Nova Science Publ; 2017. pp. 35-96
- [35] Zhou M, Chen J, Li T, Wang D. Three limit representations of the core-EP inverse. *Univerzitet u Nišu*. 2018;**32**: 5887-5894