

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

185,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.  
For more information visit [www.intechopen.com](http://www.intechopen.com)



# Aspects of Photodynamic Inactivation of Bacteria

*Faina Nakonechny and Marina Nisnevitch*

## Abstract

Increasing resistance of bacteria to antibiotics is a serious worldwide problem, and to combat resistant bacteria, new antibacterial approaches are to be developed. One alternative to traditional antibiotic therapy is photodynamic antimicrobial chemotherapy (PACT). PACT is based on excitation of photosensitizers (PS) capable of transferring the absorbed light energy to dissolved molecular oxygen causing generation of reactive oxygen species, which irreversibly damage bacterial cell components. The overall efficiency of PACT has been proven for Gram-positive and Gram-negative bacteria. The effectiveness of PACT can be increased by encapsulation of PS in liposomes providing more concentrated delivery of PS, enhanced cytotoxicity, improved pharmacokinetic properties, sustained release, and prolonged action of the PS. For continuous and reusable application, PS can be immobilized in polymers. Chemiluminescence, sonodynamic treatment, and radio-frequency irradiation allow to perform excitation of PS in the dark without external illumination, opening prospects for combating internal infections. Combination of PS with antibiotics can gain a synergistic effect, allowing in some cases to overcome the resistance of bacteria to antibiotics.

**Keywords:** photodynamic therapy (PDT), photodynamic antimicrobial chemotherapy (PACT), photosensitizer (PS), chemiluminescent antimicrobial chemotherapy (CPAT), sonodynamic antimicrobial chemotherapy (SACT), targeted drug delivery, liposomes, immobilization

## 1. Introduction

### 1.1 History of photodynamic therapy

The therapeutic properties of light were observed already in ancient Greece, Egypt, and India. However, they were not widely used for many centuries [1]. The history of modern photodynamic therapy (PDT) dates back to 1900, when Oscar Raab discovered the toxic properties of the dye acridine red on *Paramecium* spp. [2]. He and his supervisor, Hermann von Tappeiner, noticed a positive effect of illumination on the toxic activity of this dye. In his later work, von Tappeiner and his colleagues applied this approach to inactivation of bacteria [3] and to treatment of skin cancer [4]. In 1909, von Tappeiner introduced the term “Photodynamic Action” and showed that oxygen is essential for this procedure [5]. PDT has been studied and developed as an anticancer therapy for a long time and was approved by the Food and Drug Administration in the 1990s for various applications in this area of medicine [6–8]. The antimicrobial properties of this approach were unfairly

forgotten for several decades. However, interest in antibacterial PDT has been rekindled and is continuously increasing because multidrug resistance of pathogenic microorganisms has become a serious threat to public health. Photodynamic antibacterial chemotherapy (PACT) has become a promising approach for combating bacterial infections, which are resistant to modern antibiotics.

## 1.2 Photosensitizers and their mechanism of action

PACT is based on the exposure of bacteria to photosensitive compounds—photosensitizers (PSs). When a PS located in the bacteria or on the bacterial surface is exposed to light (usually visible), it transfers from its low-energy ground state to an excited singlet state. Return of the PS to its ground state is accompanied by either emission of fluorescence or transition of the PS to a longer-living, higher-energy triplet state ( $PS^*$ ) via intersystem crossing. The  $PS^*$  in turn reacts with surrounding molecules to form free radicals and hydrogen peroxide (Type I reaction) or transfers its energy to molecular oxygen to produce singlet oxygen and other highly reactive oxygen species (ROS; Type II reaction) [9, 10]. Type I and Type II reactions occur simultaneously, and the ratio at which they occur depends on both the PS type and the surrounding conditions. A detailed description of the photosensitization process can be found in the recent reviews of Castano et al. [11] and Cieplik [10]. ROSs formed in this process oxidize biomolecules, damage the cell membrane, and ultimately lead to cell death [12]. PACT usually proceeds predominantly through Type II processes. However, since Gram-negative bacteria are more susceptible to  $OH^\cdot$  radicals than to singlet oxygen, the Type I reaction may be more efficient against such microorganisms [13, 14].

## 1.3 Photosensitizers for PACT

Hundreds of compounds are currently available for mediating PDT in various areas of medicine, where some have been shown to be suitable for antimicrobial applications. PSs employed for medical uses should be a single pure compound, stable at room temperature and inexpensive. The PS must have a strong absorption peak in the visible spectrum between 600 and 900 nm and should possess a high-triplet quantum yield that will provide high production of ROS upon illumination. It should not be toxic in the dark (especially to mammalian cells), mutagenic or carcinogenic [15–18]. In addition, when talking about PACT, it is very important that the PS will display preferential association with bacteria, accumulate within the cells, or bind to the bacterial cell envelope [14, 19].

PSs can generally be assigned to several chemical classes: tetrapyrroles (which include porphyrins, chlorins, bacteriochlorins, and phthalocyanines), synthetic dyes (phenothiazinium salts, Rose Bengal, squaraines, etc.), and naturally occurring compounds (such as riboflavin or curcumin). Cyclic tetrapyrroles present the most well-known class of clinically relevant PSs used mostly for anticancer applications [20]. This structure can be found naturally in such important biomolecules such as haem, chlorophyll, and bacteriochlorophyll. Unlike other types of PSs, most tetrapyrroles (except for bacteriochlorins) are more likely to react by a Type II reaction with the creation of singlet oxygen [16], whereas bacteriochlorins act *via* a Type I mechanism. Other well-known antimicrobial agents are phenothiazinium-based synthetic dyes, including methylene blue (MB) and toluidine blue O (TBO), which also act as anticancer agents in PDT. These structures can be synthesized more easily than tetrapyrroles but possess high-dark toxicity compared to other PSs [15, 21]. Another representative of synthetic dyes, Rose Bengal (RB), has already been used successfully in antimicrobial and anticancer applications for a long


time [16]. Photodynamic active compounds isolated from plants arouse particular interest. These natural compounds include curcumin, extracted from the rhizomes of *Curcuma longa*, which was found effective in eradicating oral pathogens [22]. Another representative of this group is hypericin isolated from St. John's wort, which exhibits photodynamic activity against Gram-positive and Gram-negative bacteria. Detailed descriptions of all PS classes can be found in the reviews published by Hamblin and colleagues [15, 16].

## 2. Photosensitizer activation modes

### 2.1 Dark activity

The name photosensitizer implies the need for illumination in order to activate PS molecules and trigger their action. However, PSs possess some so-called “dark activity” even in the absence of illumination, leading to cell death in the dark [23–29]. This feature depends on the PS concentration and manifests itself in different ways for various PSs.


Shrestha demonstrated dark toxicity of RB against Gram-positive *Enterococcus faecalis*. Exposure of the cells to 10  $\mu\text{M}$  RB in the absence of illumination for 15 min led to a 0.5  $\log_{10}$  reduction in cell concentration [26]. Furthermore, a marked dark toxicity of RB against clinical isolates of Gram-negative *Pseudomonas aeruginosa* was observed by Nakonieczna [27]. Brovko compared the activity of various PSs against several types of microorganisms and noted high dark toxicity of RB, as well as of phloxine B against Gram-positive *Bacillus sp.* and *Listeria monocytogenes* (more than 5  $\log_{10}$  reduction in the bacterial concentration after 30 min of treatment with the dye) [30]. The toxicity of malachite green in the dark against the same microorganisms was very low ( $<0.1 \log_{10}$  reduction in concentration after 30 min of treatment with the dye). High concentrations ( $>500 \mu\text{g/mL}$ ) of acriflavin neutral in the absence of light were significantly toxic to *E. coli* (more than 6  $\log_{10}$  reduction in concentration after 30 min of treatment with the dye, both under illumination and



**Figure 1.** Effect of RB concentration on its cytotoxic activity. *S. aureus* cells at the initial concentration of  $10^4 \text{ CFU mL}^{-1}$  were incubated for 3 min in dark conditions at various concentrations of RB. After the incubation, bacteria were tested by viable count. Error bars present standard deviations.

in the dark). However, illumination significantly enhanced its toxic effect against other tested microorganisms [30].

In our studies, we also noted the dark toxicity of various PSs against different types of bacteria (Figures 1, 2, Table 1). Figure 1 shows the effect of various RB concentrations on *S. aureus* in the absence of light. The number of living cells decreases with increasing RB concentration in the dark. Table 1 shows a comparison between dark and light toxicity of three PSs—malachite green oxalate (MGO), RB, and safranin O. The effect of MGO in the dark was the strongest, and a 0.87  $\mu\text{M}$  concentration of MGO was sufficient for inhibiting the growth of *S. aureus*. The dark activity of RB and safranin O is noticeably weaker, and the minimal inhibitory



**Figure 2.** SACT and PACT effect of MB on *S. aureus*. In SACT experiments, the cells at  $10^8 \text{ CFU mL}^{-1}$  concentration were incubated with (a) 5  $\mu\text{M}$  RB or (b) 30  $\mu\text{M}$  MB in the ultrasonic bath for 1 h in the dark. In PACT experiments, the cells were illuminated for 15 min by  $1.6 \text{ mW cm}^{-2}$  white light under the same conditions but without sonication. After the treatment, bacteria were tested by viable count. Error bars present standard deviations.

| Photosensitizer         | MIC, $\mu\text{M}$ |              |
|-------------------------|--------------------|--------------|
|                         | Dark               | Illumination |
| Malachite green oxalate | 0.87               | 0.15         |
| Rose Bengal             | 128                | 2            |
| Safranin O              | 89                 | 23           |

**Table 1.**  
*The MIC values of water-soluble PSs in the dark and under illumination. About  $3 \times 10^4$  CFU mL<sup>-1</sup> of S. aureus were treated by malachite green oxalate, Rose Bengal, and Safranin O at doubled dilutions, illuminated at room temperature by white light of  $1.6 \text{ mW cm}^{-2}$  intensity for 1 h, and incubated overnight in the dark by shaking at 37°C.*

concentrations (MIC) for these PSs against *S. aureus* are more than 100-fold higher. **Figure 2** shows that *S. aureus* cells were completely destroyed by RB at a concentration of 5  $\mu\text{M}$  and MB at 30  $\mu\text{M}$  under illumination. These PSs also showed a cytotoxic effect when applied at the same concentrations in the dark, where MB reduced the bacterial concentration by one and RB by two orders of magnitude.

2.2 Illumination

Although PSs are known to possess a certain dark activity, illumination noticeably increases their cytotoxic effect [6, 14]. An example of the difference in antibacterial activity of different PSs with and without illumination is shown in **Table 1**. In this experiment, the MIC of three PSs was determined for the bacterium *S. aureus* in the dark and after 1 h of illumination. As a result of illumination, the MIC of the examined PSs decreased approximately 6-fold for MGO, 64-fold for RB, and 4-fold for Safranin O.

The main light sources used today for activation of PSs are lasers, light-emitting diodes (LED), and gas discharge lamps (GDL) [10, 31, 32]. There is no absolute advantage of one of these light sources over the others. The choice of light source depends on the specific application. Laser is a high-intensity monochromatic source. It can be easily coupled to a single optical fiber and installed on different lighting devices. LED lamps are cheaper and provide a wide emission spectrum. GDLs are also cheaper than lasers—both in acquisition and in maintenance and have a wide emission spectrum. However, GDLs transmit more heat to the illuminated area than lasers and LEDs, which can lead to tissue damage. In general, the emission spectrum and light intensity are more important for the excitation of a specific PS than the particular light source type [10, 31, 32].

2.3 Sonodynamic excitation of photosensitizers

Illumination is undoubtedly the easiest and most effective way to activate PSs. However, its use is restricted, due to limited penetration of visible light into tissues. There is an ongoing search for alternative methods of PS excitation in the dark in order to overcome this problem. Ultrasonic activation seems to be attractive as an alternative to illumination. As with light activation, ultrasound can be selectively focused on a specific area, thus activating only PS molecules located in the affected area. Ultrasound can also easily penetrate into tissues, which opens prospects for its application in treatment of internal lesions and infections, without the need for invasive devices [33, 34]. Ultrasonic irradiation of PSs initiates the formation of highly active cytotoxic species—ROS and free radicals—which lead to the death of pathogenic cells. It was found that some well-known PSs also have sonosensitizing properties. Among


them are porphyrins [35], RB [36, 37], chlorin e6 derivative, photodithazine [36], and curcumin [38]. Several studies found sonodynamic therapy (SDT) to be the promising treatment in various forms of cancerous tumors [39–43]. Sonodynamic therapy is also offered as treatment for atherosclerosis [44]. The applicability of sonodynamic antimicrobial chemotherapy (SACT) for the treatment of infectious diseases has been confirmed by various research groups [33, 34]. We have previously demonstrated the effectiveness of RB activated by ultrasonication for eradication of Gram-positive *S. aureus* and Gram-negative *E. coli* [29, 45, 46]. The effectiveness of SACT in inactivation of *S. aureus* by two other sensitizers—curcumin [38] and hematoporphyrin monomethyl ether [35]—was also reported. Alves et al. have recently reported on effective destruction of *Candida albicans* by photodithazine and RB in the dark under the ultrasonic excitation. A significant synergistic effect of the combination between PDT and SACT for combatting *C. albicans* biofilms was also found [36].

**Figure 2** demonstrates the effect of ultrasonic activation that we showed on the antibacterial activity of two PSs—RB (**Figure 2a**) and MB (**Figure 2b**)—against *S. aureus* compared to photodynamic activation. **Figure 2a** shows that 15 min of sonication reduces the number of living cells by almost two orders of magnitude, from  $2 \times 10^8$  to  $4 \times 10^6$  CFU mL<sup>-1</sup>. RB alone applied in the dark causes a two orders of magnitude decrease in the cell concentration. However, sonication in the presence of 5 µM RB exerts a much stronger effect, reducing the cell concentration by 5 orders of magnitude. It should be noted that RB at the same concentration under illumination by visible light of 1.6 mW cm<sup>-2</sup> fluence causes complete eradication of *S. aureus* cells, whereas light alone does not cause any significant harm to these cells. However, MB applied under sonication at the concentration causing complete destruction of *S. aureus* cells in the light did not eradicate microbial cells more than sonication alone (**Figure 2**).

## 2.4 Activation of photosensitizers by radio waves

Another possible way for activating PSs in the dark is by using nonionizing radiofrequency electromagnetic waves. The ability of radiofrequency waves to heat human tissue has been known for a long time and has already been applied for local destruction of cancerous tumors [47, 48]. The effectiveness of this method can be significantly improved by using suitable sensitizers, which can be targeted to the affected area and activated by means of radiofrequency radiation for selective destruction of cells. Tamarov et al. proposed the use of crystalline silicon-based nanoparticles as sensitizers induced by 27 MHz radiofrequency waves for effective treatment of Lewis lung carcinoma *in vivo* [48]. Another approach involved using gold nanoparticles, which were heated by an electric field using 13.56 MHz radiofrequency, and effectively destroyed human pancreatic cancer cells *in vitro* [49]. The same frequency was used in other studies to activate fullerene [50] and transferrin [51] and to eradicate cancer tumors *in vitro* and *in vivo*. A possible mechanism of radiosensitization, according to Tamarov et al. [48] and Chung et al. [51], may be thermal activation of sensitizers by hyperthermia, caused by dissipation of electromagnetic energy, which leads to thermal damage of cancer cells.

In our studies, we tested the possibility of using radiofrequency radiation to sensitize PSs in order to destroy microorganisms [29]. For this purpose, we irradiated *S. aureus* cells in physiological saline alone and in the presence of RB with radio waves at different frequencies—from 1 to 20 GHz. *S. aureus* cells in physiological saline in the dark (without RB and without radiation), *S. aureus* cells treated with radio waves (in the absence of RB), and *S. aureus* cells in the presence of RB, but not exposed to radio waves, were used as controls. Radiofrequency radiation alone did not significantly affect the survival of *S. aureus*. RB in the dark applied at the same



**Figure 3.**  
 Effect of RB at the 10  $\mu\text{M}$  concentration under activation by radio waves at various frequency ranges on eradication of *S. aureus* at the initial cell concentration of  $4.4 \times 10^4 \text{ CFU mL}^{-1}$  in the dark. Error bars present standard deviations.


concentration did not lead to any decrease in the bacterial concentration. However, exposure of *S. aureus* cells to radio waves in the presence of RB markedly reduced the number of live microorganisms. The rate of cell damage depended on the radio wave frequency. The most significant effect was observed in the frequency range of 9–12 GHz, where in the presence of RB, only 4.5% of the cells survived (**Figure 3**). For comparison, irradiation of cells treated by RB with radio waves in the frequency range of 1–3 GHz caused only a 40% reduction in the number of live cells.

To the best of our knowledge, our work was the first attempt to sensitize a PS by radio waves for destruction of bacteria. This topic naturally necessitates a broader and deeper study to understand the mechanisms of excitation and the possibilities of applying this method. The most likely mechanism of RB excitation by radio waves is conversion of electromagnetic energy into heat, which causes activation of RB, followed by energy transfer to dissolved oxygen and the formation of ROS, affecting the cells. We assume that when PSs are exposed to radiofrequency radiation, they actually behave like thermosensitizers excited by heat instead of light [29].

## 2.5 Chemiluminescent and bioluminescent excitation of photosensitizers

Another approach to overcoming the limitations of PACT in the treatment of deep infections is to replace the external light source by chemo- or bioluminescent light. Bioluminescence is a well-known phenomenon occurring in biological systems as a result of oxidation reactions of luciferins catalyzed by luciferases. This property is inherent in various microorganisms, worms, and insects, and the luciferins and luciferases of different organisms can be completely different. Bioluminescence is considered as a type of chemiluminescence, i.e., luminescence originating in the course of a chemical reaction. Bio- and chemiluminescence systems are used in various fields of medicine, pharmaceuticals, and bioanalytics [52, 53].

One of the well-studied and most effective chemical reactions involving light emission is oxidation of luminol [52, 54, 55]. Most applications of this reaction are associated with treatment of cancers [55–57]. Use of chemiluminescence as a light source for PACT has not been studied as extensively. Ferraz and colleagues



**Figure 4.**

Effect of chemiluminescent photodynamic antimicrobial treatment (CPAT) on the viability of *S. aureus* and *E. coli*. Cells were incubated with MB at 25  $\mu$ M concentration in the presence of 0.7 mM luminol. After the treatment, bacteria were tested by viable count. Error bars present standard deviations.

evaluated the potential of chemiluminescent-excited photogem in killing *S. aureus* cells [58]. Our group demonstrated the effectiveness of chemiluminescent photodynamic antimicrobial therapy (CPAT) for destruction of *S. aureus* and *E. coli* by exposing these bacteria to the photosensitizer MB in the presence of luminol [46, 59, 60]. The results presented in **Figure 4** show that the rate of growth inhibition by MB increased in the presence of luminol compared to untreated cells or to cells exposed in the dark to MB only.

The dark effect of MB discussed in the above “Dark Activity” section can be seen in **Figure 4**, where the exposure of *S. aureus* and *E. coli* to 25  $\mu$ M MB in the dark reduced the number of live cells by about 10-fold. Luminol alone had no toxic effect on the tested microorganisms. However, when combined with MB, it reduced the number of surviving bacteria by two additional orders of magnitude for *S. aureus* and 1.5 orders of magnitude in the case of *E. coli*. Thus, the use of chemiluminescence may expand the capabilities of PDT, allowing the use of PSs for the treatment of internal organs.


### 3. Encapsulation of photosensitizers in liposomes

Since PSs are usually inactive in the absence of excitation, focusing the beam of light, ultrasound or radio wave radiation on the affected area is the easiest way to achieve selective action of a PS. However, surrounding healthy tissues may also be affected by the PS, even under such focused processing. It is therefore very important to target the treatment directly to the infected site. Highly biocompatible and low immunogenic liposomes can serve as carriers for targeted delivery of PSs encapsulated into liposomes to the infected site [61–63].

Liposomes are spherical multi- or unilamellar vesicles consisting of phospholipids (e.g., phosphatidylcholines) with an internal hydrophilic cavity. They vary in composition, size, charge, and number of layers and can encapsulate and deliver both hydrophilic and hydrophobic compounds, which can be retained in the water core of liposomes or be encapsulated in the phospholipid bilayer, respectively.


A variety of methods have been developed for the production of liposomes with a controlled size and special properties. The most widely used method for producing liposomes is hydration of thin lipid films. In this case, lipids with or without active substances are dissolved in an organic solvent, which is evaporated on a rotary evaporator, producing a thin film on a flask wall. The lipid film is then rehydrated by an aqueous phase. Membrane extrusion and sonication methods are most commonly used for control of liposome size [64]. Advanced strategies for liposome preparation include charging the liposomes, attaching the ligands such as antibodies or lectins to their surface, or altering the physiological conditions such as increasing the temperature or changing the pH in the target tissues to produce heat-sensitive or pH-sensitive liposomes [65]. The works of Ghosh, Li, Bulbake, Abu Lila, and Alavi summarize the latest developments in the field of liposome design and optimization, including passive and active targeting, extended circulation, building multifunctional liposomes, and so on [62–66].

There exist several methods for PS encapsulation into liposomes (**Figure 5**). Hydrophilic PSs (e.g., MB, RB, or photofrin) are dissolved in aqueous buffer and are included into the internal cavity of liposomes. Hydrophobic compounds (such as temoporfin and bacteriochlorin a) are integrated in the phospholipid bilayer [62, 67]. Several groups have shown that encapsulation of PSs in liposomes improves their effectiveness against cancer *in vivo*. Back in 1983, Jori and colleagues reported that hematoporphyrin and its derivatives incorporated into liposomes on the basis of dipalmitoyl-phosphatidyl-choline are effective for systemic delivery of PSs to tumors in rats [68]. Enhancement of the photodynamic effects of photofrin encapsulated in a liposome carrier was later demonstrated on a human glioma implanted in rat brain [69]. A variety of PSs (temoporfin, zinc phthalocyanine, benzoporphyrin derivative monoacid, etc.) in various liposomal formulations, such



**Figure 5.**  
Schematic representation of a liposome with PS entrapped in the internal aqueous phase and within the external phospholipid bilayer.

as dipalmitoylphosphatidylcholine, dipalmitoylphosphatidylglycerol, dioleoylphosphatidylcholine, and others, were found to be effective on HT29 and Meth A tumor models *in vivo* [62]. However, the only clinically approved liposomal PS drug to date is Visudyne, developed by QLT in Vancouver, and produced by Novartis AG, Switzerland. This formulation is produced from a derivative of benzoporphyrin monoacid encapsulated in unilamellar dimyristoylphosphatidylcholine/egg phosphatidylglycerol liposomes. The liposomes in this drug not only dissolve the




**Figure 6.** MIC values of free and liposome encapsulated MB and NR determined against (a) *S. aureus* and (b) *E. coli*. Liposomes were prepared from dipalmitoyl phosphatidylcholine and dimyristoyl phosphatidylglycerol at 15 mg/mL total lipid concentration by sonication for 10 sec. Bacteria at  $3 \times 10^4$  CFU mL<sup>-1</sup> concentration were treated by MB and NR at doubled dilutions, illuminated at room temperature by white light of 1.6 mW cm<sup>-2</sup> intensity for 1 h, and incubated overnight in the dark by shaking at 37°C. Error bars present standard deviations.

lipophilic PS for intravenous administration but also contribute to its enhanced absorption in tumor tissues [62, 64].

Liposomal PS preparations are suitable for antibacterial applications. This approach ensures the delivery of the compound at a higher concentration, thus increasing the cytotoxicity of the drug. In addition, the local use of liposomal preparations provides a slow release of active components, which helps prolong their effect in infected tissues. In Gram-negative bacteria, fusion between liposomes and the outer cell membranes leads to the delivery of concentrated liposome contents directly into the cytoplasm [70–72]. In Gram-positive bacteria, the PS is probably released when liposomes interact with the external peptidoglycan and diffuse through the cell wall [72–74]. Various researchers have demonstrated the effectiveness of liposomal formulations of various PSs against Gram-positive and Gram-negative microorganisms and also against fungal infections *in vitro* and *in vivo*. Ferro et al. showed high efficacy of porphyrin incorporated into cationic liposomes against *S. aureus*, compared to the free drug [75, 76]. Tsai also showed an increase in the bactericidal efficacy of hematoporphyrin against a number of Gram-positive bacteria, including *S. aureus*, as a result of its incorporation into liposomes [77]. Yang proved the efficacy of chlorine e6 encapsulated in cationic liposomes against susceptible and drug-resistant clinical isolates of *C. albicans* both *in vitro* and for infected burn wounds *in vivo* [78].

In our studies, we tested the effect of different PSs in different liposome formulations on Gram-positive and Gram-negative bacteria. **Figure 6** presents a comparison between the MICs of free and dipalmitoyl phosphatidylcholine and dimyristoyl phosphatidylglycerol liposome-encapsulated MB and NR against *S. aureus* (**Figure 6a**) and *E. coli* (**Figure 6b**).



**Figure 7.** Chemiluminescent photodynamic antimicrobial treatment effect on the viability of *S. aureus* and *E. coli*. Cells were incubated with 25  $\mu$ M MB liposome (lip) encapsulated together with 0.7 mM luminol (LM). After the treatment, the bacteria were tested by viable count. Error bars present standard deviations.

As can be seen from the results, incorporation into liposomes significantly increased the antibacterial activity of MB and NR. Following encapsulation, the MIC of MB decreased by approximately 2-fold and that of NR by about 1.4-fold for both tested microorganisms (**Figure 6**). We tested the effect of liposome composition on the delivery of these PSs to cells and determined the conditions for efficient use of encapsulated PSs [74].


In addition, we tried to apply liposomal forms of PSs to CPAT by encapsulating not only PSs in liposomes but also luminol and introduced to activate PSs in sites inaccessible to external lighting [59]. We monitored the survival of the cells following their exposure to either liposomal MB or luminol, as well as to liposomes containing both compounds together (**Figure 7**) when the experiments were carried out in the dark.

It can be seen (**Figure 7**) that luminol itself did not lead to cell damage. MB in the liposomal form exhibited certain dark activity, similar to that in a free form discussed in the “Dark Activity” section. The addition of luminol to MB liposomes markedly increased its antibacterial activity toward *S. aureus* and *E. coli*. Liposomes were not targeted in this study. Targeting of liposomes can lead to an additional increase in the efficiency and specificity of this technique.

#### 4. Immobilization

New prospects of using PSs are opened by the immobilization of PSs onto a solid phase. This approach may allow repeated or continuous use of PSs. PSs can be immobilized by adsorption and covalent bonding onto solid supports and by ionic bonding to ion-exchange resins or incorporation into polymer films. The photodynamic properties of immobilized PSs are reported to be retained for a long time [79–83]. PSs studied in the immobilized form include RB, MB, and TBO; the porphyrin derivatives 5,10,15,20-tetrakis (p-hydroxy phenyl) porphyrin, 5,10,15,20-tetrakis (p-aminophenyl) porphyrin, and zinc (II) phthalocyanine tetrasulfonic acid; and the ruthenium salts tris (4,4'-diphenyl-2,2'-bipyridine) ruthenium (II), tris (4,7-diphenyl-1,10-phenanthroline) ruthenium (II), tris (1,10-phenanthroline)-4,7-bis (benzenesulfonate) ruthenate (II), and tris (4,40-dinonyl-1,10-phenanthroline) ruthenium (II). Solid supports applied for immobilization of PSs include polyethylene, polypropylene, polystyrene, polycarbonate, polymethyl methacrylate, polyester isophthalic resin, silicone, cationic nylon, porous silicones, poly (vinylidene difluoride), cellulose membranes, and chitosan [82–88]. Immobilized PSs demonstrated antibacterial properties against Gram-negative and Gram-positive bacteria in batch and continuous regimes and under reuse. Immobilized PSs were found more stable and resistant to photobleaching than in a free form [82, 86, 88].

Our group immobilized PSs in polymers using several techniques. The first method included mixing solutions of PSs in chloroform with solutions of polymers in the same solvent, followed by evaporation of the solvent, which yielded thin polymeric films with homogeneously incorporated PSs. This technique was applied to RB and MB immobilized onto polystyrene, polycarbonate, and polymethyl methacrylate [88–90]. In all cases, the obtained polymer films showed high antibacterial activity against Gram-positive and Gram-negative bacteria when exposed to an external source of white light. However, since this method involves using an organic solvent, it cannot be considered environmentally friendly. The second method is based on dissolution of PSs in a melted polymer under extrusion and does not require any additional chemical reagents [91]. The photosensitizers RB, Rose Bengal lactone, MB, and hematoporphyrin were immobilized in polyethylene and polypropylene using this method. The antibacterial efficiency of immobilized



**Figure 8.** Antibacterial activity of silicon-immobilized RB (5% w/w) under ultrasonic treatment in the dark. Control—*S. aureus* cells treated by ultrasound only. After the treatment, bacteria were tested by viable count. Error bars present standard deviations.

PSs obtained as polymeric strips and beads was tested against *S. aureus* and *E. coli* in batch and continuous regimes under white fluorescent light. All immobilized PSs significantly reduced the concentration of the tested microorganisms, up to their complete eradication [91].

Another immobilization technique was based on polymerization of silicon in the presence of RB as the photosensitizer. Silicon tablets produced by this method contained evenly distributed RB that was not bound to the support by covalent bonds [29]. The antibacterial activity of the immobilized RB was tested under illumination and using ultrasonic activation in the dark (**Figure 8**). **Figure 8** demonstrates the effect of immobilized RB on *S. aureus* cells when subjected to ultrasound in the dark. Silicone alone did not affect the microorganisms with and without sonication. However, the number of alive cells in samples subjected to immobilized RB under sonication decreased with sonication time and decreased by more than three orders of magnitude after 10 min of treatment.

Further development of immobilization methods and different PSs and polymers may expand the possibilities of this approach and yield the applications in various fields, such as the production of antibacterial surfaces and water disinfection.

## 5. Conclusions

Numerous studies show that photodynamic antibacterial chemotherapy is a powerful tool for killing microorganisms. Since this method requires external illumination, it can be successfully applied only to the treatment of local superficial skin and oral cavity infections. Development of new modes of PS excitation by ultrasound, radio waves, chemiluminescent, and bioluminescent light opens new prospects for their use in treating internal infections. Encapsulation of PSs in liposomes may solve the problem of using hydrophobic PSs with poor solubility in the aqueous phase. It can also provide delivery of a concentrated PS directly to the target site, thus increasing efficiency and reducing side effects of the treatment. Immobilization of PSs in a solid phase enables using them repeatedly or in

a continuous mode. It can be assumed that PSs have a good potential for various clinical and nonclinical applications.

## **Acknowledgements**

This work was supported by the Research Authority of the Ariel University, Ariel, Israel.

## **Conflict of interest**

The authors declare no conflict of interest.

## **Author details**

Faina Nakonechny and Marina Nisnevitch\*  
Department of Chemical Engineering, Biotechnology and Materials, Ariel  
University, Ariel, Israel

\*Address all correspondence to: [marinan@ariel.ac.il](mailto:marinan@ariel.ac.il)

## **IntechOpen**

© 2019 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. 

## References

- [1] Daniell MD, Hill JS. A history of photodynamic therapy. *The Australian and New Zealand Journal of Surgery*. 1991;**61**:340-348. DOI: 10.1111/j.1445-2197.1991.tb00230.x
- [2] Raab O. Über die Wirkung fluoreszierender Stoffe auf Infusoria. *Zeitschrift für Biologie*. 1900;**39**:524
- [3] Jodlbauer A, Tappeiner von H. Über die Wirkung photodynamischer (fluoreszierender) Stoffe auf Bakterien. *Münchener Medizinische Wochenschrift* (1950). 1904;**51**:1096-1097
- [4] Jesionek A, Tappeiner von H. Zur Behandlung der Hautcarcinome mit fluoreszierenden Stoffen. *Deutsches Archiv für Klinische Medizin*. 1905;**85**:223-239
- [5] Tappeiner von H. Die photodynamische Erscheinung (Sensibilisierung durch fluoreszierende Stoffe). *Ergebnisse der Physiologie*. 1909;**8**:698-741. Available from: <https://link.springer.com/article/10.1007/BF02321096>
- [6] Ormond AB, Freeman HS. Dye sensitizers for photodynamic therapy. *Materials (Basel)*. 2013;**6**:817-840. DOI: 10.3390/ma6030817
- [7] Debele TA, Peng S, Tsai HC. Drug carrier for photodynamic cancer therapy. *International Journal of Molecular Sciences*. 2015;**16**:22094-22136. DOI: 10.3390/ijms160922094
- [8] Zhang J, Jiang C, Figueiró Longo JP, Azevedo RB, Zhang H, Muehlmann LA. An updated overview on the development of new photosensitizers for anticancer photodynamic therapy. *Acta Pharmaceutica Sinica B*. 2018;**8**:137-146. DOI: 10.1016/j.apsb.2017.09.003
- [9] Macdonald IJ, Dougherty TJ. Basic principles of photodynamic therapy. *Journal of Porphyrins and Phthalocyanines*. 2001;**5**:105-129. DOI: 10.1002/jpp.328
- [10] Cieplik F, Deng D, Crielaard W, Buchalla W, Hellwig E, Al-Ahmad A, et al. Antimicrobial photodynamic therapy - what we know and what we don't. *Critical Reviews in Microbiology*. 2018;**44**:571-589. DOI: 10.1080/1040841X.2018.1467876
- [11] Castano AP, Demidova TN, Hamblin MR. Mechanisms in photodynamic therapy: Part one-photosensitizers, photochemistry and cellular localization. *Photodiagnosis and Photodynamic Therapy*. 2004;**1**:279-293. DOI: 10.1016/S1572-1000(05)00007-4
- [12] Maisch T. Anti-microbial photodynamic therapy: Useful in the future? *Lasers in Medical Science*. 2007;**22**:83-91. DOI: 10.1007/s10103-006-0409-7
- [13] Huang L, Xuan Y, Koide Y, Zhiyentayev T, Tanaka M, Hamblin MR. Type I and type II mechanisms of antimicrobial photodynamic therapy: An in vitro study on gram-negative and gram-positive bacteria. *Lasers in Surgery and Medicine*. 2012;**44**:490-499. DOI: 10.1002/lsm.22045
- [14] Liu Y, Qin R, Zaat SAJ, Breukink E, Heger M. Antibacterial photodynamic therapy: Overview of a promising approach to fight antibiotic-resistant bacterial infections. *Journal of Clinical and Translational Research*. 2015;**1**:140-167. DOI: 10.18053/jctres.201503.002
- [15] Yin R, Hamblin MR. Antimicrobial photosensitizers: Drug discovery under the spotlight. *Current Medicinal Chemistry*. 2015;**22**:2159-2185. DOI: 10.2174/0929867322666150319120134
- [16] Abrahamse H, Hamblin MR. New photosensitizers for photodynamic

- therapy. *The Biochemical Journal*. 2016;**473**:347-364. DOI: 10.1042/BJ20150942
- [17] Carrera ET, Dias HB, Corbi SCT, Marcantonio RAC, Bernardi ACA, Bagnato VS, et al. The application of antimicrobial photodynamic therapy (aPDT) in dentistry: A critical review. *Laser Physics*. 2016;**26**:123001. DOI: 10.1088/1054-660X/26/12/123001
- [18] Hamblin MR. Antimicrobial photodynamic inactivation: A bright new technique to kill resistant microbes. *Current Opinion in Microbiology*. 2016;**33**:67-73. DOI: 10.1016/j.mib.2016.06.008
- [19] Tim M. Strategies to optimize photosensitizers for photodynamic inactivation of bacteria. *Journal of Photochemistry and Photobiology. B*. 2015;**150**:2-10. DOI: 10.1016/j.jphotobiol.2015.05.010
- [20] Nyman ES, Hynninen PH. Research advances in the use of tetrapyrrolic photosensitizers for photodynamic therapy. *Journal of Photochemistry and Photobiology. B*. 2004;**73**:1-28. DOI: 10.1016/j.jphotobiol.2003.10.002
- [21] Harris F, Chatfield LK, Phoenix DA. Phenothiazinium based photosensitisers - photodynamic agents with a multiplicity of cellular targets and clinical applications. *Current Drug Targets*. 2005;**6**:615-627. DOI: 10.2174/1389450054545962
- [22] Leite DP, Paolillo FR, Parmesano TN, Fontana CR, Bagnato VS. Effects of photodynamic therapy with blue light and curcumin as mouth rinse for oral disinfection: A randomized controlled trial. *Photomedicine and Laser Surgery*. 2014;**32**:627-632. DOI: 10.1089/pho.2014.3805
- [23] Bond JS, Francis SH, Park JH. An essential histidine in the catalytic activities of 3-phosphoglyceraldehyde dehydrogenase. *The Journal of Biological Chemistry*. 1970;**245**:1041-1053. Available from: <http://www.jbc.org/content/245/5/1041.long>
- [24] Coulson AF, Yonetani T. Interaction of rose bengal with apo-hemoproteins. An essential histidine residue in cytochrome c peroxidase. *European Journal of Biochemistry*. 1972;**26**:125-131. DOI: 10.1111/j.1432-1033.1972.tb01748.x
- [25] Hsieh YH, Huang YJ, Jin JS, Yu L, Yang H, Jiang C, et al. Mechanisms of Rose Bengal inhibition on SecA ATPase and ion channel activities. *Biochemical and Biophysical Research Communications*. 2014;**454**:308-312. DOI: 10.1016/j.bbrc.2014.10.070
- [26] Shrestha A, Hamblin MR, Kishen A. Photoactivated rose bengal functionalized chitosan nanoparticles produce antibacterial/biofilm activity and stabilize dentin-collagen. *Nanomedicine*. 2014;**10**:491-501. DOI: 10.1016/j.nano.2013.10.010
- [27] Nakonieczna J, Wolnikowska K, Ogonowska P, Neubauer D, Bernat A, Kamysz W. Rose bengal-mediated photoinactivation of multidrug resistant *Pseudomonas aeruginosa* is enhanced in the presence of antimicrobial peptides. *Frontiers in Microbiology*. 2018;**9**:1949. DOI: 10.3389/fmicb.2018.01949
- [28] Goulart Rde C, Bolean M, Paulino Tde P, Thedei G Jr, Souza SL, Tedesco AC, et al. Photodynamic therapy in planktonic and biofilm cultures of *Aggregatibacter actinomycetemcomitans*. *Photomedicine and Laser Surgery*. 2010;**28**(Suppl. 1):S53-S60. DOI: 10.1089/pho.2009.2591
- [29] Nakonechny F, Barel M, David A, Koretz S, Litvak B, Ragozin E, et al. Dark antibacterial activity of Rose Bengal. *International Journal of*

Molecular Sciences. 2019;**20**:E3196.  
DOI: 10.3390/ijms20133196

[30] Brovko LY, Meyer A, Tiwana AS, Chen W, Liu H, Filipe CD, et al. Photodynamic treatment: A novel method for sanitation of food handling and food processing surfaces. *Journal of Food Protection*. 2009;**72**:1020-1024. DOI: 10.4315/0362-028X-72.5.1020

[31] Wilson BC, Patterson MS. The physics, biophysics and technology of photodynamic therapy. *Physics in Medicine and Biology*. 2008;**53**:R61-R109. DOI: 10.1088/0031-9155/53/9/R01

[32] Brancaleon L, Moseley H. Laser and non-laser light sources for photodynamic therapy. *Lasers in Medical Science*. 2002;**17**:173-186. DOI: 10.1007/s101030200027

[33] Harris F, Dennison SR, Phoenix DA. Using sound for microbial eradication - light at the end of the tunnel? *FEMS Microbiology Letters*. 2014;**356**:20-22. DOI: 10.1111/1574-6968.12484

[34] Serpe L, Giuntini F. Sonodynamic antimicrobial chemotherapy: First steps towards a sound approach for microbe inactivation. *Journal of Photochemistry and Photobiology. B*. 2015;**150**:44-49. DOI: 10.1016/j.jphotobiol.2015.05.012

[35] Zhuang D, Hou C, Bi L, Han J, Hao Y, Cao W, et al. Sonodynamic effects of hematoporphyrin monomethyl ether on *Staphylococcus aureus* in vitro. *FEMS Microbiology Letters*. 2014;**361**:174-180. DOI: 10.1111/fml.2014.361

[36] Alves F, Pavarina AC, Mima EGO, McHale AP, Callan JF. Antimicrobial sonodynamic and photodynamic therapies against *Candida albicans*. *Biofouling*. 2018;**34**:357-367. DOI: 10.1080/08927014.2018.1439935

[37] Costley D, Nesbitt H, Ternan N, Dooley J, Huang YY, Hamblin MR, et al.

Sonodynamic inactivation of gram-positive and gram-negative bacteria using a Rose Bengal-antimicrobial peptide conjugate. *International Journal of Antimicrobial Agents*. 2017;**49**:31-36. DOI: 10.1016/j.ijantimicag.2016.09.034

[38] Wang X, Ip M, Leung AW, Xu C. Sonodynamic inactivation of methicillin-resistant *Staphylococcus aureus* in planktonic condition by curcumin under ultrasound sonication. *Ultrasonics*. 2014;**54**:2109-2114. DOI: 10.1016/j.ultras.2014.06.017

[39] Sadanala KC, Chaturvedi PK, Seo YM, Kim JM, Jo YS, Lee YK, et al. Sono-photodynamic combination therapy: A review on sensitizers. *Anticancer Research*. 2014;**34**:4657-4664. Available from: <http://ar.iiarjournals.org/content/34/9/4657.full.pdf>

[40] Costley D, Mc Ewan C, Fowley C, McHale AP, Atchison J, Nomikou N, et al. Treating cancer with sonodynamic therapy: A review. *International Journal of Hyperthermia*. 2015;**31**:107-117. DOI: 10.3109/02656736.2014.992484

[41] McHale AP, Callan JF, Nomikou N, Fowley C, Callan B. Sonodynamic therapy: Concept, mechanism and application to cancer treatment. *Advances in Experimental Medicine and Biology*. 2016;**880**:429-450. DOI: 10.1007/978-3-319-22536-4\_22

[42] Pan X, Wang H, Wang S, Sun X, Wang L, Wang W, et al. Sonodynamic therapy (SDT): A novel strategy for cancer nanotheranostics. *Science China. Life Sciences*. 2018;**61**:415-426. DOI: 10.1007/s11427-017-9262-x

[43] Rengeng L, Qianyu Z, Yuehong L, Zhongzhong P, Libo L. Sonodynamic therapy, a treatment developing from photodynamic therapy. *Photodiagnosis and Photodynamic Therapy*. 2017;**19**:159-166. DOI: 10.1016/j.pdpdt.2017.06.003

- [44] Geng C, Zhang Y, Hidru TH, Zhi L, Tao M, Zou L, et al. Sonodynamic therapy: A potential treatment for atherosclerosis. *Life Sciences*. 2018;**207**:304-313. DOI: 10.1016/j.lfs.2018.06.018
- [45] Nakonechny F, Nisnevitch M, Nitzan Y, Nisnevitch M. Sonodynamic excitation of Rose Bengal for eradication of gram-positive and gram-negative bacteria. *BioMed Research International*. 2013;**2013**:684930. DOI: 10.1155/2013/684930
- [46] Nisnevitch M, Nakonechny F. Sensitivity of bacteria to photodynamic chemotherapy. In: Nisnevitch M, editor. *Prokaryotes: Physiology, Biochemistry and Cell Behavior*. New York, NY, USA: Nova Science Publishers Inc.; 2015. pp. 197-220
- [47] Mirza AN, Fornage BD, Sneige N, Kuerer HM, Newman LA, Ames FC, et al. Radiofrequency ablation of solid tumors. *Cancer Journal*. 2001;**7**:95-102
- [48] Tamarov KP, Osminkina LA, Zinovyev SV, Maximova KA, Kargina JV, Gongalsky MB, et al. Radio frequency radiation-induced hyperthermia using Si nanoparticle-based sensitizers for mild cancer therapy. *Scientific Reports*. 2014;**4**:7034. DOI: 10.1038/srep07034
- [49] Amini SM, Kharrazi S, Rezayat SM, Gilani K. Radiofrequency electric field hyperthermia with gold nanostructures: Role of particle shape and surface chemistry. *Artificial Cells, Nanomedicine, and Biotechnology*. 2018;**46**:1452-1462. DOI: 10.1080/21691401.2017.1373656
- [50] Shi J, Wang L, Gao J, Liu Y, Zhang J, Ma R, et al. A fullerene-based multi-functional nanoplatform for cancer theranostic applications. *Biomaterials*. 2014;**35**:5771-5784. DOI: 10.1016/j.biomaterials.2014.03.071
- [51] Chung HJ, Lee HK, Kwon KB, Kim HJ, Hong ST. Transferrin as a thermosensitizer in radiofrequency hyperthermia for cancer treatment. *Scientific Reports*. 2018;**8**:13505. DOI: 10.1038/s41598-018-31232-9
- [52] Merényi G, Lind J, Eriksen TE. Luminol chemiluminescence: Chemistry, excitation, emitter. *Journal of Bioluminescence and Chemiluminescence*. 1990;**5**:53-56. DOI: 10.1002/bio.1170050111
- [53] Gu X, Tang BZ. No UV irradiation needed! Chemiexcited AIE dots for cancer theranostics. *Chem*. 2017;**3**:917-927. DOI: 10.1016/j.chempr.2017.11.013
- [54] Rose AL, Waite TD. Chemiluminescence of luminol in the presence of iron(II) and oxygen: Oxidation mechanism and implications for its analytical use. *Analytical Chemistry*. 2001;**73**:5909-5920. DOI: 10.1021/ac015547q
- [55] Magalhães CM, da Silva JCGE, daSilva LP. Chemiluminescence and bioluminescence as an excitation source in the photodynamic therapy of cancer: A critical review. *ChemPhysChem*. 2016;**17**:2286-2294. DOI: 10.1002/cphc.201600270
- [56] Laptev R, Nisnevitch M, Siboni G, Malik Z, Firer MA. Intracellular chemiluminescence activates targeted photodynamic destruction of leukemic cells. *British Journal of Cancer*. 2006;**95**:189-196. DOI: 10.1038/sj.bjc.6603241
- [57] Davis RW 4th, Snyder E, Miller J, Carter S, Houser C, Klampatsa A, et al. Luminol chemiluminescence reports photodynamic therapy-generated neutrophil activity in vivo and serves as a biomarker of therapeutic efficacy. *Photochemistry and Photobiology*. 2019;**95**:430-438. DOI: 10.1111/php.13040

- [58] Ferraz RC, Fontana CR, Ribeiro AP, Trindade FZ, Bartoloni FH, Baader JW, et al. Chemiluminescence as a PDT light source for microbial control. *Journal of Photochemistry and Photobiology. B.* 2011;**103**:87-92. DOI: 10.1016/j.jphotobiol.2011.01.018
- [59] Nakonechny F, Firer MA, Nitzan Y, Nisnevitch M. Intracellular antimicrobial photodynamic therapy: A novel technique for efficient eradication of pathogenic bacteria. *Photochemistry and Photobiology.* 2010;**86**:1350-1355. DOI: 10.1111/j.1751-1097.2010.00804.x
- [60] Nakonechny F, Nisnevitch M, Nitzan Y, Firer MA. New techniques in antimicrobial photodynamic therapy: Scope of application and overcoming drug resistance in nosocomial infections. In: *Science against Microbial Pathogens: Communicating Current Research and Technological Advances*; Méndez-Vilas, A., Ed.; Formatex Research Center Publisher: Badajoz, Spain; 2011. Volume 1, p. 684-691. ISBN: 13: 978-84-939843-1-1
- [61] Derycke AS, de Witte PA. Liposomes for photodynamic therapy. *Advanced Drug Delivery Reviews.* 2004;**56**:17-30. DOI: 10.1016/j.addr.2003.07.014
- [62] Ghosh S, Carter KA, Lovell JF. Liposomal formulations of photosensitizers. *Biomaterials.* 2019;**218**:119341. DOI: 10.1016/j.biomaterials.2019.119341
- [63] Li M, Du C, Guo N, Teng Y, Meng X, Sun H, et al. Composition design and medical application of liposomes. *European Journal of Medicinal Chemistry.* 2019;**164**:640-653. DOI: 10.1016/j.ejmech.2019.01.007
- [64] Bulbake U, Doppalapudi S, Kommineni N, Khan W. Liposomal formulations in clinical use: An updated review. *Pharmaceutics.* 2017;**9**, pii: E12. DOI: 10.3390/pharmaceutics9020012
- [65] Abu Lila AS, Ishida T. Liposomal delivery systems: Design optimization and current applications. *Biological & Pharmaceutical Bulletin.* 2017;**40**:1-10. DOI: 10.1248/bpb.b16-00624
- [66] Alavi M, Karimi N, Safaei M. Application of various types of liposomes in drug delivery systems. *Advanced Pharmaceutical Bulletin.* 2017;**7**:3-9. DOI: 10.15171/apb.2017.002
- [67] Damoiseau X, Schuitmaker HJ, Lagerberg JW, Hoebeke M. Increase of the photosensitizing efficiency of the Bacteriochlorin a by liposome-incorporation. *Journal of Photochemistry and Photobiology. B.* 2001;**60**:50-60. DOI: 10.1016/S1011-1344(01)00118-X
- [68] Jori G, Tomio L, Reddi E, Rossi E, Corti L, Zorat PL, et al. Preferential delivery of liposome-incorporated porphyrins to neoplastic cells in tumour-bearing rats. *British Journal of Cancer.* 1983;**48**:307-309. DOI: 10.1038/bjc.1983.186
- [69] Jiang F, Lilge L, Grenier J, Li Y, Wilson MD, Chopp M. Photodynamic therapy of U87 human glioma in nude rat using liposome-delivered photofrin. *Lasers in Surgery and Medicine.* 1998;**22**:74-80. DOI: 10.1002/(SICI)1096-9101(1998)22:2<74::AID-LSM2>3.0.CO;2-T
- [70] Mugabe C, Halwani M, Azghani AO, Lafrenie RM, Omri A. Mechanism of enhanced activity of liposome-entrapped aminoglycosides against resistant strains of *Pseudomonas aeruginosa*. *Antimicrobial Agents and Chemotherapy.* 2006;**50**:2016-2022. DOI: 10.1128/AAC.01547-05
- [71] Sachetelli S, Khalil H, Chen T, Beaulac C, Sénéchal S, Lagacé J. Demonstration of a fusion mechanism between a fluid bactericidal liposomal formulation and bacterial cells.

Biochimica et Biophysica Acta. 2000;**1463**:254-266. DOI: 10.1016/S0005-2736(99)00217-5

[72] Nakonechny F, Nitzan Y, Nisnevitch M. Olive oil-based delivery of photosensitizers for bacterial eradication. In: Olive Oil-Constituents, Quality, Health Properties and Bioconversions; Boskou, D., Ed.; InTech Open Access Publisher: Rijeka, Croatia; 2012. p. 471-492. DOI: 10.5772/30623

[73] Storm G, Crommelin DJA. Liposomes: quo vadis? Pharmaceutical Science & Technology Today. 1998;**1**:19-31. DOI: 10.1016/S1461-5347(98)00007-8

[74] Nisnevitch M, Nakonechny F, Nitzan Y. Photodynamic antimicrobial chemotherapy by liposome-encapsulated water-soluble photosensitizers. Russian Journal of Bioorganic Chemistry. 2010;**36**:363-369. DOI: 10.1134/S106816201003012X

[75] Ferro S, Ricchelli F, Mancini G, Tognon G, Jori G. Inactivation of methicillin-resistant *Staphylococcus aureus* (MRSA) by liposome-delivered photosensitising agents. Journal of Photochemistry and Photobiology. B. 2006;**83**:98-104. DOI: 10.1016/j.jphotobiol.2005.12.008

[76] Ferro S, Ricchelli F, Monti D, Mancini G, Jori G. Efficient photoinactivation of methicillin-resistant *Staphylococcus aureus* by a novel porphyrin incorporated into a poly-cationic liposome. The International Journal of Biochemistry & Cell Biology. 2007;**39**:1026-1034. DOI: 10.1016/j.biocel.2007.02.001

[77] Tsai T, Yang YT, Wang TH, Chien HF, Chen CT. Improved photodynamic inactivation of gram-positive bacteria using hematoporphyrin encapsulated in liposomes and micelles.

Lasers in Surgery and Medicine. 2009;**41**:316-322. DOI: 10.1002/lsm.20754

[78] Yang YT, Chien HF, Chang PH, Chen YC, Jay M, Tsai T, et al. Photodynamic inactivation of chlorin e6-loaded CTAB-liposomes against *Candida albicans*. Lasers in Surgery and Medicine. 2013;**45**:175-185. DOI: 10.1002/lsm.22124

[79] Paczkowski J, Neckers DC. Photochemical properties of rose bengal. 11. Fundamental studies in heterogeneous energy transfer. Macromolecules. 1985;**18**:2412-2418. DOI: 10.1021/ma00154a013

[80] Faust D, Funken K, Horneck G, Milow B, Ortner J, Sattlegger M, et al. Immobilized photosensitizers for solar photochemical applications. Solar Energy. 1999;**65**:71-74. DOI: 10.1016/S0038-092X(98)00099-1

[81] Nowakowska M, Kępczyński M, Dąbrowska M. Polymeric photosensitizers. 5. Synthesis and photochemical properties of poly[(N-isopropylacrylamide)-co-(vinylbenzyl chloride)] containing covalently bound rose bengal chromophores. Macromolecular Chemical Physics. 2001;**202**:1679-1688. DOI: 10.1002/1521-3935(20010601)202:9<1679::AID-MACP1679>3.0.CO;2-R

[82] Bonnett R, Krysteva MA, Lalov IG, Artarsky SV. Water disinfection using photosensitizers immobilized on chitosan. Water Research. 2006;**40**:1269-1275. DOI: 10.1016/j.watres.2006.01.014

[83] Jiménez-Hernández ME, Manjón F, García-Fresnadillo D, Orellana G. Solar water disinfection by singlet oxygen photogenerated with polymer-supported Ru(II) sensitizers. Solar Energy. 2006;**80**:1382-1387. DOI: 10.1016/j.solener.2005.04.027

- [84] Villén L, Manjón F, García-Fresnadillo D, Orellana G. Solar water disinfection by photocatalytic singlet oxygen production in heterogeneous medium. *Applied Catalysis*. 2006;**69**:1-9. DOI: 10.1016/j.apcatb.2006.05.015
- [85] Manjón F, García-Fresnadillo D, Orellana G. Water disinfection with Ru(II) photosensitisers supported on ionic porous silicones. *Photochemical & Photobiological Sciences*. 2009;**8**:926-932. DOI: 10.1039/b902014d
- [86] Manjón F, Santana-Magaña M, García-Fresnadillo D, Orellana G. Singlet oxygen sensitizing materials based on porous silicone: Photochemical characterization, effect of dye reloading and application to water disinfection with solar reactors. *Photochemical & Photobiological Sciences*. 2010;**9**:838-845. DOI: 10.1039/c0pp00026d 15
- [87] Cahan R, Schwartz R, Langzam Y, Nitzan Y. Light-activated antibacterial surfaces comprise photosensitizers. *Photochemistry and Photobiology*. 2011;**87**:1379-1386. DOI: 10.1111/j.1751-1097.2011.00989.x
- [88] Valkov A, Nakonechny F, Nisnevitch M. Polymer-immobilized photosensitizers for continuous eradication of bacteria. *International Journal of Molecular Sciences*. 2014;**15**:14984-14996. DOI: 10.3390/ijms150914984
- [89] Valkov A, Nakonechny F, Nisnevitch M. Antibacterial properties of Rose Bengal immobilized in polymer supports. *Applied Mechanics and Materials*. 2015;**719**:21-24. DOI: 10.4028/www.scientific.net/AMM.719-720.21
- [90] Nakonechny F, Pinkus A, Hai S, Yehosha O, Nitzan Y, Nisnevitch M. Eradication of gram-positive and gram-negative bacteria by photosensitizers immobilized in polystyrene. *Photochemistry and Photobiology*. 2013;**89**:671-678. DOI: 10.1111/php.12022
- [91] Valkov A, Raik KA, Mualem-Sinai Y, Nakonechny F, Nisnevitch M. Water disinfection by immobilized photosensitizers. *Water*. 2019;**11**:26. DOI: 10.3390/w11010026

