

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

186,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Role of Antioxidants Supplementation in the Treatment of Male Infertility

*Houda Amor, Nyaz Shelko, Massooma Mohammed,
Peter Michael Jankowski and Mohamad Eid Hammadeh*

Abstract

Nutritional utilization of antioxidants, such as vitamins C, E, β -Carotene and micronutrients, such as folate and zinc, have been shown to be critically essential for normal semen quality and reproductive function. However, it is still, a large knowledge gap exists concerning the role of antioxidants on semen parameters and the role in treatment of male subfertility. Therefore, the current review article designed to find out the positive effect of antioxidants on semen quality, alterations in physiological functions of spermatozoa and infertility treatment. It is advisable that patients with oxidative DNA disruption should be asked to take a simple course of antioxidants prior to undertaking assisted reproduction treatment (ART). In conclusion, antioxidant may be employed as a potent antioxidant and may improve infertility treatment outcomes with ART.

Keywords: antioxidants, male infertility, semen quality, ART

1. Introduction: The impact of oxidative stress on spermatozoa

ROS include a broad category of species including: a) Oxygen free radicals, such as superoxide anion (O_2^-), hydroxyl radical (OH) and hydroxyl radical (HOO). b) Non radical species, such as hypochlorous acid (HOCl) and hydrogen peroxide (H_2O_2). c) Reactive nitrogen species and free nitrogen radicals such as nitroxyl, nitrous oxide, peroxynitrite, etc. [1–3]. These ROS are generated during normal aerobic metabolism and their level increases under stress which reflects a basic health danger. Mitochondrion is the primary cell organelle involved in ROS production. Besides, several endogenous cells and cellular components contribute towards the initiation and propagation of ROS [4, 5].

Overproduction of ROS or the deficiency of antioxidants provokes an imbalance between the per-oxidation and the anti-oxidation in the normal human body. This phenomenon is termed as Oxidative stress [6–8].

Subsequently, it leads to alterations in peroxidation of lipid membranes of sperm, disrupting the structure of membrane receptors, enzymes, transport proteins, and causes an increase in the level of DNA fragmentation of sperm [9] (Figure 1).

ROS have a significant effect on the sperm plasma membrane and subsequent functional integrity of the sperm resulting in a loss of acrosome reaction [11],




Figure 1.
Role of reactive oxygen species in male reproduction [10].

the sperm potential for fertilization [12], sperm motility and vitality [13, 14] as well as impair the sperm capacitation [15] (**Figure 1**).

Both seminal plasma and spermatozoa contain antioxidant systems able to detoxify the harmful effects of ROS. The imbalance between total antioxidant capacity and ROS generation in seminal fluid presents oxidative stress and is strongly correlated with male infertility [16].

In addition, insufficient penetration of the sperm into the oocyte in oligospermia men with elevated levels of ROS has been identified [7]. Moreover, ROS concentrations are considerably elevated in semen samples from infertile men as compared with those of healthy controls, which has suggests that infertile men may benefit highly from antioxidant supplementation [17].

Several lifestyle, stress, and environmental factors encourage excessive free radical generation and oxidative stress, including: Air pollution, cigarette smoke, alcohol intake, toxins, bacterial, fungal, or viral infections, radiation, including extensive sun bathing, intense and lengthy exercise, which provoke tissue damage, antioxidant deficiency and finally enormous intake of antioxidants, such as vitamins C and E etc.

Hammadeh presented that the oxidative stress and smoking markers, (ROS), malondialdehyde, 8-Hydroxyguanosine (8-OHdG) and cotinine were significantly higher ($P < 0.010$) in smokers as compared to non-smokers [18].

In a second study, both fertile and infertile smokers presented elevated seminal ROS level [19].

Alcohol, known as ethyl alcohol or ethanol; EtOH, promotes ROS. These ROS ultimately interact with macromolecules, among them membrane lipids, generating aldehydes such as 4-Hydroxynonenal (4-HNE) and Malondialdehyde (MDA). It is also known that aldehydes and ROS can directly interact with both proteins and DNA, ultimately leading to transcription-repression of concrete genes. In fact, the impact of ROS and aldehydes seems to serve as key factor for these alterations, partially affirmed by the fact that intake of antioxidants prevents these EtOH-induced cellular alterations [20].

The effect of oxidative stress on sperm quality has been studied extensively and estimated to be the problem in 25–87% of male infertility cases [21]. Another major driving hypothesis states that these conditions, by increasing reactive oxygen species (ROS) and nitrogen species (RONS), are capable of altering the balance of the redox status of both the steroidogenic cell population and the germ line cell

populations, causing the disruption of the hypothalamic–pituitary–testicular axis and the impairment of sperm quality [22].

Therefore, the human body has developed a very concise defense system in which antioxidants play a very important role. Antioxidants are capable of decreasing the generation of free radicals, slowing or inhibiting the oxidation and repairing the damage [23]. Mironczuk-Chodakowska also presented that ROS can serve as mediators and regulators of cell metabolism and apoptosis [23].

2. Antioxidants and the sperm quality (count, motility and morphology) and function

Sperms are especially susceptible to oxidative damages due to the presence of excessive amounts of polyunsaturated fatty acids in the plasma membrane, making them highly vulnerable for lipid peroxidation by ROS, causing decreased flexibility of the sperm membrane and decreased tail motion [24]. Besides, the cytoplasm is extruded during the final stages of spermatogenesis, leaving behind the spermatozoa alone without these important enzymes to protect them from ROS altering the sperm DNA [24].

A significant positive correlation has also been described between levels of reactive oxygen species (ROS) and percentage of spermatozoa with several types of abnormalities such as, abnormal heads, acrosome abnormalities, mid piece anomalies, cytoplasmic droplets and tail defects [25].

In previous studies, it has been demonstrated that the generation of reactive oxygen species has been linked with loss of motility and a reduced capacity for sperm–oocyte fusion [26].

In addition, seminal oxidative stress is rather negatively correlated with sperm count, function, and motility, adversely interfering with fusion required for successful fertilization [27].

Prospective studies have indicated that men with elevated levels of ROS have seven times less chance of fertilizing in comparison with men with low ROS levels. Moreover, ROS results in sperm cell damage and its high values have a negative correlation with sperm number and motility [28].

Other studies reveal that sperm exposed to elevated levels of ROS show reduction in viability and motility as confirmed through both conventional assessment and the utilization of computer-assisted sperm motility analysis [29, 30].

Antioxidant supplements are commercially available to assist treat male infertility, but research on its impact on semen quality and rates of pregnancy and live birth are rather very limited and controversial.

The male reproductive condition can be enhanced by supplementation of beneficial elements such as zinc or selenium which provide positive changes in sperm count and motility [31]. Melatonin, beta-carotene, or luteine also maintain high semen quality [32, 33]. Several studies have affirmed that higher intake of vitamin E, vitamin C, and beta-carotene, is linked with improved sperm count motility, in both fertile and infertile men [34, 35].

Also, spermatozoa are dependent on antioxidants which already present in seminal plasma as ascorbic acid (vitamin C), α -tocopherol (vitamin E), glutathione (GSH), amino acids (taurine and hypotaurine), albumin, carnitine and carotenoids [36].

A clinical trial on the impact of antioxidants on male factor infertility enrolled a population of 171 couples. The male partner presented at least one abnormal reading concentration, mobility, morphology and DNA quality; the female partners presented normal fertility test results. Men received a placebo or an antioxidant

supplement comprising of vitamins C, E, D, selenium, L-carnitine, zinc, folic acid and lycopene from three to six months [37].

In this study, no statistically significant differences in sperm concentration, mobility, morphologically normal sperm and DNA quality between the placebo and antioxidant groups after three months have been observed. Moreover, live birth rates did not differ at six months between the antioxidant (15%) and placebo (24%) groups.

3. Antioxidants and sperm DNA fragmentation and apoptosis

Spermatozoa chromatin is relatively tightly packed due to the positively charged protamine unlike histone in somatic cells [38]. This highly dense and stable structure decreases the capability of DNA disruption. Unfortunately, as spermatozoa have only a few repair mechanisms, DNA damage is generally encountered in human spermatozoa, even within the fertile donor population [39].

It has been presented that almost 40% of men searching for a fertility treatment are fertile and free of sperm oxidative DNA damage [40].

In fact, this topic was discussed in several studies. Mark et al. described that spermatozoa are especially prone to oxidative stress due to the elevated ROS level can provoke a break down of sperm phospholipids and fatty acids [41].

Others have been illustrated that an increase degeneration of ROS by morphologically abnormal spermatozoa and/or decreased antioxidant capacity of seminal plasma are possible reasons of DNA damage in these patients [42, 43].

Hammadeh et al., demonstrated that the ROS level in subfertile patients was significantly higher compared with normal subjects [18].

Besides, 8-hydroxydeoxyguanosine (8OHdG) is considered to be an important sensitive oxidative biomarker for evaluating oxidative sperm DNA damage, and its levels were found to be significantly higher in infertile patients compared with normal ones [18, 44].

Defective spermatozoa are also postulated to retain high residual cytoplasm, allowing them to produce excessive reactive oxygen species (ROS), which, provided their incomplete chromatin packaging, causes DNA damage [45]. Therefore, DNA damage in spermatozoa is primarily related to oxidative stress [45].

Sperm DNA damage could be testicular or post-testicular [46]. These damages may include disruptions in spermatogenesis (e.g., genetic or developmental abnormalities) and testicular or post-testicular damage (e.g., gonadotoxins, hyperthermia, oxidants, and endocrine abnormalities). It has been postulated that protamine deficiency (with subsequent aberrant chromatin remodeling), reactive oxygen species and abortive apoptosis may be associated with sperm DNA damage [18, 47].

Walczak-Jędrzejowska described the damaging impact of oxidative stress on sperm cells including a reduction in activity of anti-oxidative mechanisms, disruption of DNA and accelerated apoptosis [48].

Another study demonstrated that oxidative parameters in the semen of infertile men were significantly rather higher than in fertile men, and a very high correlation was observed between oxidative parameters, sperm ROS generation, and DNA fragmentation level [49].

Since sperm DNA disruption can be associated by oxidative and non-oxidative stress (caused by incomplete sperm protamination or aberrant apoptosis), the utilization of DNA fragmentation tests may not be the perfect procedure for identifying individuals with high sperm DNA damage related to oxidative stress.

Therefore, the most ideal parameters to assess DNA damage might not be DNA fragmentation but sperm-oxidation level which indirectly interferes with DNA disruptions [50, 51].

Antioxidants utilized as dietary supplements removed free radicals and decrease the degree of oxidative insult by improving the cellular redox equilibrium [52].

Several studies demonstrated positive results in the therapy of patients with moderate DNA damage exploiting oral antioxidants [42, 53, 54].

Greco and colleagues reported an enhancement in DNA damage in 76% of patients with moderate DNA damage ($\geq 15\%$) after oral intake with vitamins [53].

The beneficial impact of raised antioxidant consumption on sperm concentration and motility in the infertile patient population with no effect on sperm DNA integrity has also been observed [55].

The advantages of antioxidant therapy in male infertility are rather inconclusive with both a positive effect [53, 54] and no significant effect reported [56]. Also, Silver analyzed a cohort of fertile men and did not report any relationships between dietary antioxidant consumption (vitamins C, E or β -carotene) and sperm DNA damage [56].

Treatment with antioxidant supplements is generally related with decreased levels of sperm DNA integrity and/or increased fertility potential [57].

The impact of these antioxidants in prohibiting sperm from endogenous ROS, gentle sperm processing and cryopreservation has yet not been established [58].

It is advisable that patients with oxidative DNA disruption should be asked to take a simple course of antioxidants prior to under taking assisted reproduction treatment. The utilization of antioxidants in decreasing sperm oxidative stress has been the topic of some 20 clinical trials over the last decade for a review [59].

Systematic review involving 29 studies evaluated the impact of oral antioxidant therapy on fertility outcomes and affirmed an overall positive effect of antioxidant supplementation on basic semen parameters, advanced sperm function, results of assisted reproductive therapy (ART), and live birth rate [60].

Oral supplementation of either a single antioxidant or a combination of antioxidants such as L-carnitine, L-acetyl carnitine, N-acetyl-cysteine, Coenzyme Q10, selenium, vitamin C, vitamin E, and lycopene has been reported to enhance semen parameters and sperm DNA integrity in idiopathic infertile men [61].

4. Types of antioxidants

Seminal plasma comprises of a number of antioxidant enzymes, such as, superoxide dismutase, glutathione peroxidase, glutathione reductase, and catalase, and also include non-enzyme antioxidants, such as vitamin C, vitamin E, beta carotenes, carotenoids, flavonoids and metal binding proteins such as albumin, ferritin, coenzyme Q10 and myoglobin, which serve as antioxidant by inactivating pro-oxidant transition metal ions (**Figure 2**) [62–64].

The non-enzymatic antioxidants may protect the spermatozoa from oxidative DNA and membrane disruption by decreasing the singlet oxygen and the detrimental effect of lipid peroxidation on sperm [65].

4.1 Enzymatic natural antioxidants

The male reproductive system has an endogenous antioxidant for shielding spermatozoa from oxidative insult and these are categorized into enzymatic and non-enzymatic [1]. The enzyme system includes superoxide dismutase (SOD), glutathione peroxidase/reductase and catalase acting as a defense against lipid peroxidation in mammalian sperm. The malfunction of these enzyme activities could lead to a loss of the cell function [66].




Figure 2.
Enzymatic and non-enzymatic antioxidants.

These enzymes catalytically eliminate reactive oxygen species from biological systems. Sperm themselves predominantly channel this enzymatic antioxidant [66]. Spermatozoa comprising of low intracellular antioxidant activity include superoxide dismutase (SOD), nuclear glutathione peroxidase (GPx), peroxiredoxin (PRDX), thioredoxin (TRX), and thioredoxin reductase (TRD) [67].

4.1.1 N-acetyl-cystein (NAC)

NAC serves as a precursor of glutathione (GSH) and has thus been employed as an antioxidant in many studies and trials.

The most important studies which demonstrate positive impact with varying doses in vitro are those reported by Baker et al., Oeda et al., and Lopes et al., [68–70].

Baker et al., witnessed decreased ROS generation and enhanced motility with doses ranging from 1 to 10 mmol [68]. Oeda et al. reported preserved motility with doses from 0.1 to 15 mg/ml [69] and Lopes et al. reported decreased DNA damage with 0.1 mmol of NAC [70]. The three most relevant in vivo studies provided conflicting outcomes: Comhaire et al., did not find any enhancement in sperm parameters after 3 months’ administration of NAC (600 mg for 21 days) [71]; whilst the same dosage seemed to enhance sperm motility, volume, viscosity and oxidative status according to Ciftci et al., [72]. In another trial conducted on 468 patients with idiopathic OAT treated for 6 months with 600 mg/day of NAC or 200 mg/day of selenium or both, an enhancement in sperm count, motility and morphology was observed, with additive beneficial impact when both therapies were dispensed [73].

4.1.2 Glutathione peroxidase (GPx4)

Like peroxidase, Gpx4 is a member of the large family of peroxide reducing enzymes that employs glutathione as electron donor [74].

GPX4 exploits thiols in nuclear protein to reduce glutathione and may function as a redox regulation of sperm motility activation [75].

Sakai et al., found a correlation between undue production of glutathione peroxidase (GPx4) and infertility due to inhibition of ROS generation. GPx4 is not only an essential antioxidant enzyme required for sperm motility; it also plays a key role prior to capacitation [76]. However, while GPx4 shields sperm apoptosis, it decreases ROS which is an important factor for capacitation. Naturally occurring mutations in the human GPX4 gene have been related to the pathogenesis of oligoasthenozoospermia [77].

4.1.3 L-Glutathion (GSH)

GSH is one of the most essential endogenous antioxidant, implicated in sustaining the antioxidant balance in human tissues and directly engaged in the removal of ROS [78].

Due to its widely recognized antioxidant characteristics, GSH has been explored as a possible therapy for infertile patients. In vitro studies performed by Griveau and Le Lannou and by Lopes et al. have demonstrated a protective impact of 10 mmol of GSH against DNA damage caused by ROS [70, 79].

In vivo studies, two trials performed by (Lenzi et al.; Lenzi et al.,) revealed that GSH exerted a positive, statistically significant impact on sperm parameters [80, 81]. The first, Lenzi et al., discovered that 600 mg of GSH dispensed every other day for 2 months increased the motility and morphology in 21 men treated with varicocele or male accessory gland infection [82]. The second, performed the following year on 10 infertile men with the same attributes, led to similar outcomes, although a significant upsurge in sperm concentration was also observed [80, 81].

Tail-beat frequency and the shielding effect of 5–10 and 1–10 mmol GSH in vitro against the disruption of sperm motility by activated polymorphonuclear leukocytes has been reported [68, 83].

Parinaud et al., investigated the impact of Sperm-Fit, an antioxidant solution comprising of glucose and GSH, on sperm motility and showed that motility was well-preserved in leukocyte spermic samples, thus enhancing the possibility of recovering motile sperm after liquefaction and centrifugation [84].

Raijmakers et al., reported the presence of substantial amounts of GSH in the seminal plasma of both fertile and subfertile men. However, the median GSH concentrations were remarkable higher in fertile men in comparison with subfertile men. Furthermore, the concentrations of GSH in seminal plasma were directly correlated with sperm motility and inversely correlated with sperm morphology [85].

4.1.4 Superoxide dismutase (SOD)

There is some consensus on the helpful impacts of SOD on lipid peroxidation in vitro. A plethora of studies, published in last two decades of 20th century i.e. between 1991 and 1998, (for a review Lombardo et al.,) described enhanced motility and decreased lipid peroxidation in samples dosed with between 87.7 and 500 IU ml of SOD [86]. Similarly, Kovalski et al. reported that the incorporation of 1 mg of SOD conserved motility [87].

In the contrary, other found no decrease in lipid peroxidation after the incorporation of between 100 and 500 IU ml SOD [88]. Besides, some investigators could

not affirm the impact of SOD on semen quality and sperm fertilizing potential [89]. However, the amount of endogenous antioxidants such as superoxide dismutase and catalase were also observed to be lower in smokers in comparison with fertile non-smokers [19, 90].

A direct correlation between the SOD activity and sperm damage and sperm motility was endorsed by several researchers [7].

4.1.5 Catalase (CAT)

Catalase represents one of the most effective natural enzymes which catalyzes the conversion of H_2O_2 into H_2O and O_2 . CAT could play an essential role in male fertility and could turn out to be an excellent target for male infertility diagnosis.

In vitro investigations confirmed the positive impact of catalase in doses ranging from 0.008 to 0.1 mg/ml or from 50 to 2000 IU/ml on motility, peroxidation and DNA disruption [70, 79, 87].

However, Twigg et al., reported that the addition of 250, 500 or 2000 IU/ml of catalase to sperm samples did not provide any effect on lipid peroxidation. In the same study, no enhancement was observed in lipid peroxidation after the supplementation of albumin at a dose of 0.3–10% [88].

Other researcherse valuated the effects produced by different types of antioxidant therapies on semen quality in infertile males via measurement of CAT activity on seminal plasma through exogenous H_2O_2 degeneration. The results revealed that catalase activity improved after the administration of antioxidant treatments when compared to control samples without antioxidant administration [91–93].

Moreover, Catalase has also been shown to play a crucial role in male fertility. Its antioxidant activity prevents the upsurge of oxidative stress which can cause damage in sperm level, and accordingly, to reduced fertility. However, even though consensus prevails among most authors on the correlation of this scavenger with male fertility, it has only been investigated to a very limited degree of extent in this field [94].

4.1.6 Coenzyme Q10

A very strong relationship has been described to occur between sperm count, motility, and ubiquinol concentration in seminal fluid [95].

Lewin and Lavon employed coenzyme Q10 both in vivo and in vitro and revealed that the incorporation of 50 mmol in vitro caused a significant improvement in motility, whilst in vivo incorporation of 60 mg/day for almost four months resulted in an enhancement in the fertilization rate but did not provide any impact on motility, morphology or concentration in 17 patients with decreased fertilization rates after in vitro fertilization with intracytoplasmic sperm injection for male factor infertility [96].

A stimulating in vivo study conducted on 60 patients with idiopathic asthenozoospermia who were provided with either placebo or 200 mg daily of coenzyme Q10 showed a remarkable advancement in motility after 6 months of treatment; this enhancement was markedly decreased after a period of 3 months in treated subjects [97].

The exogenous incorporation of CoQ10 enhances both ubiquinone and ubiquinol levels in semen and can be useful in increasing sperm kinetic features in patients suffering from idiopathic asthenozoospermia [97].

CoQ10 biosynthesis is highly active in the testes and elevated levels of ubiquinol are found in sperm [98, 99]. Gvozdkova et al., revealed a direct relationship between seminal plasma CoQ10 total concentration and sperm motility [100].

4.2 Non-enzymatic/synthetic/dietary supplements

4.2.1 Vitamin A

The name vitamin A includes both preformed vitamin A (retinol and its esters) and pro vitamin A carotenoids (mostly α -carotene, β -carotene, and β -cryptoxanthin). Retinol can be oxidized to retinal in a reversible manner, which provides all of the biological implications of retinol; or even further oxidized to retinoic acid, which represents the primary active metabolite of vitamin A [101].

Pro-vitamin A carotenoids can be endogenously transformed into retinoic acid whilst other carotenoids, including lutein, zeaxanthin, and lycopene, are not precursors of vitamin A but contain antioxidant capacity. Carotenoids are naturally found in fruits and vegetable. They impart the yellow, red and orange pigment in plant [102].

The most important phytochemical in the carotenoid family is lycopene which among the carotenoids, ranks as one of the most highest quencher of singlet oxygen. A combination of carotenoids, however, is highly potent as compared to individual compounds [103].

In in vitro models, β -carotene may act both as an antioxidant or a prooxidant, depending upon the redox status of the biological environment in which it acts [104].

Comhaire et al., investigated the impact of a combination therapy of 180 mg day vitamin E and 30 mg day β -carotene on a group of 27 infertile men: ROS generation decreased significantly, although no significant enhancement in semen parameters was observed [71].

Zareba et al., evaluated the effect of regular carotenoid intake on the enhancement of sperm quality in 189 young, healthy men. They evaluated semen volume, total sperm count, motility, and morphology after a period on antioxidant rich diet and observed that β -carotene and lutein utilization elevated sperm motility [33]. However, high exploitation of some vitamins, such as vitamin A, may cause adverse reproductive outcomes [105].

4.2.1.1 Lycopene

Lycopene is one of the several constituents of the carotenoid family. Provitamin A carotenoids can be endogenously transformed into retinoic acid whilst other carotenoids, including lutein, zeaxanthin, and lycopene, which are not the precursors of vitamin A but contain antioxidant capacity. Lycopene exhibits good antioxidant activity includes singlet oxygen and free radical scavenging [106].

The presence of lycopene in human semen has been confirmed and its quantity can be enhanced after dietary intake with a natural source of Lycopene [107]. Also, very little is known about the efficacy of carotenoids, particularly that of lycopene—a potent antioxidant and singlet oxygen quencher [108].

Zini et al., preincubated washed sperm suspensions with and without lycopene followed by incubation with H_2O_2 and presented a significant decrease in sperm mobility and significant elevation of sperm DNA defragmentation in lycopene untreated samples whilst the treated samples showed a significant decrease in sperm DNA defragmentation [109].

Ghyasvand et al., performed another study to evaluate the levels of lycopene, β -carotene and retinol in serum and their relationship with sperm DNA disruption and lipid peroxidation in infertile and normo-spermic males, and concluded that lycopene, β -carotene, and retinol can decrease sperm DNA fragmentation and lipid peroxidation via their antioxidant effect [110].

4.2.1.2 Lutein and zeaxanthin

Comhaire et al., conducted a study involving 30 subfertile men which were treated with 16 mg a day of astaxanthin, a carotenoid which is not transformed into vitamin A in humans, for 3 months and did not find any, significant enhancement in concentration, motility, morphology or volume in the 11 treated men compared with the 19 control patients [111].

Intriguingly, Ming-Chieh Li, et al., found an unusual and unexpected inverse relationship of β -carotene administration from foods and of lutein and zeaxanthin administration with live birth rates. Within the observed administration ranges, total consumption of vitamins A, C and E before initiating infertility treatment with ART was not correlated with live birth rates [112].

4.2.2 Vitamin B (folic acid)

Folic Acid is a B-vitamin which is necessary for the synthesis of DNA and transfer of RNA.

The micronutrients folate and zinc are closely related with semen quality. Blood plasma and seminal plasma levels of folate are positively related with sperm concentration and count [113, 114].

Moreover, folic acid, the synthetic form of folate, effectively scavenges oxidizing free radicals and inhibits lipid peroxidation [115].

The evidence on the impact of folate on fertility is unclear. Wong et al., reported that in vivo utilization of folic acid, alone or combined with zinc sulphate (5 and 66 mg day respectively), enhanced sperm concentration and count in their formerly discussed controlled trial. The results were, however, significant only for the 103 infertile men involved in the study, whilst the 107 fertile men did not present any enhancement in sperm parameters [116].

Also, nutritional utilization of antioxidants, such as vitamins C and E, and β -carotene, and micronutrients, such as folate and zinc, have been shown to be critically essential for normal semen quality and reproductive function as confirmed by a large number of studies in both animals and humans [55].

Ebisch et al., showed that administration of zinc and folic acid enhances not only sperm quality but also the outcome of varicocele surgery [117].

Other researchers have revealed that a low folate concentration in seminal plasma is correlated with increased sperm DNA damage in infertile men [118].

Schisterman et al., designed and conducted a trial involving 2,370 couples planning infertility treatments. The men were assigned arbitrarily and received either a placebo or a daily supplement comprising of 5 milligrams of folic acid and 30 milligrams of zinc. The results indicated that the live births did not alter significantly among the two groups: 404 (34%) in the supplement group and 416 (35%) in the placebo group. Similarly, the groups did not show difference among various semen parameters, such as sperm motility, morphology and total count. However, the fraction of sperm DNA fragmentation was elevated in the supplement group (29.7%), when compared to the placebo group (27.2%). Men in the supplement group also presented an elevated proportion of gastrointestinal symptoms, when compared to the placebo group: abdominal discomfort (6% vs. 3%), nausea (4% vs. 2%) and vomiting (3% vs. 1%). The authors reported that these dietary supplements have a very little to no impact on fertility and may even provoke mild gastrointestinal symptoms [119].

4.2.3 Vitamin C

Ascorbic acid represents the main natural water-soluble antioxidant and is an essential dietary nutriment. This water-soluble antioxidant is an essential dietary nutrient.

Beside, its high potency for scavenging ROS [120], Vit-C serves as an excellent source of electrons, providing an electron to free radicals such as superoxides and hydroxyl radicals, which decreases their reactivity and damages [121].

Ascorbic acid exhibits both antioxidant and prooxidant properties depending upon the amount [122]. As an exogenous antioxidant in semen, ascorbic acid played a very crucial role in controlling the oxidative stress [47].

In an extensive study comprising of 30 infertile but healthy men, daily intake of 200 mg and 100 mg vitamin C enhanced sperm count by 112 and 140 percent respectively. Intriguingly the concentration of ascorbic acid in the seminal plasma is 10-fold higher than the serum [123].

The concentration of ascorbic acid in the seminal plasma was observed to be negatively correlated with reactive oxygen species activity in sperm of infertile men, and the reduced ascorbic acid intake was correlated with an elevated oxidative damage in the sperm of healthy men [124].

Hughes et al., reported the significance of vitamins C (300-600 mmol/l) and E (30-60 mmol/l) in protecting sperm DNA integrity in Percoll preparations; however, this combination has generated contrasting outcomes with excessive DNA damage, which is likely associated to pro-oxidant effects [125].

Donnelly et al. reported decreased H₂O₂-induced ROS production and DNA damage after the utilization of both vitamin C (300 and 600 mmol/l) and vitamin E (40 and 60 mmol/l) to normozoospermic and asthenozoospermic samples [126].

Vitamin C acts as a cofactor for various key enzymes. It assists in the metabolic processes of folic acid, tyrosine and tryptophan [127]. In cells, ascorbic acid interacts with glutathione in maintaining the reduced form of tocopherol [128].

Kini et al., reported a significant increase in the testicular GSH and SOD level but a reduction in testicular MDA level after pre-treatment of rats with Vitamin-C before cadmium chloride exposure [129]. Similar outcomes were observed by Behairy et al. [130]. In addition, vitamin C is vital in protecting from oxidative damage to the sperm and steroid cells of the Leydig cells and in the sperm chamber [131].

A potent and combined action of vitamin C and E has been observed to protect the spermatozoa against peroxidative insult and DNA fragmentation [132].

Vitamin C reduces the inter chain disulphide bridges in protamines opening the cysteine net and subsequently causing DNA decondensation in spermatozoa [54, 133, 134].

4.2.4 Vitamin D

Vitamin D plays a significant role in regulating both male and female fertility [135]. The quality of spermatozoa strongly depend upon vitamin D [136]. Vitamin D is also responsible for retaining sperm motility [137]. Low fertility rates have also been credited to low vitamin D levels in the serum of males [138]. Vitamin D deficiency, in contrast, decreases the probabilities of success when undergoing assisted reproductive technology [139-141]. Islamian et al. investigated the combined effect of vitamin D complement and docosahexaenoic acid on oxidative stress indices of semen in asthenospermic men. The outcome showed that the combined treatment with fatty acid complement, docosahexaenoic acid and vitamin E in asthenospermic men decreased oxidative stress in seminal plasma [142].

4.2.5 Vitamin E (α -tocopherol)

Vitamin E represents a fat-soluble antioxidant molecule which serves as a major chain breaking antioxidant in membranes.

Numerous studies were conducted on vitamin E. The first clinical trials showed that vitamin E supplementation increase the fertilization rates by decreasing lipid peroxidation potential [143]. Kodama et al., reported elevated sperm concentration and decreased DNA damage in 36 infertile patients who underwent 2 months of therapy with vitamin E (200 mg day), vitamin C (200 mg day) and glutathione (GSH; 400 mg day), with no significant enhancement in motility or morphology [144].

Similarly, in another study in which a co-administration of vitamin E and selenium for six months resulted in a significant increase in sperm motility and a decreased percentage of defective spermatozoa in comparison with pre-supplementation period [145, 146]. Besides, it has also been reported that vitamin E decreases the oxidative damage resulting an increase in the fertilization rate [71].

A plethora of research has been dedicated to show that antioxidants such as vitamins E and C and carnitines assist in reducing oxidative stress by quenching free radicals [42]. Greco et al. showed a significant enhancement in 38 men with increased sperm DNA fragmentation after 2 months of combined therapy comprising of 1 g of vitamin C and 1 g of vitamin E daily: clinical pregnancy and implantation rates presented a remarkable increase after the second attempt in comparison with an initial failed intracytoplasmic sperm injection attempt before treatment [53].

Also, vitamin E as an antioxidant may directly quench free radicals and together with CoQ10 shield lipid membranes from peroxidative insult [147].

Moslemi et al., showed that supplementation of vitamin E can significantly decrease lipid peroxidation in seminal plasma, enhance sperm motility and boost pregnancy occurrence [148].

However, in a Meta-analysis, four studies confirmed that it had a very little or no impact on semen parameters, whilst beneficial effects were shown in the remaining 18 studies [86].

4.2.6 Carnitine

The antioxidant activity of carnitines shields against lipid peroxidation of cells membrane, stimulating a large number of in vivo trials examining their impact on sperm parameters.

In the trial conducted by Costa and colleagues [149], 3 g day of L-carnitine was provided to 100 asthenozoospermic men. After 4 months of treatment, a significant enhancement in sperm concentration, motility and morphology was observed.

Balercia et al., evaluated various doses of L-carnitine and acetyl-L-carnitine (3 g of L-carnitine, 3 g of acetyl-L-carnitine, or 2 g L carnitine and 1 g acetyl-L-carnitine daily), showing enhanced sperm motility in asthenozoospermic subjects [150].

Also, Carnitine supplementation has been reported to enhance sperm concentration, mobility, viability, morphology, and total antioxidative capacity [151].

In contrast, Sigman et al., did not find any enhancements in semen parameters in 26 men diagnosed with asthenozoospermia and underwent 6 months of therapy with L-carnitine and L-acetyl-carnitine with 2 and 1 g day, respectively [152].

Gvozdjáková et al., demonstrated a beneficial impact of treatment with a combination of different antioxidants (carnitine with ubiquinol, vitamins E and C) on sperm density, which improved by 39.8% after 3 months of therapy and by 78.0% after 6 months of therapy and sperm motility was enhanced. Sperm pathology reduced in 25.8% after 3-month treatment [153].

4.2.7 Flavonoids

Flavonoids comprise of a very large heterogeneous group of benzopyran derivatives found naturally in fruits, vegetables, and herbs. This group of plant antioxidants has many beneficial health effects.

Flavonoids exhibit a positive health effect in cancer and neurodegenerative diseases, owing to their innate free radical-scavenging activities [154].

These flavonoid glycosides serve as extremely potent free radical scavengers [155, 156]. Some of the most popular and well-known antioxidant flavonoids also serve as prooxidants even when a transition metal is available [157].

Quercetin is one of the most abundant natural flavonoids found in a large variety of fruits and vegetables [158, 159].

The pentahydroxy flavone protects from oxidative injury and cell death by scavenging free radicals, donating hydrogen compound, quenching singlet oxygen, and shielding lipid peroxidation or chelating metal ions [160].

4.2.8 Selenium

Selenium is a very important trace mineral which is highly essential for the human body, including the immune system, cognitive function, and male and female fertility.

According to the United States Office of Dietary Supplements, it participates in the metabolism of thyroid hormone and DNA synthesis, and protects against infection and oxidative damage.

Selenium significantly contributes in the construction of the mitochondrial protective shield in sperm cells and interferes with the condition and function of sperm, and is potent in the treatment of impaired fertility [161].

Searching for a concrete proof of the advantages of *in vivo* selenium therapy is problematic, since various studies with a range of dosages have resulted in conflicting results.

Iwanier and Zachara did not find any positive impact after 3 months of therapy with selenium at a dose of 200 mg/day in 33 subfertile men [162].

In 1998, Scott and his colleagues executed a trial on a group of 64 men (comprising of 46 men diagnosed with OAT and 16 classified as subfertile), dispensing selenium alone or in combination with vitamins A, C and E at daily doses of 100 mg, 1 mg, 10 mg and 15 mg respectively. No significant enhancement was observed in sperm concentration even after 3 months, although motility was increased in treated subjects [163].

However, most of the other studies have claimed enhancements in several parameters after several months of combined therapy with selenium and other antioxidants.

Vezina et al., reported that combination of selenium and vitamin E enhanced the sperm motility, morphology and viability, and the concentration, however, did not alter significantly, as previously discussed [145].

Keskes-Ammar et al., and Safarinejad employed a combination therapy—selenium and vitamin E in the former and selenium and NAC in the latter—which enhanced the motility in both studies and improved the sperm count and morphology in the latter [73, 164].

Besides, it has been demonstrated that Selenium is a fundamental element for semen quality and plays an important function in keeping reproductive condition [31, 48].

4.2.9 Zinc

Zinc is an important micronutrient which also serves as a multifunctional co-factor for more than 80 metallo enzymes involved in DNA transcription and protein synthesis [117]. Moreover, zinc finger proteins are extensively involved in the genetic expression of steroid hormone receptors [165], and it also has antioxidant [166] and anti-apoptotic properties [167].

An interesting correlation was also revealed between the level of Zn in serum and semen in oligozoospermic infertile men, with significantly decreased levels of Zn in serum and semen of men with fertility problems [168]. Zinc levels in seminal plasma have also been positively correlated with sperm concentration and motility in the literature [169, 170].

The Zinc in seminal plasma also serves as antioxidant and anti-bacterial agent which shields the semen from heavy metals accumulation [171].

Two studies were published in which ZnSO_4 was utilized: in the first, 500 mg/day of Zinc sulphate (ZnSO_4) was dispensed to 100 asthenozoospermic men for 3 months, and a significant upturn in sperm count, motility and membrane integrity was observed [172]. The second study, published 10 years later, employed a combination of drugs and Zinc sulphate (ZnSO_4) has been postulated as an infertility treatment, although only a few studies have shown its impact on semen parameters when applied in vivo [173].

Wong et al. observed an elevated sperm count and enhanced semen concentration in a controlled trial with 103 infertile and 107 fertile men who had consumed 5 mg folic acid and 66 mg ZnSO_4 per day for 6 months, either alone or in combination but no enhancements in any of the sperm parameters, most notably concentration were observed [116].

Due to its key role in the processes of DNA compaction, administration of this micronutrient i.e. Zn was very successful in enhancing sperm morphology and DNA integrity in patients suffering from prostate abnormalities [174, 175].

High seminal Zn levels may provide harmful effect on the spermatozoa-zona pellucida-induced acrosome reaction in normozoospermic men [176].

Zinc, is extensively involved in processes of reproduction, not only in the hormone metabolism but also in sperm formation and in the regulation of sperm viability and motility [31].

4.2.10 Manganese (Mn)

Manganese (Mn^{2+}) is one of the most abundant element which is widely distributed in soil, air, water, and food [177].

It is known by its ability to quench the superoxide anions and hydroxyl radicals and due to its chain breaking capacity [178].

The structural flexibility induced by Mn^{2+} is also vital for enzyme dynamics, since Mn is crucial for RNA polymerization [179].

Therefore, Manganese is an essential metal which serves as a co-factor for several enzymes and plays several important biological functions [180]. Mn^{2+} may stimulate the enzymes of glutathione cycle and interact with the total thiols (TSH), glutathione reduced (GSH), glutathione oxidized (GSSG) contents in human spermatozoa. It reduces the generation of thiobarbituric acid reactive substances. In several organisms, elevated intracellular manganese shields against oxidative damage via unknown pathways [181].

Few studies have evaluated the impact of Mn exposure on male reproductive health. Lafond et al., showed that decreased Mn levels in seminal plasma from men with diminished sperm densities [182]. Besides, Mn^{2+} is a very potent stimulator of sperm motility through the stimulation of adenylate cyclase activity [183].

In addition, Manganese accelerates the progressive motility of human washed sperm in a time and dose dependent manner [184]. Moreover, it is proposed that anti-oxidative activity of Mn^{2+} stabilizes the plasma membrane, thereby sustaining the membrane integrity and viability [185].

Also, Mn is a necessary element for humans, but although it has several crucial functions for normal reproductive health, overexposure to Mn^{2+} may be harmful to reproductive health [176, 186].

4.2.11 Pentoxifylline (PTX)

PTX is a derivative of xanthine and a methylxanthine, it is a vasodilating compound which increases red blood cell deformability, prevents from inflammatory reactions and decreases blood viscosity by avoiding platelet aggregation [187].

Pentoxifylline serves as a phosphodiesterase inhibitor and shields the cells from lipid peroxidation by H_2O_2 , therefore it may be helpful to minimize H_2O_2 induced embryo damage and improve IVF outcome [188].

Controversial outcomes were presented in various studies. Some studies reported that the pentoxifylline exerts a beneficial effect on sperm parameters, by decreasing the superoxide anion generation [189–191] or by minimizing lipid peroxidation [191, 192]. However, Twigg et al., did not find any enhancement in lipid peroxidation after the utilization of 3.6 mmol/l of pentoxifylline [88].

Safarinejad carried out a randomized controlled trial on a population comprising of men with idiopathic Oligoasthenoteratozoospermie (OAT). He investigated the response of semen parameters to supplementation with 400 mg of PTX twice daily for a 24-week therapy phase followed by a 12-week therapy-free period. The outcomes of that study revealed a significant enhancement in seminal parameters such as concentration, motility, and morphology [91].

It was also reported that PTX exhibited a positive impact on ICSI outcomes, including fertilization, embryo quality, and pregnancy rates, in asthenozoospermic patients [187].

In conclusion, antioxidants intake separately or combined for at least three months is advisable for patients who planning to undergoing ART therapy.

Conflict of interest

The authors declare that they have no conflict of interest.

IntechOpen

Author details

Houda Amor¹, Nyaz Shelko^{1,2}, Massooma Mohammed³, Peter Michael Jankowski¹ and Mohamad Eid Hammadeh^{1*}

1 Department of Obstetrics and Reproductive Medicine, University of Saarland, Homburg/Saar, Germany

2 Community Health Departments, Technical College of Health, Sulaimani Polytechnic University, Kurdistan, Iraq

3 Gynecology and Obstetrics, Medical College, University of Sulaimani, Kurdistan, Iraq

*Address all correspondence to: mehammadeh@yahoo.de;
mohamad.eid.hammadeh@uks.eu

IntechOpen

© 2021 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. 

References

- [1] S.C Sika, "Relative impact of oxidative stress on male reproductive functions." *Curr. Med. Chem.*, 2001. Volume 8, Number 7, 2001, pp. 851-862(12), doi: <https://doi.org/10.2174/0929867013373039>
- [2] V. Darley-USmar, H. Wiseman, and B. Halliwell, "Nitric oxide and oxygen radicals: a question of balance," *FEBS Letters*. 1995, doi: 10.1016/0014-5793(95)00764-Z.
- [3] A. R. Talebi, Sperm nuclear maturation: A basic and clinical approach. 2011.
- [4] Y. Liu, G. Fiskum, and D. Schubert, "Generation of reactive oxygen species by the mitochondrial electron transport chain," *J. Neurochem.*, 2002, doi: 10.1046/j.0022-3042.2002.00744.x.
- [5] M. Inoue et al., "Mitochondrial Generation of Reactive Oxygen Species and its Role in Aerobic Life," *Curr. Med. Chem.*, 2005, doi: 10.2174/0929867033456477.
- [6] A. Agarwal, A. Hamada, and S. C. Esteves, "Insight into oxidative stress in varicocele-associated male infertility: Part 1," *Nature Reviews Urology*. 2012, doi: 10.1038/nrrol.2012.197.
- [7] A. Agarwal, G. Virk, C. Ong, and S. S. du Plessis, "Effect of Oxidative Stress on Male Reproduction," *World J. Mens. Health*, 2014, doi: 10.5534/wjmh.2014.32.1.1.
- [8] P. P. Mathur and S. C. D'Cruz, "The effect of environmental contaminants on testicular function," *Asian Journal of Andrology*. 2011, doi: 10.1038/aja.2011.40.
- [9] P. Gałęcka, E.; Mrowicka, M.; Malinowska, K.; Gałęcki, "Role of free radicals in the physiological processes," *Woln. Rod. tlenu i azotu w Fizjol. Pol. Merk. Lek*, pp. 24, 446-448., 2008.
- [10] S. Fatima, "Role of Reactive Oxygen Species in Male Reproduction," in *Novel Prospects in Oxidative and Nitrosative Stress*, 2018.
- [11] T. Ichikawa, T. Oeda, H. Ohmori, and W. B. Schill, "Reactive oxygen species influence the acrosome reaction but not acrosin activity in human spermatozoa," *Int. J. Androl.*, 1999, doi: 10.1046/j.1365-2605.1999.00145.x.
- [12] J. G. ALVAREZ, J. C. TOUCHSTONE, L. BLASCO, and B. T. STOREY, "Spontaneous Lipid Peroxidation and Production of Hydrogen Peroxide and Superoxide in Human Spermatozoa Superoxide Dismutase as Major Enzyme Protectant Against Oxygen Toxicity," *J. Androl.*, 1987, doi: 10.1002/j.1939-4640.1987.tb00973.x.
- [13] K. Tremellen, "Oxidative stress and male infertility - A clinical perspective," *Human Reproduction Update*. 2008, doi: 10.1093/humupd/dmn004.
- [14] R. Mahfouz et al., "Semen characteristics and sperm DNA fragmentation in infertile men with low and high levels of seminal reactive oxygen species," *Fertil. Steril.*, 2010, doi: 10.1016/j.fertnstert.2009.12.030.
- [15] R. J. Aitken and B. J. Curry, "Redox regulation of human sperm function: From the physiological control of sperm capacitation to the etiology of infertility and DNA damage in the germ line," *Antioxidants Redox Signal.*, 2011, doi: 10.1089/ars.2010.3186.
- [16] R. K. Sharma, F. F. Pasqualotto, D. R. Nelson, A. J. Thomas, and A. Agarwal, "The reactive oxygen species - Total antioxidant capacity score is a new measure of oxidative stress to predict male infertility," *Hum. Reprod.*, 1999, doi: 10.1093/humrep/14.11.2801.

- [17] A. Agarwal, G. Ahmad, and R. Sharma, "Reference values of reactive oxygen species in seminal ejaculates using chemiluminescence assay," *J. Assist. Reprod. Genet.*, 2015, doi: 10.1007/s10815-015-0584-1.
- [18] M. E. Hammadeh, M. F. Hamad, M. Montenarh, and C. Fischer-Hammadeh, "Protamine contents and P1/P2 ratio in human spermatozoa from smokers and non-smokers," *Hum. Reprod.*, 2010, doi: 10.1093/humrep/deq226.
- [19] E. A. Taha, A. M. Ez-Aldin, S. K. Sayed, N. M. Ghandour, and T. Mostafa, "Effect of smoking on sperm vitality, DNA integrity, seminal oxidative stress, zinc in fertile men," *Urology*, 2012, doi: 10.1016/j.urology.2012.07.002.
- [20] H. H. Byung. et al., "Contribution of reactive oxygen species to cerebral amyloid angiopathy, vasomotor dysfunction, and microhemorrhage in aged Tg2576 mice," *Proc. Natl. Acad. Sci. U. S. A.*, 2015.
- [21] R. Smits, K. D'Hauwers, J. Inthout, Di. Braat, and K. Fleischer, "Impact of a nutritional supplement (Impryl) on male fertility: Study protocol of a multicentre, randomised, double-blind, placebo-controlled clinical trial (SUPpleMent Male fERtility, SUMMER trial)," *BMJ Open*, 2020, doi: 10.1136/bmjopen-2019-035069.
- [22] A. Ferlin and C. Foresta, "New genetic markers for male infertility," *Current Opinion in Obstetrics and Gynecology*. 2014, doi: 10.1097/GCO.0000000000000061.
- [23] I. Mironczuk-Chodakowska, A. M. Witkowska, and M. E. Zujko, "Endogenous non-enzymatic antioxidants in the human body," *Advances in Medical Sciences*. 2018, doi: 10.1016/j.advms.2017.05.005.
- [24] C. Lewis and A. T. Ford, "Infertility in male aquatic invertebrates: A review," *Aquatic Toxicology*. 2012, doi: 10.1016/j.aquatox.2012.05.002.
- [25] C. I. Kobayashi and T. Suda, "Regulation of reactive oxygen species in stem cells and cancer stem cells," *Journal of Cellular Physiology*. 2012, doi: 10.1002/jcp.22764.
- [26] A. Agarwal, R. A. Saleh, and M. A. Bedaiwy, "Role of reactive oxygen species in the pathophysiology of human reproduction," *Fertility and Sterility*. 2003, doi: 10.1016/S0015-0282(02)04948-8.
- [27] S. L., "Human Physiology from Cell to Systems. 6th ed," Cengage Learn. Boston, MA, USA, 2011.
- [28] A. Mehrotra, D. K. Katiyar, A. Agarwal, V. Das, and K. K. Pant, "Role of total antioxidant capacity and lipid peroxidation in fertile and infertile men," *Biomed. Res.*, 2013.
- [29] T. Y. Shi et al., "Effects of reactive oxygen species from activated leucocytes on human sperm motility, viability and morphology," *Andrologia*, 2012, doi: 10.1111/j.1439-0272.2011.01252.x.
- [30] L. S. De Castro et al., "Sperm oxidative stress is detrimental to embryo development: A dose-dependent study model and a new and more sensitive oxidative status evaluation," *Oxid. Med. Cell. Longev.*, 2016, doi: 10.1155/2016/8213071.
- [31] F. Atig, M. Raffa, B. A. Habib, A. Kerkeni, A. Saad, and M. Ajina, "Impact of seminal trace element and glutathione levels on semen quality of Tunisian infertile men," *BMC Urol.*, 2012, doi: 10.1186/1471-2490-12-6.
- [32] R. J. Aitken and S. D. Roman, "Antioxidant systems and oxidative stress in the testes," *Oxidative medicine and cellular longevity*. 2008, doi: 10.4161/oxim.1.1.6843.

- [33] P. Zareba et al., "Semen quality in relation to antioxidant intake in a healthy male population," *Fertil. Steril.*, 2013, doi: 10.1016/j.fertnstert.2013.08.032.
- [34] M. Akmal, J. Q. Qadri, N. S. Al-Waili, S. Thangal, A. Haq, and K. Y. Saloom, "Improvement in human semen quality after oral supplementation of vitamin C," *J. Med. Food*, 2006, doi: 10.1089/jmf.2006.9.440.
- [35] D. P. D. A. F. Braga, G. Halpern, R. D. C. S. Figueira, A. S. Setti, A. Iaconelli, and E. Borges, "Food intake and social habits in male patients and its relationship to intracytoplasmic sperm injection outcomes," *Fertil. Steril.*, 2012, doi: 10.1016/j.fertnstert.2011.10.011.
- [36] S. E. M. Lewis et al., "The impact of sperm DNA damage in assisted conception and beyond: Recent advances in diagnosis and treatment," *Reproductive BioMedicine Online*. 2013, doi: 10.1016/j.rbmo.2013.06.014.
- [37] A. Z. Steiner et al., "The effect of antioxidants on male factor infertility: the Males, Antioxidants, and Infertility (MOXI) randomized clinical trial," *Fertil. Steril.*, 2020, doi: 10.1016/j.fertnstert.2019.11.008.
- [38] S. Sepaniak, T. Forges, H. Gerard, B. Foliguet, M. C. Bene, and P. Monnier-Barbarino, "The influence of cigarette smoking on human sperm quality and DNA fragmentation," *Toxicology*, 2006, doi: 10.1016/j.tox.2006.03.001.
- [39] L. Simon, G. Brunborg, M. Stevenson, D. Lutton, J. McManus, and S. E. M. Lewis, "Clinical significance of sperm DNA damage in assisted reproduction outcome," *Hum. Reprod.*, 2010, doi: 10.1093/humrep/deq103.
- [40] P. Cohen-Bacrie, S. Belloc, Y. J. R. Ménéz, P. Clement, J. Hamidi, and M. Benkhalifa, "Correlation between DNA damage and sperm parameters: a prospective study of 1,633 patients," *Fertil. Steril.*, 2009, doi: 10.1016/j.fertnstert.2008.01.086.
- [41] V. Mak, K. Jarvi, M. Buckspan, M. Freeman, S. Hechter, and A. Zini, "Smoking is associated with the retention of cytoplasm by human spermatozoa," *Urology*, 2000, doi: 10.1016/S0090-4295(00)00700-7.
- [42] A. Agarwal, K. P. Nallella, S. S. R. Allamaneni, and T. M. Said, "Role of antioxidants in treatment of male infertility: An overview of the literature," *Reproductive BioMedicine Online*. 2004, doi: 10.1016/S1472-6483(10)61641-0.
- [43] S. I. Moskovtsev, J. Willis, A. Azad, and J. B. M. Mullen, "Sperm DNA integrity: Correlation with sperm plasma membrane integrity in semen evaluated for male infertility," *Arch. Androl.*, 2005, doi: 10.1080/014850190512770.
- [44] S. Loft et al., "Oxidative DNA damage in human sperm influences time to pregnancy," *Hum. Reprod.*, 2003, doi: 10.1093/humrep/deg202.
- [45] R. J. Aitken and G. N. De Iuliis, "On the possible origins of DNA damage in human spermatozoa," *Molecular Human Reproduction*. 2009, doi: 10.1093/molehr/gap059.
- [46] F. D. Henkel RR, "Sperm DNA fragmentation: origin and impact on human reproduction.," *J Reprod Biotech Fertil.*, pp. 2: 88-108., 2011.
- [47] A. Zini, O. Albert, and B. Robaire, "Assessing sperm chromatin and DNA damage: Clinical importance and development of standards," *Andrology*. 2014, doi: 10.1111/j.2047-2927.2014.00193.x.
- [48] R. Walczak-Jedrzejowska, J. K. Wolski, and J. Slowikowska-Hilczer, "The role of oxidative stress and

- antioxidants in male fertility,” *Cent. Eur. J. Urol.*, 2013, doi: 10.5173/cej.2013.01.art19.
- [49] G. Aktan, S. Doğru-Abbasoğlu, C. Küçükgergin, A. Kadioğlu, G. Özdemirler-Erata, and N. Koçak-Toker, “Mystery of idiopathic male infertility: Is oxidative stress an actual risk?,” *Fertil. Steril.*, 2013, doi: 10.1016/j.fertnstert.2012.11.045.
- [50] A. Noblanc et al., “DNA oxidative damage in mammalian spermatozoa: Where and why is the male nucleus affected?,” *Free Radic. Biol. Med.*, 2013, doi: 10.1016/j.freeradbiomed.2013.07.044.
- [51] S. Vorilhon et al., “Accuracy of human sperm DNA oxidation quantification and threshold determination using an 8-OHdG immuno-detection assay,” *Hum. Reprod.*, 2018, doi: 10.1093/humrep/dey038.
- [52] W. Y. Wong, C. M. G. Thomas, J. M. W. M. Merkus, G. A. Zielhuis, and R. P. M. Steegers-Theunissen, “Male factor subfertility: Possible causes and the impact of nutritional factors,” *Fertility and Sterility*. 2000, doi: 10.1016/S0015-0282(99)00551-8.
- [53] E. Greco, M. Iacobelli, L. Rienzi, F. Ubaldi, S. Ferrero, and J. Tesarik, “Reduction of the incidence of sperm DNA fragmentation by oral antioxidant treatment,” *J. Androl.*, 2005, doi: 10.2164/jandrol.04146.
- [54] Y. J. R. Ménéz et al., “Antioxidants to reduce sperm DNA fragmentation: An unexpected adverse effect,” *Reprod. Biomed. Online*, 2007, doi: 10.1016/S1472-6483(10)60887-5.
- [55] B. Eskenazi, S. A. Kidd, A. R. Marks, E. Slotter, G. Block, and A. J. Wyrobek, “Antioxidant intake is associated with semen quality in healthy men,” *Hum. Reprod.*, 2005, doi: 10.1093/humrep/deh725.
- [56] E. W. Silver, B. Eskenazi, D. P. Evenson, G. Block, S. Young, and A. J. Wyrobek, “Effect of antioxidant intake on sperm chromatin stability in healthy nonsmoking men,” *J. Androl.*, 2005, doi: 10.2164/jandrol.04165.
- [57] P. Piomboni, L. Gambera, F. Serafini, G. Campanella, G. Morgante, and V. De Leo, “Sperm quality improvement after natural anti-oxidant treatment of asthenoteratospermic men with leukocytospermia,” *Asian J. Androl.*, 2008, doi: 10.1111/j.1745-7262.2008.00356.x.
- [58] A. Zini, M. San Gabriel, and A. Baazeem, “Antioxidants and sperm DNA damage: A clinical perspective,” *Journal of Assisted Reproduction and Genetics*. 2009, doi: 10.1007/s10815-009-9343-5.
- [59] P. Gharagozloo and R. J. Aitken, “The role of sperm oxidative stress in male infertility and the significance of oral antioxidant therapy,” *Human Reproduction*. 2011, doi: 10.1093/humrep/der132.
- [60] A. Majzoub and A. Agarwal, “Systematic review of antioxidant types and doses in male infertility: Benefits on semen parameters, advanced sperm function, assisted reproduction and live-birth rate,” *Arab J. Urol.*, 2018, doi: 10.1016/j.aju.2017.11.013.
- [61] M. Arafa et al., “Efficacy of antioxidant supplementation on conventional and advanced sperm function tests in patients with idiopathic male infertility,” *Antioxidants*, 2020, doi: 10.3390/antiox9030219.
- [62] M. G. GERMAN, J. B., & TRABER, “Nutrients and oxidation: Actions, transport, and metabolism of dietary antioxidants,” *Handb. Vitamins*. 3rd Ed, Rucker RB, JW Suttie, DB McCormick,

LJ Machlin (Eds), Marcel Dekker Inc., New York, USA, pp. 569-588., 2001.

[63] L. C. GM, . "Potential for *Ginkgo biloba* as a functional food," Dr. Diss., 2004.

[64] S. Jube and D. Borthakur, "Recent Advances in Food Biotechnology Research," in Food Biochemistry and Food Processing, 2007.

[65] S. E. M. Lewis, E. S. L. Sterling, I. S. Young, and W. Thompson, "Comparison of individual antioxidants of sperm and seminal plasma in fertile and infertile men," Fertil. Steril., 1997, doi: 10.1016/S0015-0282(97)81871-7.

[66] N. Sheikh, I. Amiri, M. Farimani, R. Najafi, and J. Hedeie, "Correlation between sperm parameters and sperm DNA fragmentation in fertile and infertile men," Iran. J. Reprod. Med., 2008.

[67] C. O'flaherty, "Peroxiredoxin 6: The protector of male fertility," Antioxidants. 2018, doi: 10.3390/antiox7120173.

[68] H. W. G. Baker, J. Brindle, D. S. Irvine, and R. J. Aitken, "Protective effect of antioxidants on the impairment of sperm motility by activated polymorphonuclear leukocytes," Fertil. Steril., 1996, doi: 10.1016/S0015-0282(16)58109-6.

[69] T. Oeda, R. Henkel, H. Ohmori, and W. B. Schill, "Scavenging effect of N-acetyl-L-cysteine against reactive oxygen species in human semen: A possible therapeutic modality for male factor infertility?," Andrologia, 1997, doi: 10.1111/j.1439-0272.1997.tb00305.x.

[70] S. Lopes, A. Jurisicova, J. G. Sun, and R. F. Casper, "Reactive oxygen species: Potential cause for DNA fragmentation in human spermatozoa," Hum. Reprod., 1998, doi: 10.1093/humrep/13.4.896.

[71] F. H. Comhaire, A. B. Christophe, A. A. Zalata, W. S. Dhooge, A. M. A. Mahmoud, and C. E. Depuydt, "The effects of combined conventional treatment, oral antioxidants and essential fatty acids on sperm biology in subfertile men," Prostaglandins Leukot. Essent. Fat. Acids, 2000, doi: 10.1054/plef.2000.0174.

[72] H. Ciftci, A. Verit, M. Savas, E. Yeni, and O. Erel, "Effects of N-acetylcysteine on Semen Parameters and Oxidative/Antioxidant Status," Urology, 2009, doi: 10.1016/j.urology.2009.02.034.

[73] M. R. Safarinejad and S. Safarinejad, "Efficacy of Selenium and/or N-Acetyl-Cysteine for Improving Semen Parameters in Infertile Men: A Double-Blind, Placebo Controlled, Randomized Study," J. Urol., 2009, doi: 10.1016/j.juro.2008.10.015.

[74] L. Flohé and F. Ursini, "Peroxidase: A term of many meanings," Antioxidants and Redox Signaling. 2008, doi: 10.1089/ars.2008.2059.

[75] R. J. Aitken, "Possible redox regulation of sperm motility activation," Journal of Andrology. 2000, doi: 10.1002/j.1939-4640.2000.tb02113.x.

[76] S. O., U. T., I. H., U. T., and A. S., "Role of glutathione peroxidase 4 in conjunctival epithelial cells," Investig. Ophthalmol. Vis. Sci., 2015.

[77] M. Diaconu et al., "Failure of phospholipid hydroperoxide glutathione peroxidase expression in oligoasthenozoospermia and mutations in the PHGPx gene," Andrologia, 2006, doi: 10.1111/j.1439-0272.2006.00729.x.

[78] X. Shan, T. Y. Aw, and D. P. Jones, "Glutathione-dependent protection against oxidative injury," Pharmacology and Therapeutics. 1990, doi: 10.1016/0163-7258(90)90045-4.

- [79] J. E. GRIVEAU and D. LE LANNOU, "Effects of antioxidants on human sperm preparation techniques," *Int. J. Androl.*, 1994, doi: 10.1111/j.1365-2605.1994.tb01247.x.
- [80] A. lenzi, F. Culasso, L. Gandini, F. Lombardo, and F. Dondero, "Andrology: Placebo-controlled, double-blind, cross-over trial of glutathione therapy in male infertility," *Hum. Reprod.*, 1993, doi: 10.1093/oxfordjournals.humrep.a137909.
- [81] A. lenzi et al., "Andrology: Glutathione treatment of dyspermia: Effect on the lipoperoxidation process," *Hum. Reprod.*, 1994, doi: 10.1093/oxfordjournals.humrep.a138391.
- [82] L. A., C. F., G. L., L. F., and D. F., "Placebo-controlled, double-blind, cross-over trial of glutathione therapy in male infertility," *Hum. Reprod.*, 1993.
- [83] C. Y. Hong, M. F. Lee, L. J. Lai, and C. P. Wang, "Effect of lipid peroxidation on beating frequency of human sperm tail," *Andrologia*, 1994, doi: 10.1111/j.1439-0272.1994.tb00757.x.
- [84] J. Parinaud, D. Le Lannou, G. Vieitez, J. F. Griveau, P. Milhet, and G. Richoille, "Enhancement of motility by treating spermatozoa with an antioxidant solution (Sperm-Fit®) following ejaculation," *Hum. Reprod.*, 1997, doi: 10.1093/humrep/12.11.2434.
- [85] M. T. M. Raijmakers et al., "Glutathione and glutathione S-transferases A1-1 and P1-1 in seminal plasma may play a role in protecting against oxidative damage to spermatozoa," *Fertil. Steril.*, 2003, doi: 10.1016/S0015-0282(02)04404-7.
- [86] F. Lombardo, A. Sansone, F. Romanelli, D. Paoli, L. Gandini, and A. lenzi, "The role of antioxidant therapy in the treatment of male infertility: An overview," *Asian Journal of Andrology*. 2011, doi: 10.1038/aja.2010.183.
- [87] N. N. Kovalski, E. De Lamirande, and C. Gagnon, "Reactive oxygen species generated by human neutrophils inhibit sperm motility: Protective effect of seminal plasma and scavengers," *Fertil. Steril.*, 1992, doi: 10.1016/s0015-0282(16)55332-1.
- [88] J. Twigg, N. Fulton, E. Gomez, D. Stewart Irvine, and R. John Aitken, "Analysis of the impact of intracellular reactive oxygen species generation on the structural and functional integrity of human spermatozoa: Lipid peroxidation, DNA fragmentation and effectiveness of antioxidants," *Hum. Reprod.*, 1998, doi: 10.1093/humrep/13.6.1429.
- [89] Y. Y. Hsieh, Y. L. Sun, C. C. Chang, Y. S. Lee, H. Der Tsai, and C. S. Lin, "Superoxide dismutase activities of spermatozoa and seminal plasma are not correlated with male infertility," *J. Clin. Lab. Anal.*, 2002, doi: 10.1002/jcla.10029.
- [90] M. F. Elshal, I. H. El-Sayed, M. A. Elsaied, S. A. El-Masry, and T. A. Kumosani, "Sperm head defects and disturbances in spermatozoal chromatin and DNA integrities in idiopathic infertile subjects: Association with cigarette smoking," *Clin. Biochem.*, 2009, doi: 10.1016/j.clinbiochem.2008.11.012.
- [91] M. R. Safarinejad, "Effect of pentoxifylline on semen parameters, reproductive hormones, and seminal plasma antioxidant capacity in men with idiopathic infertility: A randomized double-blind placebo-controlled study," *Int. Urol. Nephrol.*, 2011, doi: 10.1007/s11255-010-9826-4.
- [92] A. Nadjarzadeh et al., "Effect of Coenzyme Q10 supplementation on antioxidant enzymes activity and oxidative stress of seminal plasma: A double-blind randomised clinical trial," *Andrologia*, 2014, doi: 10.1111/and.12062.

- [93] B. Tartibian and B. H. Maleki, "The effects of honey supplementation on seminal plasma cytokines, oxidative stress biomarkers, and antioxidants during 8 weeks of intensive cycling training," *J. Androl.*, 2012, doi: 10.2164/jandrol.110.012815.
- [94] N. Rubio-Riquelme, N. Huerta-Retamal, M. J. Gómez-Torres, and R. M. Martínez-Espinosa, "Catalase as a molecular target for male infertility diagnosis and monitoring: An overview," *Antioxidants*. 2020, doi: 10.3390/antiox9010078.
- [95] R. Alleva, A. Scaramucci, F. Mantero, S. Bompadre, *L. Leoni*, and G. P. Littarru, "The protective role of ubiquinol-10 against formation of lipid hydroperoxides in human seminal fluid," 1997, doi: 10.1016/S0098-2997(97)00040-X.
- [96] A. Lewin and H. Lavon, "The effect of coenzyme Q10 on sperm motility and function," 1997, doi: 10.1016/S0098-2997(97)00036-8.
- [97] G. Balercia et al., "Coenzyme Q10 and male infertility," *Journal of Endocrinological Investigation*. 2009, doi: 10.3275/6301.
- [98] A. Mancini, G. Conte, D. Milardi, L. De Marinis, and G. P. Littarru, "Relationship between sperm cell ubiquinone and seminal parameters in subjects with and without varicocele," *Andrologia*, 1998, doi: 10.1111/j.1439-0272.1998.tb01374.x.
- [99] G. P. Littarru and L. Tiano, "Clinical aspects of coenzyme Q10 in relationship with its bioenergetic and antioxidant properties," in *Mitochondrial Medicine: Mitochondrial Metabolism, Diseases, Diagnosis and Therapy*, 2008.
- [100] A. Gvozdjakova et al., "Importance of the assessment of coenzyme Q10, alpha-tocopherol and oxidative stress for the diagnosis and therapy of infertility in men," *Bratislava Med. J.*, 2013, doi: 10.4149/BLL_2013_129.
- [101] S. NW, "Vitamin A In: Erdman JWMI, Zeisel SH, ed. *Present Knowledge in Nutrition*. 10th Edition ed," John Wiley Sons Ltd, pp. 149-184, 2012.
- [102] A. V. Rao, M. R. Ray, and L. G. Rao, "Lycopene," *Advances in Food and Nutrition Research*. 2006, doi: 10.1016/S1043-4526(06)51002-2.
- [103] A. V. Rao and S. Agarwal, "Role of lycopene as antioxidant carotenoid in the prevention of chronic diseases: A review," *Nutrition Research*. 1999, doi: 10.1016/S0271-5317(98)00193-6.
- [104] P. Palozza, S. Serini, F. Di Nicuolo, E. Piccioni, and G. Calviello, "Prooxidant effects of β -carotene in cultured cells," *Molecular Aspects of Medicine*. 2003, doi: 10.1016/S0098-2997(03)00031-1.
- [105] K. J. Rothman, L. L. Moore, M. R. Singer, U.-S. D. T. Nguyen, S. Mannino, and A. Milunsky, "Teratogenicity of High Vitamin A Intake," *N. Engl. J. Med.*, 1995, doi: 10.1056/nejm199511233332101.
- [106] E. J. Johnson, "The role of carotenoids in human health," *Nutrition in clinical care : an official publication of Tufts University*. 2002, doi: 10.1046/j.1523-5408.2002.00004.x.
- [107] A. Goyal, M. Chopra, B. A. Lwaleed, B. Birch, and A. J. Cooper, "The effects of dietary lycopene supplementation on human seminal plasma," *BJU Int.*, 2007, doi: 10.1111/j.1464-410X.2007.06804.x.
- [108] A. V. Rao and L. G. Rao, "Carotenoids and human health," *Pharmacological Research*. 2007, doi: 10.1016/j.phrs.2007.01.012.

- [109] A. Zini, M. San Gabriel, and J. Libman, "Lycopene supplementation in vitro can protect human sperm deoxyribonucleic acid from oxidative damage," *Fertil. Steril.*, 2010, doi: 10.1016/j.fertnstert.2009.04.004.
- [110] T. Ghyasvand, M. T. Goodarzi, I. Amiri, J. Karimi, and M. Ghorbani, "Serum levels of Lycopene, beta-carotene, and retinol and their correlation with sperm DNA damage in Normospermic and infertile men," *Iran. J. Reprod. Med.*, 2015, doi: 10.29252/ijrm.13.12.787.
- [111] F. H. Comhaire, Y. El Garem, A. Mahmoud, F. Eertmans, and F. Schoonjans, "Combined conventional/antioxidant 'Astaxanthin' treatment for male infertility: A double blind, randomized trial," *Asian J. Androl.*, 2005, doi: 10.1111/j.1745-7262.2005.00047.x.
- [112] L. M.-C. et al., "Men's Intake of Vitamin C and β -Carotene Is Positively Related to Fertilization Rate but Not to Live Birth Rate in Couples Undergoing Infertility Treatment," *J. Nutr.*, 2019.
- [113] L. M. Wallock, T. Tamura, C. A. Mayr, K. E. Johnston, B. N. Ames, and R. A. Jacob, "Low seminal plasma folate concentrations are associated with low sperm density and count in male smokers and nonsmokers," *Fertil. Steril.*, 2001, doi: 10.1016/S0015-0282(00)01697-6.
- [114] J. R. Wallock L, Tamura T, Ames B, Johnston K, Gretz D, "Improvements in folate indices in blood and seminal plasma following vitamin supplementation in smokers and nonsmokers [abstract].," *FASEB J*, p. 13:A890., 1999.
- [115] R. Joshi, S. Adhikari, B. S. Patro, S. Chattopadhyay, and T. Mukherjee, "Free radical scavenging behavior of folic acid: Evidence for possible antioxidant activity," *Free Radic. Biol. Med.*, 2001, doi: 10.1016/S0891-5849(01)00543-3.
- [116] W. Y. Wong, H. M. W. M. Merkus, C. M. G. Thomas, R. Menkveld, G. A. Zielhuis, and R. P. M. Steegers-Theunissen, "Effects of folic acid and zinc sulfate on male factor subfertility: A double-blind, randomized, placebo-controlled trial," *Fertil. Steril.*, 2002, doi: 10.1016/S0015-0282(01)03229-0.
- [117] I. M. W. Ebisch, C. M. G. Thomas, W. H. M. Peters, D. D. M. Braat, and R. P. M. Steegers-Theunissen, "The importance of folate, zinc and antioxidants in the pathogenesis and prevention of subfertility," *Human Reproduction Update*. 2007, doi: 10.1093/humupd/dml054.
- [118] J. C. Boxmeer et al., "Low folate in seminal plasma is associated with increased sperm DNA damage," *Fertil. Steril.*, 2009, doi: 10.1016/j.fertnstert.2008.06.010.
- [119] E. F. Schisterman et al., "Effect of Folic Acid and Zinc Supplementation in Men on Semen Quality and Live Birth among Couples Undergoing Infertility Treatment: A Randomized Clinical Trial," *JAMA - J. Am. Med. Assoc.*, 2020, doi: 10.1001/jama.2019.18714.
- [120] T. Brzozowski et al., "Comparison of nitric oxide-releasing NSAID and vitamin C with classic NSAID in healing of chronic gastric ulcers; involvement of reactive oxygen species," *Med. Sci. Monit.*, 2001.
- [121] E. Levine M., Eбенуwa I., Violet P.-C. Chapter 17—Vitamin C. In: Prasad A.S., Brewer G.J., "Essential and Toxic Trace Elements and Vitamins in Human Health," *Acad. Press*. Cambridge, MA, USA, pp. 241-262, 2020.
- [122] M. Y. Seo and S. M. Lee, "Protective effect of low dose of ascorbic acid on hepatobiliary function

in hepatic ischemia/reperfusion in rats,” *J. Hepatol.*, 2002, doi: 10.1016/S0168-8278(01)00236-7.

[123] R. A. Jacob, F. S. Pianalto, and R. E. Agee, “Cellular ascorbate depletion in healthy men,” *J. Nutr.*, 1992, doi: 10.1093/jn/122.5.1111.

[124] C. G. Fraga, P. A. Motchnik, M. K. Shigenaga, H. J. Helbock, R. A. Jacob, and B. N. Ames, “Ascorbic acid protects against endogenous oxidative DNA damage in human sperm,” *Proc. Natl. Acad. Sci. U. S. A.*, 1991, doi: 10.1073/pnas.88.24.11003.

[125] C. M. Hughes, S. E. M. Lewis, V. J. McKelvey-Martin, and W. Thompson, “The effects of antioxidant supplementation during Percoll preparation on human sperm DNA integrity,” *Hum. Reprod.*, 1998, doi: 10.1093/humrep/13.5.1240.

[126] E. T. Donnelly, N. McClure, and S. E. M. Lewis, “The effect of ascorbate and α -tocopherol supplementation in vitro on DNA integrity and hydrogen peroxide-induced DNA damage in human spermatozoa,” *Mutagenesis*, 1999, doi: 10.1093/mutage/14.5.505.

[127] K. Iqbal, A. Khan, and M. Muzaffar Ali Khan Khattak, “Biological Significance of Ascorbic Acid (Vitamin C) in Human Health-A Review,” 2004.

[128] A. E. Wagner et al., “Free radical scavenging and antioxidant activity of ascorbigen versus ascorbic acid: Studies in vitro and in cultured human keratinocytes,” *J. Agric. Food Chem.*, 2008, doi: 10.1021/jf802403d.

[129] K. P. . Kini R.D., Tripathi Y.B., Raghuveer C.V., Pai S., Ramaswamy C., “Role of vitamin C as an antioxidant in cadmium chloride induced testicular damage,” *Int. J. Appl. Biol. Pharm*, pp. 2: 484-488, 2011.

[130] A. Behairy et al., “The modulatory role of vitamin C in boldenone undecylenate induced testicular oxidative damage and androgen receptor dysregulation in adult male rats,” *Antioxidants*, 2020, doi: 10.3390/antiox9111053.

[131] E. Mangoli et al., “Vitamin C attenuates negative effects of vitrification on sperm parameters, chromatin quality, apoptosis and acrosome reaction in neat and prepared normozoospermic samples,” *Taiwan. J. Obstet. Gynecol.*, 2018, doi: 10.1016/j.tjog.2018.02.006.

[132] U. R. Acharya, M. Mishra, J. Patro, and M. K. Panda, “Effect of vitamins C and E on spermatogenesis in mice exposed to cadmium,” *Reprod. Toxicol.*, 2008, doi: 10.1016/j.reprotox.2007.10.004.

[133] D. Giustarini, I. Dalle-Donne, R. Colombo, A. Milzani, and R. Rossi, “Is ascorbate able to reduce disulfide bridges? A cautionary note,” *Nitric Oxide - Biol. Chem.*, 2008, doi: 10.1016/j.niox.2008.07.003.

[134] E. T. Donnelly, N. McClure, and S. E. M. Lewis, “Antioxidant supplementation in vitro does not improve human sperm motility,” *Fertil. Steril.*, 1999, doi: 10.1016/S0015-0282(99)00267-8.

[135] M. J. Berridge, “Vitamin d deficiency: Infertility and neurodevelopmental diseases (attention deficit hyperactivity disorder, autism, and schizophrenia),” *American Journal of Physiology - Cell Physiology*. 2018, doi: 10.1152/ajpcell.00188.2017.

[136] I. M. Boisen, L. Bøllehuus Hansen, L. J. Mortensen, B. Lanske, A. Juul, and M. Blomberg Jensen, “Possible influence of vitamin D on male reproduction,” *Journal of Steroid Biochemistry and Molecular Biology*. 2017, doi: 10.1016/j.jsbmb.2016.09.023.

- [137] M. B. Jensen et al., "Vitamin D deficiency and low ionized calcium are linked with semen quality and sex steroid levels in infertile men," *Hum. Reprod.*, 2016, doi: 10.1093/humrep/dew152.
- [138] M. Tartagni et al., "Males with low serum levels of vitamin D have lower pregnancy rates when ovulation induction and timed intercourse are used as a treatment for infertile couples: Results from a pilot study," *Reprod. Biol. Endocrinol.*, 2015, doi: 10.1186/s12958-015-0126-9.
- [139] S. Ozkan et al., "Replete vitamin D stores predict reproductive success following in vitro fertilization," *Fertil. Steril.*, 2010, doi: 10.1016/j.fertnstert.2009.05.019.
- [140] M. M. Pacis, C. N. Fortin, S. M. Zarek, S. L. Mumford, and J. H. Segars, "Vitamin D and assisted reproduction: should vitamin D be routinely screened and repleted prior to ART? A systematic review," *J. Assist. Reprod. Genet.*, 2015, doi: 10.1007/s10815-014-0407-9.
- [141] P. Skowrońska et al., "The role of vitamin D in reproductive dysfunction in women – a systematic review," *Ann. Agric. Environ. Med.*, 2016, doi: 10.5604/12321966.1226865.
- [142] J. P. Islamian and H. Mehrali, "Lycopene as a carotenoid provides radioprotectant and antioxidant effects by quenching radiation-induced free radical singlet oxygen: An overview," *Cell Journal*. 2015, doi: 10.22074/cellj.2015.485.
- [143] E. Geva, B. Bartoov, N. Zabludovsky, J. B. Lessing, L. Lerner-Geva, and A. Amit, "The effect of antioxidant treatment on human spermatozoa and fertilization rate in an in vitro fertilization program," *Fertil. Steril.*, 1996, doi: 10.1016/S0015-0282(16)58514-8.
- [144] H. Kodama, H. Kasai, R. Yamaguchi, T. Tanaka, and J. Fukuda, "Increased oxidative deoxyribonucleic acid damage in the spermatozoa of infertile male patients," *Fertil. Steril.*, 1997, doi: 10.1016/S0015-0282(97)00236-7.
- [145] D. Vézina, F. Mauffette, K. D. Roberts, and G. Bleau, "Selenium-vitamin E supplementation in infertile men: Effects on semen parameters and micronutrient levels and distribution," *Biol. Trace Elem. Res.*, 1996, doi: 10.1007/BF02784546.
- [146] S. Sinclair, "Male infertility: Nutritional and environmental considerations," *Alternative Medicine Review*. 2000.
- [147] C. Ross et al., "A systematic review of the effect of oral antioxidants on male infertility," *Reproductive BioMedicine Online*. 2010, doi: 10.1016/j.rbmo.2010.03.008.
- [148] M. K. Moslemi and S. Tavanbakhsh, "Selenium-vitamin E supplementation in infertile men: Effects on semen parameters and pregnancy rate," *Int. J. Gen. Med.*, 2011, doi: 10.2147/IJGM.S16275.
- [149] M. Costa, D. Canale, M. Filicori, S. D'Addio, and A. Lenzi, "L-carnitine in idiopathic asthenozoospermia: a multicenter study," *Andrologia*, 1994, doi: 10.1111/j.1439-0272.1994.tb00780.x.
- [150] G. Balercia, F. Regoli, T. Armeni, A. Koverech, F. Mantero, and M. Boscaro, "Placebo-controlled double-blind randomized trial on the use of L-carnitine, L-acetylcarnitine, or combined L-carnitine and L-acetylcarnitine in men with idiopathic asthenozoospermia," *Fertil. Steril.*, 2005, doi: 10.1016/j.fertnstert.2005.03.064.
- [151] X. Zhou, F. Liu, and S. Zhai, "Effect of L-carnitine and/or L-acetyl-carnitine

in nutrition treatment for male infertility: A systematic review,” 2007, doi: 10.6133/apjcn.2007.16.s1.70.

[152] M. Sigman, S. Glass, J. Campagnone, and J. L. Pryor, “Carnitine for the treatment of idiopathic asthenospermia: a randomized, double-blind, placebo-controlled trial,” *Fertil. Steril.*, 2006, doi: 10.1016/j.fertnstert.2005.10.055.

[153] A. Gvozdjaková, J. Kucharská, J. Dubravicky, V. Mojto, and R. B. Singh, “Coenzyme Q10, α -tocopherol, and oxidative stress could be important metabolic biomarkers of male infertility,” *Dis. Markers*, 2015, doi: 10.1155/2015/827941.

[154] C. A. Rice-evans, N. J. Miller, P. G. Bolwell, P. M. Bramley, and J. B. Pridham, “The relative antioxidant activities of plant-derived polyphenolic flavonoids,” *Free Radic. Res.*, 1995, doi: 10.3109/10715769509145649.

[155] K. C. Huang, “The Pharmacology of Chinese Herbs, Second Edition By K. C. Huang (University of Louisville). CRC Press, Inc., Boca Raton, FL. 1998. xxi + 512 pp. 17.5 × 25 cm. \$129.00. ISBN 0-8493-1665-0,” *J. Nat. Prod.*, 1998, doi: 10.1021/np980247y.

[156] Y. C. S. Attele A.S., Wu J.A., “Multiple constituents and multiple actions,” *Biochem. Pharm.*, 1999.

[157] B. Halliwell, “Are polyphenols antioxidants or pro-oxidants? What do we learn from cell culture and in vivo studies?,” *Archives of Biochemistry and Biophysics*. 2008, doi: 10.1016/j.abb.2008.01.028.

[158] and T. A. Ahmad AH; Rahal A, “Optimising drug potential of plants,” *Proc. Symp. Recent Trends Dev. Herb. Drugs Challenges Oppor. 6th Annu. Conf. ISVPT*, pp. 23-25, 2006.

[159] Mahima et al., “Immunomodulatory and therapeutic

potentials of herbal, traditional/ indigenous and ethnoveterinary medicines,” *Pakistan J. Biol. Sci.*, 2012, doi: 10.3923/pjbs.2012.754.774.

[160] J. Lee, E. R. Hahm, and S. V. Singh, “Withaferin A inhibits activation of signal transducer and activator of transcription 3 in human breast cancer cells,” *Carcinogenesis*, 2010, doi: 10.1093/carcin/bgq175.

[161] V. Badmaev, M. Majeed, and R. A. Passwater, “Selenium: a quest for better understanding [published erratum appears in *Altern Ther Health Med* 1996 Sep;2(5):101],” *Altern Ther Heal. Med.*, 1996.

[162] K. IWANIER and B. A. ZACHARA, “Selenium Supplementation Enhances the Element Concentration in Blood and Seminal Fluid But Does Not Change the Spermatozoal Quality Characteristics in Subfertile Men,” *J. Androl.*, 1995, doi: 10.1002/j.1939-4640.1995.tb00561.x.

[163] R. Scott, A. Macpherson, R. W. S. Yatest, B. Hussain, and J. Dixon, “The effect of oral selenium supplementation on human sperm motility,” *Br. J. Urol.*, 1998, doi: 10.1046/j.1464-410X.1998.00683.x.

[164] L. Keskes-Ammar et al., “Sperm oxidative stress and the effect of an oral vitamin E and selenium supplement on semen quality in infertile men,” *Arch. Androl.*, 2003, doi: 10.1080/01485010390129269.

[165] A. E. Favier, “The role of zinc in reproduction - Hormonal mechanisms,” *Biol. Trace Elem. Res.*, 1992, doi: 10.1007/BF02784623.

[166] M. P. Zago and P. I. Oteiza, “The antioxidant properties of zinc: Interactions with iron and antioxidants,” *Free Radic. Biol. Med.*, 2001, doi: 10.1016/S0891-5849(01)00583-4.

[167] F. Chimienti, M. Aouffen, A. Favier, and M. Seve, “Zinc

Homeostasis-regulating Proteins: New Drug Targets for Triggering Cell Fate,” *Curr. Drug Targets*, 2005, doi: 10.2174/1389450033491082.

[168] H. Mohan, J. Verma, I. Singh, P. Mohan, S. Marwah, and P. Singh, “Inter-relationship of Zinc Levels in Serum and Semen in Oligospermic Infertile Patients and Fertile Males,” *Indian J. Pathol. Microbiol.*, 1997.

[169] H. Fuse, T. Kazama, S. Ohta, and Y. Fujiuchi, “Relationship between zinc concentrations in seminal plasma and various sperm parameters,” *Int. Urol. Nephrol.*, 1999, doi: 10.1023/A:1007190506587.

[170] S. E. Chia, C. N. Ong, L. H. Chua, L. M. Ho, and S. K. Tay, “Comparison of zinc concentrations in blood and seminal plasma and the various sperm parameters between fertile and infertile men,” *J. Androl.*, 2000, doi: 10.1002/j.1939-4640.2000.tb03275.x.

[171] N. Batra, B. Nehru, and M. P. Bansal, “Influence of lead and zinc on rat male reproduction at ‘biochemical and histopathological levels,”” *J. Appl. Toxicol.*, 2001, doi: 10.1002/jat.796.

[172] A. E. Omu, H. Dashti, and S. Al-Othman, “Treatment of asthenozoospermia with zinc sulphate: Andrological, immunological and obstetric outcome,” *Eur. J. Obstet. Gynecol. Reprod. Biol.*, 1998, doi: 10.1016/S0301-2115(97)00262-5.

[173] A. E. Omu, M. K. Al-Azemi, E. O. Kehinde, J. T. Anim, M. A. Oriowo, and T. C. Mathew, “Indications of the mechanisms involved in improved sperm parameters by zinc therapy,” *Med. Princ. Pract.*, 2008, doi: 10.1159/000112963.

[174] C. Foresta et al., “Role of zinc trafficking in male fertility: From germ to sperm,” *Hum. Reprod.*, 2014, doi: 10.1093/humrep/deu075.

[175] A. Fallah, A. Mohammad-Hasani, and A. H. Colagar, “Zinc is an essential element for male fertility: A review of zn roles in men’s health, germination, sperm quality, and fertilization,” *Journal of Reproduction and Infertility*. 2018.

[176] D. Y. Liu, B. S. Sie, M. L. Liu, F. Agresta, and H. W. G. Baker, “Relationship between seminal plasma zinc concentration and spermatozoa-zona pellucida binding and the ZP-induced acrosome reaction in subfertile men,” *Asian J. Androl.*, 2009, doi: 10.1038/aja.2009.23.

[177] D. GB, “Manganese Clin. Toxicol,” pp. 37: 293-307, 1999.

[178] R. K. Anand and U. Kanwar, “Role of some trace metal ions in placental membrane lipid peroxidation,” *Biol. Trace Elem. Res.*, 2001, doi: 10.1385/bter:82:1-3:061.

[179] M. M. Poranen et al., “Structural explanation for the role of Mn²⁺ in the activity of ϕ 6 RNA-dependent RNA polymerase,” *Nucleic Acids Res.*, 2008, doi: 10.1093/nar/gkn632.

[180] J. P. Prestifilippo et al., “Acute effect of manganese on hypothalamic luteinizing hormone releasing hormone secretion in adult male rats: Involvement of specific neurotransmitter systems,” *Toxicol. Sci.*, 2008, doi: 10.1093/toxsci/kfn135.

[181] A. R. Reddi et al., “The overlapping roles of manganese and Cu/Zn SOD in oxidative stress protection,” *Free Radic. Biol. Med.*, 2009, doi: 10.1016/j.freeradbiomed.2008.09.032.

[182] J. L. Lafond, B. Sele, and A. Favier, “Concentration of selected metals in normal and pathological human seminal plasma,” *J. Trace Elem. Electrolytes Health Dis.*, 1988.

[183] S. Lapointe, I. Ahmad, M. M. Buhr, and M. A. Sirard, “Modulation of

Postthaw Motility, Survival, Calcium Uptake, and Fertility of Bovine Sperm by Magnesium and Manganese,” *J. Dairy Sci.*, 1996, doi: 10.3168/jds.S0022-0302(96)76592-X.

[184] Magnus, I. Brekke, T. Åbyholm, and K. Purvis, “Effects of manganese and other divalent cations on progressive motility of human sperm,” *Syst. Biol. Reprod. Med.*, 1990, doi: 10.3109/01485019008986875.

[185] A. Bansal and G. Bilaspuri, “Effect of manganese on bovine sperm motility, viability, and lipid peroxidation in vitro,” *Anim. Reprod.*, 2008.

[186] J. G. Anderson, P. T. Cooney, and K. M. Erikson, “Inhibition of DAT function attenuates manganese accumulation in the globus pallidus,” *Environ. Toxicol. Pharmacol.*, 2007, doi: 10.1016/j.etap.2006.08.006.

[187] A. Ghasemzadeh, F. Karkon-Shayan, S. Yousefzadeh, M. Naghavi-Behzad, and K. Hamdi, “Study of pentoxifylline effects on motility and viability of spermatozoa from infertile asthenozoospermic males,” *Niger. Med. J.*, 2016, doi: 10.4103/0300-1652.193857.

[188] X. Zhang, R. K. Sharma, A. Agarwal, and T. Falcone, “Antioxidant effect of pentoxifylline in reducing oxidative stress induced embryotoxicity,” *Fertil. Steril.*, 2004, doi: 10.1016/j.fertnstert.2004.07.878.

[189] M. GAVELLA, V. LIPOVAC, and T. MAROTTI, “Effect of pentoxifylline on superoxide anion production by human sperm,” *Int. J. Androl.*, 1991, doi: 10.1111/j.1365-2605.1991.tb01099.x.

[190] M. Gavella and V. Lipovac, “Pentoxifylline-mediated reduction of superoxide anion production by human spermatozoa,” *Andrologia*, 1992, doi: 10.1111/j.1439-0272.1992.tb02606.x.

[191] H. Okada, N. Tatsumi, M. Kanzaki, M. Fujisawa, S. Arakawa, and S. Kamidono, “Formation of reactive oxygen species by spermatozoa from asthenospermic patients: Response to treatment with pentoxifylline,” *J. Urol.*, 1997, doi: 10.1016/S0022-5347(01)64697-4.

[192] K. A. McKinney, S. E. M. Lewis, and W. Thompson, “The effects of pentoxifylline on the generation of reactive oxygen species and lipid Peroxidation in human spermatozoa,” *Andrologia*, 1996, doi: 10.1111/j.1439-0272.1996.tb02752.x.