We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists



186,000

200M



Our authors are among the

TOP 1% most cited scientists





WEB OF SCIENCE

Selection of our books indexed in the Book Citation Index in Web of Science™ Core Collection (BKCI)

Interested in publishing with us? Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected. For more information visit www.intechopen.com



Chapter

Flavonoids: Understanding Their Biosynthetic Pathways in Plants and Health Benefits

InnocentMary IfedibaluChukwu Ejiofor and Maria-Goretti Chikodili Igbokwe

Abstract

Flavonoids are polyphenolic compounds and are one of the most abundant secondary metabolites present in plants. They are found in almost all vegetables and fruits. Flavonoids are of essence to plants and to man as well, due to their Medicinal and Pharmaceutical importance. Explicit understanding of the biosynthetic pathway of flavonoids is very essential. This will provide a stepwise explanation of the processes and mechanisms through which different forms of flavonoids are synthesized in plants, including the enzyme(s) responsible for each step. The importance in plants, medicine and pharmacy, of all the product(s) of each step will be emphasized.

Keywords: flavonoid, biosynthesis, plant, phenylpropanoid

1. Introduction

Flavonoids represent an important class of natural products; mainly, they are of the family of secondary plant metabolites having a multi-phenolic structure, found commonly in fruits, vegetables and certain beverages. They have various favorable biochemical, and antioxidant effects associated with various diseases such as cancer, Alzheimer's disease (AD), atherosclerosis and other reported pharmacological effects [1–3]. Flavonoids are associated with a wide spectrum of health-promoting effects and are crucial component in various nutraceutical, pharmaceutical, medicinal and cosmetic applications. This broad spectrum of health-promoting effect is due to their antioxidative, anti-inflammatory, anti-mutagenic and anti-carcinogenic properties coupled with their ability for cellular enzyme functions modulation [4].

In discussing the understanding of biosynthesis of flavonoids and their health benefits, we will be looking at it based on the sub headings of the enzymes involved in the biosynthetic pathway.

2. Phenylalanine ammonia lyase

Lyases are group of enzymes that catalyzes the removal of a functional group or a moiety from a compound by cleaving a carbon–carbon, carbon–oxygen, phosphorous-oxygen, and carbon-nitrogen bonds by mechanism of reaction other than

Bioactive Compounds - Biosynthesis, Characterization and Applications

hydrolysis, or oxidation. This removal of a functional group or a moiety often leads to the formation of a new double bond or ring structure. The bonds cleaving usually occur by means of elimination reaction [5]. In the synthesis of flavonoids, an ammonia-lyase, Phenylalanine ammonia-lyase (PAL) is the primary enzyme in the pathway for the synthesis of phenol [6]. At the initial synthesis of flavonoids, the conversion of Phenylalanine to trans-cinnamic acid is catalyzed by Phenylalanine ammonia lyase through a mechanism of non-oxidative deamination [7] as shown below. This Phenylalanine that is deaminated is a product of another pathway, shikimic acid pathway. Shikimic acid pathway produces three amino acid; L-Tyrosine, L-Phenylalanine and L-Tryptophan. The phenylpropanoid pathway takes its root from the L-Phenylalanine generated from the shikimic acid pathway.



Cinnamic acids belong to a group of aromatic carboxylic acids (C6–C3), which occurs naturally in plant kingdom. In the biochemical process that leads to the formation of lignin, which is the naturally occurring polymeric material that is responsible for providing mechanical support to plant cell wall, cinnamic acids are produced [8]. In all green plants cinnamic acids occur [9]. They are covalently bound to cell walls in minute quantities [10]. They are also found in the reproductive organs of flowering plants [11].

Coffee beans, tea, cocoa, apples, pears, berries, citrus, brassicas vegetables, spinach, beetroot, artichoke, potato, tomato, celery, faba beans, grape and cereals also contain cinnamic acids [12]. Cinnamic acids, with quinic acid, usually appears as conjugates known as the chlorogenic acids. With other acids, sugars or lipids they can also form esters. They can also with aromatic and aliphatic amines, form amides. Some cinnamic-related molecules have been shown in literature those possess the following pharmacological properties; anticancer [13], antituberculosis [14], antimalarial [15], antifungal [16], antimicrobial [17], antiatherogenic [18] and antioxidant [17] activities.

Also, various surveys directed towards the synthetic procedure for cinnamic acid preparation and related atoms have been shown in the literature [19–21]. Medicinal chemists have done the alteration of potency, permeability, solubility or other parameters of a preferred drug or pharmacophore with the aid of cinnamic acids [22].

Cinnamic acid exists in two isomeric forms; trans and cis. Most often available in nature and commercially is the trans form. Cinnamic acid can be obtained from cinnamon bark and balsam resins such as storax.

In the flavors, dyes, and pharmaceuticals production, trans-Cinnamic acid is utilized. The principal use of trans-cinnamic acid is in manufacturing of its methyl, ethyl, and benzyl esters, which are an essential component of perfumes. Also, in the production of the sweetener aspartame, the acid serves as a precursor [23].

3. Cinnamate-4-hydroxylase (C4H)

Hydroxylases are enzymes which add hydroxyl group to organic compounds. C4H found in plants is a cytochrome P450 that catalyzes trans-cinnamic acid conversion to

p-coumaric acid and is the first hydroxylation step of lignin, flavonoids and hydroxycinnamic acid ester biosynthetic pathway correlating with lignifications [24–27]. Generally, in the synthesis of flavonoids, Cinnamate-4-hydroxylase catalyzes the addition of hydroxyl group to the trans-cinnamic acid generated from the deamination of the L-Phenylalanine, leading to the production of p-coumaric acid as shown below.



p-coumaric acid also known as 4-Hydroxycinnamic acid or p-Hydroxycinnamic acid, is a hydroxycinnamic acid, and also organic compound which is a hydroxy derivative of cinnamic acid. p-coumaric acid exists naturally in three isomers, namely; ortho-, meta- and para- coumaric acid [28].



As can be observed from above, these isomers differ from each other differ by the position of the hydroxy group substitution on the phenyl group. p-coumaric acid (4-hydroxy-cinnamic acid) occurs widely in the cell walls of graminaceous plants and is the most abundant of the three isomers [29]. It decreases low-density lipoprotein (LDL) peroxidation [30, 31], antimicrobial activities [32, 33] and plays a vital role in human health. Coumaric acids have been shown to possess radical-scavenging effect [34–40] which reduces stomach cancer risk by suppressing carcinogenic nitrosamines formation [41–43].

p-coumaric acid effectively suppressed endothelial cell migration, tube formation, and rat aorta ring sprouting [44]. It reduces intracellular and mitochondrial reactive oxygen species production [44]. In vivo p-coumaric acid significantly suppressed tumor growth in vivo by blocking angiogenesis. p-coumaric acid is found in various edible plants, such as carrots, tomatoes and cereals [44].

4. 4-coumarate CoA ligase

Ligases are a class of enzymes capable of catalyzing the merging of two large compounds by forming a new chemical bond, mostly accompanied by a small chemical group's hydrolysis on one of the large compounds or commonly causing the linkage of two mixtures together. Ligases are classified under EC 6 primary class of enzymes. They are also further group into six subclasses which are known as ligases that create carbon–oxygen bonds, carbon-sulfur bonds, carbon-nitrogen bonds, carbon–carbon bonds, phosphoric–ester bonds, and nitrogen–metal bonds [45, 46]. Coenzyme A (CoA) is a type of coenzyme that contains pantothenic acid, adenosine 3-phosphate 5-pyrophosphate, and cysteamine; which take part in the transfer of acyl groups, notably in transacetylations [47]. Coenzymes can be defined as organic molecules or compounds that many enzymes required to elicit a catalytic effect [48].



4-coumarate-CoA ligase (4CL) is essential to the general phenylpropanoid pathway and takes part in monolignol biosynthesis through the production of p-coumaroyl-CoA, a precursor for the biosynthesis of p-coumaryl alcohol and coniferyl alcohol in conifers. Essentially, p-coumaroyl-CoA is also involved in the production of other metabolites of plant as a precursor, including stilbenes and flavonoids [49].

4-coumarate-CoA ligase causes the joining of Coenzyme A to *p*-coumaric acid, leading to the formation of *p*- coumaroyl-CoA.



Coumaroyl-coenzyme A is a molecule or compound present in plants. It is the THIOESTER of coenzyme-A and coumaric acid. Coumaroyl-coenzyme A is a basic or fundamental intermediate in the biosynthesis of various natural products found in plants [50].

5. Chalcone synthase

Synthases are enzymes that catalyze the formation of a particular compound. Chalcone synthase catalyzes the production of chalcone, in phenylpropanoid metabolic pathway.

Chalcone synthase (CHS) or naringenin-chalcone synthase is a member of the plant polyketide synthase superfamily, which also includes stilbene synthase (STS), acridone synthase, pyrone synthase, bibenzyl synthase, and p-coumaroyl triacetic acid synthase [51]. Polyketides are a ubiquitous group of secondary metabolites which contain either alternating carbonyl and methylene groups (-CO-CH2-)

or are derived from precursors which have alternating carbonyl and methylene groups [52].

Chalcone synthases, the most well-known representatives of this family, make available the materials needed to initiate various sets of metabolites (flavonoids). These metabolites have important diverse role to play in flowering plants, like the provision of floral pigments, antibiotics, UV protectants and insect repellents [53].

In the production of chalcone, one molecule of p-coumaroyl-CoA and three malonyl-CoA molecules condensation is required. This is catalyzed by the enzyme, chalcone synthase. This process starts with the transfer of a coumaroyl moiety from a p-coumaroyl-CoA which is the starter molecule to an active site cysteine (Cys164) [54]. The next that follows is the series of condensation reactions involving three acetate units obtained from three malonyl-CoA molecules, each proceeding through an acetyl-CoA carbanion derived from malonyl-CoA decarboxylation, extends the polyketide intermediate. Following generation of the thioester-linked tetraketide, a regiospecific intramolecular Claisen condensation forms a new ring system to yield chalcone [55, 56].



Malonyl-CoA is the starting molecule for the synthesis of fatty acid and its elongation. Malonyl-CoA is one of the foundations for the biosynthesis of some phytoalexins, flavonoids, and many malonylated compounds [57]. In plants and also in animals, malonyl-CoA is almost entirely obtained from acetyl-CoA by acetyl-CoA carboxylase [58].

Chalcone is an essential and resourceful molecule. It is a biogenetic precursor for flavonoids and isoflavonoids, which are bountiful in consumable plants. Chalcone contains two aromatic rings, linked together by a three-carbon- α , β unsaturated carbonyl system, i.e., 1,3-diphenyl-2-propen-1-one derivative. Structurally, chalcones are one of the most divergent forms of flavonoids. Chalcone derivatives exhibit a wide range of therapeutic activities which include anticancer [58–62] antioxidants [63–67], anti-inflammatory [68–73], antihypertensive [74], antimalarial [75], antiulcer [76, 77], antiviral [78–81], antiprotozoal [82], cardiovascular activity [83] and mutagenic properties [84], and many other pharmacological properties.

6. Chalcone isomerase

Isomerases are class of EC 5 primary enzymes that catalyze the reactions that involve the rearrangement of a molecule structure [46].

The first detected flavonoid pathway enzyme was Chalcone isomerase. It catalyzes the stereospecific cyclization of chalcones to (2S)-flavanones, which were found to be the exclusive substrates for the reactions to the formation of other classes of flavonoids [85].

Although this type of isomerization reaction can go on spontaneously, the rate of turnover can be increased by 10⁷ fold in the presence of CHI [86]. The CHIs present in plants can be divided into four types (type I to type IV). This

division depends on its phylogenetic relationships [87]. Type I and type II are considered bona fide catalysts with representative CHI enzymatic activity. Type I CHIs are mostly found in vascular plants; they are responsible for forming general flavonoids [88, 89].

In comparison with Type I CHIs, type II CHIs have wider substrate acceptability, besides making use of naringenin chalcone as substrate, they also undertake the conversion of isoliquiritigenin to isoflavonoid which appear to be the specific metabolites in legume [90, 91]. Both type III and type IV CHIs do not participate in chalcone cyclization activity, unlike type I and type II CHI proteins. Due to this, they are termed CHI-like proteins (CHIL). Type III CHIs, which is extensively dispersed in land plants and green algae, have been shown to be fatty acid-binding proteins that function to influence the synthesis and storage of fatty acid in plants [92]. Nonetheless, the action of type IV CHIs which completely lose the bona fide CHI activity is not well known, yet new studies have revealed type CHI-fold proteins might serve as the enhancer of colouration of flowers and production of flavonoid in diverse plant species [93]. All CHIs have a similar backbone arrangement and type III CHIs are thought to be the common forebear of bona fide CHIs [92, 94]. CHI, also regarded as chalcone flavonone isomerase.



Flavanones are primarily found in about 42 larger plant families, specifically in Compositae, Leguminosae, and Rutaceae. Depending on the type of plants, flavanones can be found in all of the parts above and below ground, from vegetative parts to generative organs: branches, bark, stem, leaves, roots, flowers, fruits, seeds, rhizomes, peels, and others [95].

Flavanones show strong antioxidant and radical scavenging activity [96–103] and appear to be associated with a reduced risk of certain chronic diseases [98, 99] the prevention of some cardiovascular disorders [104–107] and certain kinds of cancer [108–112]. Flavanones also exhibit antiviral [113], antimicrobial, [114] and anti-inflammatory activities, [115] beneficial effects on capillary fragility, [116] and an ability to inhibit human platelet aggregation, [117] anti-ulcer [118, 119] and anti-allergenic [120] properties.

From flavanones other classes of flavonoids are biosynthesized with the aid of specific enzymes.

a. Flavones

Flavones are biosynthesized from flavanone with the aid of flavone synthase as catalyst. This enzyme catalyzes a double bond formation between C2 and C3 of flavanones.

Two FNS (I and II) enzyme systems have been described in dicots for flavone biosynthesis. FNSI is a soluble 2-oxoglutarate-dependent dioxygenase (2-ODD),

while FNSII is a cytochrome P-450-dependent monooxygenase enzyme system. FNSI and most FNSII enzymes convert flavanones to flavones directly [121].

Flavones are one of the largest subgroups of flavonoids [121]. Flavones are involved in various interactions with microbes, insects, and other plants [122–124]. In addition to their extensive functions in plants' biochemistry and physiology, flavones are also essential for human nutrition and health [121, 125]. Their pharmacological effects, such as antioxidant, antiviral, anti-inflammatory activities and potential, have made these compounds increasingly popular as dietary constituents or supplements [121].



b.Flavonol

Flavonols are a major class of the family of flavonoids, molecules that have interesting biological activity such as antioxidant, antimicrobial, hepatoprotective, anti-inflammatory, and vasodilatation effects, and they have been considered as potential anticancer agents [126, 127]. Examples of flavonol include fisetin, quercetin, kaempferol, myricetin etc.

The biosynthetic pathway for the synthesis of flavanols is shown below.



c. Isoflavones

Isoflavones are a polyphenol class usually found in legumes, including soybeans, chickpeas, fava beans, pistachios, peanuts, and other fruits and nuts [128]. Soybeans are the richest source of isoflavones, and soy foods and ingredients contain varying concentrations of isoflavones [129]. Isoflavones can be biosynthesized from flavanone with the aid of isoflavone synthase as a catalyst, as shown below.



Common isoflavones include daidzin, genistin, biochanin A, and formononetin [130]. Isoflavones exhibit antioxidant, anticancer, antimicrobial, anti-inflammatory, antiosteoporotic, and estrogenic properties [131–136]. Several studies have also shown that isoflavonoids may contribute to other multiple additional health benefits by reducing cardiovascular risk, osteoporosis, and decreasing the intensity of bone resorption [137].

d.Anthocyanidins

Anthocyanidins are a group of phytochemicals, which are natural pigments responsible for blue, red, purple and orange colors present in many fruits and vegetables and many fruits- and vegetable-based food products. Over and above 500 different anthocyanidins are known and described in the literature [138, 139]. This flavonoid group dominates teas, honey, fruits, vegetables, nuts, olive oil, cocoa and cereals. They can also be found in berries (e.g. black currant, blueberries, strawberries, elderberries), their juices, as well as red wine [140]. Cyanidin, pelargonidin, delphinidin, malvidin, petunidin and peonidin are the most common anthocyanidins present in fruits and vegetables.

The number and position of the hydroxyl and methoxyl moiety is the determinant of different types of anthocyanidins [141]. Anthocyanidins have been reported to have some essential pharmacological role in cardiovascular disease, cholesterol decomposition, visual acuity, as well as antioxidant efficacy, and cytotoxicity [142]. Anthocyanidins can be synthesized as shown below.



e. Flavan-3-ol

The most common flavonoids in the diet, flavan-3-ols are considered functional ingredients of beverages, fruits and vegetables, food grains, herbal remedies, dietary supplements, and dairy products. Flavan-3-ols have been reported to exhibit several pharmacological effects by acting as an antioxidant, anticarcinogen, cardio-preventive, antimicrobial, anti-viral, and neuroprotective agents [143]. Flavan-3-ol can be synthesized as shown below, from dihydroflavanol.



Flavonoids are an essential group of secondary metabolites that play so much role for the benefit of plants in which they exist or in their surroundings and for the health benefit of humanity. Understanding the biosynthetic processes of flavonoids and their pharmacological effects will aid proper utilization of flavonoids for health benefits and also draw more attention to the development of a synthetic laboratory process for the synthesis of flavonoid classes.

Author details

InnocentMary IfedibaluChukwu Ejiofor^{*} and Maria-Goretti Chikodili Igbokwe Department of Pharmacognosy and Traditional Medicine, Faculty of Pharmaceutical Sciences, Nnamdi Azikiwe University, Awka, Anambra State, Nigeria

*Address all correspondence to: ii.ejiofor@unizik.edu.ng

IntechOpen

© 2021 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/3.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

References

[1] Burak M & Imen Y. Flavonoids and their antioxidant properties. Turkiye Klin Tip Bil Derg, 1999, 19, 296-304.

[2] Ovando C, Hernandez D, Hernandez E, et al. Chemical studies of anthocyanins: a review. Food Chem, 2009, 113, 859-871.

[3] Lee Y, Yuk D, Lee J, et al. Epigallocatechin-3-gallate prevents lipopolysaccharide-induced elevation of β -amyloid generation and memory deficiency. Brain Res, 2009, 1250, 164-174.

[4] Panche A. N., Diwan A. D., and Chandra S. R. Flavonoids: an overview. Journal of Nutritional Science, 2016, 5.

[5] Creative Enzymes. Lyase Introduction. (https://www.creative enzymes.com/ resource/lyase-introduction 22.html, accessed on 01/02/2021

[6] G.W. Cheng, P.J. Breen, Activity of phenylalanine ammonia-lyase (PAL) and concentrations of anthocyanins and phenolics in developing strawberry fruit, J. Amer. Soc. Hort. Sci. 116 (1991) 865-869.

[7] Yuri Dyakov, Vitaly Dzhavakhiya, Timo Korpela, 2007, Comprehensive and Molecular Phytopathology, Elsevier Science.

[8] Xu Z., Zhang D, Hu J, Zhou X, Ye X, Reichel K, Stewart N, Syrenne R, Yang X, Gao P, et al. Comparative genome analysis of lignin biosynthesis gene families across the plant kingdom. BMC Bioinform, 2009, 10, S3.

[9] Lichtenthaler, H.K.; Schweiger, J. Cell wall bound ferulic acid, the major substance of the blue-green fluorescence emission of plants. J. Plant Physiol. 1998, 152, 272-282.

[10] Kroon P.A., Williamson G., Hydroxycinnamates in plants and food: current and future perspectives. J. Sci. Food Agric. 1999, 79, 355-361.

[11] Martin-Tanguy J, Cabanne F, Perdrizet E. Martin, C. The distribution of hydroxycinnamic acid amides in flowering plants. Phytochemistry 1978, 17, 1927-1928.

[12] Clifford M.N. Chlorogenic acids and other cinnamates—nature, occurrence and dietary burden. J. Sci. Food Agric. 1999, 79, 362-372.

[13] De P., Baltas M., Bedos-Belval F. Cinnamic acid derivatives as anticancer agents-a review. Curr. Med. Chem. 2011, 18, 1672-1703.

[14] De P., Veau D., Bedos-Belval F., Chassaing S., Baltas M. Cinnamic derivatives in tuberculosis. In Understanding Tuberculosis—New Approaches to Fighting against Drug Resistance; Cardona, P.-J., Ed.; InTech Publishing: Rijeka, Croatia, 2012; Chapter 15.

[15] Wiesner J., Mitsch A., Wißner P., Jomaa H., Schlitzer M. Structure– activity relationships of novel antimalarial agents. Part 2: Cinnamic acid derivatives. Bioorg. Med. Chem. Lett. 2001, 11, 423-424.

[16] Tawata S.; Taira S., Kobamoto N., Zhu J., Ishihara M.; Toyama S. Synthesis and antifungal activity of cinnamic acid esters. Biosci. Biotechnol. Biochem. 1996, 60, 909-910.

[17] Sova M. Antioxidant and antimicrobial activities of cinnamic acid derivatives. Mini Rev. Med. Chem. 2012, 12, 749-767.

[18] Lapeyre C., Delomenède M.,Bedos-Belval F., Duran H.,Nègre-Salvayre A., Baltas M. Design,synthesis, and evaluation ofpharmacological properties of cinnamic

derivatives as antiatherogenic agents. J. Med. Chem. 2005, 48, 8115-8124.

[19] Simonyan A. Activity of cinnamic acid derivatives and new methods for their synthesis (review). Pharm. Chem. J. 1993, 27, 92-100.

[20] Wall V.M., Eisenstadt A., Ager D. J., Laneman S.A. The Heck reaction and cinnamic acid synthesis by heterogeneous catalysis. Platin. Metals Rev. 1999, 43, 138-144.

[21] Sharma P. Cinnamic acid derivatives: A new chapter of various pharmacological activities. J. Chem. Pharm. Res. 2011, 3, 403-423.

[22] Guzman J. D. Natural Cinnamic Acids, Synthetic Derivatives and Hybrids with Antimicrobial Activity. Molecules 2014, 19, 19292-19349.

[23] American Chemical Society. Molecule of the Week Archive, *trans*-Cinnamic acid, 2014. https://www.acs. org/content/acs/en/molecule-of-theweek/archive/c/trans-cinnamicacid. html, accessed on 10/01/2021.

[24] Koopmann E., Logemannm, E. & Hahlbrock K. Regulation and functional expression of cinnamate 4-hydroxylase from parsley. Plant Physiol., 1999, 119, 49-56.

[25] Blount J. W. et al. Altering Expression of Cinnamic Acid 4-Hydroxylase in Transgenic Plants Provides Evidence for a Feedback Loop at the Entry Point into the Phenylpropanoid Pathway. Plant Physiol., 2000, 122, 107-116.

[26] Blee K. A. et al. Antisense and sense expression of cDNA coding for CYP73A15, a class II cinnamate
4-hydroxylase, leads to a delayed and reduced production of lignin in tobacco.
Phytochemistry, 2001, 57, 1159-1166.

[27] Ro D. K., Mah N., Ellis B. E. & Douglas C. J. Functional characterization

and subcellular localization of poplar (Populus trichocarpa£Populus deltoids) cinnamate 4-hydroxylase. Plant Physiol., 2001, 126, 317-329.

[28] p-Coumaric acid' 2021. *Wikipedia*. Available at https://en.wikipedia.org/ wiki/P-Coumaric_acid, Accessed : 02 February 2021.

[29] Huseyin Boz. p-Coumaric acid in cereals: presence, antioxidant and antimicrobial effects, International Journal of Food Science and Technology 2015, 50, 2323-2328.

[30] Garrait G., Jarrige J. F., Blanquet S., Beyssac E., Cardot J.M. & Alric M. Gastrointestinal absorption and urinary excretion of trans-cinnamic and p-coumaric acids in rats. Journal of Agriculture and Food Chemistry, 2006, 54, 2944-2950.

[31] Zilic S., Sukalovic V. H-T., Dodig D., Maksimovic V., Maksimovic M. & Basic Z. Antioxidant activity of small grain cereals caused by phenolics and lipid soluble antioxidants. Journal of Cereal Science, 2011, 54, 417-424.

[32] Lou Z., Wang H., Rao S., Sun J., Ma C. & Li J. p-Coumaric acid kills bacteria through dual damage mechanisms. Food Control, 2012, 25, 550-554.

[33] Sanchez-Maldonado A. F., Schieber A. & G€anzle M.G. Structure-function relationships of the antibacterial activity of phenolic acids and their metabolism by lactic acid bacteria. Journal of Applied Microbiology, 2011, 111, 1176-1184.

[34] Halliwell B. & Ahluwalia S. Hydroxylation of p-coumaric acid by horseradish peroxidase: the role of superoxide and hydroxylradicals. Biochemical Journal, 1976, 153, 513-518.

[35] Castelluccio C., Paganga G., Melikian N. et al. Antioxidant potential of intermediates in phenylpropanoid metabolism in higher plants. FEBS Letters, 1995, 368, 188-192.

[36] Zang L.Y., Cosma G., Gardner H., Shi X., Castranova V. & Vallyathan V. Effect of antioxidant protection by p-coumaric acid on low-density lipoprotein cholesterol oxidation. American Journal of Physiology – Cell Physiology, 2000, 279, 954-960.

[37] Rao R. S. P. and Muralikrishna G. Non-starch polysaccharidephenolic acid complexes from native and germinated cereals and millet. Food Chemistry, 2004, 84, 527-531.

[38] Kovacova M. & Malinova E. Ferulic and coumaric acids, total phenolic compounds and their correlation in selected oat genotypes. Czech Journal of Food Science, 2007, 25, 325-332.

[39] Gani A., Wani S.M., Masoodi F.A. & Hameed G. Wholegrain cereal bioactive compounds and their health benefits: a review. Journal of Food Process and Technology, 2012, 3, 146.

[40] Yoon S.-A., Kang S.-I., Shin H.-S. et al. p-Coumaric acid modulates glucose and lipid metabolism via AMP-activated protein kinase in L6 skeletal muscle cells. Biochemical and Biophysical Research Communications, 2013, 432, 553-557.

[41] Biswick T., Park D., Shul Y.G. & Choy J.H. p- Coumaric acid-zinc basic salt nanohybrid for controlled release and sustained antioxidant activity. Journal of Physics and Chemistry of Solids, 2010, 71, 647-664.

[42] Shabir G., Anwar F., Sultana B. et al. Antioxidant and antimicrobial attributes and phenolics of different solvent extracts from leaves, flowers and bark of gold mohar [*Delonix regia* (Bojerex Hook.) Raf.]. Molecules, 2011, 16, 7302-7319.

[43] Navaneethan D. & Rasool M. p-Coumaric acid, a common dietary polyphenol, protects cadmium chlorideinduced nephrotoxicity in rats. Renal Failure, 2014, 36, 244-251.

[44] Kong C. S., Jeong C. H., Choi J. S., Kim K. J. and Jeong J. W. Antiangiogenic Effects of P-Coumaric Acid in Human Endothelial Cells. Phytother. Res. 2012.

[45] Lehninger AL, Nelson DL, Cox MM, editors. Lehninger's Principles of Biochemistry, 4 ed., New York, NY, USA, 2004.

[46] Manubolu M., Goodla L., Pathakoti K. and Malmlo"f K. (2018). Enzymes as direct decontaminating agents—mycotoxins. In C. S. Nunes and V. Kumar (Eds). *Enzymes in Human and Animal Nutrition* (pp. 312-330). USA: Elsevier Academic press.

[47] National Cancer Institute Thesaurus. Coenzyme A. https://ncit.nci.nih. gov/ncitbrowser/ConceptReport. jsp?dictionary=NCI_Thesaurus&ns=NCI_ Thesaurus&code=C384, Accessed : 02 February 2021.

[48] Palmer T. and Bonner P. L. (2007). Enzymes Biochemistry, Biotechnology, Clinical Chemistry (pp. 204-220). United Kingdom: Woodhead Publishing Limited.

[49] Boudet A. Evolution and current status of research in phenolic compounds. Phytochemistry, 2007, 68, 2722-2735.

[50] Vogt, T. Phenylpropanoid Biosynthesis. Molecular Plant, 2010, 3(1), 2-20.

[51] Sanchez IJF. (2008) Polyketide synthase in *Cannabis sativa* L. PhD thesis, Leiden University, Leiden, The Netherlands.

[52] IUPAC, Compendium of Chemical Terminology, 2nd ed. (the "Gold Book") (1997). Online corrected version:(2006) "Polyketides.

[53] Hahlbrock K., Scheel D. (1989) Physiology and molecular biology of phenylpropanoid metabolism. Annu Rev Plant Physiol Plant Mol Biol 40:347-369.

[54] Lanz T., Tropf S., Marner FJ, Schröder J, Schröder G. The role of cysteines in polyketide synthases: sitedirected mutagenesis of resveratrol and chalcone synthases, two key enzymes in different plant-specific pathways. J Biol Chem, 1991, 266, 9971-9976.

[55] Hahlbrock K., Zilg H., Grisebach H. Stereochemistry of the enzymatic cyclisation of 4,2', 4'-trihydroxychalcone to 7,4'-dihydroxyflavanone by isomerases from Mung Bean seedlings. Eur J Biochem, 1970. 15, 13-18.

[56] Jez JM, Austin MB, Ferrer JL, Bowman ME, Schröder J, Noel JP. Structural control of polyketide formation in plant-specific polyketide synthases. Chem Biol, 2000, 40, 1-12.

[57] Chen H., Kim H. U., Weng H., and Browse J. Malonyl-CoA Synthetase, Encoded by ACYL ACTIVATING ENZYME13, is Essential for Growth and Development of Arabidopsis. The Plant Cell, 2011, 23, 2247-2262.

[58] Wu W., Ye H., Wan L., Han X., Wang G., Hu J., Tang M., Duan X., Fan Y., He S., Huang L. Millepachine, a novel chalcone, induces G 2/M arrest by inhibiting CDK1 activity and causing apoptosis via ROS-mitochondrial apoptotic pathway in human hepatocarcinoma cells in vitro and in vivo. Carcinogenesis, 2013, 34, 1636-1643.

[59] Shibata S., Inoue H., Iwata S., Ma R., Yu L., Ueyama H., Takayasu J., Hasegawa T., Tokuda H., Nishino A., Nishino H. (1991) Inhibitory efects of licochalcone A isolated from Glycyrrhiza infata root on infammatory ear edema and tumor promotion in mice. Planta Med, 1991, 57, 221-224. [60] Edwards ML, Stemerick DM,Sunkara PS. Chalcones: a new class of antimitotic agents. J Med Chem, 1990, 33, 1948-1954.

[61] Wang Y., Chan FL, Chen S., Leung LK (2005) The plant polyphenol butein inhibits testosterone-induced proliferation in breast cancer cells expressing aromatase. Life Sci, 2005 77, 39-51.

[62] Yit CC, Das NP. Cytotoxic efect of butein on human colon adenocarcinoma cell proliferation. Cancer Lett, 1994, 82, 65-72.

[63] Anto RJ, Sukumaran K., Kuttan G., Rao MNA, Subbaraju V., Kuttan R. Anticancer and antioxidant activity of synthetic chalcones and related compounds. Cancer Lett, 1995, 97, 33-37.

[64] Hatano T., Takagi M., Ito H., Yoshida T. (1997) Phenolic constituents of liquorice. VII. A new chalcone with a potent radical scavenging activity and accompanying phenolics from liquorice. Chem Pharm Bull, 1997, 45, 1485-1492.

[65] Koorbanally NA,

RandrianarivelojosiaM., MulhollandDA, van Uford LQ, van den Berg AJ. Chalcones from the seed of *Cedrelopsis* grevei (Ptaeroxylaceae). Phytochemistry, 2003, 62:1225-1229.

[66] Cioffi G., Escobar LM, Braca A., De Tommasi N. (2003) Antioxidant chalcone glycosides and favanones from *Maclura* (Chlorophora) *tinctoria*. J Nat Prod, 2003, 66, 1061-1064.

[67] Doan TN, Tran DT. Synthesis, antioxidant and antimicrobial activities of a novel series of chalcones, pyrazolic chalcones, and allylic chalcones. Pharmacol Pharm, 2011, 2, 282.

[68] Ballesteros JF, Sanz MJ, Ubeda A., Miranda MA, Iborra S., Paya M., Alcaraz MJ. Synthesis and pharmacological evaluation of 2'-hydroxychalcones and favones as inhibitors of inflammatory mediators generation. J Med Chem, 1995, 38, 2794-2797.

[69] Hirai S., Kim YI, Goto T., Kang MS, Yoshimura M., Obata A., Yu R., Kawada T. Inhibitory effect of naringenin chalcone on inflammatory changes in the interaction between adipocytes and macrophages. Life Sci, 2007, 81, 1272-1279.

[70] Funakoshi TM, Okamoto K., Izumi R., Tago K., Yanagisawa K., Narukawa Y., Kiuchi F., Kasahara T., Tamura H. Anti-infammatory activity of favonoids in Nepalese propolis is attributed to inhibition of the IL-33 signaling pathway. Int Immunopharmacol, 2015, 25, 189-198.

[71] Yang EB, Zhang K., Cheng LY, Mack P. Butein, a specifc protein tyrosine kinase inhibitor. Biochem Biophys Res Commun, 1998, 245, 435-438.

[72] Zhao F., Nozawa H., Daikonnya A., Kondo K., Kitanaka S. Inhibitors of nitric oxide production from hops (*Humulus lupulus* L.). Biol Pharm Bull, 2003, 26, 61-65.

[73] Piramal Life Sciences Ltd. Preparation of heterocyclic benzene compounds as cytokine inhibitors for treating inflammatory disorders, 2011, WO2011/121505.

[74] Won SJ, Liu CT, Tsao LT, Weng JR, Ko HH, Wang JP, Lin CN. Synthetic chalcones as potential anti-infammatory and cancer chemopreventive agents. Eur J Med Chem, 2005, 40:103-112.

[75] Dominguez JN, Leon C., Rodrigues J., de Dominguez NG, Gut J., Rosenthal PJ. Synthesis and antimalarial activity of sulfonamide chalcone derivatives. Il Farmaco, 2005, 60, 307-311.

[76] KyogokuK., HatayamaK., YokomoriS., SazikiR., NakaneS., SasajimaM., SawadaJ., Ohzeki M., Tanaka I. Anti-ulcer efect of isoprenyl favonoids. II. Synthesis and anti-ulcer activity of new chalcones related to sophoradin. Chem Pharm Bull, 1979, 27, 2943-2953.

[77] Ming DS, Lopez A., Hillhouse BJ, French CJ, Hudson JB, Towers GN. Bioactive Constituents from Iryanthera megistophylla. J Nat Prod, 2002, 65, 1412-1416.

[78] Sasajima M., Nakane S., Saziki R., Saotome H., Hatayama K., Kyogoku K., Tanaka I. Studies on the anti-ulcer efects of isoprenyl favonoids (1). The anti-ulcer efects of isoprenyl chalcone extracted from Sophora subprostrata. Folia Pharmacol Japonica, 1978, 74, 897-905.

[79] Ishitsuka H., Ninomiya YT, Ohsawa C., Fujiu M., Suhara Y. Direct and specifc inactivation of rhinovirus by chalcone Ro09-0410. Antimicrob Agents Chemother, 1982, 22, 617-621.

[80] Kaul TN, Middleton E Jr, Ogra PL. Antiviral efect of flavonoids on human viruses. J Med Vir, 1985, 15, 71-79.

[81] Mustafa KA, Kjaergaard HG, Perry NB, Weavers RT. Hydrogenbonded rotamers of 2',4',6'-trihydroxy-3'formyldihydrochalcone, an intermediate in the synthesis of a dihydrochalcone from *Leptospermum recurvum*. Tetrahedron, 2003, 59, 6113-6120.

[82] Chen M, Christensen SB, Blom J, Lemmich E, Nadelmann L, Fich K, Theander TG, Kharazmi A (1993) Licochalcone A, a novel antiparasitic agent with potent activity against human pathogenic protozoan species of Leishmania. Antimicrob Agents Chemother, 1993, 37, 2550-2556.

[83] Cheng ZJ, Lin CN, Hwang TL, Teng CM. Broussochalcone A, a potent antioxidant and efective suppressor of inducible nitric oxide synthase in lipopolysaccharide-activated

macrophages. Biochem Pharmacol, 2001, 61, 939-946.

[84] Rashid KA, Mullin CA, Mumma RO. Structure-mutagenicity relationships of chalcones and their oxides in the Salmonella assay. Mutat Res, 1986, 169, 71-79.

[85] Forkmann G. and Heller W. (1999).
Polyketides and Other Secondary
Metabolites Including Fatty Acids and
Their Derivatives. In D. Barton, K.
Nakanishi and O. Meth-Cohn (Eds)
Comprehensive Natural Products
Chemistry, Vol. 1 (pp. 714-743). Elsevier
Science Ltd.

[86] Cheng A., Zhang X., Han X., Zhang Y., Gao S., Liu C., et al. Identification of chalcone isomerase in the basal land plants reveals an ancient evolution of enzymatic cyclization activity for synthesis of flavonoids. New Phytol, 2018, 217, 909-924.

[87] Ralston L., Subramanian, S., Matsuno, M., and Yu, O. Partial reconstruction of flavonoid and isoflavonoid biosynthesis in yeast using soybean type I and type II chalcone isomerases. Plant Physiol, 2005, 137, 1375-1388.

[88] Jez J. M., Bowman M. E., Dixon R. A., and Noel J. P. Structure and mechanism of the evolutionarily unique plant enzyme chalcone isomerase. Nat. Struct. Biol, 2000, 7, 786-791.

[89] Jez J. M., Bowman M. E., and Noel J. P. Role of hydrogen bonds in the reaction mechanism of chalcone isomerase. J. Biol. Chem, 2002, 41, 5168-5176.

[90] Dixon R. A., Blyden E. R., Robbins M. P., Tunen A. V., and Mol J. N. Comparative biochemistry of chalcone isomerases. Phytochemistry, 1988, 27, 2801-2808.

[91] Ngaki M. N., Louie G. V., Philippe R. N., Manning G., Pojer F., Bowman M.

E., et al. Evolution of the chalconeisomerase fold from fatty-acid binding to stereospecific catalysis. Nature, 2012, 485, 530-533.

[92] Morita Y., Takagi K., Fukuchi-Mizutani M., Ishiguro K., Tanaka Y., Nitasaka E., et al. (2014). A chalcone isomerase like protein enhances flavonoid production and flower pigmentation. Plant J, 2014, 78, 294-304.

[93] Das A. B., Goud V. V., Das C. (2019) Phenolic Compounds as Functional Ingredients in Beverages. In A. Grumezescu, A. M. Holban (Eds). The Science of Beverages; Valueadded Ingredients and Enrichments of Beverages. Elsevier Science & Technology.

[94] Wang H, Cao G, Prior RL. Total antioxidant capacity of fruits. J Agric Food Chem 1996, 44(3), 701-5.

[95] Cao G, Sofic E, Prior RL. Antioxidant and prooxidant behavior of flavonoids: structure-activity relationships. Free Radic Biol Med, 1997, 22(5), 749-60.

[96] Pietta PG. Flavonoids as antioxidants. J Nat Prod, 2000, 63 (7):1035-42.

[97] Rice-Evans C. Flavonoid antioxidants. Curr Med Chem, 2001, 8 (7), 797-807.

[98] Kaur C, Kapoor HC. Antioxidants in fruits and vegetables-the millennium's health. Int J Food Sci Tech, 2001, 36(7), 703-25.

[99] Burda S, Oleszek W. Antioxidant and antiradical activities of flavonoids. J Agric Food Chem 2001;49(6):2774-9.

[100] Sa'nchez-Moreno C, Plaza L, De Ancos B, Cano MP. Quantitative bioactive compounds assessment and their relative contribution to the antioxidant capacity of commercial orange juices. J Sci Food Agric, 2003, 83(5), 430-9. [101] Zhang HY, Yang DP, Tang GY.Multipotent antioxidants: from screening to design. Drug Discov Today, 2006, 11(15-16):749-54.

[102] Drewnowski A, Gomez-Carneros C. Bitter taste, phytonutrients and the consumer: a review. Am J Clin Nutr, 2000, 72 (6), 1424-35.

[103] Kris-Etherton PM, Lefevre M, Beecher GR, Gross MD, Keen CL, Etherton TD. Bioactive compounds in nutrition and healthresearch methodologies for establishing biological function: the antioxidant and antiinflammatory effects of flavonoids on atherosclerosis. Ann Rev Nutr, 2004, 24, 511-38.

[104] Hertog MG. Flavonoid intake and long-term risk of coronary heart-disease and cancer in the seven countries study. Arch Intern Med, 1995, 155(4), 381-6.

[105] Knekt P, Jarvinen R, Reunanen A, Maatela J. Flavonoid intake and coronary mortality in Finland: a cohort study. Brit Med J, 1996, 312(7029), 478-81.

[106] Yochum L., Kushi LH, Meyer K., Folsom AR. Dietary flavonoid intake and risk of cardiovascular disease in postmenopausal women. Am J Epidemiol, 1999, 149(10), 943-9.

[107] Gross M. Flavonoids and cardiovascular disease. Pharm Biol, 2004, 42(Suppl.), 21-35.

[108] Steinmetz KA, Potter JD. Vegetables, fruit, and cancer prevention: a review. J Am Diet Assoc, 1996, 96(10), 1027-39.

[109] Le Marchand L, Murphy SP, Hankin JH, Wilkens LR, Kolonel LN. Intake of flavonoids and lung cancer. J Natl Cancer Inst, 2000, 92(2), 154-60.

[110] Kris-Etherton PM, Hecker KD, Bonanome A, Coval SM, Binkoski AE, Hilpert KF, et al. Bioactive compounds in foods: their role in the prevention of cardiovascular disease and cancer. Am J Med, 2002, 113(Suppl. 9B), 71-88.

[111] Nichenametla SN, Taruscio TG, Barney DL, Exon JH. A review of the effects and mechanism of polyphenolics in cancer. Crit Rev Food Sci, 2006, 46(2), 161-83.

[112] Moon YJ, Wang X, Morris ME. Dietary flavonoids: effects on xenobiotic and carcinogen metabolism. Toxicol In Vitro, 2006, 20 (2), 187-210.

[113] Asres K, Seyoum A, Veeresham C, Bucar F, Gibbons S. Naturally derived anti-HIV agents. Phytother Res, 2005, 19(7), 557-81.

[114] Cushnie TPT, Lamb AJ. Antimicrobial activity of flavonoids. Int J Antimicrob Agent, 2005, 26(5), 343-56.

[115] Kim HP, Son KH, Chang HW, Kang SS. Anti-inflammatory plant flavonoids and cellular action mechanisms. J Pharmacol Sci, 2004, 96(3), 229-45.

[116] Benavente-Garcı'a O, Castillo J, Marı'n FR, Ortun~o A, Del Rı'o JA.mUses and properties of Citrus flavonoids.
J Agric Food Chem, 1997, 45(12), 4505-15.

[117] Tijburg LBM, Mattern T., Folts JD,
Weisgerber UM, Katan MB. Tea
flavonoids and cardiovascular diseases:
a review. Crit Rev Food Sci, 1997,
37(8), 771-85.

[118] Borrelli F., Izzo AA. The plant kingdom as a source of anti-ulcer remedies. Phytother Res, 2000, 14(8), 581-91.

[119] Wightman JD. Red berries and their health benefits. In: Nutraceutical Beverages: Chemistry, Nutrition, and Health Effects (ACS Symp. Ser. 871), 2004, pp. 123-132.

[120] Middleton E, Kandaswami C. Effects of flavonoids on immune and

inflammatory cell functions. Biochem Pharmacol, 1992, 43(6), 1167-79.

[121] Martens S., & Mithöfer A. Flavones and flavone synthases. Phytochemistry, 2005, 66(20), 2399-2407.

[122] Fisher R. F. & Long S. R. Rhizobiumplant signal exchange. Nature, 1992, 357, 655-660.

[123] Simmonds M. S. Flavonoid-insect interactions: recent advances in our knowledge. Phytochemistry, 2003, 64, 21-30.

[124] Soriano I., Asenstorfer R., Schmidt O. and Riley I. Inducible flavone in oats (*Avena sativa*) is a novel defense against plant-parasitic nematodes. Phytopathology, 2004, 94, 1207-1214.

[125] Pandey A. et al. AtMYB12 expression in tomato leads to large scale differential modulation in transcriptome and flavonoid content in leaf and fruit tissues. Sci. Rep. 2015, -Uk 5.

[126] Cushnie T. P. T., & Lamb A. J. Antimicrobial activity of flavonoids. International Journal of Antimicrobial Agents, 2005, 26(5), 343-356.

[127] Erlund I. (2004). Review of the flavonoids quercetin, hesperetin, and naringenin. Dietary sources, bioactivities, bioavailability, and epidemiology. Nutrition Research, 24(10), 851-874.

[128] United States Department of Agriculture. (2008). A Database for the Isoflavone Content of Selected Foods, Release 2.0. September

[129] Thrane M., Paulsen P. V., Orcutt M.
W. and Krieger T. M. (2017). Soy Protein: Impacts, Production, and Applications.
In S. Nadathur, J. Wanasundara, L.
Scanlin (Eds). *Sustainable Protein Sources* (pp. 23-45). Elsevier Inc.

[130] COT (committee on toxicity), Committee on toxicity of chemicals in food, consumer products and the environment. Phytoestrogens and health; 2003, London, UK: Crown Copyright. Available at http://www. food.gov.uk/science/ouradvisors/ toxicity/COTwg/wg_phyto

[131] Jang C.H.; Lim J.K.; Kim J.H.;
Park C.S.; Kwon D.Y.; Kim Y.S.;
Shin D.H.; Kim J.S. Change of Isoflavone Content during Manufacturing of Cheonggukjang, a Traditional Korean Fermented Soyfood. Food Sci.
Biotechnol. 2006, 15, 643-646.

[132] Iwasaki M., Inoue M., Otani T., Sasazuki S., Kurashi N., Miura T., Yamoto S., Tsugane, S. Plasma isoflavone level and subsequent risk of breast cancer among Japanese women: A nested casecontrol study from Japan Public Health Center-base prospective study group. J. Clin. Oncol. 2008, 26, 1677-1683.

[133] Bezek S., Ujhazy E., Mach M., Navarova J., Dubovicky M.Developmental origin of chronic diseases: Toxicological implication. Interdisc.Toxicol. 2008, 1, 29-31.

[134] Ørgaard A., Jensen L. The effects of soy isofl avones on obesity. Exp. Biol. Med. 2008, 233, 1066-1080.

[135] Sabudak T., Guler N., Trifolium L. A review on its phytochemical and pharmacological profile. Phytother. Res. 2009, 23, 439-446.

[136] Choi E.J., Kim G.H. Antiproliferative activity of daidzein and genistein may be related to $ER\alpha/c$ -erbB-2 expression in human breast cancer cells. Mol. Med. Rep. 2013, 7, 781-784.

[137] Rochfort S., Panozzo J. Phytochemicals for health, the role of pulses. J. Agric. Food Chem. 2007, 55, 7381-7994.

[138] de Pascual-Teresa S., Moreno DA, Garcia-Viguera C. Flavanols and anthocyanins in cardiovascular health:

Bioactive Compounds - Biosynthesis, Characterization and Applications

a review of current evidence. Int J Mol Sci, 2010, 11, 1679-1703.

[139] de Pascual-Teresa S., Sanchez-Ballesta MT. Anthocyanins: from plant to health. Phytochem Rev, 2008, 7, 281-299.

[140] Roy HJ, Lundy S., Eriksen C., Kalicki B. Anthocyanins. Pennigton Nutrition Series, 2009.

[141] de Pascual-Teresa S., Santos-Buelga C., RivasGonzalo JC. Quantitative analysis of flavan-30ls in Spanish foodstuffs and beverages. J Agric Food Chem, 2000, 48, 5331-5337.

[142] Bagchi D., Sen CK, Bagchi M., Atalay M. Antiangiogenic, antioxidant, and anti-carcinogenic properties of a novel anthocyanin-rich berry extract formula. Biochem, 2004, 69(1), 75-80.

[143] Aron P. M., & Kennedy J. A. Flavan-3-ols: Nature, occurrence and biological activity. Molecular Nutrition & Food Research, 2008, 52(1), 79-104.

