

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

186,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Introductory Chapter: Biodiversity of Mexico

Levente Hufnagel and Ferenc Mics

1. Introduction

1.1 Flora, fauna and vegetation

Mesoamerica (starting from the southern states of Mexico) differs from Central America, which is a geopolitical name. The expression “Middle America” is in use as well, which involves all areas south from the border of the US including the Islands of the Caribbean [1]. In this chapter the biodiversity of Mexico is presented and the literature on its protection is analyzed.

Mexico being the largest country in the region is very rich in species in itself (Tables 1–4). Lot of species occur even in the dry northern areas. In the Chihuahuan Desert 826 plant species are noted by Villarreal-Quintanilla et al. [7], out of which 560 are endemic, 165 are quasi-endemic and 176 are microendemic. 116 taxa can be originated from a non-arid habitat. The most species-rich are Cactaceae with 141, Asteraceae with 106, Boraginaceae with 34 and Brassicaceae with 31 species. On the California Peninsula 723 endemic species are noted by Riemann and Exequiel [8], claiming that the great number of endemic species is due to the heterogeneity of the environment. The flora and fauna are very interesting because the area of Mexico involves the border of *Neotropis* and *Nearctis* (Mexican Transition Zone), which is not exactly a border but rather a wide transition zone, its accurate definition is yet to be created. The determination is based on the distribution of the endemic genera characteristic to one or the other area. The several results obtained regarding this vary hugely. The determination of the location and width of the transition zone is different among authors without a consensus, which requires further floristic examinations [9]. Vegetation varies depending on the topography that has a great role in the formation of the great number of endemic species as well [10].

| Group | Number of species worldwide | Number of species in Mexico | Worldwide/Mexico % | Number of endemic species | Endemic species % |
|-----------------|-----------------------------|-----------------------------|--------------------|---------------------------|-------------------|
| Vascular plants | 248 428 | 18000–30000 | 7–12 | 10000–15000 | 33–50 |
| Amphibians | 4222 | 284 | 7 | 169 | 60 |
| Reptiles | 6458 | 717 | 11 | 368 | 51 |
| Birds | 9040 | 1050 | 12 | 125 | 12 |
| Mammals | 4629 | 450 | 10 | 140 | 31 |

Table 1.
Biodiversity of Mexico according to Mittermeier et al. [2].

| Vegetation | % area | Number of species | Endemisms % |
|--------------------------|--------|-------------------|-------------|
| Cloud forest | 0.7 | 3000 | 30 |
| Rainforest | 4.4 | 5000 | 5 |
| Pine and oak | 12.9 | 7000 | 70 |
| Xerophyl and steppe | 34.8 | 6000 | 60 |
| Wetlands | 1.43 | 1000 | 15 |
| Tropical deciduous | 7.9 | 6000 | 40 |
| Agricultural and ruderal | ND | 2000 | 20 |

Table 2.
Flora of Mexico according to vegetation types based on Flores-Villela and Gerez [3].

| Biom/ecoregions | Area (km ²) | Estimated number of species | Remaining intact vegetation % |
|---|-------------------------|-----------------------------|-------------------------------|
| Tropical and Subtropical Moist Broadleaf Forests | | | |
| Chiapas moist forests | 5759 | 3000–4500 | 67 |
| Chimalpas montane forests | 2076 | 2000–3000 | 49 |
| Pantanos de Centla | 17152 | 1500–3000 | 11 |
| Petén-Veracruz moist forests | 148604 | 5000–8000 | 31 |
| Sierra de los Tuxtlas | 3890 | 2400–3500 | 11 |
| Sierra Madre de Chiapas moist forests | 11218 | 3500–4700 | 34 |
| Veracruz moist forests | 68949 | 4500–7000 | 20 |
| Veracruz montane forests | 4942 | 2200–3500 | 55 |
| Yucatán moist forests | 69485 | 1300–1900 | 64 |
| Tropical and Subtropical Dry Broadleaf Forests | | | |
| Bajío dry forests | 37282 | 2900–5000 | 0,64 |
| Balsas dry forests | 62249 | 2500–5100 | 2 |
| Central American dry forests | 67777 | 2800–400 | 12 |
| Chiapas Depression dry forests | 13974 | 1500–3500 | 7 |
| Jalisco dry forests | 26051 | 1000–2500 | 26 |
| Revillagigedo Islands dry forests | 210 | ND | ND |
| Sierra de la Laguna dry forests | 3975 | 500–1000 | 0,008 |
| Sinaloan dry forests | 77364 | 1700–2500 | 13 |
| Sonoran-Sinaloan transition subtropical dry forest | 50326 | ND | ND |
| Southern Pacific dry forests | 42283 | 2500–5100 | 15 |
| Veracruz dry forests | 6616 | 900–2000 | 5 |
| Tropical and Subtropical Coniferous Forests | | | |
| Central American pine-oak forests | 110948 | 4000–6000 | 42 |
| Sierra de la Laguna pine-oak forests | 1061 | 700–1200 | 4 |

| Biom/ecoregions | Area (km ²) | Estimated number of species | Remaining intact vegetation % |
|--|-------------------------|-----------------------------|-------------------------------|
| Sierra Juárez and San Pedro Mártir pine-oak forests | 4000 | ND | ND |
| Sierra Madre de Oaxaca pine-oak forests | 14299 | 2500–3700 | 55 |
| Sierra Madre del Sur pine-oak forests | 60976 | 3600–5000 | 43 |
| Sierra Madre Occidental pine-oak forests | 222700 | ND | ND |
| Sierra Madre Oriental pine-oak forests | 65600 | ND | ND |
| Trans-Mexican volcanic belt pine-oak forests | 91800 | 4000–6000 | 26 |
| Tropical and Subtropical Grasslands, Savannas, and Shrublands | | | |
| Western Gulf coastal grasslands | 77425 | 2150–2250 | 3 |
| Deserts and Xeric Shrublands | | | |
| Baja California desert | 45940 | 1500–2200 | 8 |
| Central Mexican matorral | 59195 | 2500–4500 | 0,011 |
| Chihuahuan desert | 501896 | 3300–3600 | 50 |
| Gulf of California xeric shrub | 22573 | 900–1900 | 29 |
| Meseta Central matorral | 124975 | 3000–4500 | 4 |
| San Lucan xeric scrub | 3685 | ND | 12 |
| Sonoran desert | 260000 | 2600–3000 | 37 |
| Tamaulipan matorral | 16300 | 1500–2500 | 9 |
| Tamaulipan mezquital | 141500 | 1700–2500 | 23 |
| Tehuacan Valley matorral | 9842 | | 0 |
| Flooded Grasslands and Savannas | | | |
| Central Mexican wetlands | 259 | 100–600 | ND |
| Montane Grasslands and Shrublands | | | |
| Zacatonal | 306 | 150–500 | ND |
| Mangroves | | | |
| Alvarado mangroves | 4534 | 20–400 | 1,12 |
| Marismas Nacionales-San Blas mangroves | 2034 | 20–400 | |
| Mayan Corridor mangroves | 4079 | 20–400 | |
| Mexican South Pacific Coast mangroves | 1168 | 20–400 | |
| Petenes mangroves | 1971 | 20–400 | |
| Ría Lagartos mangroves | 3457 | 20–400 | |
| Tehuantepec-El Manchon mangroves | 2685 | 20–400 | |
| Usumacinta mangroves | 3118 | 20–400 | |
| Mediterranean Forests, Woodlands, and Scrub | | | |
| California chaparral and woodlands | 121000 | 1550–1750 | 1 |

Table 3.
Bioms and ecoregions of Mexico according to Kier et al. [4] and Dinerstein et al. [5].

| State | Birds | Mammals |
|---------------------|-------|---------|
| Veracruz | 664 | 101 |
| Chiapas | 628 | 90 |
| Oaxaca | 687 | 116 |
| Jalisco | 481 | 107 |
| Guerrero | 476 | 72 |
| Puebla | 367 | 76 |
| San Luis Potosí | 469 | 93 |
| Michoacán | 460 | 79 |
| Chihuahua | 329 | 95 |
| Edo. México | 281 | 55 |
| Tamaulipas | 444 | 90 |
| Sonora | 456 | 100 |
| Durango | 308 | 81 |
| Nayarit | 407 | 72 |
| Nuevo León | 252 | 63 |
| Hidalgo | 236 | 59 |
| Morelos | 274 | 46 |
| Sinaloa | 460 | 69 |
| Coahuila | 275 | 80 |
| Tabasco | 370 | 47 |
| Baja California | 292 | 95 |
| Quintana Roo | 340 | 51 |
| Distrito Federal | 222 | 44 |
| Yucatán | 343 | 58 |
| Baja California Sur | 258 | 77 |
| Colima | 318 | 51 |
| Zacatecas | 154 | 75 |
| Guanajuato | 256 | 45 |
| Campeche | 281 | 50 |
| Querétaro | 181 | 36 |
| Aguascalientes | 89 | 33 |
| Tlaxcala | 89 | 21 |

Table 4.
Bird and mammalian fauna of Mexico according to CONABIO [6] (National Commission for knowledge and use of biodiversity) by states. In coastal states marine mammals are included.

The climate formed by the emerging mountains (Sierra Madre Oriental, Sierra Madre Occidental, Trans-Mexico Volcanic Belt) also affects evolutionary processes resulting in the development of new species [11, 12]. The richness of species and the species composition of the community depends on the heterogeneity and diversity of the environment [10]. As Moonlight et al. [13] presents on the example of *Begonia* genus regarding the DNA sequence data of the plastis diversification is fast and new species develop in the hetrogeneous environment. According to their results there were two indipendent colonization events from Africa towards the Neotropis. Two

different clads were reconstructed, which diversified around the middle of miocene in South America and radiation occurred once towards Central America and Mexico.

Today tropical deciduous forest is typical on the western side but it is fragmented due to human activity and only 3% is protected. Agricultural areas are concentrated in areas where the climate is seasonal that is why the reduction is faster than in the case of rainforests [14]. 10% and 19.7% of tropical forest and shrubland, respectively are protected. On the east side 28% of the rainforests are protected (Estado [15]). Examinations of flora and fauna are still not complete, our knowledge on the wildlife of the area is deficient [16, 17]. New species are still identified in Mexico such as *Tryonia* (Caenogastropoda: Cochliopidae) species discovered by Hershler et al. [18] in the creeks of the Chihuahuan Desert (Durango State). Some of these probably have become extinct since they were failed to be found again. The cause of their extinction is the destruction and disappearance of their habitat. According to forecasts the average annual temperature will rise by 3,7–3,8°C to 2090 in Mexico, the amount of precipitation will reduce by 18,2% and AAI (Annual Aridity Index) will rise by 26% [19]. These will result in a 25% decrease in the value of ENS (effective number of species), if the forecasts prove to be accurate [20]. Characteristic plants of Mexico the cacti (Cactaceae) can also become rare by then due to land transforming activity of man, 31% of the species are already endangered. Climate change may cause other species to become endangered. Cacti are drought tolerant but because of climate change species may become endangered due to the reduction or extinction of pollinators and animals dispersing seeds. In the case of cacti often ants carry the seeds (Myrmecochory) only to limited distances. Many species have a small area of distribution (75–100 km²) as well, making them particularly vulnerable [21]. A remarkable example of the destruction of human activity to habitat is the wall that is being built on the border of the US and Mexico, which parts that have already been set up initiated the reduction of size, quality and connections of habitat in the otherwise diverse area [22]. Migration of many amphibian, reptile and mammal species are hindered by the barrier and human activity. This becomes relevant when certain species look for new habitat due to climate change. Obstructing their movement endanger species or certain populations. The genetic diversity of species reduce with the extinction of populations endangering the entire species [23].

2. Environmental protection and biodiversity conservation

Since the nineties Mexico have been involved in the work of international environmental organizations with increasing activity and now have signed 44 international agreements. It is an active stakeholder of organizations such as the CBD (Convention on Biological Diversity) and CITES (Convention on International Trade in Endangered Species of Wild Fauna and Flora), CEC (Commission for Environmental Cooperation), Trilateral Committee, The Ramsar Convention on Wetlands, EMSA (Mesoamerican Biological Corridor and the Mesoamerican Strategy of Environmental Sustainability). The NDP (National Development Plan) can be considered as a legal framework within the country providing basic regulations for the federal government by giving guidance for the work of the government with the development of clear strategies, setting regional aims and measures to be implemented, coordinating institutional and regional programs involving several areas [24].

The program, in which Mexico also participates, aiming to preserve biodiversity is the Payments for Ecosystem Services (PES) that is an economic framework to plan and introduce payment schemes that provide market remuneration for ecosystem services [25]. This tool seems to be eligible for the protection of rainforests in the developing countries. 2.6 million hectares are involved in the program in

Mexico, which in terms of money means 450 million USD and this is one of the biggest among such programs in the world [26]. With these territories approximately 25% of the biodiversity that needs protection is now included in the program [27]. According to Honey-Rosés et al. [28] 3–16% more forest were managed to be preserved along with those habitats that these forests include. Deforestation in those areas that are involved in the program is carried out at a slower rate, than in those that are not involved. This can be beneficial to the population and to the owners of the forests as well since it can lead to other income sources, such as with the development of ecotourism. In the protected areas it is also important to involve local residents and educate them on the importance of wildlife protection since mostly they only experience difficulties in accessing their resources [29, 30].

In order to preserve at least some part of biodiversity a system of protected areas is required. Determination of the most valuable protected areas can be carried out by ecological modeling. Torres-Miranda et al. [31] used red oaks and their distribution area to estimate (section *Lobatae*; 75 species) those areas that are worth the most concerning protection. These species representing ecosystems prove to be useful indicators for conservation professionals. These species occur in various habitats often as dominant species and have an important part in preserving diversity. Based on a computer simulation (Complementarity analyses) 12 current areas under protection should be enlarged and 26 new should be established with a total of 512 500 ha area to ensure adequate protection for habitats. Certain species, especially arthropods bond to certain plant communities. With the protection of fast reducing oak forests species bonded to them can also be preserved [32].

An experiment in the Tehuacán Valley showed that biodiversity is preserved with greater success in areas under forest management than in areas that are not involved. Although, preserving rare species is limited even in this system. In average 59% of plant species and 94% of the genetic variety of dominant species (*Polaskia chichipe*, *Escontria chiotilla*, *Myrtillocactus schenckii*) was managed to be preserved in the examined area. In areas that are not involved the remaining natural flora decrease fast due to the increasing intensity of agriculture [33]. The number of species that can be preserved depends on whether the areas in question are private or community properties, also on the availability of natural resources, ecosystem services to people, as well as on the culture of certain communities. Ornamental or herbal species and those that are used for human or animal nutrition tend to be preserved even if the area gets involved in agriculture. Lanes dividing plots and islands within the agricultural area promote later regeneration [34]. Traditional methods of agriculture have less harmful effects on biodiversity than the current intensive mechanized agriculture. In the state of Oaxaca indigenous people have been carrying out agricultural production for centuries, which lead to a mosaic landscape with patches of forest and agricultural land. Nowadays due to urbanization more and more people give up farming and move to cities. On lands left behind a secondary forest forms, which surprisingly lead to the decrease of biodiversity.

3. Agricultural aspects

Traditional agriculture plays a part in sustaining biodiversity, since the landscape renews regularly. Traditional farming also has to be considered in the decision making process related to protected area [35]. Larios et al. [36] also claim that traditional farming has a great role in the preservation of biodiversity. According to a survey carried out in the area of the Tehuacán Valley 281 vascular plants were identified even in the gardens out of which 34% were endemic. Though abundance of cultivated plants was the largest. The highest value (199) was found in gardens lying near the cloud forests

in the mountains. The lowest value (141) was found in those that are located near deciduous forests. This can be explained by the tendency that owners cultivate plants in their own land to compensate the rarity of useful species in the nearby forests.

An agricultural effort to protect the diversity of the species is the production of shade-grown coffee [37]. With the production of shade-grown coffee most of biodiversity can be preserved since a proportion of the original vegetation survives. Coffee produced with this method has a high price, which can encourage more and more people to choose this cultivation method. Cultivation area is increasing unfortunately to the detriment of the primary forest, thus in its current form this is not the appropriate method to preserve biodiversity [38]. For the protection of marine ecosystems aquaculture is spreading in Mexico as well regarding both fishes and marine invertebrates. The development of the technology means income for the economy and wildlife can also be protected since the importance of illegal fishing decreases. Aquacultural production can mean a legal income source for the local people, while overfishing and the decrease of biodiversity can be avoided [39].

Phytoremediation plays an important part in the region as well in the neutralization of industrial pollutants, therefore the research of those organizations are important, which can be used for this purpose [40]. An abandoned mine in the state of Hidalgo was recultivated and the area was reforested. 56 species representing 29 families were managed to be planted. Samples of AMF (Arbuscular Mycorrhizal Fungi) from tree roots *Glomus* (Glomeraceae: Glomerales) and *Acaulospora* (Acaulosporaceae: Diversisporales) species were the most common. These have a great role in phytoremediation due to increasing the trees tolerance against heavy metals in the extreme environment, as well as decreasing the distribution of toxic substances in the environment [41]. Harmful effects of industrial pollutants can be reduced by phytoremediation, therefore it serves the protection of wildlife. Regarding the big biodiversity of the area it is likely that new species will be found to be suitable for this purpose. There is another example of a microbial scale biodiversity research with direct economic benefits. Diversity of bacteria and fungi living around cultivated *Agave tequilana* roots in the soil, rhizosphere and phyllosphere, in the endosphere of the root and the leaf was compared by Coleman-Derr et al. [42] with similar microbes of wild *Agave salmiana* and *Agave deserti* populations. *Agave tequilana* can be cultivated in areas where no other crop can survive and can be used for bioethanol production. Symbiotic microorganisms influence plant health and accommodation to stress due to this the rate of growth as well [43]. Manipulation of microbiome may increase the rate of growth and therefore, the amount of ethanol that can be produced [44]. The composition of the microorganism community based on the analysis of the traceable DNA changes depending on the compartment, which was obtained by the amplification of ITS2 and 16S regions. Geographical distribution also affected composition. In the case of cultivated plants alpha diversity was low, which can be explained with agricultural practices. The community is dominated by the genera of Enterobacteriaceae family (*Pantoea*, *Leclercia*, *Trabusiella*), therefore soft rot disease became often, which cost millions. Genetic diversity of plants is also low due to vegetative reproduction. That is why the bacteria *Pantoea agglomerans* could develop avoiding strategies during the evolution against the plants defense mechanisms [42]. Apart from agave, oil pressed from the seed of *Jatropha* (Euphorbiaceae) species, especially from the seed of *J. curcas*, is also appropriate for the production of biodiesel [45]. 50 of the 186 species occur in Mexico as well, most of which are endemic. They are distributed from the rainforests to the deserts everywhere. Their distribution is limited by the frost in the mountains. Other species may be eligible for agricultural use, which require further research [46].

Area, estimated number of species and percentage of remaining intact vegetation. The boundaries of ecoregions are not the same as national borders.

IntechOpen

Author details

Levente Hufnagel^{1*} and Ferenc Mics²

¹ Research Institute of Multidisciplinary Ecotheology, John Wesley Theological College, Budapest, Hungary

² Department of Environmental Security, John Wesley Theological College, Budapest, Hungary

*Address all correspondence to: leventehufnagel@gmail.com

IntechOpen

© 2021 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. 

References

- [1] Winkler, K. (2011): Middle America, not Mesoamerica, is the accurate term for biogeography. – *The Condor* 113(1): 5-6.
- [2] Mittermeier, R.A., Myers, N., Mittermeier, C.G. (1999): Hotspots: Earth's biologically richest and Most endangered terrestrial ecoregions. – *Journal of Mammalogy* 83(2): 630-633.
- [3] Flores-Villela, O., Gerez, P. (1994): Biodiversidad y conservación en México: vertebrados, vegetación y uso del suelo. – Comisión Nacional para el Conocimiento y Uso de La Biodiversidad y Universidad Nacional Autónoma de México, México.
- [4] Kier, G., Mutke, J., Dinerstein, E., Ricketts, T.H., Küper, Kreft, H., Barthlott, W. (2005): Global patterns of plant diversity and floristic knowledge. – *Journal of Biogeography* 32(7): 1107-1116.
- [5] Dinerstein, E., Vynne, C., Sala, E., Joshi, A.R., Fernando, S., Lovejoy, T.E., Mayorga, J., Olson, D., Asner, G.P., Baillie, J.E.M., Burgess, N.D., Burkart, K., Noss, R.F., Baccini, A., Birch, T., Hahn, N., Joppa, L.N., Wikramanayake, E. (2019): A global Deal for nature: Guiding principles, milestones, and targets. – *Science Advances* 5(4): eaaw2869.
- [6] CONABIO. (1998): La diversidad biológica en México: Estudio de País, 1998. – Comisión Nacional para el Conocimiento y Uso de la Biodiversidad. México.
- [7] Villarreal-Quintanilla, J.A., Bartolomé-Hernández, J.A., Estrada-Castillón, E., Ramírez-Rodríguez, H., Martínez-Amador, S. J. (2017): The endemic element of the Chihuahuan Desert vascular flora. – *Acta Botanica Mexicana* 118: 65-96.
- [8] Riemann, H., Exequiel, E. (2007): Endemic regions of the vascular Flora of the peninsula of Baja California, Mexico. – *Journal of Vegetation Science* 18(3): 327-336.
- [9] Villaseñor, J.L., Ortiz, E., Delgadillo-Moya, C., Juárez, D. (2020): The breadth of the Mexican transition zone as defined by its flowering plant generic flora. – *PLoS ONE* 15(6): e0235267.
- [10] López-González, C., Presley, S.J., Lozano, A., Stevens, R.D., Higgins, C.L. (2015): Ecological biogeography of Mexican bats: The relative contributions of habitat heterogeneity, beta diversity, and environmental gradients to species richness and composition patterns. – *Ecography* 38: 261-272.
- [11] Fernández, J.A. (2012): Phylogenetics and biogeography of the microendemic rodent *Xerospermophilus perotensis* (Perote ground squirrel) in the oriental basin of Mexico. – *Journal of Mammalogy* 93(6): 1431-1439.
- [12] Fernández, J.A., Cervantes, F.A., Hafner, M.S. (2012): Molecular systematics and biogeography of the Mexican endemic kangaroo rat, *Dipodomys phillipsii* (Rodentia: Heteromyidae). – *Journal of Mammalogy* 93(2): 560-571.
- [13] Moonlight, P.W., Richardson, J.E., Tebbitt, M.C., Thomas, D.C., Hollands, R., Peng, C.I., Hughes, M. (2015): Continental-scale diversification patterns in a megadiverse genus: The biogeography of neotropical begonia. – *Journal of Biogeography* 42: 1137-1149.
- [14] Chazdon, R.L., Harvey, C.A., Martínez-Ramos, M., Balvanera, P., Schondube, J.E., Stoner, K.E., Cabadilla, L.D.A., Flores-Hidalgo, M. (2011): Seasonally dry tropical Forest biodiversity and conservation value in agricultural landscapes of Mesoamerica. – In: Dirzo,

- R., Young, H.S., Mooney, H.A., Ceballos, G. (eds.) *Seasonally Dry Tropical Forests*. Island Press, Washington, DC.
- [15] Estado de la Región. (2008): Estado de la Región en Desarrollo Humano Sostenible un informe desde Centroamérica y para Centroamérica. – Programa Estado de la Región. San José, Costa Rica.
- [16] Bastida-Zavala, J.R., del Socorro García-Madrigal, M., Rosas-Alquicira, E.F., López-Pérez, R.A., Benítez-Villalobos, F., Meraz-Hernando, J.F., Torres-Huerta, A.M., Montoya-Márquez, A., Barrientos-Luján, N.A. (2013): Marine and coastal biodiversity of Oaxaca, Mexico. – *Check List* 9(2): 329-390.
- [17] de León, G.P.P., García-Prieto, L., Mendoza-Garfia, B. (2011): Describing parasite biodiversity: The case of the helminth Fauna of wildlife vertebrates in Mexico. – In: Grillo, O., Venora, G. *Changing Diversity in Changing Environment*. BoD – Books on Demand.
- [18] Hershler, R., Liu, H.P., Landye, J.J. (2011): New species and records of springsnails (Caenogastropoda: Cochliopidae: Tryonia) from the Chihuahuan Desert (Mexico and United States), an imperiled biodiversity hotspot. – *Zootaxa* 3001: 1-32.
- [19] Sáenz-Romero, C., Rehfeldt, G.E., Crookston, N.L., Duval, P., St-Amant, R., Beaulieu, J., Richardson, B.A. (2010): Spline models of contemporary, 2030, 2060 and 2090 climates for Mexico and their use in understanding climate-change impacts on the vegetation. – *Climatic Change* 102: 595-623.
- [20] Silva-Flores, R., Pérez-Verdín, G., Wehenkel, C. (2014): Patterns of tree species diversity in relation to climatic factors on the Sierra Madre occidental, Mexico. – *PLOS One* 9(8): e105034.
- [21] Mosco, A. (2017): Niche characteristics and potential distribution of *Thelocactus* species, a Mexican genus of globular cacti. – doi:10.1101/124511. PPR:PPR32150.
- [22] Peters, R., Ripple, W.J., Wolf, C., Moskwik, M., Carreón-Arroyo, G., Ceballos, G., Córdova, A., Dirzo, R., Ehrlich, P.R., Flesch, A.D., List, R., Lovejoy, T.E., Noss, R.F., Pacheco, J., Sarukhán, J.K., Soulé, M.E., Wilson, E.O., Miller, J.R.B., and 2556 scientist signatories from 43 countries. (2018): *Nature Divided, Scientists United: US–Mexico Border Wall Threatens Biodiversity and Binational Conservation*. – *BioScience* 68(10): 740-743.
- [23] Lasky, J.R., Jetz, W., Keitt, T.H. (2011): Conservation biogeography of the US–Mexico border: A transcontinental risk assessment of barriers to animal dispersal. – *Diversity and Distributions* 17: 673-687.
- [24] Dávila, P., Benítez, H., Barrios, Y., Cruz-Angón, A., Álvarez-Girard, N. (2011): Definition and insertion of the GSPC in the political context of Mexico. – *Botanical Journal of the Linnean Society* 166(3): 326-330.
- [25] Fodor, R.K. (2014): Ökoszisztéma szolgáltatások egy újfajta keretrendszerben. – In: Lukovics, M., Zuti, B. *A területi fejlődés dilemmái*. SZTE Gazdaságtudományi Kar, Szeged.
- [26] Alix-Garcia, J.M., Sims, K.R.E., Yañez-Pagans, P. (2015): Only one tree from each seed? Environmental effectiveness and poverty alleviation in Mexico's payments for ecosystem. – *American Economic Journal: Economic Policy* 7(4): 1-40.
- [27] Costedoat, S., Corbera, E., Ezzine-de-Blas, D., Honey-Rosés, J., Baylis, Castillo-Santiago, M.A. (2015): How Effective Are Biodiversity Conservation

- Payments in Mexico? – PLoS ONE 10(3): e0119881.
- [28] Honey-Rosés, J., Baylis, K., Ramírez, M.I. (2011): A spatially explicit estimate of avoided forest loss. – *Conservation Biology* 25(5): 1032-1043.
- [29] Durand, L., Vázquez, L.B. (2011): Biodiversity conservation discourses. A case study on scientists and government authorities in sierra de Huautla biosphere reserve, Mexico. – *Land Use Policy* 28: 76-82.
- [30] Mendez-Lopez, M.E., García-Frapolli, E., Pritchard, D.J., Gonzalez, M.C.S., Ruiz-Mallen, I., Porter-Bolland, L., Reyes-Garcia, V. (2014): Local participation in biodiversity conservation initiatives: A comparative analysis of different models in south East Mexico. – *Journal of Environmental Management* 145: 321-329.
- [31] Torres-Miranda, A., Luna-Vega, I., Oyama, K. (2011): Conservation biogeography of red oaks (*Quercus*, section *Lobatae*) in Mexico and Central America. – *American Journal of Botany* 98(2): 290-305.
- [32] Sanborn, A.F., Phillips, P.K. (2013): Biogeography of the cicadas (Hemiptera: Cicadidae) of North America, North of Mexico. – *Diversity* 5: 166-239.
- [33] Moreno-Calles, A., Casas, A., Blancas, J., Torres, I., Masera, O., Caballero, J., Garcia-Barrios, L., Perez-Negron, E., Rangel-Landa, S. (2010): Agroforestry systems and biodiversity conservation in arid zones: The case of the Tehuacan Valley, Central Mexico. – *Agroforestry Systems* 80: 315-331.
- [34] Vallejo, M., Casas, A., Blancas, J., Moreno-Calles, A.I., Solís, L., Rangel-Landa, S., Dávila, P., Téllez, O. (2014): Agroforestry systems in the highlands of the Tehuacán Valley, Mexico: Indigenous cultures and biodiversity conservation. – *Agroforestry Systems* 88: 125-140.
- [35] Robson, J.P., Berkes, F. (2011): Exploring some of the myths of land use change: Can rural to urban migration drive declines in biodiversity? – *Global Environmental Change* 21: 844-854.
- [36] Larios, C., Casas, A., Vallejo, M., Moreno-Calles, A.I., Blancas, J. (2013): Plant management and biodiversity conservation in Náhuatl homegardens of the Tehuacán Valley, Mexico. – *Journal of Ethnobiology and Ethnomedicine* 9:74.
- [37] Ruben, R., Zuniga, G. (2010): How standards compete: Comparative impact of coffee certification schemes in northern Nicaragua. – *Supply Chain Management: An International Journal* 16(2): 98-109.
- [38] Tejeda-Cruz, C., Silva-Rivera, E., Barton, J.R., Sutherland, W.J. (2010): Why shade coffee does not guarantee biodiversity conservation. – *Ecology and Society* 15(1): 13.
- [39] Solís-Marín, F.A., Honey-Escandón, M.B.I., Herrero-Perezrul, M.D., Benitez-Villalobos, F., Díaz-Martínez, J.P., Buitrón-Sánchez, B.E., Palleiro-Nayar, J.S., Durán-González, A. (2013): The echinoderms of Mexico: Biodiversity, distribution and current state of knowledge. – In: Alvarado, J.J., Solís-Marín, F.A. (eds.) *Echinoderm Research and Diversity in Latin America*. – Springer-Verlag, Berlin.
- [40] Wong-Arguelles, C., Alonso-Castro, A.J., Ilizaliturri-Hernandez, C.A., Carranza-Alvarez, C. (2020): Credibility of In Situ Phytoremediation for Restoration of Disturbed Environments. – In: Hakeem, K., Bhat, R., Qadri, H. *Bioremediation and Biotechnology*. Springer, Cham.
- [41] del Pilar Ortega-Larrocea, M., Xoconostle-Cazares, B.,

Maldonado-Mendoza, I.E., Carrillo-Gonzalez, R., Hernandez-Hernandez, J., Garduño, M.D., Lopez-Meyer, M., Gomez-Flores, L., del Carmen A. Gonzalez-Chavez, M. (2010): Plant and fungal biodiversity from metal mine wastes under remediation at Zimapan, Hidalgo, Mexico. – *Environmental Pollution* 158: 1922-1931.

[42] Coleman-Derr, D., Desgarennes, D., Fonseca-Garcia, C., Gross, S., Clingenpeel, S., Woyke, T., North, G., Visel, A., Partida-Martinez, L.P., Tringe, S.G. (2016): Plant compartment and biogeography affect microbiome composition in cultivated and native agave species. – *New Phytologist* 209: 798-811.

[43] Panke-Buisse, K., Poole, A.C., Goodrich, J.K., Ley, R.E., Kao-Kniffin, J. (2015): Selection on soil microbiomes reveals reproducible impacts on plant function. – *ISME Journal* 9: 980-989.

[44] Turner, T.R., James, E.K., Poole, P.S. (2013): The plant microbiome. – *Genome Biology* 14: 209.

[45] Folaranmi, J. (2013): Production of Biodiesel (B100) from *Jatropha* Oil Using Sodium Hydroxide as Catalyst. – *Journal of Petroleum Engineering* Article ID: 956479.

[46] Fresnedo-Ramírez, J., Orozco-Ramírez, Q. (2013): Diversity and distribution of genus *Jatropha* in Mexico. – *Genetic Resources and Crop Evolution* 60: 1087-1104.