

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

185,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Arsenic Speciation Techniques in Soil Water and Plant: An Overview

Mohammed Zia Uddin Kamal and Md. Yunus Miah

Abstract

There are more than 100 different arsenic with different characteristics in the soil-water-plant ecosystem. The identification and quantification of individual arsenic species is essential for understanding the distribution, environmental fate and behavior, metabolism and toxicity of arsenic. Due to the hazardous nature of arsenic, people have a high interest in the measurement of arsenic species. The reaction of the formation of arsenic speciation in the soil-water-plant environment is briefly studied. There is little information on methods used to quantify arsenic forms and species in contaminated soil, water and plant. The purpose of this article is to understand the available sample pretreatment, extraction, separation, detection and method validation techniques for arsenic speciation analysis of arsenic species in soil, water and plant. The performances of various sample preparation and extraction processes, as well as effective separation techniques, that contribute greatly to excellent sensitivity and selectivity in arsenic speciation when coupling with suitable detection mode, and method validity are discussed. The outlines of arsenic speciation techniques are discussed in view of the importance to the completeness and accuracy of analytical data in the soil-water-plant samples. To develop cheap, fast, sensitive, and reproducible techniques with low detection limits, still needed to confine research on arsenic speciation present in environmental matrices.

Keywords: Arsenic speciation, extraction, separation, detection, techniques, soil water and plant

1. Introduction

Arsenic (As) is a geogenic toxic metalloid found ubiquitously in environmental systems such as lithosphere (earth crusts, soil, rock, and sediment), hydrosphere (surface water, aquifers, deep wells, and oceans), atmosphere and biosphere (food chain and ecosystems) [1]. Arsenic is considered as the 12th most abundant elements in the earth's crust. Elevated arsenic having been introduced in the ecosystem either by natural routes involve in weathering and other biogeochemical processes or via anthropogenic activities, including mining, and smelting, excessive agricultural utilization of As-based fertilizers and pesticides and irrigation with As-laden groundwater [2–4]. This problem becomes serious concern because once arsenic is released in the soil and water resources, it is bioaccumulated by the terrestrial and aquatic biota, and subsequently enters in the human or animal food chain [5, 6]. In highly arsenic contaminated ($\geq 0.01 \text{ mg L}^{-1}$) area, the migration of arsenic from soil to water and plant is a serious problem, becoming a major threat to sustainable agriculture practices and food security [7, 8]. Empirical data shows that the

concentration of arsenic in contaminated soils lies between 10 mg kg^{-1} and as high as $30,000 \text{ mg/kg}$ [9]. In addition, the reported concentrations of arsenic in all natural waters is between $<0.5 \text{ } \mu\text{g L}^{-1}$ and more than $5000 \text{ } \mu\text{g L}^{-1}$, although maximum permissible contaminant total As limits in drinking water by WHO is $10 \text{ } \mu\text{g L}^{-1}$ [1, 10]. Moreover, considering toxicity, the Joint Food and Agriculture Organization and the World Health Organization (FAO/WHO) Expert Committee on Food Additives proposed that the maximum inorganic arsenic content in food such as polished rice is 0.2 mg kg^{-1} [11–13]. Thus, exposure to arsenic (As) in soil-water-plant becomes global public health and the environment concern due to the wide distribution in ecosystem and its close association with numerous adverse effects.

There are more than 100 different arsenic compounds in the soil-water-plant ecosystem [14, 15]. It is well known that the toxicity, bioavailability, physiological and metabolic processes and mobility of arsenic vary greatly depending on the chemical species and oxidative states rather than its total content [16, 17]. Arsenic (As) speciation analysis may specify not only the determination of total As contents but also considering its specific ionic forms in the aqueous solution and the sequential extracted As related to various mineral phases [18]. According to the IUPAC recommendations, “speciation of an element” is defined as “the distribution of an element amongst defined chemical species in a system” rather than fractionation. While speciation analysis is defined as “analytical activity of measuring the quantities of one or more individual chemical species in a sample” [19].

Arsenic exists multiple oxidation states (+III, +V, 0, –III) and various inorganic and organic chemical species. In environmental assessment, it is far from enough to know the total arsenic content in actual samples, because the toxicity of As element is predestined by distinct arsenical species [20]. Generally speaking, trivalent arsenic compounds are usually more toxic than pentavalent arsenic compounds [4] and inorganic species are more toxic than the organic ones. Again, trivalent organic arsenicals can be more toxic than trivalent inorganic arsenicals [21]. The United States Environmental Protection Agency (USEPA) priority pollutants list represents inorganic As is the first category of toxins [22], classified as Group I carcinogens based on human epidemiological data. In addition, the organic species toxicity usually decreases with the increase of methylation. For example, monomethylarsonic acid (MMA) and dimethylarsinic acid (DMA) are less toxic, arsenobetaine (AsB), arsenocholine (AsC) and other arsenosugars are even considered non-toxic [21]. However, in certain environments, such as in aquatic biomass, AsB can be converted into toxic inorganic arsenicals and enter the food chain [23, 24]. Depending on the source, metals or metalloids may enter the environment, where they may be transformed into another compound. Therefore, As speciation is essential for understanding its distribution, transformation in the environment, toxicity, metabolism, bioavailability and health effects in the natural system [15].

There is a huge difference between the toxicity and distribution of the arsenic species observed in the environment, which heights the importance of detecting and quantifying a single compound. Recently, various techniques have been developed to figure out arsenic species in environmental and biological samples, including soil, water and plants. The establishment of the new arsenic speciation analysis program not only improves our understanding of arsenic biogeochemistry, toxicity and metabolism but also provides a lot of information about exposure biomarkers and arsenic cycling in the natural environment. However, it is still a challenge to completely isolate the target arsenic compound from background interference [25]. Therefore, a quick and simple method is needed to analyze the arsenic species in different matrices. In addition, optimizing the extraction of target arsenic is also crucial for accurate quantification [26]. Determining the exact species of arsenic in biological and environmental samples helps to more accurately assess the

environmental impact and health risks caused by arsenic exposure. Appropriate sample pretreatment techniques are necessary to reduce the influence of matrix, to enrich the target species and/or separation of As species for accurate identification. The newly developed As speciation protocols must achieve suitable detection mode, excellent selectivity and sensitivity in various environmental and biological species. Various non-chromatographic and chromatographic methods are involved in the selective separation of the arsenic species.

To date, several study on overall arsenic speciation analysis have been done [15, 27, 28]. Nevertheless, there is still a big knowledge gap in the speciation of arsenic. This overview includes arsenic speciation analysis, species detection systems, key extraction/separation techniques and mechanisms used in the accuracy assessment of speciation methods, and focuses on important strategies for specific arsenic speciation. This study will provide sentinels on comprehensively discuss in the latest developments in arsenic speciation analysis and challenges for further research.

2. Reactions of arsenic speciation on environment

Arsenic is introduced into the environment either naturally or anthropogenically; once released, it cannot be degraded or destroyed. The environmental transformation of arsenic depends on the availability of arsenic in the geological source, as well as their oxidation state, speciation and other environmental factors [29, 30]. There are different forms of arsenic containing mineral in the Earth's crust. For example, 60% are in arsenate form, 20% are in the form of sulfides and sulfonates, and the remaining 20% are in the form of arsenites, arsenides, silicates, oxides, and elemental As [31]. In the soil and water environments, As can exist in four different oxidation states (As^{3+} , As^{5+} , As^0 and As^{3-}), in inorganic as well as in organic forms [4]. The most widespread arsenic species detected in the environment and biological systems are shown in **Table 1**.

In natural environment, inorganic arsenic contains two oxygen anions, arsenite As (III) and arsenate As (V) but there are many organic arsenic compounds including monomethyl arsonic acid (MMA) and dimethyl arsinic acid (DMA) is the most common. According to intake and mobility, the toxicity of arsenic compounds decreases in the following sequential order: arsines > inorganic arsenites > organic trivalent compounds (arsenoxides), inorganic arsenates > organic pentavalent compounds > arsonium compounds > elemental arsenic. Arsenobetaine and arsenocholine are considered nontoxic [4]. At the same time, arsenic species exhibit various reaction behaviors and metabolic transformations in soil-water and plant ecosystems. For arsenic risk assessment of environmental samples and detection of appropriate speciation analysis, it is necessary to understand the main forms and metabolic transformations of arsenic compounds.

2.1 Arsenic speciation in soil

The various species or chemical forms of As in soil include- free ionic forms, precipitated as solids, adsorbed on soil organic or inorganic constituents, exchangeable, and structural constituent of primary and secondary minerals [32, 33]. There are both inorganic and organic forms (species) of arsenic in the soil. The most common inorganic species are arsenate (AsV) and arsenite (AsIII), while the most common organic species are monomethylarsonic acid (MMA) and dimethylarsinic acid (DMA). The order of toxicity of arsenic species is $\text{AsIII} > \text{AsV} > \text{MMA} > \text{DMA}$ [34]. In general, minor amount of naturally occurring arsenic in soil exists as a form of amorphous iron and aluminum oxides.

Name	Abbreviation	Chemical structure
Inorganic form		
Arsenite (Arsenous acid)	As (III)	H_3AsO_3 , $\text{H}_2\text{AsO}_3^{3-}$, HAsO_3^{2-} , AsO_3^{3-}
Arsenate (arsenic acid)	As (V)	H_3AsO_4 , H_2AsO_4^- , HAsO_4^{2-} , AsO_4^{3-}
Organic form		
Monomethylarsenic acid	MMA	$\text{CH}_3\text{AsO}(\text{OH})_2$
Monomethylarsonic acid	MMA(V)	$\text{CH}_3\text{AsO}(\text{OH})_2$
Monomethylarsonous acid	MMA(III)	$\text{CH}_3\text{As}(\text{OH})_2$
Dimethylarsinic acid	DMA (V)	$(\text{CH}_3)_2\text{AsO}(\text{OH})$
Dimethylarsinous acid	DMA (III)	$(\text{CH}_3)_2\text{AsOH}$
Dimethylarsinoyl ethanol	DMAE	$(\text{CH}_3)_2\text{AsOCH}_2\text{CH}_2\text{OH}$
Trimethylarsine oxide	TMAO	$(\text{CH}_3)_3\text{AsO}$
Trimethylarsoniopropionate	TMAP	$(\text{CH}_3)_3\text{As}^+\text{CH}_2\text{CH}_2\text{COO}^-$
Tetramethylarsonium ion	TETRA, TMA	$(\text{CH}_3)_4\text{As}^+$
Arsenobetaine	AsB	$(\text{CH}_3)_3\text{As}^+\text{CH}_2\text{COOH}$
Arsenobetaine 2	AsB-2	$(\text{CH}_3)_3\text{As}^+\text{CH}_2\text{CH}_2\text{COO}^-$
Arsenocholine	AsC	$(\text{CH}_3)_3\text{As}^+\text{CH}_2\text{CH}_2\text{OH}$
Trimethylarsine	TMA (III)	$(\text{CH}_3)_3\text{As}$
Arsines	AsH_3 , MeAsH_2 , Me_2AsH	$(\text{CH}_3)_x\text{AsH}_{3-x}$ ($x = 0-3$)
Ethylmethylarsines	$\text{Et}_x\text{AsMe}_{3-x}$	$(\text{CH}_3\text{CH}_2)_x\text{As}(\text{CH}_3)_{3-x}$ ($x = 0-3$)
Dimethylarsinyolacetic acid	DMAA	$(\text{CH}_3)_2\text{AsOCH}_2\text{COOH}$
Phenylarsine oxide	PAO	$\text{C}_6\text{H}_5\text{AsO}$
Phenylarsonic acid	PAA	$\text{C}_6\text{H}_5\text{AsO}(\text{OH})_2$
Arsenosugars $\text{C}_7\text{H}_{14}\text{AsO}_3\text{CH}_2\text{CH}(\text{OH})\text{CH}_2\text{R}$		
Arsenosugar 1 (glycerol sugar)	—	$\text{R} = \text{OH}$
Arsenosugar 2 (phosphate sugar)	—	$\text{R} = \text{OP}(\text{O})(\text{O}^-)\text{OCH}_2\text{CH}(\text{OH})\text{CH}_2\text{OH}$
Arsenosugar 3 (sulphonate sugar)	—	$\text{R} = \text{SO}_3^-$
Arsenosugar 4 (sulphate sugar)	—	$\text{R} = \text{OSO}_3^-$

Table 1.
Arsenic species commonly identified in the environment and biological systems.

Arsenic can be transformed in the soils through various mechanisms, such as oxidation, reduction, adsorption, dissolution, precipitation, and volatilization. The inorganic species of As, As(III) and As(V), are present in different forms (e.g., fully protonated As acids or arsenous acid) [35]. The main and thermodynamically stable forms of As(V/III) in soil may include H_2AsO_4^- , HAsO_4^{2-} and H_3AsO_3 . The existence of different As forms in soil largely depends on the texture, organic matter, pH value and redox potential of the surrounding environment. Arsenic exists in aerobic soil (oxidized conditions) in the form of arsenate (As^{V}) and is rapidly adsorbed on clay minerals and Fe/Mn oxides/hydroxides [2]. However, in reducing soil environment such as paddy fields, the arsenite (As^{III}) form of arsenic dominates and its

solubility, mobility, and toxicity are about 60 times that of As(V) [2]. In addition, under anaerobic conditions in the presence of sulfides, arsenic may precipitate in the form of arsenic sulfide and release excess arsenic into the environment [36]. Anaerobic bacteria degrade into less toxic volatile forms, such as dimethyl arsenic acid (DMAA) and monomethyl arsenic acid (MMAA) [37]. The oxidation and reduction of arsenic species takes place biologically and chemically in soil and water [38]. In addition, higher concentrations of arsenic were observed in alluvial soils and organic soils, while lower concentrations of arsenic were found in sandy soils [39]. Clay played a leading role in arsenic fixation. Arsenate is mainly adsorbed on clay particles in soils with neutral pH. Soil pH plays a major role in the types of arsenic compounds present in the soil. Under acidic conditions, arsenic tends to form compounds with aluminum and iron (AlAsO_4 , FeAsO_4), whereas under alkaline conditions (limestone soils) Under acidic conditions, arsenic tends to form compounds with aluminum and iron.

Arsenate exists in the form of oxygen anions at neutral pH, while arsenite has a neutral charge at pH 7.0. It leads to the formation of $\text{Ca}_3(\text{AsO}_4)_2$, $\text{Mn}_3(\text{AsO}_4)_2$, AlAsO_4 and FeAsO_4 [40]. However, when the pH value is higher than 8.5, as the pH value increases, the adsorption capacity of As^{V} decreases, while the case of As^{III} is the opposite. But at pH around 4, the adsorption for As^{V} on FeOOH is maximum, whereas for As^{III} the optimum pH is 7–8.5 [41]. Arsenic is more soluble under high or low pH values. In reducing environment, as arsenate is reduced to arsenite, it binds less strongly to the hydroxide solids, which increase the bioavailability of arsenic [42]. On the contrary, due to the larger arsenic sorption affinity, organic matter has formed organo-arsenic complex and reduced the solubility of arsenic in the soil [43]. Naturally, arsenic can be released into the soil environment through the hydrolysis and oxidation process of primary sulfide mineral (i.e. arsenopyrite) and absorbed by ferric hydroxide. Meanwhile, the retention and mobilization of arsenic in soil is largely controlled by Ferric hydroxide. Such as iron oxides has stronger arsenic adsorption capacity than manganese oxides. Moreover, Phosphate plays significant role in absorption of arsenic from contaminant soil. Williams et al. [44] reported that in iron oxides dominated acidic soil, approximately 60% of the adsorbed pentavalent arsenic and 70% of the trivalent arsenic were displaced by H_2PO_4 . Soil microbial activities can affect the adsorption/desorption, solubility, bioavailability, mobility and soil–plant transfer of arsenic by changing the chemical speciation of As in soil [45, 46]. Due to the action of microorganisms or the past use of methylated arsenic compounds, dimethyl sulfoxide, or sodium salts of MMA and DMA as pesticides, methylated As species, (i.e. MMA and DMA) might be accumulate in soil [45]. Microorganisms in soil may interconvert As species As^{V} and As^{III} and further transform into MMA and DMA.

2.2 Arsenic speciation in water

The presence of arsenic in water is either dissolved or in particulate form. Arsenic pollution in groundwater mainly occurs due to release of geothermal water, desorption and reductive dissolution of iron oxides and oxidation of sulfide minerals [37]. High levels of arsenic in groundwater have been observed in many countries, such as more than $3000 \mu\text{g L}^{-1}$ because As has been released geogenically either by oxidation of arsenopyrite, or by reductive dissolution of arsenic rich ferrous oxyhydroxides in reducing aquifer environment [47, 48]. The most common forms of arsenic in natural waters are arsenite and arsenate. However, the main species found in natural water are forms of inorganic arsenic, namely H_2AsO_4^- , H_3AsO_3 , HAsO_4^- and $\text{As}_3\text{O}_4^{3-}$. The change in the arsenic solubility in sulfidic water promotes the formation of amorphous metal-sulfide complex thioarsenic

compounds [49]. Various species of aquatic micro and higher organisms plays important role on biomethylation process of arsenic, which reduces As^{5+} to soluble As^{3+} species [37]. Arsenate is the main form of arsenic in seawater. Dimethyl arsenic acid (DMMA) and methylarsenic acid (MMA) are also present in small amounts in seawater [50]. Moreover, the determination of arsenic in saline waters bear much importance due to gaining knowledge on because salinity induced inorganic arsenic specifically arsenite transformation to arsine gas [51].

2.3 Arsenic speciation in plants

Arsenic is not essential elements for plants development, although very small amounts of As in plants may have a positive effect on plant species. The concentration of As in plants is usually less than 1.0 mg kg^{-1} dry weight (DW) [52]. The mechanism by which plants absorb arsenic varies depending on the chemical form of arsenic.

2.3.1 Transformation of inorganic arsenic species in plants

Plants absorb inorganic arsenic through two mechanisms. The first mechanism involves the use of a high-affinity PO_4 transporter through phosphate (PO_4) transport pathway [53, 54] which uptake As (V) from soil solution and subsequently to aerial parts of the plants [55]. While, the second route for plant roots to absorb arsenic is through the aquaporin channels, which uptake As (III) (silicic acid analog) and methylated As species (MMA and DMA) [56]. In rice root cells, As (III) uses generally Si transporter owing to its similarities with silicic acid. Once in plant cell, As (V) is reduced to As (III) with the help of As reductase, ACR2 [57]. The detoxification of As (III) is achieved by forming complexes with thiol- rich peptides [58]. The form of As in the phloem is considered to be very crucial for the redistribution of As in various tissues in the plant [59].

2.3.2 Organic arsenic species transformation in plant

The methylated arsenic species, such as MMA and DMA, contribute less to the total arsenic in the soil. The organic arsenic substances MMA and DMA have taken up by the intrinsic protein of Oncokin 26. Rabb et al. [60] showed that the absorption efficiency of inorganic As species (AsIII and AsV) in roots is much higher than that of methylated As species (DMA and MMA), but the translocation efficiency of inorganic species in plant stems is much lower than that of methylation As species. The decrease in the As complex formed with ligands (such as glutathione) in the root may be the reason for the better translocation of methylated As species [60]. In rice, As (III) is found to be the most abundant species, followed by DMA. As (V), MMA and two other unidentified As also have found in lower concentrations [61]. A speciation analysis revealed the As (V) as a predominant species in rice straw followed by As (III) and DMA [62]. Meanwhile, in rice grain, As (III) and DMA are the dominant species.

3. Soil, water and plant sample preparation and extraction of arsenic species

3.1 Soil, water and plant sample preparation for arsenic speciation

Sample preparation and storage procedures are considered to be a key prerequisite for maintaining the concentration and chemical structure of the original species

in the sample during the analysis process to obtain accurate As speciation information. Impractical As speciation data may arise due to losses during sampling, unrepresentative samples [63], contamination, mutual conversion between species, inefficient extraction of the analyte, and the possibility of precipitation and wall effects from the sample container [64]. To obtain reliable arsenic speciation data in soil, water and plant samples, two main strategies should be considered. First, determine appropriate species preservation practices to keep the chemical species of interest unchanged throughout the analysis process through avoiding changes in oxidation state, changes caused by microbial activity, and losses caused by volatilization or adsorption. Secondly, the species can be quantitatively converted into appropriate derivatives for further separation, accumulation and quantification [65]. Microbial transformation of arsenical compounds in contaminated samples is observed through a change in valence (i.e. oxidation/reduction) or chemical form (i.e. solid, liquid and gas) and formed biomethylate arsenic and both volatile (e.g., methylarsines) and nonvolatile (e.g., methylarsonic acid and dimethylarsinic acid) compounds [66]. To avoid degradation of arsenic speciation, biological samples should be kept in low temperature. To reduce analyte loss, drying in oven used for the stabilization of samples particularly freeze-drying [67].

To avoid arsenic speciation loss during sampling, the soil-plant-water should be collected in polyethylene flip-top bottle/plastic with lock and/or seal lead. Immediately after collection all of the samples should be kept in freezer until sample preparation for analysis. The soil samples were air-dried, gently crushed and sieved through a 2 mm sieve and used for analysis. Meanwhile the plant sample placed in a oven drier at 40°C until constant weight and then grind and sieve the sample and stored in brown glass bottles in a desiccator in order to avoid exposure to light and moisture until required for analysis. Sample preparation for solid samples generally may include procedures such as mincing, freeze drying, milling, grinding, homogenization, and sieving, followed by extraction. For achieving the best extraction efficiency and reproducibility of arsenic speciation in soil and plant sample, the tested sample must dried and homogenized before extraction. Because, particle size plays a crucial role in the extraction efficiency of As [68]. After sampling the fresh plant sample, it should be kept in freezing (−80°C) to avoid species interchange. Moreover, dry and grind plant and soil sample can store at −20°C up to one year [69].

The most reliable method for preserving natural water samples is, therefore, acidification to pH 2, refrigeration and deoxygenation [70]. Preservation of natural water in polypropylene bottles in refrigerator arsenic species in water is stable under neutral conditions for a period of 4 months [71]. To increase the stability of dissolved As redox species (As (III) and As(V)) of water sample, the samples to be filtered and stored in darkened polythene containers. For longer preservation, water samples are acidified with HCl [72], HNO₃ [73], H₂SO₄ [74] and H₃PO₄ [75], ascorbic acid [76] ethylene diamine tetraacetic acid [65]. Filtering the sample removes most of the colloidal material and microorganisms; acidification prevents oxidation and precipitation of Fe and Mn hydroxides and EDTA sequesters Fe and inhibits precipitation. Using 10 mM H₃PO₄ as a preserving agent combined with keeping samples at 6°C in dark, As species remain stable for 3 months, even with high concentrations of Fe and Mn.

3.2 Extraction procedures for arsenic speciation

The great challenge of As speciation, as it has been highlighted, is to maintain the original characteristics of species during extraction step. Extraction is the first step for speciation of target As species from their matrix (water, soil, sediment,

plant, biological tissue or fluid). Determining an appropriate sample preparation method that provides high extraction efficiency for the arsenic species of interest and prevents inter-conversion between arsenic species can be challenging. To achieve maximum extraction competence of arsenic speciation from solid or liquid matrix, the extraction protocols must have three criteria's, such as (i) the extracting solution must solubilize only the specific form, (ii) reduction of native As (V) to As (III) may not occur during the extraction, and (iii) oxidation of native As (III) to As (V) should not occur [77]. Extraction procedures employ a range of approaches including solid–liquid extraction [78] liquid– liquid extraction, solid phase extraction (SPE) [79] and solid phase microextraction (SPME) [80]. Enhanced techniques such as soxhlet, [69] sonication, [81] pressurized liquid extraction (PLE), [82] microwave-assisted extraction (MWA) [83] and supercritical fluid extraction (SFE) [84] have also been utilized for the determination of As speciation in soil–plant–water sample. Soil–plant–water sample preparation and extraction methods applied for the arsenic speciation analysis are presented in **Table 2**.

3.2.1 Solvent extraction

The solvent extraction technique is often used to determine organic arsenic compounds, especially arsenic compounds in biological samples. The extraction of arsenic substances is usually achieved through mild extraction solvents (ie water, methanol, methanol–water solvent system) and/or rarely uses acetonitrile–water and sequential extraction [15, 103]. Methanol/water mixture 1/1 (v/v) is widely used for the quantification of water-soluble As compounds in environmental samples, followed by centrifugation and filtration [104], while methanol–chloroform or hexane is used in non-polar species [15]. Moreover, extraction with water–methanol (1:1v⁻¹) had offered easy oxidation of As (III) in basic medium such as soil and the best efficiency was achieved after 20 min of extraction [105]. Extraction efficiency of arsenic species in soil–plant–water samples varied according with the changing the ratio of methanol– water solvent. Nevertheless, the methanol:water extraction solvent ratio of 1:1 provides the best processing and extraction efficiency for the extraction of arsenic species from plant samples, while 1 M phosphoric acid is suitable for soil samples [15]. At the same time, Rahman et al., [98] noticed that addition of extracting agent NH₄H₂PO₄ in edible part of spinach had shown similar extraction efficiency of As (III) and As (V) by water, 50% vv⁻¹water/methanol solution on shaking and microwave techniques. However, As(III) was extracted twice as much by the protein extract, indicating that it is a good extractor. Zheng and Hintelmann [106] pointed out that methanol/water mixture is an effective extractant for organic species, and its efficiency for inorganic species drops sharply. The solvent extraction reagent, tetramethylammonium hydroxide (TMAH) in alkaline medium, is also useful for the determination of AB, DMA and inorganic arsenic form a wide variety of biological matrices. In addition, sequential extraction procedure using different solvents (i.e (NH₄)₂SO₄, (NH₄)₄H₂PO₄, NH₄-oxalate buffer, KOH and hot water) can effectively extract organic and inorganic arsenic species, namely arsenous acid, arsenic acid, monomethylarsonic acid, dimethylarsinic acid, arsenobetaine, trimethylarsine oxide and glycerol-ribose in both soil and plant [107, 108]. Larios et al. [109] found that orthophosphoric acid followed by graphite block heating at 90°C for 60 min was provided the best conditions for As speciation in plants grown in contaminated environment. The applied extraction solvent led to an extraction efficiency of 80% for samples without species interconversion and recovery of 95% for leaves As speciation of Arsenic (V), As (III), DMA and MMA.

Matrix	Arsenic species	Sample preparation/extraction	Extraction solution	Detection	References
Soil	AS total, AS ^{III} , AS ^V	Shaking/mixing	10 M HCl	HGAAS; XRF	[85]
Soil	AS total, AS ^{III} , AS ^V	Shaking/mixing	10 mM phosphate	HPLC	[86]
Soil	AS total, AS ^{III} , AS ^V	Micro wave heating + Shaking	1 M H ₃ PO ₄ + 0.5 M ascorbic acid (C ₆ H ₈ O ₆)	LC=UV- HG- AFS	[87]
Soil	AS ^{III} , AS ^V , MMA ^V , DMA ^V	Ultrasonic, shaking, Microwave heat	500 mM Phosphate solution	HPLC-HG-AFS	[88]
Soil	AS ^{III} , AS ^V	Filtration. Shaking and water bath heat	[BMIM][PF ₆]	IL-LLME-ETAAS	[89]
Soil	AS total	Microwave heat	HNO ₃ + HClO ₄ + H ₂ SO ₄	HGAAS	[90]
Soil	AS total, AS ^{III} , AS ^V	Microwave heat	1 M H ₃ PO ₄ + 0.1–1 M ascorbic acid	LC–HG–AFS	[91]
Soil	AS ^{III} , AS ^V	Shaking/mixing	2.5 mM CaCl ₂	LC–HG–AFS	[21]
Soil	AS ^V	Shaking/mixing	1 M HCl	XRF	[92]
Soil and plant (chickpea)	AS ^{III} , AS ^V , MMA ^V , DMA ^V	Shaking/mixing	Buffer solution (1.5 mM NaH ₂ PO ₄ + 0.2 mM Na ₂ EDTA + 3 mM NaNO ₃ ,+ 10 mM CH ₃ COONa + 1% C ₂ H ₅ OH; pH 6.0)	ICP-MS	[93]
Plant	AS ^{III} , AS ^V , MMA ^V , DMA ^V	Shaking/mixing	0.3 M H ₃ PO ₄	HPLC-ICP-MS	[94]
Plant	AS ^{III} , AS ^V , MMA ^V , DMA ^V	Shaking/mixing	1% HCOOH	HPLC-ICP-MS-ESI-MS	[95]
Plant	AS ^{III} , AS ^V , MMA ^V , DMA ^V	Sonication	2 mM NaH ₂ PO ₄ + 0.2 mM Na ₂ EDTA (pH 6.0)	HPLC-ICP-MS-ESI-MS.	[95]
Soil and plant	AS ^{III} , AS ^V , MMA ^V , DMA ^V , AsC	Shaking/mixing + sonication	CH ₃ OH/H ₂ O 1 + 1 v/v	HPLC-ICP-MS	[96]
Soil and plant	AS ^{III} , AS ^V , MMA ^V , DMA ^V ,	Shaking/mixing + sonication	(a) CH ₃ OH/H ₂ O 1 + 1 v/v; (b) 0.1 M HCl	HPLC, AAS and XANES	[97]
Plant	AS ^{III} , AS ^V ,	MW-heating	0.33 M sucrose, 50 mM MES, 5 mM EDTA, 5 mM Lascorbate	HPLC-ICP-MS	[98]
Plant	Total As, AS ^{III} , AS ^V ,	MW-heating	Methanol–water (1: 1) /HNO ₃	HPLC-ICP-MS	[99]

Matrix	Arsenic species	Sample preparation/extraction	Extraction solution	Detection	References
Plant	As ^{III} , As ^V , MMA ^V , DMA ^V ,	MW-heating	1% (v/v) HNO ₃	HPLC-ICP-MS	[100]
Plant	As ^{III} , As ^V , MMA ^V , DMA ^V ,	MW-heating	0.01 mol/L TMAH	ETAAS	[101]
Surface/drinking water	As ^{III} , As ^V , MMA ^V , DMA ^V	Filtration	EDTA	HPLC-ICP-MS	[102]
Sea water	As ^{III} , As ^V , MMA, DMA, AsB, TMAO	Shaking/mixing + ultra-sonication	1% (v/v) HNO ₃	HPLC-ICP-MS	[26]

Table 2.
Several soil-water-plant sample preparation/extraction methods for determination of arsenic speciation.

3.2.2 Enzymatic hydrolysis

Biomolecular hydrolysis of complex matrix, enzymes are able to break down specific bonds of the substrate at neutral pH and room temperature, and they allow selective release of the analyte from the sample matrix without chemical changes. Enzymes can digest various matrix components, enzyme-assisted reactions usually require several hours of incubation. Microwave-assisted extraction (MAE) is used in combination with the enzyme extraction of pronase E and lipase to effectively extract AsB, As(III), DMA, MMA, and As(V) from seafood, rice, and plants [110, 111]. Viscozyme, was considered the most effective multi-enzyme mixture (consisted of a wide range of carbohydrases, including arabanase, cellulase, glucanase, hemicellulase, and xylanase) useful to extracted arsenic species from algae and terrestrial plant materials [112].

3.2.3 Microwave-assisted extraction

Microwave extraction is a common technique for extracting biological and environmental matrices, which is much faster than traditional Soxhlet extraction procedures. The extraction procedure using dilute acids or organic solvents at low temperatures can be easily achieved in a focused microwave oven. Generally microwave extraction is used to determine inorganic arsenic in food and provided good arsenic speciation extraction efficiencies (generally >90%) for samples of rice and wheat [113]. The method is based on extracting samples with trifluoroacetic acid/H₂O₂ and measuring arsenate by anion exchange HPLC-ICP-MS using aqueous malonic acid as the mobile phase. By using 2 M trifluoroacetic acid assisted with microwave heating for 6 h at 100°C to hydrolyze rice samples, the conversion between As^{III} and As^V was also observed and recovered 83, 88, 100, and 93% of fortified arsenite (100 ng As g⁻¹), arsenate (100 ng As g⁻¹), methylarsonic acid (MMA, 50 ng As g⁻¹), and dimethylarsinic acid (DMA, 200 ng As g⁻¹), respectively [114].

3.2.4 Solid phase extraction

Solid phase extraction (SPE) method is an efficient extraction technique for arsenic speciation from complex environmental and biological matrices. The principle mechanism of SPE is partitioning sorbent and sample matrix phase and may include simple adsorption, chelation, ion exchange or ion-pair solid phase extraction. In recent years, the techniques gaining popularity for As speciation because of its simple operation offers acceptable recovery and pre-concentration efficiency, lower reagent consumption and offer effective combination ability with different on-line and off-line As detection systems.

3.2.4.1 Conventional sorbent

Several conventional sorbent (i.e ion exchange resin, glass and modified mesoporous silica) based protocols have been developed for inorganic As speciation. To avoid inter-conversion of arsenic species, extraction with anion exchange cartridges prior to the inductively coupled plasma sector field mass spectrometric (ICP-SF-MS) becomes an efficient technique. During on-site separation and speciation of inorganic arsenic (As (III) and As (V)) from high arsenic- ground-water and ferrihydrite removal anaerobic arsenics species, anion-exchange resin (AG 1-X8) adsorbed As(V) in acetate form, while no adsorption to As(V)/As(III) in chloride form [115]. A dual-sorbent SPE protocol, in which the sorbent is composed of strong basic anion exchange (SBAE) resin and hydrate iron oxide particles

integrated HY resin, has been adopted successfully for the retention of inorganic arsenic species As (V) and As(III) simultaneously [116]. On-line continuous leaching extraction method is also effective for speciation of bio-accessible As species in plant [108].

3.2.4.2 Functional nanomaterials extractant

The modern technological invention of nanomaterials such as Nanofibers [117], magnetic nanoparticles [118], metal hydroxide precipitation [119], and nano-TiO₂ colloid [120] has offered selective and efficient extraction techniques for As speciation from different matrix. Like ammonium pyrroline- dithiocarbonate (APDC) have excellent selectivity of As (III) from ground water samples [117]. Moreover, yttrium hydroxide precipitate layer coated cellulose fiber is used as extracting material, [119] of As (III) and As (V) at acidic condition. Multi-wall carbon nanotubes (MWCNTs) modified with branched cationic polyethyleneimine (BPEI) is also proved to be excellent adsorbent with favorable selectivity toward adsorption of As(V) [121] in combined with sequential injection technique. Nano particle TiO₂ colloid has effectively extracted ultra-trace As from environmental water sample without agglomeration [120]. Besides, As (III) and As(V) from aqueous solution can be effectively extracted by hematite-coated Fe₃O₄ particles. Moreover, due to the fact simple and rapid separation capacity of As species, magnetic extraction techniques also gaining much popularity day by day.

3.2.4.3 Multi-sorbent based SPE procedure

A combined SPE procedure for arsenic speciation developed by using three molecular recognition technology (MRT) gel resins, which includes strong base anion exchange (SBAE) and two hybrid (HY) resins, HY-Fe and HY-AgCl, This methods has constructed for simultaneously extraction of four water-soluble arsenic species: arsenite, arsenate, monomethylarsonic acid (MMA) and dimethylarsinic acid (DMA) and retain in the SPE columns and separately eluted by using different elution [115].

3.2.5 Liquid-liquid extraction (LLE)

A liquid-liquid extraction generally used dodecane modified with 4% dodecanol containing Aliqua t336 as the extractant has been developed for the extraction of arsenic species in environmental matrices [122]. Here only As (V) is quantitatively transported to organic phase but no transport of As(III) takes place. A rapid in-situ liquid-liquid micro-extraction procedure has been developed for successfully determination of As (III) and As(V) in water samples, food salts, and total As in biological samples [123].

4. Derivatization of total arsenic

The vital challenge in element speciation is to determine each form independently without interference from other species. Because arsenic could complex with certain derivatizing agents, that hampered the detection. The derivitization process consists of two steps for example. i) reduction of AsV to AsIII and ii) convert to arsine (AsH₃) [124]. During measurement, the inert g N₂ is pushed by hydride

generation (HG) step, reaches the atomic absorption spectrophotometer or ICP-MS and finally produced arsines [125]. The main limitation of derivitization is that it only limits the formation of volatile arsines materials, but it is difficult to separate un-derivatized arsenic species (eg. AsB, AsC, or arsenosugars), using conventional reversed phase liquid chromatography and almost impossible by spectrophotometry or mass spectrometry [126]. In addition, the derivitization process requires control condition. In addition, the derivitization process requires control condition. This technique largely depends on the type and concentration of the sample matrix. To overcome obstacle, sodium borohydride is now commonly used for the hydrides synthesis [18]. Arsenic speciation after derivatization can be overcome by combining couple technique with detection techniques such as LC-MS/MS retention in liquid chromatography and ionization in mass spectrometry. Under such circumstances, the hyphenated technique is the most reliable because it includes several facilities like high sensitivity, good reproducibility, short analysis time and reduced risk for species transformation [18].

5. Separation techniques of arsenic speciation

Usually, two main techniques, namely chromatography (gas and liquid) and capillary electrophoresis are used to separate arsenic from various complex matrices [65]. Based on the complexity of As compounds, sometimes two technologies are introduced simultaneously or cumulatively.

5.1 Liquid chromatography

Liquid chromatography generally provides excellent possibilities for the separation of environmental and biological samples [18]. Various commonly used liquid chromatography techniques are high performance liquid chromatography (HPLC), ion chromatography and ion interaction chromatography [127]. Chromatographic separation can be performed by using ion exchange columns to separate metal ions directly or by adsorption (reverse phase or normal phase) liquid chromatography (if the metal species are complexed with organic ligands). Liquid chromatography is connected to many other detection systems, such as ICP-MS, HG-AFS, HG-AAS and GF-AAS [65]. Several liquid chromatography techniques can be used for the organic and inorganic As species, as follows:

5.1.1 Anion exchange liquid chromatography

Anion exchange chromatography is commonly used for speciation analysis of arsenic in environmental and biological samples. The anionic nature of arsenic species is different (at neutral pH, arsenic acid As(V), monomethylarsonic acid (MMA) and dimethylarsonic acid (DMA) are deprotonated, but As(III) exists) to make anion exchange chromatography suitable for their separation. Anion exchange chromatography has been used to analyze many arsenic compounds, including As(III), As(V), MMA, DMA, arsenobetaine (AsB), arsenocholine (AsC), oxoarsenic sugar (oxoAsS), thiosulfate Arsenic sugar (thioAsS) and benzene arsenic [27, 28]. The most commonly used column for arsenic speciation analysis is a strong anion exchange column, such as PRP-X100. Generally, the As form of the matrix separated by anion exchange chromatography techniques is detected by inductively coupled plasma mass spectrometry (ICP-MS) and electrospray ionization tandem mass spectrometry (ESI-MS/MS).

5.1.2 Cation exchange liquid chromatography

Cation exchange chromatography works similarly to anion exchange, except that the stationary phase is negatively charged to interact with the cation analyte. However, in cation exchange liquid chromatography, the separation speed of As species is relatively fast. The retention of arsenicals is directly related to the strength of its cationic charge: positively charged analytes have stronger retention. Cation exchange chromatography is commonly used for speciation analysis of positively charged As compounds, such as AsB, AsC, trimethylarsenic oxide (TMAO) and tetramethylarsenic ion (TMA) [15].

5.1.3 Reverse phase liquid chromatography

Reversed-phase chromatography is the most common HPLC separation technique used to separate compounds that are less hydrophobic or polar. In particular, reversed-phase liquid chromatography is particularly suitable for the analysis of arsenic lipids, including arsenic-containing hydrocarbons, fatty acids, phospholipids, phosphatidylcholines, fatty alcohols, and phosphatidylethanolamines of biological samples [24].

5.1.4 Ion pair chromatography

Ion pair chromatography can separate ions and neutral analytes using popular reversed-phase chromatography. It has been widely used for arsenic speciation analysis of various substrates. The reagent of ion pair chromatography reagent comprises with two groups a charged group for interaction with the analyte and a hydrophobic region for interrelates with the stationary phase. Usually, tetraalkylammonium, tetrabutylammonium and tetraethylammonium are used as the ion pair reagents for the separation of anionic and neutral arsenic species, and alkyl sulfonates, such as hexanesulfonic acid and 1-pentane sulfonic acid, for cationic and neutral arsenic species. Two most commonly used organic modifiers, methanol and acetonitrile are added to the mobile phase to decrease retention time [15].

5.1.5 Hydrophilic interaction liquid chromatography

Hydrophilic Interaction Chromatography (HILIC) is an important substitute to RP-HPLC separations of polar compounds. Although the stationary phase is polar, HILIC can separate neutral, cationic and anionic species simultaneously. HILIC has great potential to separate more arsenic species in a single analysis. This separation technique is more useful for organoarsenicals. Xie et al. [128] successfully detected nine kinds of organoarsenicals (I.e MMA, DMA, AsB, AsC, TMAO, phenylarsonic acid (PAA), phenylarsine oxide (PAO), 4-hydroxy-3-nitrophenylarsonic acid (Roxarsone), and 4-aminophenylarsonic acid (p-arsanilic acid, ASA) using a zwitterionic HILIC column followed by ICP-MS/ ESI-MS detection.

5.1.6 Size exclusion chromatography

Size-exclusion chromatography (SEC), also known as molecular sieve chromatography, is a chromatographic method in which molecules in solution are separated by their size, and in some cases molecular weight. SEC is very effective for analysis of arsenic interactions with large molecules or macromolecular

complexes such as arsenic-protein binding, humic acid-arsenic complexes and industrial polymers. SEC commonly used to separate protein-bound arsenic from free arsenic [129]. This separation method is expensive and useful for biological samples.

5.1.7 Multidimensional chromatography

Combining a variety of chromatographic columns and separation modes, try to separate a series of arsenic substances. Multidimensional separation has been performed offline or online. These usually involve a cation exchange column and an anion exchange column connected by a switching valve. This combination allows separation of cationic and anionic arsenic species. Applications were demonstrated for the determination of water-soluble arsenic species [20].

5.2 Capillary electrophoresis

Capillary electrophoresis separates As species according to the electrophoretic mobility related to the charge, viscosity, and atomic radius of the molecule, which is controlled by the composition, concentration, and pH of the buffer. This method is applicable for all type of soil-plant-water samples. Capillary electrophoresis can be used in many different detection systems but the most common is ICP-MS [15]. Although, Although capillary electrophoresis separation is simple, cost-effective, fast analysis and a certain degree of matrix independence, the additional complexity of coupling with the detection system makes CE a less common As speciation analysis method.

6. Detection systems of arsenic species

Several sensitive and element techniques can be used for the detection of arsenic species. Various detection techniques are: atomic mass spectrometry, molecular mass spectrometry, optical spectrometry, X-ray methods and others (voltammetry, potentiometry, conductometry and spectrophotometry), which provide different level of specificity, cost effectiveness and detection limits [21]. Detection methods applied for the arsenic speciation analysis of soil-plant-water samples are assembled in **Table 3**.

ICP-MS is the most commonly used technique for the detection of multiple arsenic species because of its high sensitivity, high selectivity and wide dynamic range. The coupling of chromatography to ICP-MS has several benefits due to the compatibility of the mobile phase with the behavior of the plasma torch and the carefully determined quality inspection interference. Various techniques have been developed to eliminate or reduce isobaric interference in the detection of arsenic by mass-to-charge ratio. Recently, compared with the traditional single quadrupole ICP-MS, the combination of ICP and triple quadrupole tandem mass spectrometry (ICP-QQQ) helps to eliminate isobaric interference, reduce background, and improve selectivity [156]. Quantification is performed by preparing standard solutions of commercially available substances, such as iAs(III), iAs(V), MA, DMA, and AB. It is generally believed that arsenate is used to calibrate anionic substances, and arsenobetaine is used to calibrate cationic substances [157]. DMA is considered to be the most suitable calibration standard for arsenic lipid quantification [158]. The sensitivity of ICPMS to detect arsenic is limited by its relatively high ionization potential. In order to compensate for this effect, various methods have been used,

Matrix	Arsenic species	Detection techniques	Coupled with	References
Atomic mass spectrometry				
Soil	AS ^{III} , AS ^V	ICP = MS	—	[130]
Water	AS ^{III} , AS ^V , MMA ^V , DMA ^V			[131]
Soil-water-plant	Total As, AS ^{III} , AS ^V ,			[132]
Water	AS ^{III} , AS ^V	ICP-SFMS	—	[133]
Water	AS ^{III} , AS ^V , MMA ^V , DMA ^V , AsB AsC	ICP = MS	HPLC	[134]
Plant	AS ^{III} , AS ^V , MMA ^V , DMA ^V , AsB			[135]
Soil	AS ^{III} , AS ^V			[136]
Soil	AS ^{III} , AS ^V , MMA ^V , DMA ^V	ICP-SFMS	HPLC	[137]
Plant	AS ^{III} , AS ^V , MMA ^V , DMA ^V			[138]
Water	AS ^{III} , AS ^V , MMA ^V , DMA ^V			[139]
Soil	MMA ^V , DMA ^V	ICP = MS	GC	[140]
Soil	AS ^{III} , AS ^V , MMA ^V , DMA ^V	ICP = MS	CE	[137]
Molecular mass spectrometry				
Soil	PA and AA	ESI-qMS	HPLC	[141]
Soil -water	PA and AA	ESI-TOF-MS	=	[142]
Soil-plant	AS ^{III} , AS ^V , MMA ^V , DMA ^V , MMMTA, DMMTA, DMDTA	ESI-qTOF-MS	=	[143]
Soil-plant	AS ^{III} , AS ^V , MMA ^V , DMA ^V , MMMTA, DMMTA, DMDTA	ESI-qTOF-MS	HPLC	[143]
Soil	AS ^{III} , AS ^V , N-AHPAA, 3-AHPAA	ESI-triple quad-MS	=	[144]
Plant	Arsenolipids	ESI-triple quad-MS	HPLC	[145]
Water	AS ^{III} , AS ^V , MMA ^V , DMA ^V , TMAO	ESI-Orbitrap-MS	=	[146]
Plant	Arsenic peptides	ESI-IT-MS	HPLC	[147]
Plant	PA and AA	EI-MS	GC	[141]
Optical spectroscopy				
Plant	AS total, AS ^{III} , AS ^V	GF-AAS	=	[148]
Soil	AS total, AS ^{III} , AS ^V	HG-AAS	=	[85]
Water	AS ^{III} , AS ^V , MMA ^V , DMA ^V	HG-AAS	HPLC	[127]
Soil-plant	AS ^{III} , AS ^V , MMA ^V , DMA ^V			[69]
Soil	AS total, AS ^{III} , AS ^V	HG-AFS	=	[87]
Soil	AS ^{III} , AS ^V , MMA ^V , DMA ^V	HG-AFS	HPLC	[88]
Plant	AS ^{III} , AS ^V , MMA ^V , DMA ^V			[149]

Matrix	Arsenic species	Detection techniques	Coupled with	References
X-ray methods				
Soil and Plant	AS ^{III} , AS ^V , MMA ^V , DMA ^V	XANES		[150]
Soil	AS ^{III} , AS ^V	EXAFS		[151]
Soil	AS ^{III} , AS ^V	STXM		[152]
Soil	AS ^V	XPS		[153]
Others				
Water	AS ^{III} , AS ^V ,	Voltammetry	=	[154]
Water	AS ^V	Potentiometry	=	[155]
Water	AS ^{III} , AS ^V	Spectrophotometry	HPLC	[126]

Aqueous phenylarsonic acid (PA); o-arsanilic acid (AA); N-acetyl-4-hydroxy-m-arsanilic acid (N-AHPAA), 3-amino-4-hydroxyphenylarsonic acid (3-AHPAA).

Table 3.
Examples of detection systems for arsenic speciation analysis of soil-water-plant samples.

including adding supplemental methanol or ethanol solution to the spray chamber [159] or after the column via the T-piece [91], and the use of correction response factors. Finally, internal standardization was used to overcome the non-spectral matrix effects and instrumental drift [160].

Recently, molecular mass spectrometry is considered as a forward-looking technique for arsenic speciation analysis, especially for the detection of new organic arsenic species, such as thioarsenosugar [21, 161] and arsenolipids [21, 60, 162]. In this detection technique, the purified part of the extractable sample is introduced by electrospray ionization (ESI), and then mass spectrometry is combined with liquid chromatography. Generally, for the As forms, a simple single quadrupole mass analyzer is used, while tandem mass spectrometry is used for precise structure determination, whether it is a “spatial” triple quadrupole or a quadrupole time combination, or a “time” and Orbitrap system [21]. However, it has been recognized that ESI-MS analysis lacks selectivity for complex matrices, and quantification is more difficult than ICP-MS [163]. Therefore, the most powerful setting for arsenic speciation analysis that combines atomic and mass spectrometers is used as the detector of the same chromatographic system [21, 60, 145].

The optical spectroscopy technique such as atomic absorption spectroscopy (AAS) and atomic fluorescence spectroscopy (AFS) is popular to researcher as an attractive alternative to mass spectrometry. Due to the low purchase and operation cost, high speed, low consumption of organic solvents, high enrichment coefficient, combined with hydride generation provides high sensitivity and reduced matrix effect, this technology has been applied to the determination of arsenic species in environmental samples. Moreover, hydride generation systems (HG-AAS and HG-AFS) facilitate a direct measurement of the more As. Graphite furnace atomic absorption spectroscopy can be an independent facility and does not require AsH₃ because of the low level of interference [18]. In fact, the optical spectroscopy is an effective technique, when combined with different separation techniques and chemical modifiers, iAs(III), iAs(V), MA, DMA and TMAO, can be identified, and significant hydride generation of arsenosugars [164] and thioarsenates can be observed [21]. Nevertheless, HGAAS and HG-AFS are mainly used for water samples [150], sediment extracts and soil, [165] and plants [97] mainly contain

inorganic arsenic. These techniques are also applicable to biological substrates and more stubborn arsenic Analysis [166].

X-ray method is an important technique for morphological analysis of environmental samples, which can record raw data about the chemical environment of arsenic atoms in situ without sample preparation. X-ray atomic absorption spectroscopy (XAS) is generally divided into two regions: X-ray absorption near edge structure (XANES) and extended X-ray absorption fine structure (EXAFS). These technologies are mainly used to directly detect solid samples, including sediments, [167] soil [97, 168] and plants [96, 97, 169]. Both XANES and EXAFS have studied abiotic matrices to measure arsenic redox status and geochemical correlation.

7. Accuracy evaluation of speciation methods

In order to obtain precise analytical information about the bioavailability and toxicity of arsenic in the environmental process, it is necessary to carefully consider any possible sources of error during analysis and validate the data. To avoid or minimize the impact of species changes and ensure the reliability and quality of speciation data, mass balance ratio, extraction efficiency, column recovery of arsenic species during separation and standard reference materials quality need to be tracked. The main difficulty of specific analysis of arsenic may occur in the sample preparation stage and species stability. Mass balance data provides information about the distribution of elements in each analysis step (extraction, separation, and species detection) and quantitatively determine the fate of arsenic during speciation [170]. The extraction efficiency can provide some important information about the extraction procedure, the polarity of the extracted species, and help to select the best extraction solvent and separation system. It helps to establish a non-toxic, effective and simple extraction procedure for arsenic speciation analysis. Column recovery is an important aspect of any separation technique. It is critical to eliminate loss and to ensure there is no cross-contamination between analyses. The column recovery compares the total arsenic concentration with the sum of the detected substances, which can provide information about the elution and retention of the analyte. In fact, depending on the type of sample and the concentration of the arsenic species, the column recovery rate of the arsenic species has great variability [171]. The column recovery also affected by the extraction solvent of the column. It is difficult to evaluate the mutual transformation of arsenic species in the actual sample in the column, which is related to the individual arsenic standard. However, the lack of available standards for new arsenic species is the main challenge in studying the inter-conversion of arsenic compounds during separation [15]. For accurate method validation of arsenic speciation, the use of standard reference materials (SRM) and certified reference materials (CRM) is essential. With reference or certified values available, SRM and CRM can be used to test and verify the accuracy of the method. In order to verify the arsenic speciation analysis methods of environmental samples, different types of soil and sediments, natural waters, marine and terrestrial plants and other biological samples are used as reference samples. It should be noted that a single SRM or CRM could not be used to verify method calibration and results [15]. SRM 1640 (NIST) is commonly used to check calibration curves for trace elements in water. The type of CRM used depends on the sample matrix and the type of arsenic studied [1].

8. Conclusion

Arsenic pollution is a universal problem. The form of arsenic in soil, water, and plants play an important role in understanding arsenic exposure, metabolism and environmental arsenic cycle, and food chain. A crucial requirement for obtaining reliable speciation information is to maintain the concentration of the original chemical species in the sample prior to analysis. In order to determine the total element concentration, the main considerations for sample collection and storage are to prevent contamination and minimize the loss of trace analytes. Research on simple and efficient extraction procedures that use less or non-toxic solvents is very urgent for better arsenic speciation. In the case of speciation analysis, the concentration of individual species of the element must be constant through sample handling and processing. Therefore, the time between the extraction procedure and the analysis must be as short as possible to avoid inter-conversion between species. The selection of extraction and sample preparation methods must be complementary and compatible with the separation method in order to perform qualitative and quantitative analysis of arsenic species and its concentration. It may require a combination of multiple extraction methods and multiple separation techniques to achieve a comprehensive arsenic speciation analysis. Several techniques have been used to study arsenic speciation, each with its advantages and disadvantages. However, research efforts are still needed to develop cheap, fast, sensitive, and reproducible methods for arsenic species that can work at low detection limits. However, research efforts are still needed to develop cheap, fast, sensitive, and reproducible methods for arsenic species that can work at low detection limits. In addition, in order to find a unified analysis protocol i.e. at least for the more common matrices, for the prevalent and unidentified arsenic species, advanced investigations and routine measurements are necessary.

Conflict of interests

The authors declare that they have no conflicting interests.

IntechOpen

IntechOpen

Author details

Mohammed Zia Uddin Kamal* and Md. Yunus Miah
Department of Soil Science, Bangabandhu Sheikh Mujibur Rahman Agricultural
University (BSMRAU), Gazipur, Bangladesh

*Address all correspondence to: zia_ssc@yahoo.com; zia@bsmrau.edu.bd

IntechOpen

© 2021 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. 

References

- [1] Sadee B, Foulkes ME, Hill SJ. Coupled techniques for arsenic speciation in food and drinking water: a review. *Journal of Analytical Atomic Spectrometry*. 2015; 30(1): 102-18.
- [2] Khalid S, Shahid M, Niazi NK, Rafiq M, Bakhat HF, Imran M, Abbas T, Bibi I, Dumat C. Arsenic behaviour in soil-plant system: Biogeochemical reactions and chemical speciation influences. In *Enhancing cleanup of environmental pollutants 2017* (pp. 97-140). Springer, Cham.
- [3] Chandrakar V, Naithani SC, Keshavkant S. Arsenic-induced metabolic disturbances and their mitigation mechanisms in crop plants: A review. *Biologia*. 2016 May 1; 71(4): 367-77.
- [4] Abbas G, Murtaza B, Bibi I, Shahid M, Niazi NK, Khan MI, Amjad M, Hussain M. Arsenic uptake, toxicity, detoxification, and speciation in plants: physiological, biochemical, and molecular aspects. *International journal of environmental research and public health*. 2018; 15(1): 59.
- [5] Huq SI, Alam MD. A handbook on analyses of soil, plant and water. BACER-DU, University of Dhaka, Bangladesh. 2005; 246.
- [6] Chakraborty S, Alam MO, Bhattacharya T, Singh YN. Arsenic accumulation in food crops: a potential threat in Bengal delta plain. *Water Quality, Exposure and Health*. 2014 Dec 1; 6(4): 233-46.
- [7] Rahman MM, Asaduzzaman M, Naidu R. Arsenic exposure from rice and water sources in the Noakhali district of Bangladesh. *Water Quality, Exposure and Health*. 2011 Jun 1; 3(1): 1-0.
- [8] Rahman MA, Kadohashi K, Maki T, Hasegawa H. Transport of DMAA and MMAA into rice (*Oryza sativa* L.) roots. *Environmental and Experimental Botany*. 2011 Aug 1; 72(1): 41-6.
- [9] Rauf MA, Hakim MA, Hanafi MM, Islam MM, Rahman GK, Panaullah GM. Bioaccumulation of arsenic (As) and phosphorous by transplanting Aman rice in arsenic-contaminated clay soils. *Australian Journal of Crop Science*. 2011 Jan; 5(12): 1678-84.
- [10] Punshon T, Jackson BP, Meharg AA, Warczack T, Scheckel K, Guerinot ML. Understanding arsenic dynamics in agronomic systems to predict and prevent uptake by crop plants. *Science of the Total Environment*. 2017 Mar 1; 581:209-20.
- [11] FAO/WHO (Session TS). Joint FAO/WHO Food Standards Programme Codex Committee on Contaminants in Foods, 8th Session. Codex. 2014 Dec; 150(2014/8). World Health Organization, Geneva
- [12] US Food and Drug Administration. Questions & Answers: Arsenic in Rice and Rice Products; US. FDA: Silver Spring, MD, USA, 2015
- [13] Bakhat HF, Zia Z, Fahad S, Abbas S, Hammad HM, Shahzad AN, Abbas F, Alharby H, Shahid M. Arsenic uptake, accumulation and toxicity in rice plants: possible remedies for its detoxification: a review. *Environmental Science and Pollution Research*. 2017 Apr 1; 24(10): 9142-58.
- [14] Sun Y, Liu G, Cai Y. Thiolated arsenicals in arsenic metabolism: occurrence, formation, and biological implications. *Journal of Environmental Sciences*. 2016 Nov 1; 49:59-73.
- [15] Reid MS, Hoy KS, Schofield JR, Uppal JS, Lin Y, Lu X, Peng H, Le XC. Arsenic speciation analysis: A review with an emphasis on chromatographic

separations. *TrAC Trends in Analytical Chemistry*. 2020 Feb 1; 123:115770.

[16] Moe B, Peng H, Lu X, Chen B, Chen LW, Gabos S, Li XF, Le XC. Comparative cytotoxicity of fourteen trivalent and pentavalent arsenic species determined using real-time cell sensing. *Journal of Environmental Sciences*. 2016 Nov 1; 49:113-24.

[17] Ebert F, Leffers L, Weber T, Berndt S, Mangerich A, Beneke S, Bürkle A, Schwerdtle T. Toxicological properties of the thiolated inorganic arsenic and arsenosugar metabolite thio-dimethylarsinic acid in human bladder cells. *Journal of Trace Elements in Medicine and Biology*. 2014 Apr 1; 28(2): 138-46.

[18] Rukh S, Akhtar MS, Memon M, Mehmood A, Imran M. An overview of arsenic extraction and speciation techniques in soil and water. *Amer. Chem. Sci. J*. 2015; 6:1-5.

[19] Templeton DM, Ariese F, Cornelis R, Danielsson LG, Muntau H, van Leeuwen HP, Lobinski R. Guidelines for terms related to chemical speciation and fractionation of elements. Definitions, structural aspects, and methodological approaches (IUPAC Recommendations 2000). *Pure and applied chemistry*. 2000 Aug 31; 72(8): 1453-70.

[20] Gong Z, Lu X, Ma M, Watt C, Le XC. Arsenic speciation analysis. *Talanta*. 2002 Aug 16; 58(1): 77-96.

[21] Ardini F, Dan G, Grotti M. Arsenic speciation analysis of environmental samples. *Journal of Analytical Atomic Spectrometry*. 2020; 35(2): 215-37.

[22] Ng JC, Wang J, Shraim A. A global health problem caused by arsenic from natural sources. *Chemosphere*. 2003 Sep 1; 52(9): 1353-9.

[23] Caumette G, Koch I, Reimer KJ. Arsenobetaine formation in plankton: a

review of studies at the base of the aquatic food chain. *Journal of Environmental Monitoring*. 2012; 14(11): 2841-53.

[24] Taylor V, Goodale B, Raab A, Schwerdtle T, Reimer K, Conklin S, Karagas MR, Francesconi KA. Human exposure to organic arsenic species from seafood. *Science of the Total Environment*. 2017 Feb 15; 580:266-82.

[25] Jackson BP. Fast ion chromatography-ICP-QQQ for arsenic speciation. *Journal of analytical atomic spectrometry*. 2015; 30(6): 1405-7.

[26] Park MK, Choi M, Kim L, Choi SD. An improved rapid analytical method for the arsenic speciation analysis of marine environmental samples using high-performance liquid chromatography/inductively coupled plasma mass spectrometry. *Environmental monitoring and assessment*. 2019 Aug; 191(8): 1-4.

[27] Liu Q, Lu X, Peng H, Popowich A, Tao J, Uppal JS, Yan X, Boe D, Le XC. Speciation of arsenic—a review of phenylarsenicals and related arsenic metabolites. *TrAC Trends in Analytical Chemistry*. 2018 Jul 1; 104:171-82.

[28] Reis VA, Duarte AC. Analytical methodologies for arsenic speciation in macroalgae: a critical review. *TrAC Trends in Analytical Chemistry*. 2018 May 1; 102:170-84.

[29] Pfeifer H R, Beatrizotti G, Berthoud J, Rossa M D, Girardet A, Jäggli M, Lavanchy J C, Reymond D, Righetti G, Schlegel C, Schmit V, Temgoua E. Natural arsenic-contamination of surface and ground waters in Southern Switzerland (Ticino). *Bull. Appl. Geol*. 2002; 7:81-103.

[30] Jang YC, Somanna Y, Kim H. Source, distribution, toxicity and remediation of arsenic in the environment—a review. *Int J Appl Environ Sci*. 2016; 11(2): 559-81.

- [31] Onishi, H, & Wedepohl KH. Handbook of geochemistry. 1969; Vol. II-2. Springer: Berlin, Germany.
- [32] Shahid M, Shamshad S, Rafiq M, Khalid S, Bibi I, Niazi NK, Dumat C, Rashid MI. Chromium speciation, bioavailability, uptake, toxicity and detoxification in soil-plant system: a review. *Chemosphere*. 2017 Jul 1; 178:513-33.
- [33] Niazi NK, Singh B, Shah P. Arsenic speciation and phytoavailability in contaminated soils using a sequential extraction procedure and XANES spectroscopy. *Environmental science & technology*. 2011 Sep 1; 45(17): 7135-42.
- [34] Baig JA, Kazi TG, Shah AQ, Kandhro GA, Afridi HI, Khan S, Kolachi NF. Biosorption studies on powder of stem of *Acacia nilotica*: removal of arsenic from surface water. *Journal of hazardous materials*. 2010 Jun 15; 178(1-3): 941-8.
- [35] Sadiq M. Arsenic chemistry in soils: an overview of thermodynamic predictions and field observations. *Water, air, and soil pollution*. 1997 Jan; 93(1): 117-36.
- [36] Koyama T. Arsenic in soil-plant system. *Nippon Dojo Hiryogaku Zasshi*. 1975; 46:491-502.
- [37] Bhattacharya P, Jacks G, Frisbie S H, Smith E, Naidu R, Sarkar B. Arsenic in the environment: a global perspective. in Sarkar, B. (Eds.). *Heavy Metals in the Environment*. Marcel Dekker. Inc. New York: 2002. p 147-215.
- [38] Rhine E D, Phelps C D, Young L Y. Anaerobic arsenite oxidation by novel denitrifying isolates. *Environ. Microbiol*. 2006; 8: 899-908.
- [39] Mandal B K, Suzuki K T. Arsenic round the world: a review. *Talanta*. 2002; 58: 201-235.
- [40] Sadiq M, Zaidi TH, Mian AA. Environmental behavior of arsenic in soils: Theoretical. *Water, Air, & Soil Pollution*. 1983 Nov; 20(4): 369-77.
- [41] Mahimairaja S, Bolan NS, Adriano DC, Robinson B. Arsenic contamination and its risk management in complex environmental settings. *Advances in Agronomy*. 2005 Jan 1; 86:1-82.
- [42] Marquez EB, Gurian PL, Barud-Zubillaga A, Goodell PC. Correlates of arsenic mobilization into the groundwater in El Paso, Texas. *Air Soil Water Res* 2011; 4:19-29
- [43] Rahaman S, Sinha AC, Mukhopadhyay D. Effect of water regimes and organic matters on transport of arsenic in summer rice (*Oryza sativa* L.). *Journal of Environmental Sciences*. 2011 Apr 1; 23(4): 633-9.
- [44] Williams LE, Barnett MO, Kramer TA, Melville JG. Adsorption and transport of arsenic (V) in experimental subsurface systems. *Journal of Environmental Quality*. 2003 May; 32(3): 841-50.
- [45] Mishra S, Mattusch J, Wennrich R. Accumulation and transformation of inorganic and organic arsenic in rice and role of thiol-complexation to restrict their translocation to shoot. *Sci. Rep*. 2017, 7, 40522.
- [46] Ayangbenro AS, Babalola OO. A new strategy for heavy metal polluted environments: a review of microbial biosorbents. *International journal of environmental research and public health*. 2017 Jan; 14(1):94.
- [47] Shahid N, Zia Z, Shahid M, Faiq Bakhat H, Anwar S, Mustafa Shah G, Rizwan Ashraf M. Assessing Drinking Water Quality in Punjab, Pakistan. *Polish J. Environ. Stud*. 2015, 24, 2597-2606.

- [48] Polizzotto ML, Harvey CF, Li G, Badruzzman B, Ali A, Newville M, Sutton S, Fendorf S. Solid-phases and desorption processes of arsenic within Bangladesh sediments. *Chemical Geology*. 2006 Apr 16; 228(1-3): 97-111.
- [49] Wilkin RT, Wallschläger D, Ford RG. Speciation of arsenic in sulfidic waters. *Geochemical Transactions*. 2003; 4(1): 1-7.
- [50] Cabon JY, Cabon N. Speciation of major arsenic species in seawater by flow injection hydride generation atomic absorption spectrometry. *Fresenius' journal of analytical chemistry*. 2000 Oct; 368(5): 484-9.
- [51] Nriagu J O. Arsenic in the environment. Wiley. New York: 1994. p. 448
- [52] Adriano DC. Arsenic. In *Trace Elements in Terrestrial Environments*; Springer: Berlin, Germany, 2001; pp. 219-261.
- [53] Catarcha P, Segura MD, Franco-Zorrilla JM, García-Ponce B, Lanza M, Solano R, Paz-Ares J, Leyva A. A mutant of the *Arabidopsis* phosphate transporter PHT1; 1 displays enhanced arsenic accumulation. *Plant Cell*, 2007; 19:1123-1133.
- [54] Shin H, Shin HS, Dewbre GR, Harrison MJ. Phosphate transport in *Arabidopsis*: Pht1; 1 and Pht1;4 play a major role in phosphate acquisition from both low- and high-phosphate environments. *Plant Journal*, 2004; 39:629-642.
- [55] Wu Z, Ren H, McGrath SP, Wu P, Zhao FJ. Investigating the contribution of the phosphate transport pathway to arsenic accumulation in rice. *Plant Physiology*, 2011; 157:498-508.
- [56] Ma JF, Yamaji N, Mitani N, Xu XY, Su YH, McGrath SP, Zhao FJ. Transporters of arsenite in rice and their role in arsenic accumulation in rice grain. *Proceedings of the National Academy of Sciences*. 2008 Jul 22; 105(29): 9931-5.
- [57] Kumar S, Dubey RS, Tripathi RD, Chakrabarty D, Trivedi PK. Omics and biotechnology of arsenic stress and detoxification in plants: current updates and prospective. *Environment international*. 2015 Jan 1; 74:221-30.
- [58] Liu WJ, Wood BA, Raab A, McGrath SP, Zhao FJ, Feldmann J. Complexation of arsenite with phytochelatins reduces arsenite efflux and translocation from roots to shoots in *Arabidopsis*. *Plant Physiology*. 2010 Apr 1; 152(4): 2211-21.
- [59] Ye WL, Wood BA, Stroud JL, Andralojc PJ, Raab A, McGrath SP, Feldmann J, Zhao FJ. Arsenic speciation in phloem and xylem exudates of castor bean. *Plant physiology*. 2010 Nov 1; 154(3): 1505-13.
- [60] Raab A, Williams PN, Meharg A, Feldmann J. Uptake and translocation of inorganic and methylated arsenic species by plants. *Environmental Chemistry*. 2007 Jul 13; 4(3): 197-203.
- [61] Huang JH, Fecher P, Ilgen G, Hu KN, Yang J. Speciation of arsenite and arsenate in rice grain—Verification of nitric acid based extraction method and mass sample survey. *Food chemistry*. 2012 Jan 15; 130(2): 453-9.
- [62] Abedin MJ, Feldmann J, Meharg AA. Uptake kinetics of arsenic species in rice plants. *Plant physiology*. 2002 Mar 1; 128(3): 1120-8.
- [63] Feldmann J. R. Cornelis, J. Caruso, H. Crews and K. Heumann. *Handbook of element speciation techniques and methodology*. John Wiley & Sons, 2005. Chichester, ISBN: 0-471-49214-0. 634+ xii pp.49
- [64] Moreda-Piñero, E. Alonso-Rodríguez, A. Moreda-Piñero, C.

- Moscoso-Pérez, S. Muniategui-Lorenzo, P. López-Mahía, D. Prada-Rodríguez and P. Bermejo-Barrera, *Anal. Chim. Acta*, 2010, 679, 63-73.
- [65] Rajaković LV, Todorović ŽN, Rajaković-Ognjanović V, Onjia A. Analytical methods for arsenic speciation analysis. *Journal of the Serbian Chemical Society*. 2013; 78(10): 1461-79.
- [66] Bentley R, Chasteen TG. Microbial methylation of metalloids: arsenic, antimony, and bismuth. *Microbiology and molecular biology reviews*. 2002 Jun; 66(2): 250.
- [67] Mester Z, Sturgeon RE. Sample preparation for trace element analysis. Elsevier; 2003 Dec 16.
- [68] Amaral CD, Nóbrega JA, Nogueira AR. Sample preparation for arsenic speciation in terrestrial plants—a review. *Talanta*. 2013 Oct 15; 115:291-9.
- [69] Mir KA, Rutter A, Koch I, Smith P, Reimer KJ, Poland JS. Extraction and speciation of arsenic in plants grown on arsenic contaminated soils. *Talanta*. 2007 Jun 15; 72(4): 1507-18.
- [70] Aggett J, Kriegman MR. Preservation of arsenic (III) and arsenic (V) in samples of sediment interstitial water. *Analyst*. 1987; 112(2): 153-7.
- [71] Segura M, Muñoz J, Madrid Y, Cámara C. Stability study of As (III), As (V), MMA and DMA by anion exchange chromatography and HG-AFS in wastewater samples. *Analytical and bioanalytical chemistry*. 2002 Oct; 374(3): 513-9.
- [72] Mc Cleskey RB, Nordstrom DK, Meast AS. Preservation of water samples for arsenic (III/V) determinations: An evaluation of the literature and new analytical results. *Applied Geochemistry*. 2004; 9:995-1009.
- [73] Hall GM. Stability of inorganic arsenic (III) and arsenic (V) in water samples. *Journal of Analytical Atomic Spectrometry*. 1999; 14(2): 205-13.
- [74] Nikolaidis NP, Dobbs GM, Chen J, Lackovic JA. Arsenic mobility in contaminated lake sediments. *Environmental pollution*. 2004 Jun 1; 129(3): 479-87.
- [75] Daus B, Weiss H, Mattusch J, Wennrich R. Preservation of arsenic species in water samples using phosphoric acid—Limitations and long-term stability. *Talanta*. 2006 Apr 15; 69(2): 430-4.
- [76] Edwards M, Patel S, McNeill L, Chen HW, Frey M, Eaton AD, Antweiler RC, Taylor HE. Considerations in As analysis and speciation. *Journal-American Water Works Association*. 1998 Mar; 90(3): 103-13.
- [77] Melanie G, Zagury JG, Deschenes L, Blouin J. Comparison of four extraction procedures to assess arsenate and arsenite species in contaminated soils. *Environmental Pollution*. 2009; 158:1890-1898.
- [78] Van Herreweghe S, Swennen R, Vandecasteele C, Cappuyns V. Solid phase speciation of arsenic by sequential extraction in standard reference materials and industrially contaminated soil samples. *Environmental pollution*. 2003 Apr 1; 122(3): 323-42.
- [79] Baig JA, Kazi TG, Shah AQ, Arain MB, Afridi HI, Kandhro GA, Khan S. Optimization of cloud point extraction and solid phase extraction methods for speciation of arsenic in natural water using multivariate technique. *Analytica Chimica Acta*. 2009 Sep 28; 651(1): 57-63.
- [80] Wu J, Mester Z, Pawliszyn J. Speciation of organoarsenic compounds by polypyrrole-coated capillary in-tube

solid phase microextraction coupled with liquid chromatography/electrospray ionization mass spectrometry. *Analytica chimica acta*. 2000 Dec 1; 424(2): 211-22.

[81] Pizarro I, Gómez M, Cámara C, Palacios MA. Arsenic speciation in environmental and biological samples: extraction and stability studies. *Analytica Chimica Acta*. 2003 Oct 24; 495(1-2): 85-98.

[82] Carabias-Martínez, R., Rodríguez-Gonzalo, E., Revilla-Ruiz, P. and Hernández-Méndez, J., 2005. Pressurized liquid extraction in the analysis of food and biological samples. *Journal of Chromatography A*, 1089(1-2), pp. 1-17.

[83] Foster S, Maher W, Krikowa F, Apte S. A microwave-assisted sequential extraction of water and dilute acid soluble arsenic species from marine plant and animal tissues. *Talanta*. 2007 Feb 15; 71(2):537-49.

[84] Dietz C, Sanz J, Sanz E, Munoz-Olivas R, Cámara C. Current perspectives in analyte extraction strategies for tin and arsenic speciation. *Journal of Chromatography A*. 2007 Jun 15; 1153(1-2): 114-29.

[85] Chappell J, Chiswell B, Olszowy H. Speciation of arsenic in a contaminated soil by solvent extraction. *Talanta*. 1995 Mar 1; 42(3): 323-9.

[86] Georgiadis M, Cai Y, Solo-Gabriele HM. Extraction of arsenate and arsenite species from soils and sediments. *Environmental Pollution*. 2006 May 1; 141(1): 22-9.

[87] Garcia-Manyes S, Jiménez G, Padró A, Rubio R, Rauret G. Arsenic speciation in contaminated soils. *Talanta*. 2002 Aug 16; 58(1): 97-109.

[88] Yuan CG, He B, Gao EL, Lü JX, Jiang GB. Evaluation of extraction

methods for arsenic speciation in polluted soil and rotten ore by HPLC-HG-AFS analysis. *Microchimica Acta*. 2007 Jun 1; 159(1-2): 175-82.

[89] Wu J, Zhang SD, Zhu XS. Study on the analytical method of inorganic arsenic species in environmental samples. *Applied Ecology and Environmental Research*. 2019 Jan 1; 17(4):7943-55.

[90] Rahman MS, Abdul Mazid Miah M, Khaled HM, Islam A, Panaullah GM. Arsenic concentrations in groundwater, soils, and irrigated rice in Southwestern Bangladesh. *Communications in soil science and plant analysis*. 2010 Aug 4; 41(16): 1889-95.

[91] Ruiz-Chancho MJ, Taleshi MS, Goessler W, Francesconi KA. A method for screening arsenolipids in fish oils by HPLC-ICPMS. *Journal of Analytical Atomic Spectrometry*. 2012; 27(3): 501-4.

[92] Gutiérrez-Ruiz ME, Cenicerós-Gómez AE, Villalobos M, Romero F, Santiago P. Natural arsenic attenuation via metal arsenate precipitation in soils contaminated with metallurgical wastes: II. Cumulative evidence and identification of minor processes. *Applied geochemistry*. 2012 Nov 1; 27(11): 2204-14.

[93] Tripathi P, Singh PC, Mishra A, Srivastava S, Chauhan R, Awasthi S, Mishra S, Dwivedi S, Tripathi P, Kalra A, Tripathi RD. Arsenic tolerant *Trichoderma* sp. reduces arsenic induced stress in chickpea (*Cicer arietinum*). *Environmental Pollution*. 2017 Apr 1; 223:137-45.

[94] Márquez-García B, Pérez-López R, Ruíz-Chancho MJ, López-Sánchez JF, Rubio R, Abreu MM, Nieto JM, Córdoba F. Arsenic speciation in soils and *Erica andevalensis* Cabezudo & Rivera and *Erica australis* L. from São Domingos Mine area, Portugal. *Journal*

of Geochemical Exploration. 2012 Aug 1; 119:51-9.

[95] Zhang X, Uroic MK, Xie WY, Zhu YG, Chen BD, McGrath SP, Feldmann J, Zhao FJ. Phytochelatins play a key role in arsenic accumulation and tolerance in the aquatic macrophyte *Wolffia globosa*. Environmental Pollution. 2012 Jun 1; 165:18-24.

[96] Yang F, Xie S, Wei C, Liu J, Zhang H, Chen T, Zhang J. Arsenic characteristics in the terrestrial environment in the vicinity of the Shimen realgar mine, China. Science of the Total Environment. 2018 Jun 1; 626:77-86.

[97] Bergqvist C, Herbert R, Persson I, Greger M. Plants influence on arsenic availability and speciation in the rhizosphere, roots and shoots of three different vegetables. Environmental pollution. 2014 Jan 1; 184:540-6.

[98] Rahman F, Chen Z, Naidu R. A comparative study of the extractability of arsenic species from silverbeet and amaranth vegetables. Environmental Geochemistry and Health. 2009 Apr; 31(1): 103-13.

[99] Paik MK, Kim MJ, Kim WI, Yoo JH, Park BJ, Im GJ, Park JE, Hong MK. Determination of arsenic species in polished rice using a methanol-water digestion method. Journal of the Korean Society for Applied Biological Chemistry. 2010 Oct; 53(5): 634-8.

[100] Kim JY, Kim WI, Kunhikrishnan A, Kang DW, Kim DH, Lee YJ, Kim YJ, Kim CT. Determination of arsenic species in rice grains using HPLC-ICP-MS. Food Science and Biotechnology. 2013 Dec 1; 22(6): 1509-13

[101] López-García I, Briceño M, Hernández-Córdoba M. Non-chromatographic screening procedure for arsenic speciation analysis in

fish-based baby foods by using electrothermal atomic absorption spectrometry. Analytica chimica acta. 2011 Aug 5; 699(1): 11-7.

[102] Sun J, Yang Z, Lee H, Wang L. Simultaneous speciation and determination of arsenic, chromium and cadmium in water samples by high performance liquid chromatography with inductively coupled plasma mass spectrometry. Analytical Methods. 2015; 7(6): 2653-8.

[103] Pereira ÉR, Kopp JF, Raab A, Krupp EM, del Campo Menoyo J, Carasek E, Welz B, Feldmann J. Arsenic containing medium and long chain fatty acids in marine fish oil identified as degradation products using reversed-phase HPLC-ICP-MS/ESI-MS. Journal of Analytical Atomic Spectrometry. 2016; 31(9): 1836-45.

[104] Rubio R, Ruiz-Chancho MJ, López-Sánchez JF. Sample pre-treatment and extraction methods that are crucial to arsenic speciation in algae and aquatic plants. TrAC Trends in Analytical Chemistry. 2010 Jan 1; 29(1): 53-69.

[105] He B, Fang Y, Jiang G, Ni Z. Optimization of the extraction for the determination of arsenic species in plant materials by high-performance liquid chromatography coupled with hydride generation atomic fluorescence spectrometry. Spectrochimica Acta Part B: Atomic Spectroscopy. 2002 Nov 11; 57(11): 1705-11.

[106] Zheng J, Hintelmann H. HPLC-ICP-MS for a comparative study on the extraction approaches for arsenic speciation in terrestrial plant, *Ceratophyllum demersum*. Journal of Radioanalytical and Nuclear Chemistry. 2009 Apr; 280(1): 171-9.

[107] Mrak T, Šlejkovec Z, Jeran Z. Extraction of arsenic compounds from lichens. Talanta. 2006 Mar 15; 69(1): 251-8.

- [108] Amaral CD, Nóbrega JA, Nogueira AR. Sample preparation for arsenic speciation in terrestrial plants—a review. *Talanta*. 2013 Oct 15; 115:291-9.
- [109] Larios R, Fernández-Martínez R, LeHecho I, Rucandio I. A methodological approach to evaluate arsenic speciation and bioaccumulation in different plant species from two highly polluted mining areas. *Science of the Total Environment*. 2012 Jan 1; 414:600-7.
- [110] Guzmán Mar JL, Hinojosa Reyes L, Mizanur Rahman GM, Kingston HS. Simultaneous extraction of arsenic and selenium species from rice products by microwave-assisted enzymatic extraction and analysis by ion chromatography-inductively coupled plasma-mass spectrometry. *Journal of Agricultural and Food Chemistry*. 2009 Apr 22; 57(8): 3005-13.
- [111] Sadee BA, Foulkes ME, Hill SJ. An evaluation of extraction techniques for arsenic in staple diets (fish and rice) utilising both classical and enzymatic extraction methods. *Food Additives & Contaminants: Part A*. 2016 Mar 3; 33(3): 433-41.
- [112] Mattusch J, Cimpean M, Wennrich R. Enzyme-assisted extraction of arsenic species from plant material. *International Journal of Environmental and Analytical Chemistry*. 2006 Aug 10; 86(9): 629-40.
- [113] Raber G, Stock N, Hanel P, Murko M, Navratilova J, Francesconi KA. An improved HPLC–ICPMS method for determining inorganic arsenic in food: application to rice, wheat and tuna fish. *Food chemistry*. 2012 Sep 1; 134(1): 524-32.
- [114] Heitkemper DT, Vela NP, Stewart KR, Westphal CS. Determination of total and speciated arsenic in rice by ion chromatography and inductively coupled plasma mass spectrometry. *Journal of Analytical Atomic Spectrometry*. 2001; 16(4): 299-306.
- [115] Chen ML, Ma LY, Chen XW. New procedures for arsenic speciation: A review. *Talanta*. 2014 Jul 1; 125:78-86.
- [116] Issa NB, Rajaković-Ognjanović VN, Marinković AD, Rajaković LV. Separation and determination of arsenic species in water by selective exchange and hybrid resins. *Analytica chimica acta*. 2011 Nov 7; 706(1): 191-8.
- [117] Jia X, Gong D, Wang J, Huang F, Duan T, Zhang X. Arsenic speciation in environmental waters by a new specific phosphine modified polymer microsphere preconcentration and HPLC–ICP-MS determination. *Talanta*. 2016 Nov 1; 160:437-43.
- [118] Huang C, Xie W, Li X, Zhang J. Speciation of inorganic arsenic in environmental waters using magnetic solid phase extraction and preconcentration followed by ICP-MS. *Microchimica Acta*. 2011 Apr 1; 173(1-2): 165-72.
- [119] An MI, Zhang X, Yang T, Chen M, Wang J. Uptake and speciation of inorganic arsenic with cellulose fibre coated with yttrium hydroxide layer as a novel green sorbent. *Chinese Journal of Chemistry*. 2012 Sep; 30(9): 2225-31.
- [120] Qian S, Huang Z, Fu J, Kuang J, Hu C. Preconcentration of ultra-trace arsenic with nanometre-sized TiO₂ colloid and determination by AFS with slurry sampling. *Analytical Methods*. 2010; 2(8): 1140-3.
- [121] Chen M, Lin Y, Gu C, Wang J. Arsenic sorption and speciation with branch-polyethyleneimine modified carbon nanotubes with detection by atomic fluorescence spectrometry. *Talanta*. 2013 Jan 30; 104:53-7.

- [122] Güell R, Fontàs C, Salvadó V, Anticó E. Modelling of liquid–liquid extraction and liquid membrane separation of arsenic species in environmental matrices. *Separation and purification technology*. 2010 May 11; 72(3): 319-25.
- [123] Majidi B, Shemirani F. In situ solvent formation microextraction in the presence of ionic liquid for preconcentration and speciation of arsenic in saline samples and total arsenic in biological samples by electrothermal atomic absorption spectrometry. *Biological trace element research*. 2011 Oct; 143(1): 579-90.
- [124] Shraim A, Chiswell B, Olszowy H. Speciation of arsenic by hydride generation–atomic absorption spectrometry (HG–AAS) in hydrochloric acid reaction medium. *Talanta*. 1999 Dec 1; 50(5): 1109-27.
- [125] Anawar HM. Arsenic speciation in environmental samples by hydride generation and electrothermal atomic absorption spectrometry. *Talanta*. 2012 Jan 15; 88:30-42.
- [126] Cathum SJ, Brown CE, Obenauf A, Punt M. Speciation of arsenic using chelation solvent extraction and high performance liquid chromatography. *CLEAN–Soil, Air, Water*. 2007 Feb; 35(1): 71-80.
- [127] Akter KF, Chen Z, Smith L, Davey D, Naidu R. Speciation of arsenic in ground water samples: A comparative study of CE-UV, HG-AAS and LC-ICP-MS. *Talanta*. 2005 Dec 15; 68(2): 406-15.
- [128] Xie D, Mattusch J, Wennrich R. Separation of Organoarsenicals by Means of Zwitterionic Hydrophilic Interaction Chromatography (ZIC®-HILIC) and Parallel ICP-MS/ESI-MS Detection. *Engineering in Life Sciences*. 2008 Dec; 8(6): 582-8.
- [129] Shen S, Li XF, Cullen WR, Weinfeld M, Le XC. Arsenic binding to proteins. *Chemical reviews*. 2013 Oct 9; 113(10): 7769-92.
- [130] Migoni D, Papadia P, Cannito F, Fanizzi FP. Sequential Extraction Analysis of Arsenic in Soil Samples Collected in an Agricultural Area of Brindisi, Apulia (Italy), in the Proximity of a Coal-Burning Power Plant. *Applied Sciences*. 2021 Jan; 11(5): 2115.
- [131] Watts MJ, O'reilly J, Marcilla AL, Shaw RA, Ward NI. Field based speciation of arsenic in UK and Argentinean water samples. *Environmental geochemistry and health*. 2010 Dec 1; 32(6): 479-90.
- [132] Sadee BA, Foulkes ME, Hill SJ. A study of arsenic speciation in soil, irrigation water and plant tissue: a case study of the broad bean plant, *Vicia faba*. *Food chemistry*. 2016 Nov 1; 210:362-70.
- [133] Sugár É, Tatár E, Záray G, Mihucz VG. Field separation-based speciation analysis of inorganic arsenic in public well water in Hungary. *Microchemical Journal*. 2013 Mar 1; 107:131-5.
- [134] Hong S, Choi SD, Khim JS. Arsenic speciation in environmental multimedia samples from the Youngsan River Estuary, Korea: A comparison between freshwater and saltwater. *Environmental Pollution*. 2018 Jun 1; 237:842-50.
- [135] Nogueira R, Melo EA, Figueiredo JL, Santos JJ, Nascimento Neto AP. Arsenic Speciation in Fish and Rice by HPLC-ICP-MS Using Salt Gradient Elution. *Journal of the Brazilian Chemical Society*. 2018 Aug; 29(8): 1593-600.
- [136] Rahman MM, Chen Z, Naidu R. Extraction of arsenic species in soils using microwave-assisted extraction detected by ion chromatography

coupled to inductively coupled plasma mass spectrometry. *Environmental geochemistry and health*, 2009; 31(1): 93-102.

[137] Casiot C, Donard OF, Potin-Gautier M. Optimization of the hyphenation between capillary zone electrophoresis and inductively coupled plasma mass spectrometry for the measurement of As-, Sb-, Se-and Te-species, applicable to soil extracts. *Spectrochimica Acta Part B: Atomic Spectroscopy*. 2002 Jan 9; 57(1): 173-87

[138] Qu H, Mudalige TK, Linder SW. Arsenic speciation in rice by capillary electrophoresis/inductively coupled plasma mass spectrometry: enzyme-assisted water-phase microwave digestion. *Journal of agricultural and food chemistry*. 2015 Apr 1; 63(12): 3153-60.

[139] Richardson DD, Kannamkumarath SS, Wuilloud RG, Caruso JA. Hydride generation interface for speciation analysis coupling capillary electrophoresis to inductively coupled plasma mass spectrometry. *Analytical chemistry*. 2004 Dec 1; 76(23): 7137-42.

[140] Duester L, Diaz-Bone RA, Kösters J, Hirner AV. Methylated arsenic, antimony and tin species in soils. *Journal of Environmental Monitoring*. 2005; 7(12): 1186-93.

[141] Arroyo-Abad U, Mattusch J, Möder M, Elizalde-González MP, Matysik FM. Identification of degradation products of phenylarsonic acid and o-arsanilic acid in contact with suspensions of soils of volcanic origin. *Talanta*. 2012 Sep 15; 99:310-5.

[142] Arroyo-Abad U, Mattusch J, Möder M, Elizalde-González MP, Wennrich R, Matysik FM. Identification of roxarsone metabolites produced in the system: Soil-chlorinated water-light by using HPLC-ICP-MS/ESI-MS,

HPLC-ESI-MS/MS and High Resolution Mass Spectrometry (ESI-TOF-MS). *Journal of Analytical Atomic Spectrometry*. 2011; 26(1):171-7.

[143] Dai J, Chen C, Gao AX, Tang Z, Kopittke PM, Zhao FJ, Wang P. Dynamics of Dimethylated Monothioarsenate (DMMTA) in Paddy Soils and Its Accumulation in Rice Grains. *Environmental Science & Technology*. 2021 Jun 10.

[144] Huang K, Peng H, Gao F, Liu Q, Lu X, Shen Q, Le XC, Zhao FJ. Biotransformation of arsenic-containing roxarsone by an aerobic soil bacterium *Enterobacter* sp. CZ-1. *Environmental Pollution*. 2019 Apr 1; 247:482-7.

[145] Yu X, Xiong C, Jensen KB, Glabonjat RA, Stiboller M, Raber G, Francesconi KA. Mono-acyl arsenosugar phospholipids in the edible brown alga Kombu (*Saccharina japonica*). *Food Chemistry*. 2018 Feb 1; 240:817-21.

[146] Vriens B, Ammann AA, Hagendorfer H, Lenz M, Berg M, Winkel LH. Quantification of methylated selenium, sulfur, and arsenic in the environment. *PLOS one*. 2014 Jul 21; 9(7): 102906.

[147] Bluemlein an CG, Jiang GB, He B. Evaluation of the extraction methods for arsenic speciation in rice straw, *Oryza sativa* L., and analysis by HPLC-HG-AFS. *Journal of Analytical Atomic Spectrometry*. 2005; 20(2): 103-10.

[148] de Oliveira RM, Antunes AC, Vieira MA, Medina AL, Ribeiro AS. Evaluation of sample preparation methods for the determination of As, Cd, Pb, and Se in rice samples by GF AAS. *Microchemical Journal*. 2016 Jan 1; 124:402-9.

[149] Yuan CG, Jiang GB, He B. Evaluation of the extraction methods for arsenic speciation in rice straw, *Oryza sativa* L., and analysis by

HPLC-HG-AFS. *Journal of Analytical Atomic Spectrometry*. 2005; 20(2): 103-10.

[150] Huynh T, Harris HH, Zhang H, Noller BN. Measurement of labile arsenic speciation in water and soil using diffusive gradients in thin films (DGT) and X-ray absorption near edge spectroscopy (XANES). *Environmental Chemistry*. 2015 Feb 17; 12(2): 102-11.

[151] Voegelin A, Weber FA, Kretzschmar R. Distribution and speciation of arsenic around roots in a contaminated riparian floodplain soil: Micro-XRF element mapping and EXAFS spectroscopy. *Geochimica et Cosmochimica Acta*. 2007 Dec 1; 71(23): 5804-20.

[152] Wang D, Root RA, Chorover J. Biochar-templated surface precipitation and inner-sphere complexation effectively removes arsenic from acid mine drainage. *Environmental Science and Pollution Research*. 2021 Apr 18; 1-5.

[153] Frau F, Rossi A, Ardau C, Biddau R, Da Pelo S, Atzei D, Licheri C, Cannas C, Capitani G. Determination of arsenic speciation in complex environmental samples by the combined use of TEM and XPS. *Microchimica Acta*. 2005 Nov; 151(3): 189-201.

[154] Bullen JC, Torres-Huerta A, Salaün P, Watson JS, Majumdar S, Vilar R, Weiss DJ. Portable and rapid arsenic speciation in synthetic and natural waters by an As (V)-selective chemisorbent, validated against anodic stripping voltammetry. *Water research*. 2020 May 15; 175:115650.

[155] Giuffrè O, Aiello D, Chillè D, Napoli A, Foti C. Binding ability of arsenate towards Cu 2+ and Zn 2+: thermodynamic behavior and simulation under natural water conditions. *Environmental Science: Processes & Impacts*. 2020; 22(8): 1731-42.

[156] Jackson BP, Liba A, Nelson J. Advantages of reaction cell ICP-MS on doubly charged interferences for arsenic and selenium analysis in foods. *Journal of analytical atomic spectrometry*. 2015; 30(5): 1179-83.

[157] Stucker VK, Silverman DR, Williams KH, Sharp JO, Ranville JF. Thioarsenic species associated with increased arsenic release during biostimulated subsurface sulfate reduction. *Environmental science & technology*. 2014 Nov 18; 48(22): 13367-75.

[158] Sele V, Sloth JJ, Holmelid B, Valdersnes S, Skov K, Amlund H. Arsenic-containing fatty acids and hydrocarbons in marine oils—determination using reversed-phase HPLC–ICP-MS and HPLC–qTOF-MS. *Talanta*. 2014 Apr 1; 121:89-96.

[159] Khan M, Francesconi KA. Preliminary studies on the stability of arsenolipids: Implications for sample handling and analysis. *Journal of Environmental Sciences*. 2016 Nov 1; 49:97-103.

[160] Maher WA, Foster S, Krikowa F, Duncan E, St John A, Hug K, Moreau JW. Thio arsenic species measurements in marine organisms and geothermal waters. *Microchemical Journal*. 2013 Jul 1; 111:82-90.

[161] Lai VW, Kanaki K, Pergantis SA, Cullen WR, Reimer KJ. Arsenic speciation in freshwater snails and its life cycle variation. *Journal of Environmental Monitoring*. 2012; 14(3): 743-51.

[162] Raab A, Newcombe C, Pitton D, Ebel R, Feldmann J. Comprehensive analysis of lipophilic arsenic species in a brown alga (*Saccharina latissima*). *Analytical chemistry*. 2013 Mar 5; 85(5): 2817-24.

[163] Feldmann J, Raab A, Krupp EM. Importance of ICPMS for speciation

analysis is changing: future trends for targeted and non-targeted element speciation analysis. *Analytical and bioanalytical chemistry*. 2018 Jan;410(3):661-7.

[164] Schmeisser E, Goessler W, Kienzl N, Francesconi KA. Volatile analytes formed from arsenosugars: determination by HPLC–HG-ICPMS and implications for arsenic speciation analyses. *Analytical chemistry*. 2004 Jan 15; 76(2): 418-23.

[165] Yang H, He M. Distribution and speciation of selenium, antimony, and arsenic in soils and sediments around the area of Xikuangshan (China). *CLEAN–Soil, Air, Water*. 2016 Nov; 44(11): 1538-46.

[166] Kristan U, Kanduč T, Osterc A, Šlejkovec Z, Ramšak A, Stibilj V. Assessment of pollution level using *Mytilus galloprovincialis* as a bioindicator species: The case of the Gulf of Trieste. *Marine pollution bulletin*. 2014 Dec 15; 89(1-2):455-63.

[167] Rivera N, Hesterberg D, Kaur N, Duckworth OW. Chemical speciation of potentially toxic trace metals in coal fly ash associated with the Kingston fly ash spill. *Energy & Fuels*. 2017 Sep 21; 31(9): 9652-9.

[168] Cui JL, Zhao YP, Li JS, Beiyuan JZ, Tsang DC, Poon CS, Chan TS, Wang WX, Li XD. Speciation, mobilization, and bioaccessibility of arsenic in geogenic soil profile from Hong Kong. *Environmental pollution*. 2018 Jan 1; 232:375-84.

[169] Whaley-Martin KJ, Koch I, Reimer KJ. Determination of arsenic species in edible periwinkles (*Littorina littorea*) by HPLC–ICPMS and XAS along a contamination gradient. *Science of the total environment*. 2013 Jul 1; 456:148-53.

[170] Madsen AD, Goessler W, Pedersen SN, Francesconi KA.

Characterization of an algal extract by HPLC-ICP-MS and LC-electrospray MS for use in arsenosugar speciation studies. *Journal of Analytical Atomic Spectrometry*. 2000; 15(6): 657-62.

[171] Maher WA, Ellwood MJ, Krikowa F, Raber G, Foster S. Measurement of arsenic species in environmental, biological fluids and food samples by HPLC-ICPMS and HPLC-HG-AFS. *Journal of Analytical Atomic Spectrometry*. 2015; 30(10): 2129-83.