

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

185,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.  
For more information visit [www.intechopen.com](http://www.intechopen.com)



# The Effect of Neonicotinoid Insecticides on the Structure and Stability of Bio-Macromolecules

*Valéria Verebová and Jana Staničová*

## Abstract

Insecticides are among the most widely used pesticides in the world. They are preparations of chemical and biological origin used to control insects, which means its killing or preventing its destructive activity. Majority are used in forestry, agriculture, and households. Neonicotinoids represent the class of insecticides that is most frequently used in the world and replaced by more dangerous pyrethroids, organophosphates, and carbamates. In recent years, the focus has been mainly on the ecological and environmental risks caused by the use of neonicotinoids. These insecticides pose a very high risk to bees and also to soil and aquatic organisms. It is therefore highly topical to address the impact of neonicotinoids on biological systems on individual bio-macromolecules (DNA and serum albumins). Monitoring the impact of neonicotinoids on the structure and stability of biological macromolecules may contribute to reducing the use of these insecticides, as well as to considering and adjusting the tolerances of insecticides and their residues in food.

**Keywords:** insecticide, neonicotinoids, DNA, serum albumin, structure, stability

## 1. Introduction

The most widely applied pesticides in common practice are insecticides. They are preparations of chemical and biological origin used to control insects, which means their killing or preventing its destructive activity (**Table 1**).

Insecticides are mostly used in forestry, agriculture, and households. The use of insecticides has increased agricultural productivity, quantity, quality, and prolonged the lifetime of food and fodder plants. Insecticides have also revolutionized the fight against endemic diseases in developing countries. Unfortunately, these compounds pose the risk to humans and animals due to the presence of their residues in the food [4]. Insecticides are divided according to their mode of action into the ovicides (destroy eggs), larvicides (destroy larvae), and imagocides (destroy adult insects) [5]. The effect of insecticides is either immediate or slow acting and the insects die after a longer period of time [6].

### 1.1 Neonicotinoids

Neonicotinoids are a class of insecticides that belong to the most widely used in the world [7] because they allow for a rational approach to agricultural pest control [8, 9].

Insecticides class	Mechanism of action on pests	
Organochlorines	Dichlorodiphenyl-trichloroethane (DDT)	Destroying the delicate balance of Na <sup>+</sup> and K <sup>+</sup> ions in the axons of the neuron, which prevents the normal transmission of nerve impulses
	Hexachloro-cyclohexane (HCH)	Similar to DDT, only the effect is faster
	Cyclodienes	Act on the gamma-aminobutyric (GABA) acid receptor, which causes increased permeability of neurons to Cl <sup>-</sup> ions
	Polychloroterpenes	Similar to cyclodienes
Organophosphates	Inhibition of nervous system enzymes, resulting in the accumulation of acetylcholine at neuron–neuron or neuron-muscle junctions	
Carbamates	Inhibition of aliesterase, which promotes hydrolysis of aliphatic ester bonds	
Formamidines	Inhibition of monoamine oxidase, which is responsible for the degradation of the neurotransmitters norepinephrine and serotonin	
Dinitrophenols	Inhibition of oxidative phosphorylation	
Pyrethroids	Influencing the peripheral and central nervous system	
Nicotinoids	Postsynaptic acetylcholine receptor blockade	
Neonicotinoids	Strong bond to nicotinic acetylcholine receptors in the central nervous system	

**Table 1.**  
*The major classes of insecticides and mechanism of action on pests [1–3].*

Such insecticides tend to be referred to as “bio-rational” [10]. They replaced by more dangerous pyrethroids, organophosphates, and carbamates [11]. Discovery of imidacloprid and its subsequent introduction to the market in 1991 ushered in the era of neonicotinoids [12], which in 2014 accounted for more than 25% of the world’s insecticide market [13]. We know the neonicotinoids of four generations. The first generation consists of chloropyridyls, which include imidacloprid, acetamiprid, thiacloprid, nitenpyram, cycloxaprid and paichongding. These are divided into three classes according to pharmacophore groups: N-nitroimine, N-cyanoimine, and nitromethylene [14]. Chlorothiazoles, as thiamethoxam, imidaclothiz and clothianidin, form the second-generation of neonicotinoids. Furanyls (e.g., dinotefuran) belong to the third generation. Fourth generation is made up of sulfoximines, such as sulfoxaflor [7, 15–18]. Neonicotinoids are used on all types of crops in temperate and tropical regions as well as in forestry, gardens, urban parks, and as veterinary control products ectoparasites in domestic animals [19]. The flexibility of their use is due to their system properties, which allow their application in the form of direct sprays on crops, soil granules or seed coating [20]. They have unique biological and chemical properties such as broad-spectrum insecticidal activity, low application rates and mode of action [21]. Due to their octanol/water partition coefficient and dissociation constant (pKa) values, they readily enter plant tissues and translocate to all parts of the plant regardless of the method of application. This will automatically become toxic to any insect and potentially other organisms feeding on plants [22, 23]. However, some of their characteristics increase their negative impact on the environment and non-target organisms [23, 24]. Recent studies show that neonicotinoids are already ubiquitous in the environment because of their versatile use, high mobility and relatively long half-life in water and soil [25, 26]. In the USA, for example, their presence was confirmed in twelve out of nineteen different

fruits monitored and vegetables, eleven of which contained more than two neonicotinoids. Value of thiamethoxam even exceeded the maximum residue limit. The most frequent occurrence was paradoxically found in foods primarily intended for infants and toddlers (prevalence ranging 6–31%) [27]. Neonicotinoid contamination has also been demonstrated in drinking water [28, 29], vegetables and fruit [30], milk [31], and in honey [32].

### *1.1.1 Mechanism of neonicotinoid action*

Neonicotinoid's mode of action is by blocking the nicotinacetylcholine receptor (nAChR) leading to paralysis and death of the pests [33, 34]. nAChR is an ion channel responsible for immediate neurotransmission and belongs to the group of neurotransmitter ion channels along with gamma-aminobutyric acid, glycine, 5-HT<sub>3</sub> and serotonin receptors. It consists of ten  $\alpha$ , four  $\beta$ ,  $\gamma$ , and  $\delta$  subunits, which combine to form three basic types of receptors (muscle, neuronal, and ganglionic) with different structures. Different combinations of subunits result in differences in sensitivity to acetylcholine and other pharmacological systems. The most potent nAChR agonist is the nicotinic derivative epibatidine [35].

Neonicotinoids contain a negatively charged, electronegative cyano or nitro group that reacts with the positively charged nAChR site of the insect. In vertebrates, this interaction is blocked by the protonation of nitrogen in their organism [35].

### *1.1.2 Neonicotinoid toxicity*

In recent years, the focus has been mainly on ecological and environmental risks caused by the use of neonicotinoids.

N-nitromine neonicotinoids show a very high risk to wild bees and honey bees [36]. Exposure to already low doses of insecticides occurs sublethal effects such as reduced immunity, disorientation and behavioral changes and reproduction of bees [37–40]. Based on these facts, the use of imidacloprid, thiamethoxam, clothianidin have been completely banned in the European Union since May 2018 [41].

N-cyanoimine insecticides in turn pose a high risk to soil and aquatic organisms [36, 42, 43]. Because of their long persistence in soil and high water solubility, they tend to pass into groundwater, surrounding rivers [44], lakes, and seas [45, 46]. Studies focusing on neonicotinoid content in watercourses have shown concentrations ranging from 0 to 380 ng/L depending on the region [47–49]. Because of adverse effect of neonicotinoids on aquatic organisms such as in food intake [50, 51], changes in swimming and nesting [52, 53], growth inhibition [54], changes in reproduction [53] and acute and chronic mortality [53, 55, 56], this contamination poses a major risk to aquatic ecosystem and for the supply water for its consumption [34, 57].

Several other studies of the adverse impact of neonicotinoids on non target organisms have demonstrated the development of immunosuppression in birds, bats, fish and amphibians [58]. Neonicotinoids can also adversely affect mammalian nAChRs, leading to neurobehavioral deficits and increased glial fibrillary acidic protein expression in the hippocampus [59–61], further affect the reproductive cycle, liver function and have genotoxic and neurological effects [11, 61–65].

Neonicotinoids are capable of disrupting the endocrine system. The study focused on exposure to thiacloprid in rats showed an increase in triiodothyronine and thyroxine hormones [66]. They are further associated with the development of oxidative stress [67], which in various cases leads to changes in the levels of ovarian

and antioxidant hormones [64], but also to increased germ cell apoptosis and DNA fragmentation in rat testes [68].


Next, we will discuss the most widely used representatives of first-generation neonicotinoid insecticides belonging to the group of N-cyanoimines (thiacloprid, acetamiprid) and N-nitroimines (imidacloprid).

### 1.1.3 Thiacloprid


Thiacloprid ((Z)-3-(6-chloro-3-pyridylmethyl)-1,3-thiazolidine-2-ylidenecyanamide) is an insecticide belonging to the class of neonicotinoids (**Figure 1**), which is used for protection of vegetables, orchards, tea, maize and oilseed rape seeds [69].

Due to quite good solubility of thiacloprid in water and its low potential in groundwater, contamination of mainly surface water bodies of water, but its presence has also been detected in drinking water [28]. Thiacloprid occurs as a white to yellow powder [70] and it is polar, slightly soluble in water (0.185 g/L) and in organic solvents: dimethyl sulfoxide (150 g/L), acetone (64 g/L), ethyl acetate (9.4 g/L), acetonitrile (52 g/L). Partition coefficient octanol/water has a  $\log P = 1.26$  (pH = 7, 20°C), indicating poor solubility in fats, low absorption and distribution in the body and its pH is in the range of 4–9 stable [71].

In general, thiacloprid shows higher toxicity to aquatic organisms compared to other neonicotinoids studied [35]. This toxicity is probably related to its resistance to degradation in water at neutral and acidic pH values. WHO classifies it as moderately hazardous (Class II) [72] while the Environmental Protection Agency in the United States (US EPA) characterizes it as a potential carcinogen, based on the occurrence of thyroid tumors in male rats and uterine and ovarian tumors in rats, and mice [73]. Thiacloprid is metabolized and excreted in the urine within 24 hours after oral administration. The target organ is primarily the liver; a toxic effect on the liver has also been observed in dog prostate [41, 74]. Other studies have shown that it causes fetal resorption, skeletal retardation, changes in motor activity in rats, thyroid adenomas, and uterine adenocarcinogens in mice [74]. In fish, after exposure to TCLs, there is growth retardation, delayed fetal development, and changes in antioxidant enzyme levels [75]. Moreover observed were reduced cell proliferation associated with higher levels of chromosomal aberrations in bovine lymphocytes [76] and genotoxic and cytotoxic effects on human [77] and bovine lymphocytes [78].



**Figure 1.**  
*Structural formula of thiacloprid.*



**Figure 2.**  
*Structural formula of acetamiprid.*



### 1.1.4 Acetamiprid


Acetamiprid ((E)-N1-[(6-chloro-3-pyridyl)methyl]-N2-cyano-N1-methylacetamidine) is a new broad-spectrum neonicotinoid (**Figure 2**) which is used to control sucking insects by interfering with insects' nervous system [79, 80]. It protects crops such as leafy vegetables, citrus fruits, pome fruits, grapes, cotton, cole crops, and ornamental plants. Acetamiprid plays a key role in commercial cherry farming due to its effectiveness against the larvae of the cherry fruit fly [20].

Acetamiprid occurs as a white crystalline substance, it is polar and water soluble (4.2 g/L), therefore it can be transported to surface waters and may be toxic to aquatic organisms and life. It is also soluble in organic solvents: acetone, chloroform and dichloromethane (200g/L) and not very frothy in hexane (0.005g/L). Partition coefficient octanol/water has a logP = 0.80 (pH = 7, 25°C) and its dissociation constant is pKa = 0.7 [81].

Acetamiprid contains a 6-chloro-pyridine motif in its molecule, as does the animal alkaloid epibatidine from the poisonous South American frog *Epipedobates tricolor* [82]. An important property of acetamiprid as a representative of N-cyanoimine neonicotinoids is that, unlike N-nitroimines such as clothianidin, dinotefuran, imidacloprid, thiamethoxam or nitenpyram, it is of little toxicity to bees [83]. It does not accumulate in soil, is mobile and rapidly degrades by aerobic mechanisms [9] and microorganisms are involved in its degradation [84]. Its half-life in soil ranges from <1 to 8.2 days [85] and its content in vegetables and fruits is low [86, 87]. The application of acetamiprid in greenhouses and agricultural farms is safe and not associated with major health risks [88]. Human poisonings are known only in cases where acetamiprid is used as a suicide agent. Two such cases of acute suicidal poisoning are described in the medical literature. In both cases, nausea and vomiting, muscle weakness, hypothermia, convulsions and other clinical manifestations, including tachycardia, hypotension, ECG changes, hypoxia and thirst, occurred. These symptoms are partly similar to acute organophosphate intoxication. Supportive treatment of the clinical symptoms was sufficient and both patients were discharged without complications 2 days after ingestion of acetamiprid [89]. It is not yet sufficiently clear whether acetamiprid is genotoxic to mammals [90].

### 1.1.5 Imidacloprid

Imidacloprid (N-{1-[6-chloro-3-pyridyl)methyl]-4,5-dihydroimidazol-2-yl} nitramide) is a systemic insecticide (**Figure 3**) that acts as an insect neurotoxin, used for pest control in agriculture. In the year 1999, it was the most widely applied insecticide in the world [91]. Imidacloprid can be used by soil injection, broadcast foliar, application to the skin of the plant, tree injection, ground application as a liquid or granular formulation, or as a pesticide-coated seed treatment [92]. It is extra effective against sucking insects and mining pests such as mealybugs, aphids, thrips, and rice leafhoppers and against whitefly [93].



**Figure 3.**  
Structural formula of imidacloprid.

Imidacloprid is a colorless crystalline substance with slight characteristic odor. It is weakly soluble in water (0.61 g/L) but better soluble in organic solvents: dichloromethane (67 g/L), isopropanol (23 g/L) and toluene (0.69 g/l). Partition coefficient octanol/water has a  $\log P = 0.57$  ( $\text{pH} = 7$ ,  $21^\circ\text{C}$ ) and its dissociation constants are  $\text{pK}_{\text{a}1} = 1.56$  and  $\text{pK}_{\text{a}2} = 11.12$  [94].

Imidacloprid has been classified by WHO as moderately hazardous (Class-II) based on animal studies [95]. The use of imidacloprid has a devastating impact on biodiversity, in particular on rivers and watercourses, not only on crustaceans [96], mollusks [97] and non-target species (insects), but also on soil organisms [36], as well as on bird populations [98]. Acute intoxication by imidacloprid and its metabolites resulted in the fast appearance of neurotoxicity symptoms, such as hyperactivity, hyperresponsiveness and trembling and led to hypoactivity and hyporesponsiveness [99]. It has been mentioned having harmful effects (oral toxicity) on honeybees in fields [100]. There is known case of imidacloprid poisoning with suicidal intent that developed various manifestations including hypokalaemia, central nervous system depression, respiratory arrest and paroxysmal atrial fibrillation [101]. Imidacloprid is metabolized by photodegradation from soil surface and water [102]. It is proven that plant metabolites of imidacloprid, the imidazolidine derivative, the olefin metabolite and nitroso-derivative were more toxic to aphids than imidacloprid itself [103].

## **2. Interaction of neonicotinoids with bio-macromolecules**

Monitoring the interaction of neonicotinoid insecticides with biological macromolecules (DNA and serum albumins), determining the binding constant and the interaction mode provides a more comprehensive view of their distribution, toxicity and metabolism in the organism. It is also of great importance in adjusting the tolerance of these insecticides and their residues in food. The presented summarized results could contribute positively to the reduction of the use of the group of insecticides discussed by us.

### **2.1 Methods and analysis**

The interaction of neonicotinoids with bio-macromolecules can be studied most often using UV-Vis, fluorescence and IR spectroscopy, circular dichroism, monitoring of bio-macromolecules melting, viscosity assays and last but not least molecular docking.

#### *2.1.1 UV-Vis measurements*

UV-Vis spectroscopy is a very effective method for investigating structural changes in the formation of DNA or protein and ligand/DNA or ligand/protein complexes. These measurements have been made on absorption spectrophotometer using quartz cuvettes of 1 cm path length at laboratory temperature. Usually the concentration of DNA or protein is constant while the concentration of neonicotinoid varies. A common practice is to add the same concentration of insecticide to the reference sample (with respect to the absorption maximum of the neonicotinoid). By monitoring the changes in absorbance intensity and the shift of the DNA/protein absorption maximum, it can be predicted how the neonicotinoid binds to the DNA/protein structure.

#### *2.1.2 Fluorescence measurements*

Fluorescence measurements were performed on spectrofluorimeter in a 1 cm quartz cuvette. Upon interaction, an extinguishing mechanism can be noted.

In general, extinguishing can be qualified as dynamic and static. The quenching mechanism is accurately described by the Stern-Volmer Eq. (1)

$$\frac{F_0}{F} = 1 + K_q \tau_0 = 1 + K_{sv} [Q] \quad (1)$$

where  $F_0$  and  $F$  are the fluorescence intensities in the absence and in the presence of the quencher (Q) respectively.  $K_q$  and  $K_{sv}$  are the bio-macromolecule quenching rate constant, and Stern-Volmer constant respectively.  $\tau_0$  is the average lifetime of the bio-macromolecule without quencher and  $[Q]$  is the concentration of quencher [104, 105].

Subsequently the binding parameters such as binding constant ( $K_a$ ) and the number of binding sites ( $n$ ) can be calculated according to Hill Eq. (2).

$$\log \left[ \frac{F_0 - F}{F} \right] = \log K_a + n \log [Q] \quad (2)$$

The number of binding sites and association constant have been obtained by the plot of  $\log[(F_0 - F)/F]$  versus  $\log[Q]$  [106–108].

### 2.1.3 Melting studies

The values of thermodynamic parameters can be determined from the Van't Hoff equation. This equation has been used in the form (3) for the interaction of neonicotinoid insecticides with proteins

$$\ln K_a = \frac{-\Delta H}{RT} + \frac{\Delta S}{R} \quad (3)$$

while  $\Delta H$  is the enthalpy change,  $\Delta S$  the entropy change,  $T$  is the temperature, and  $R$  is the universal gas constant. The binding studies from fluorescence measurements were usually realized at three different temperatures. By plotting  $\ln K_a$  versus  $1/T$  it can be possible to determine the values of  $\Delta H$  and  $\Delta S$ . Gibbs free energy change  $\Delta G$  could be calculated using Eq. (4) [109, 110].

$$\Delta G = \Delta H - T\Delta S = -RT \ln K_a \quad (4)$$

The thermal stability of DNA was studied by an absorption spectrophotometer with a Peltier module. A special type of Van't Hoff Eq. (5) has fitted melting curves

$$A = A_{\min} + \frac{A_{\max} - A_{\min}}{1 + e^{\left[ \frac{\Delta H}{R} \times \left( \frac{1}{T} - \frac{1}{T_m} \right) \right]}} \quad (5)$$

where  $A$  is the absorbance,  $A_{\min}$ ,  $A_{\max}$  are the minimal and maximal measured absorbance respectively,  $H$  is the enthalpy of transition,  $T$ ,  $T_m$  are actual and melting temperature [111].



### 2.1.4 CD studies

CD spectroscopy was used to record changes in the secondary structure of optically active substances, including proteins and DNA. The CD results of complexes neonicotinoid/protein have been expressed in terms of mean residue ellipticity (MRE) (6)

$$MRE = \frac{\text{Observed CD (mdeg)}}{10 c n l} \quad (6)$$

where  $c$  is the concentration of protein,  $n$  is the number of amino acid residues of protein, and  $l$  is the path length. The  $\alpha$ -helix contents of free protein and neonicotinoid/protein complexes can be calculated from MRE values at 208 nm using the following Eq. (7) [112].

$$\alpha - \text{helix}(\%) = \frac{-MRE_{208nm} - 4,000}{33,000 - 4,000} \times 100 \quad (7)$$

CD spectra of DNA are very sensitive to the interaction mode of DNA with small molecules like neonicotinoid insecticides [113]. The characteristic spectrum of DNA in the B-form helix is characterized by a negative band at 245 nm resulting from right-handed helicity and a positive band at 278 nm resulting from base stacking [114]. By forming a neonicotinoid/DNA complex, a decrease in the intensity of the negative band at 245 nm was usually observed with accompanying red shift, while the intensity of the positive band at 276 nm increased. This increase in intensity was associated with a slight red shift. These changes of CD spectra represent the transformation from a B-form DNA structure to an A-conformation [115].

### 2.1.5 FT-IR spectra analysis

Infrared spectra have been collected approximately after 2h incubation of solutions containing neonicotinoid insecticides and bio-macromolecule at various ratios. The amount of DNA or protein was constant. The intensity and shifting changes in infrared spectra could be applied to accurately specify the site of incorporation of neonicotinoids into the structure of bio-macromolecules (DNA and proteins) [116].

### 2.1.6 Viscosity tests

Viscosity tests can provide further information about the nature of the interaction neonicotinoid insecticides and DNA. This type of experiment consists in measuring the relative viscosity ( $t/t_0$ ) of ethidium bromide/DNA and neonicotinoid/DNA complexes. A significant increase in viscosity was detected during the interaction of a typical intercalator ethidium bromide and DNA. The task was to compare the changes in DNA viscosity after adding neonicotinoids to the solution. If the same increase in viscosity was observed as for ethidium bromide, then the intercalation mode of binding between insecticides and DNA could be predicted [117].

### 2.1.7 Molecular docking studies

The trend of today's research is to confirm the results obtained from experiments by molecular docking studies. Serum albumin and DNA crystallographic

data can be used from the Brookhaven Protein Data Bank. The species selected were 1AO6 (HSA), 4F5S (BSA), and 453D (DNA) [116, 118]. The molecular structures of neonicotinoid insecticides could be taken from The PubChem database. Most often AutoDockVina has been used for molecular docking and molecular simulation of the interaction between neonicotinoids and bio-macromolecules [119].

## 2.2 Interaction of neonicotinoids with DNA

Neonicotinoid insecticides are highly reactive compounds that form complexes with a variety of cellular biomolecules, including DNA. DNA plays an important role in cell proliferation, protein synthesis and genetic transcription. It is necessary to protect DNA from the harmful effects of insecticides because they will damage the genetic structure of cells and disrupt metabolic processes [120]. The study of the interaction between insecticides and nucleic acids is considered to be important to allow screening for carcinogenic properties of pesticides, especially insecticides [121, 122]. Studies addressing the specific mode of interaction and binding sites of neonicotinoid insecticides with DNA are few. Several techniques have been used to study the binding properties between neonicotinoid insecticides and DNA, including UV-visible absorption, fluorescence, circular dichroism, Fourier transform infrared spectroscopy, coupled with DNA melting investigations and viscosity measurements in physiological buffer. To predict the possible binding site and binding mode was used the molecular docking study [120–122].

Neonicotinoid insecticides belong to small molecules known to bind to the double helix of DNA by two dominant modes, especially such as groove binding and intercalation. Groove binding presents docking the thin ribbon-like molecules in the DNA minor groove, in close proximity to the sugar-phosphate backbone. Conversely, their intercalation into the helix includes the insertion of a drug – usually a planar aromatic cation – into the base reservoir of the helix [123].

It has been demonstrated that thiacloprid interacts with DNA, which influences on the length and denaturation of DNA. The binding strength of thiacloprid and DNA is expressed by a binding constant  $K_a$  whose value is  $9.3 \cdot 10^3$  L/mol at laboratory temperature [117]. The presence of DNA significantly affects the emission spectrum of thiacloprid. The quenching of its fluorescence intensity was analyzed using the Stern-Volmer method [114]. The quenching constant  $K_{SV}$  was determined to be  $2.8 \cdot 10^4$  L/mol and correlation coefficient for  $K_{SV}$  was  $R = 0.998$ . Not considerable changes in viscosity behavior were observed as a result of its addition to the DNA solution. The presence of thiacloprid causes a decrease in melting temperature  $T_m$  and Van't Hoff enthalpy  $\Delta H$  of DNA. These changes in thermodynamic parameters lead to destabilization of the DNA molecule. In addition, it was recorded a two-phase character melting curve of the complex DNA-thiacloprid with an expression destabilization of AT regions in DNA [117]. It is generally assumed small molecules that consist of at least two aromatic rings coupled by a no rigid bond enabling their torsional flexibility, like thiacloprid bind preferentially to the minor groove of DNA, specially to the regions rich in AT-base pairs [124, 125]. Considering all the above results, it can be concluded that thiacloprid does not interact with DNA via an intercalating binding mode. Increasing of the length and influencing of DNA denaturation points to the groove-binding mode of interaction. Probably incorporation of this insecticide occurs into DNA minor groove by hydrophobic or hydrogen bonds [117].

Similarly, the effect of acetamiprid on DNA structure and stability was investigated. As shown by the UV-visible spectra, after the addition of acetamiprid to DNA solution, the absorption peak of DNA at 260 nm (associated with strong purine and pyrimidine base absorption in DNA [126]) increased markedly and there was

significant blue shift. The fluorescence emission spectra of acetamiprid upon addition of DNA revealed that increasing concentration of DNA gradually decreased fluorescence emission peak of acetamiprid and a new band developed at approximately 370 nm. These facts can be attributed to the formation of the acetamiprid-DNA complex. The quenching mechanism between acetamiprid and DNA was analyzed by use of the Stern–Volmer method. The obtained  $K_{SV}$  value was  $1.86 \cdot 10^4$  L/mol and the relevant correlation coefficient for  $K_{SV}$  was  $R = 0.997$ . The binding constant value  $K_a$  was determined  $5.27 \cdot 10^3$  L/mol. Both constants were found at laboratory temperature [116]. Small molecules intercalating into the DNA causes stabilization of base stacking and leads to significant increase of  $T_m$ , whereas non-intercalation binding leads to no obvious increase in melting temperature [127]. In acetamiprid binding,  $T_m$  DNA increased by approximately 3°C. It can therefore be deduced that acetamiprid interacted with DNA via no classic rather than classical intercalation. This result was supported by evidence obtained from molecular docking, namely, the intercalation of acetamiprid into the double helix DNA from one side. Measurement of the relative viscosity of DNA shows a significant decrease in its viscosity, which excludes the classical intercalation mode of binding. The model in which acetamiprid binds to DNA in a partial intercalation manner explained this phenomenon. This means that partial intercalation can occur where acetamiprid can act as a ‘wedge’ that pushes apart one side of the base pair stack (but in contrast to the classical intercalation model, does not completely separate the stack), and thus induce static bending or kinking in the double helix [128]. Thermodynamic data for the interaction of acetamiprid and DNA were calculated. The Gibbs free energy value  $\Delta G$  has a negative sign, indicating that the process is happening spontaneously [116]. Positive values of enthalpy  $\Delta H$  and entropy  $\Delta S$  changes suggest that hydrophobic interactions play an important function in the formation of the bond between acetamiprid and DNA and stabilize the complex [129]. Significant changes, in position and intensity, were observed in FTIR spectra for the GC base pair than for the AT base pair. It can be predicted that the specific binding site of acetamiprid to DNA is a site rich in GC base pairs. In addition, it is confirmed that acetamiprid binding leads to a change in the secondary structure of DNA from the B conformation to the A [116].

The interaction of acetamiprid with DNA is the basis for the development of the DNA probe, which is used in practice to detect the presence of acetamiprid in environmental samples and agricultural products. The results of the detection of acetamiprid using the DNA probe are in almost complete agreement with the results obtained using the HPLC technique [130].

At present, the interaction of imidacloprid with DNA is poorly studied, as its mode of incorporation into DNA, exact binding site, and binding constant are not known. Nevertheless, it is proven that imidacloprid can induce oxidative stress and DNA damage in zebrafish [131] and bees [132]. The study of the interaction between imidacloprid (also other neonicotinoid insecticides) and DNA is important for us to understand the insecticidal mechanism of neonicotinoids and their side effects, such as carcinogenesis, teratogenesis and mutagenesis [133]. Therefore, the mechanism of the reaction between imidacloprid and DNA is necessary to examine in detail.

### **2.3 Interaction of neonicotinoids with serum albumins**

The binding study of neonicotinoids with proteins has toxicological importance [134]. The results of these interactions may cast some light on the future study of the interaction between neonicotinoid insecticides and other proteins such as enzymes and have toxicological to ecotoxicology importance. Therefore, it is essential to investigate the effect of neonicotinoids on the structural and optical properties of serum albumins, especially human serum albumin (HSA), the thermodynamic aspects in



the binding process, and characters of the binding sites. The binding of insecticides, but also pesticides in general, to proteins has been exploited in the construction of pesticide biosensors. Biosensor assays can bring measures of the toxic effects of chemicals on the target organism and of the molecular mechanisms that are the basis of toxicity [135]. Another important direction in studies of the biological properties of neonicotinoids is directly related to protection of the health of humans and agricultural animals, and includes study of the interactions between these compounds and proteins, enzymes, and receptors in blood plasma and tissues. The structure of neonicotinoid insecticides contains sections and groups capable of forming electrostatic, hydrophobic, and hydrogen bonds as well as other types of bonds typical of endogenous ligands in their complexes with proteins. It is the reason many insecticides can play the important role of exogenous ligands and change their own properties within the composition of protein complexes, as is true for natural bio-regulators such as hormones. These alternations can involve metabolic parameters and biological effects of the pesticides in the human and animal bodies. The mechanism for the interaction of neonicotinoid with serum albumin probably includes “recognition” and initial binding of the ligand because of its polar group. Followed by adaptation of the ligand-binding site of the HSA molecule for binding by means of conformational transitions. This mechanism is terminated by subsequent interaction of the hydrophobic core of the ligand with the nonpolar side chains of serum albumin in the cavity of the binding site. The new insights into the interaction of serum albumins and neonicotinoid insecticides, about the forms in which neonicotinoids exist in the body and the approaches developed for studying interactions of these compounds with the major transport protein HSA are a necessary basis for estimating the biological effects of pesticides in this class when they enter human blood [136].

It has been shown that thiacloprid interacts with HSA, its binding properties have been characterized at the molecular level under physiological conditions. The intensity of HSA absorption maximum decreased after the addition of thiacloprid, and its little red shift occurred at the same time. This indicated the probable formation of a complex between thiacloprid and HSA. With a gradual increase in thiacloprid concentration, a period decrease in fluorescence intensity was also observed. The Stern-Volmer plots were used for analyses of the quenching mechanism. The obtained  $K_{SV}$  value was  $3.304 \cdot 10^4$  L/mol and the binding constant  $K_a$  was found  $3.07 \cdot 10^4$  L/mol. Both constants were determined at laboratory temperature. The linear Van't Hoff equation was applied to track changes in the thermodynamic parameters. The entropy, enthalpy and Gibbs free energy values indicate that the coupling of thiacloprid to HSA is an exothermic process due to the positive value of  $\Delta S$  and negative values of  $\Delta H$  and  $\Delta G$  [134]. It is well known that thermodynamic parameters play an important role in determining the type of binding by which a ligand (insecticide) binds to an HSA. Positive value of  $\Delta H$  and  $\Delta S$  is the result of hydrogen bonding. Negative values of these quantities ( $\Delta H$ ,  $\Delta S$ ) are corresponded with hydrogen bonds and van der Waals interaction in a low dielectric solution. The electrostatic interaction in aqueous solution between ionic species is associated with a positive change of  $\Delta S$  and very small negative change of  $\Delta H$ , almost zero [137]. Hydrophobic force and electrostatic force interactions are characterized by negative value of  $\Delta H$  and positive  $\Delta S$  values. In view of the above facts, it is not easy to interpret the results obtained from the thermodynamic analysis of the thiacloprid-HSA interaction. It is difficult to explain this interaction mode by a single intermolecular force. It has been published that the interaction force acting between small molecules and proteins is generally not just a single force. Probably may be there are variety of forces existing in the interaction forces which contain the electrostatic force. The hydrophobic molecule embedded in the internal hydrophobic region of proteins can be responsible for the fluorescence quenching [138, 139]. One

can hypothesize that the incorporation of thiacloprid into HSA can be realized by hydrophobic interaction, as evidenced by the positive  $\Delta S$  value, but it cannot be excluded the influence of electrostatic interaction. The given conclusions are consistent with molecular modeling. This suggests that thiacloprid could be located on the surface of the binding pocket of subdomain IIA in the HSA molecule. It also confirms that the hydrogen bonding also plays a significant role [134]. Study of HSA secondary structure revealed the decrease of the percentage  $\alpha$ -helix structure the effect of thiacloprid binding on the amino acid residue of the main polypeptide chain of HSA. Thiacloprid upset their hydrogen bonds [140].

Hemoglobin is essential protein in the blood plasma. It is working as a transporter of oxygen. Bovine hemoglobin (BHb) is used as a model protein to study the binding properties of drugs and insecticides [141]. The results obtained from the study of the influence of thiacloprid on this model protein show that their interaction is a static process. Two types of bonds play an important role in this binding, namely, hydrogen bonding and hydrophobic interactions. Increasing concentration of thiacloprid causes a marked decrease in BHb fluorescence intensity. The strength of cross-linking is characterized by the binding constant  $K_a$ , the magnitude of which was determined to be  $8.04 \cdot 10^4$  L/mol at laboratory temperature. Tracking changes in thermodynamic parameters suggests that hydrogen bonding forces are most important for a given interaction. This is evident from the fact that negative values of the enthalpy change  $\Delta H$  and entropy change  $\Delta S$  were calculated. Changes in the secondary structure of BHb in the presence of thiacloprid show that there is a 3.7% decrease in  $\alpha$ -helix content. Molecular modeling was used to determine the amino acid residues involved in thiacloprid-BHb binding. The thiacloprid pyridine ring interacts with Leu105, Pro95, Trp37 by hydrophobic interactions. The incorporation of thiacloprid into BHb is not exclusively hydrophobic, as several ionic and polar residues (Thr137, Tyr35) are present near the bound ligand. These polar residues stabilize the neonicotinoid insecticide via H-bonding and electrostatic interactions. Trp37 is involved in the formation of H-bonds with side chain imino groups [142].

Similarly, the effect of imidacloprid on HSA structure was investigated. The absorbance of the imidacloprid-HSA complex decreased with increasing imidacloprid concentration. A shift of the absorption maximum to the red region was also observed as in the thiacloprid interaction [135]. The HSA molecule contains 585 amino acid residues forming a single polypeptide of known sequence [118]. The protein response to conformational transitions, subunit association, ligand binding, or denaturation are changes in tryptophan emission spectra [143]. The study of the intrinsic fluorescence of Trp HSA is important for a better understanding of the specific changes that occur in the macromolecule [135]. Tyrosine fluorescence in HSA is mostly quenched due to the presence of nearby amino acids or efficient energy transfer from tyrosine fluorescence to Trp214 [144]. Imidacloprid caused a decrease in the fluorescence intensity of tryptophan residues. Fluorescence resonance energy transfer determined the distance between the donor (HSA) and acceptor (imidacloprid). Its size is 2.10 nm [135]. It is generally accepted that the average distances between the donor fluorophore and the acceptor fluorophore are 2–8 nm and indicate that energy transfer from HSA to imidacloprid occurs with a high probability [145]. It represents static quenching. During the measurement, a change in the structure of the surroundings of the Trp and Tyr residues was observed. Imidacloprid affected the physiological function of HSA. The fluorescence maximum of HSA shifted to higher wavelengths. The above results suggest that imidacloprid formed a specific bond in subdomain IIA, close to the tryptophan residue at position 214 of the HSA polypeptide chain like thiacloprid [135, 136]. The binding constant  $K_a$  was determined to the value  $1.51 \cdot 10^4$  L/mol [135]. Thermodynamic parameters allowed predicting that



imidacloprid incorporates into HSA via hydrophobic interactions. These occur in the presence of an aromatic pyridine imidacloprid ring. Its nitroimine group could provide further enhancement of the affinity of the above neonicotinoid insecticide for HSA [135, 136].

A study looking at the interaction of imidacloprid with bovine serum albumin (BSA) gave the same results as in the interaction with HSA. From the fluorescence quenching spectra, the binding constant  $K_a$  was calculated, the magnitude of which reaches a value of  $3.42 \cdot 10^4$  L/mol. Static quenching of emission without energy transfer by radiation within a single BSA molecule was also detected. Variations in thermodynamic parameters ( $\Delta H > 0$ ,  $\Delta G < 0$ ,  $\Delta S > 0$ ) made it possible to predict the type of interaction. It is probably a hydrophobic interaction. Among the effects, that imidacloprid has on BSA can include increasing the polarity of the microenvironment in which Trp and Tyr are found, increasing the compaction of peptide bonds, and modifying the conformation of BSA [146].

The effect of acetamiprid on the structure of serum albumin is currently not sufficiently studied (not enough relevant results have been published). Considering the above results obtained by studying the interaction of similar neonicotinoid insecticides (thiacloprid and imidacloprid), it can be assumed that acetamiprid also affects the structure and conformation of serum albumins as well as their thermodynamic parameters. Therefore, it is essential to study and characterize in detail the incorporation of this insecticide into albumin.

### 3. Conclusion

In conclusion, we have to state that the topic discussed by us is still insufficiently studied and it is necessary to further continuously address the issue of the interaction of neonicotinoid insecticides with bio-macromolecules. Using of neonicotinoid insecticides leads to serious environmental problems, including contamination of soil and ground water, which can cause them to accumulate in the human and animal bodies and subsequently damage DNA. Several studies published in this chapter show that molecules of thiacloprid, acetamiprid, and imidacloprid are able to incorporate into important biological macromolecules and disturb their structure and function. All of them are honey bee killers and harmful to pollinators. Only thiacloprid is allowed to use in EU, imidacloprid and acetamiprid are prohibited which are considered to be one of the causes of bee colony decline in the world [41, 147]. On the other hand, they can still stay in the US market by Environmental Protection Agency decision [148].

However, the published results clearly indicate that neonicotinoid insecticides such as thiacloprid, acetamiprid, and imidacloprid, should be used very sparingly and cautiously in practice, especially in densely populated countries of the world where insecticides are overused.

### Acknowledgements

This work was also supported with grant from the Slovak Research Grant Agency VEGA No. 1/0242/19.

### Conflict of interest

The authors declare no conflict of interest among themselves.

IntechOpen

### **Author details**

Valéria Verebová<sup>1\*</sup> and Jana Staničová<sup>1,2</sup>

1 Department of Chemistry, Biochemistry and Biophysics, University of Veterinary Medicine and Pharmacy, Košice, Slovakia

2 First Faculty of Medicine, Charles University in Prague, Prague, Czech Republic

\*Address all correspondence to: [valeria.verebova@uvlf.sk](mailto:valeria.verebova@uvlf.sk)

### **IntechOpen**

© 2021 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. 

## References

- [1] Metcalf RL, Luckmann WH. Introduction to insect pest management. John Willey & Sons; 1994. 650 p.
- [2] Grundlingh J, Dargan PI, El-Zanfaly M, Wood DM. 2,4-dinitrophenol (DNP): a weight loss agent with significant acute toxicity and risk of death. *Journal of Medical Toxicology*. 2011;7(3):205-212. DOI: 10.1007/s13181-011-0162-6
- [3] Radcliffe ED, Hutchison WD, Cancelado RE. Integrated Pest Management. Cambridge University Press, Cambridge UK; 2009. 530 p.
- [4] Elbert A, Buchholz A, Ebbinghaus-Kintscher U, Erdelen C, Nauen R, Schnorbach HJ. The biological profile of thiacloprid-A new chloronicotinyl insecticide. *Pflanzenschutz- Nachrichten Bayer*. 2001;54:185-208.
- [5] Van Emden HF, Peakall DB. Beyond Silent Spring. Integrated pest management and chemical safety. Springer; 1996; XVIII: 320 p.
- [6] Sánchez-Bayo F, Tennekes HA, Goka K. Impact of Systemic Insecticides on Organisms and Ecosystems, Insecticides. 2013; IntechOpen. DOI: 10.5772/52831. Available from: <https://www.intechopen.com/books/insecticides-development-of-safer-and-more-effective-technologies/impact-of-systemic-insecticides-on-organisms-and-ecosystems> [Accessed: 2021-06-15] DOI: 10.5772/52831
- [7] Simon-Delso N, Amaral-Rogers V, Belzunces LP, Bonmatin JM, Chagnon M, Downs C, Furlan L, Gibbons DW, Giorio C, Girolami V, Goulson D, Kreutzweiser DP, Krupke CH, Liess M, Long E, McField M, Mineau P, Mitchell EAD, Morrissey CA, Noome DA, Pisa L, Settele J, Stark JD, Tapparo A, Van Dyck H, Van Praagh J, Van der Sluijs JP, Whitehorn PR, Wiemers M. Systemic insecticides (neonicotinoids and fipronil): trends, uses, mode of action and metabolites. *Environmental Science and Pollution Research*. 2015;22(1):5-34. DOI: 10.1007/s11356-014-3470-y
- [8] Ishaaya I, Kontsedalov S, Mazirov D, Horowitz AR. Biorational agents--mechanism and importance in IPM and IRM programs for controlling agricultural pests. *Mededelingen Rijksuniversiteit te Gent Fakulteit van de Landbouwkundige en Toegepaste Biologische Wetenschappen*. 2001;66(2a):363-374
- [9] Foster SP, Denholm I, Thompson R. Variation in response to neonicotinoid insecticides in peach-potato aphids, *Myzus persicae* (Hemiptera: Aphididae). *Pest Management Science*. 2003;59(2): 166-173. DOI: 10.1002/ps.570
- [10] Ishaaya I, Kontsedalov S, Horowitz AR. Biorational insecticides: mechanism and cross-resistance. *Archives of Insect Biochemistry and Physiology*. 2005;58(4):192-129. Available from: Biorational insecticides: Mechanism and cross-resistance (wiley.com) [Accessed: 2020-03-12]
- [11] Han W, Tian Y, Shen X. Human exposure to neonicotinoid insecticides and the evaluation of their potential toxicity: An overview. *Chemosphere*. 2018;192:59-65. DOI: 10.1016/j.chemosphere.2017.10.149
- [12] Tomizawa M, Casida JE. Neonicotinoid insecticides: highlights of symposium on strategic molecular designs. *Journal of Agricultural and Food Chemistry*. 2010;59(7):2883-2886. DOI: 10.1021/jf103856c
- [13] Bass Ch, Denholm I, Williamson MS, Nauen R. The global status of insect resistance to

neonicotinoid insecticides. *Pesticide Biochemistry and Physiology*. 2015;**121**:78-87. DOI: 10.1016/j.pestbp.2015.04.004

[14] Casida JE. Neonicotinoid metabolism: compounds, substituents, pathways, enzymes, organisms, and relevance. *Journal of Agricultural and Food Chemistry*. 2010;**59**(7):2923-2931. DOI: 10.1021/jf102438c

[15] Elbert A, Overbeck H, Iwaya K, Tsuboi S. Imidacloprid, a novel systemic nitromethylene analog insecticide for crop protection. *The Brighton Crop Protection Conference: Pests and Diseases*. 1990; Brighton. p. 21-28

[16] Maienfisch P, Angst M, Brandl F, Fischer W, Hofer D, Kayser H, Kobel W, Rindlisbacher A, Senn R, Steinemann A, Widmer H. Chemistry and biology of thiamethoxam: a second generation neonicotinoid. *Pest Management Science*. 2001;**57**:906-913. DOI: 10.1002/ps.365

[17] Meredith RH, Heatherington PJ, Morris E. Clothianidin – a new chloronicotinyl seed treatment for use on sugar beet and cereals: field trial experiences from Northern Europe. *The Brighton Crop Protection Conference: Pests and Diseases*. Brighton. 2002; p. 691-696

[18] Wang X, Anadon A, Wu Q, Qiao F, Ares I, Martiney-Larranaga MR, Yuan Z, Martinez MA. Mechanism of neonicotinoid toxicity: Impact on oxidative stress and metabolism. *Annual Review of Pharmacology and Toxicology*. 2018;**58**:471-507. DOI: 10.1146/annurev-pharmtox-010617-052429

[19] Sanchez-Bayo F. Systemic insecticides and their environmental repercussions. Dellasala DA, Goldstein MI (eds.) *Encyclopedia of the Anthropocene*. Elsevier, Oxford. 2018; p. 111-117. Available from: <https://doi.org/10.1016/B978-0-12-409548-9.09895-X> [Accessed: 2020-09-21]

[20] Jeschke P, Nauen R, Schindler M, Elbert A. Overview of the status and global strategy for neonicotinoids. *Journal of Agricultural and Food Chemistry*. 2011;**59**(7):2897-2908. DOI: 10.1021/jf101303g

[21] Maienfisch P, Huerlimann H, Rindlisbacher A, Gsell L, Dettwiler H, Haettenschwiler J, Sieger E, Walti M. The discovery of thiamethoxam: a second-generation neonicotinoid. *Pest Management Science*. 2001;**57**(2):165-176. DOI: 10.1002/1526-4998(200102)57:2<165::AID-PS289>3.0.CO;2-G

[22] Bromilow RH, Chamberlain K. Principles governing uptake and transport of chemicals. Trapp S, McFarlane JC (eds.) *Plant contamination: modelling and simulation of organic chemical process*. CRC Press, Boca Raton, Florida, 1995; 37-68

[23] Bonmatin JM, Giorio C, Girolami V, Goulson D, Kreutzweiser DP, Krupke C, Liess M, Long E, Marzaro M, Mitchell EAD, Noome DA, Simon-Delso N, Tapparo A. Environmental fate and exposure; neonicotinoids and fipronil. *Environmental Science and Pollution Research*. 2015;**22**(1):35-67. DOI: 10.1007/s11356-014-3332-7

[24] Magalhaes LC, Hunt TE, Siegfries BD. Efficacy of neonicotinoid seed treatments to reduce soybean aphid populations under field and controlled conditions in Nebraska. *Journal of Economic Entomology*. 2009;**102**(1):187-195. DOI: 10.1603/029.102.0127

[25] Goulson D. An overview of the environmental risks posed by neonicotinoid insecticides. *Journal of Applied Ecology*. 2013;**50**(4):977-987. DOI: 10.1111/1365-2664.12111



- [26] Hladik M, Main AR, Goulson D. Environmental risks and challenges associated with neonicotinoid insecticides. *Environmental Science and Technology*. 2018;**52**:3329-3335. DOI: 10.1021/acs.est.7b06388
- [27] United States Department of Agriculture (USDA). Pesticide Data Program: Annual Summary. Calendar Year 2014. 2016. Available from: <https://www.ams.usda.gov/sites/default/files/media/2014%20PDP%20Annual%20Summary.pdf> [Accessed: 2020-11-24]
- [28] Seccia S, Fidente P, Barbini DA, Morrica P. Multiresidue determination of nicotinoid insecticide residues in drinking water by liquid chromatography with electrospray ionization mass spectrometry. *Analytica Chimica acta*. 2005;**553**(1-2):21-26. DOI: 10.1016/j.aca.2005.08.006
- [29] Klarich KL, Pflug NC, DeWald EM, Hladik ML, Kolpin DW, Cwiertny DM, LeFevre GH. Occurrence of neonicotinoid insecticides in finished drinking water and fate during drinking water treatment. *Environmental Science and Technology Letters*. 2017;**4**(5):168-173. DOI: 10.1021/acs.estlett.7b00081
- [30] Xie W, Han C, Qian Y, Ding H, Chen X, Xi J. Determination of neonicotinoid pesticides residues in agricultural samples by solid-phase extraction combined with liquid chromatography-tandem mass spectrometry. *Journal of Chromatography A*. 2011;**1218**(28): 4426-4433. DOI: 10.1016/j.chroma.2011.05.026
- [31] Seccia S, Fidente P, Montesano D, Morrica P. Determination of neonicotinoid insecticides residues in bovine milk samples by solid-phase extraction clean-up and liquid chromatography with diode-array detection. *Journal of Chromatography A*. 2008;**1214**(1-2):115-120. DOI: 10.1016/j.chroma.2008.10.088
- [32] Mitchell EAD, Mulhauser B, Mulot M, Mutabazi A, Glauser G, Aebi A. A worldwide survey of neonicotinoids in honey. *Science*. 2017;**358**(6359):109-111. DOI: 10.1126/science.aan3684
- [33] Matsuda K, Buckingham SD, Kleier D, Rauh JJ, Grauso M, Sattelle DB. Neonicotinoids: insecticides acting on insect nicotinic acetylcholine receptors. *Trends in Pharmacological Sciences*. 2001;**22**(11):573-580. DOI: 10.1016/s0165-6147(00)01820-4
- [34] Morrissey CA, Mineau P, Devries JH, Sanchez-Bayo F, Liess M, Cavallaro MC, Liber K. Neonicotinoid contamination of global surface waters and associated risk to aquatic invertebrates: a review. *Environment International*. 2015;**74**:291-303. DOI: 10.1016/j.envint.2014.10.024
- [35] Tomizawa M, Casida JE. Neonicotinoid insecticide toxicology: mechanism of selective action. *Annual Review of Pharmacology and Toxicology*. 2005;**45**:247-268. DOI: 10.1146/annurev.pharmtox.45.120403.095930
- [36] E Silva CL, Brennan N, Brouwer JM, Commandeur D, Verweij RA, Van Gestel CAM. Comparative toxicity of imidacloprid and thiacloprid to different species of soil invertebrates. *Ecotoxicology*. 2017;**26**(4):555-564. DOI: 10.1007/s10646-017-1790-7
- [37] Whitehorn PR, O'Connor S, Wackers FL, Goulson D. Neonicotinoid pesticide reduces bumble bee colony growth and queen production. *Science*. 2012;**336**(6079):351-352. DOI: 10.1126/science.1215025
- [38] Sandrock C, Tanadini LG, Pettis JS, Biesmeijer JC, Potts SG, Neumann P. Sublethal neonicotinoid insecticide exposure reduces solitary bee reproductive success. *Agricultural and Forest Entomology*. 2014;**16**(2):119-128. DOI: 10.1111/afe.12041



- [39] Williams GR, Troxler A, Retschnig G, Roth K, Zaney O, Shutler D, Neumann P, Gauthier L. Neonicotinoid pesticides severely affect honey bee queens. *Scientific Report*. 2015;**5**:14621. DOI: 10.1038/srep14621
- [40] Brandt A, Gorenflo A, Siede R, Meixner M, Büchler R. The neonicotinoids thiacloprid, imidacloprid, and clothianidin affect the immunocompetence of honey bees (*Apis mellifera* L.). *Journal of Insect Physiology*. 2016;**86**:40-47. DOI: 10.1016/j.jinsphys.2016.01.001
- [41] European Food Safety Authority (EFSA), Abdourahime H, Anastassiadou M, Arena M, Auteri D, Barmaz S, Brancato A, Brocca D, Bura L, Cabrera LC, Chiusolo A, Civitella C, Marques DC, Crivellente F, Ctverackova L, De Lentdecker C, Egsmose M, Fait G, Ferreira L, Gatto V, Greco L, Ippolito A, Istace F, Jarrah S, Kardassi D, Leuschner R, Lostia A, Lythgo C, Magrans JO, Medina P, Messinetti S, Mineo D, Miron I, Nave S, Molnar T, Padovani L, Morte JMP, Pedersen R, Raczyk M, Reich H, Ruocco S, Saari KE, Sacchi A, Santos M, Serafimova R, Sharp R, Stanek A, Streissl F, Sturma J, Szentes C, Tarazona J, Terron A, Theobald A, Vagenende B, Vainovska P, Van Dijk J, Verani A, Villamar-Bouza L. Peer review of the pesticide risk assessment of the active substance thiacloprid. *EFSA Journal*. 2019;**17**(3):e05595. DOI: 10.2903/j.efsa.2019.5595
- [42] Ellis C, Park KJ, Whitehorn P, David A, Goulson D. The neonicotinoid insecticide thiacloprid impacts upon bumblebee colony development under field conditions. *Environmental Science and Technology*. 2017;**51**(3):1727-1732. DOI: 10.1021/acs.est.6b04791
- [43] Zhang P, Sun H, Min L, Ren C. Biochars change the sorption and degradation of thiacloprid in soil: insights into chemical and biological mechanisms. *Environmental pollution*. 2018;**236**:158-167. DOI: 10.1016/j.envpol.2018.01.030
- [44] Chrétien F, Giroux I, Thériault G, Garnon P, Corriveau J. Surface runoff and subsurface tile drain losses of neonicotinoids and companion herbicides at edge-of-field. *Environmental pollution*. 2017;**224**:255-264. DOI: 10.1016/j.envpol.2017.02.002
- [45] Smith R, Middlebrook R, Turner R, Huggins R, Vardy S, Warne M. Large-scale pesticide monitoring across Great Barrier Reef catchments-paddock to reef integrates monitoring, modelling and reporting program. *Marine Pollution Bulletin*. 2012;**64**(4-9):117-127. DOI: 10.1016/j.marpolbul.2011.08.010
- [46] Gonzalez-Rey M, Tapie N, Le Menach K, Dévier MH, Budzinski H, Bebianno MJ. Occurrence of pharmaceutical compounds and pesticides in aquatic systems. *Marine Pollution Bulletin*. 2015;**96**(1-2):384-400. DOI: 10.1016/j.marpolbul.2015.04.029
- [47] Yamamoto A, Terao T, Hisatomi H, Kawasaki H, Arakawa R. Evaluation of river pollution of neonicotinoids in Osaka City (Japan) by LC/MS with dopant-assisted photoionization. *Journal of Environmental Monitoring*. 2012;**14**(8):2189-2194. DOI: 10.1039/c2em30296a
- [48] Main AR, Headley JV, Peru KM, Michel NL, Cessna AJ, Morrissey CA. Widespread use and frequent detection of neonicotinoid insecticides in wetlands of Canada's Prairie Pothole Region. *Plos One*. 2014;**9**(3):e92821. DOI: 10.1371/journal.pone.0092821
- [49] Sanchez-Bayo F, Hyne RV. Detection and analysis of neonicotinoids in river waters-development of a passive sampler for three commonly used insecticides. *Chemosphere*. 2014;**99**:143-151. DOI: 10.1016/j.chemosphere.2013.10.051

- [50] Alexander AC, Culp JM, Liber K, Cessna AJ. Effects of insecticide exposure on feeding inhibition in mayflies and oligochaetes. *Environmental Toxicology and Chemistry*. 2007;**26**(8):1726-1732. DOI: 10.1897/07-015r.1
- [51] Agatz A, Cole TA, Preuss TG, Zimmer E, Brown CD. Feeding inhibition explains effects of imidacloprid on the growth, maturation, reproduction, and survival of *Daphnia magna*. *Environmental Science and Technology*. 2013;**47**(6):2909-2917. DOI: 10.1021/es304784t
- [52] Barletta M, Lima ARA, Costa MF. Distribution, sources and consequences of nutrients, persistent organic pollutants, metals, and microplastics in South America estuaries. *Science of the Total environment*. 2019;**651**(1):1199-1218. DOI: 10.1016/j.scitotenv.2018.09.276
- [53] Raby M, Nowierski M, Perlov D, Zhao X, Hao C, Poirier DG, Sibley PK. Acute toxicity of 6 neonicotinoid insecticides to freshwater invertebrates. *Environmental Toxicity and Chemistry*. 2018;**37**(5):1430-1445. DOI: 10.1002/etc.4088
- [54] Cavallaro MC, Morrissey CA, Headley JV, Peru KM, Liber K. Comparative chronic toxicity of imidacloprid, clothianidin, and thiamethoxam to *Chironomus dilutus* and estimation of toxic equivalency factors. *Environmental Toxicity and Chemistry*. 2017;**36**(2):372-382. DOI: 10.1002/etc.3536
- [55] Stoughton SJ, Liber K, Culp J, Cessna A. Acute and chronic toxicity of imidacloprid to the aquatic invertebrates *Chironomus tentans* and *Hyalella azteca* under constant- and pulse-exposure conditions. *Archives of Environmental Contamination and Toxicology*. 2008;**54**(4):662-673. DOI: 10.1007/s00244-007-9073-6
- [56] Roessink I, Merga LB, Zweers HJ, Van den Brink PJ. The neonicotinoid imidacloprid shows high chronic toxicity to mayfly nymphs. *Environmental Toxicity and Chemistry*. 2013;**32**(5):1096-1100. DOI: 10.1002/etc.2201
- [57] Basley K, Goulson D. Neonicotinoids thiamethoxam and clothianidin adversely affect the colonisation of invertebrate populations in aquatic microcosms. *Environmental Science and Pollution research*. 2018;**25**(10):9593-9599. DOI: 10.1007/s11356-017-1125-5
- [58] Mason R, Tennekes H, Sánchez-Bayo F, Jepsen PU. Immune suppression by neonicotinoid insecticides at the root of global wildlife declines. *Journal of Environmental Immunology and Toxicology*. 2013;**1**(1):3-12. DOI: 10.7178/jeit.1
- [59] Abou-Donia MB, Goldstein LB, Bullman S, Tu T, Khan WA, Dechkovskaia AM, Abdel-Rahman AA. Imidacloprid induces neurobehavioral deficits and increases expression of glial fibrillary acidic protein in the motor cortex and hippocampus in offspring rats following in utero exposure. *Journal of Toxicology and Environmental Health, Part A*. 2008;**71**(2):119-130. DOI: 10.1080/15287390701613140
- [60] Li P, Ann J, Akk Gustav. Activation and modulation of human  $\alpha 4\beta 2$  nicotinic acetylcholine receptors by the neonicotinoids, clothianidin and imidacloprid. *Journal of Neuroscience Research*. 2011;**89**(8):1295-1301. DOI: 10.1002/jnr.22644
- [61] Kimura-Kuroda J, Komuta Y, Kuroda Y, Hayashi M, Kawano H. Nicotine-like effects of the neonicotinoid insecticides acetamiprid and imidacloprid on cerebellar neurons from neonatal rats. *Plos One*. 2012;**7**(2):e32432. DOI: 10.1371/journal.pone.0032432

- [62] Cimino AM, Boyles AL, Thayer KA, Perry MJ. Effects of neonicotinoid pesticide exposure on human health: a systematic review. *Environmental Health Perspectives*. 2016;**125**(2):155-162. DOI: 10.1289/EHP515
- [63] Gu Y, Li Y, Huang X, Zheng J, Yang J, Diao H, Yuan Y, Xu Y, Liu M, Shi H, Xu W. Reproductive effects of two neonicotinoid insecticides on mouse sperm function and early embryonic development in vitro. *Plos One*. 2013;**8**(7):e70112. DOI: 10.1371/journal.pone.0070112
- [64] Kapoor U, Srivastava MK, Srivastava LP. Toxicological impact of technical imidacloprid on ovarian morphology, hormones and antioxidant enzymes in female rats. *Food and Chemical Toxicology*. 2011;**49**(12):3086-3089. DOI: 10.1016/j.fct.2011.09.009
- [65] Toor HK, Sangha GK, Khera KS. Imidacloprid induces histological and biochemical alterations in liver of female albino rats. *Pesticide Biochemistry and Physiology*. 2013;**105**(1):1-4. DOI: 10.1016/j.pestbp.2012.10.001
- [66] Şekeroğlu V, Şekeroğlu ZA, Demirhan ES. Effects of commercial formulations of deltamethrin and/or thiacloprid on thyroid hormone levels in rat serum. *Toxicology and Industrial Health*. 2014;**30**(1):40-46. DOI: 10.1177/0748233712448114
- [67] Aydin B. Effects of thiacloprid, deltamethrin and their combination on oxidative stress in lymphoid organs, polymorphonuclear leukocytes and plasma of rats. *Pesticide Biochemistry and Physiology*. 2011;**100**(2):165-171. DOI: 10.1016/j.pestbp.2011.03.006
- [68] Bal R, Naziroğlu M, Türk G, Yilmaz Ö, Kuloğlu T, Etem E, Baydas G. Insecticide imidacloprid induces morphological and DNA damage through oxidative toxicity on the reproductive organs of developing male rats. *Cell Biochemistry and Function*. 2012;**30**(6):492-499. DOI: 10.1002/cbf.2826
- [69] Sharma N, Banerjee H, Pal S, Sharma KK. Persistence of thiacloprid and deltamethrin residues in tea grown at different locations of North-East India. *Food Chemistry*. 2018;**253**:88-92. DOI: 10.1016/j.foodchem.2018.01.132
- [70] O'Neil MJ. The Merck Index: An encyclopedia of chemicals, drugs and biologicals. Whitehouse Station: Merck Research Laboratories. Division of Merck and Co. 2006. p. 2708
- [71] National Center for Biotechnology Information. PubChem Compound Summary for CID 115224. Thiacloprid. 2021. Available from: <https://pubchem.ncbi.nlm.nih.gov/compound/Thiacloprid> [Accessed: 2020-10-11]
- [72] World Health Organization & International Programme on Chemical Safety. The WHO recommended classification of pesticides by hazard and guidelines to classification 2009. World Health Organization. 2010. p. 78. Available from: <https://apps.who.int/iris/handle/10665/44271> [Accessed: 2021-06-15]
- [73] U.S. Environmental Protection Agency. Office of Pesticide Programs. Chemicals evaluated for carcinogenic potential annual cancer report 2018. p. 40. Available from: <https://www.epa.gov/pesticide-registration/chemicals-evaluated-for-carcinogenic-potential-annual-cancer-report-2018> [Accessed: 2020-09-10]
- [74] EPA U.S. Thiacloprid. Pesticide Tolerances. Federal Register. 2013;**70**:8410-8416. Available from: <https://www.federalregister.gov/documents/2013/03/14/2013-05410-thiacloprid-pesticide-tolerances> [Accessed: 2020-08-13]
- [75] Velisek J, Stara A. effect of thiacloprid on early life stages of common carp (*Cyprinus carpio*).



- Chemosphere. 2018;**194**:481-487. DOI: 10.1016/j.chemosphere.2017.11.176
- [76] Galdíková M, Šivíková K, Holečková B, Dianovský J, Drážovská M, Schwarzbacherová V. The effect of thiacloprid formulation on DNA/chromosome damage and changes in GST activity in bovine peripheral lymphocytes. *Journal of Environmental Science and Health, Part B*. 2015;**50**(10):698-707. DOI: 10.1080/03601234.2015.1048102
- [77] Kocaman AY, Rencüzoğullari E, Topaktaş M. In vitro investigation of the genotoxic and cytotoxic effects of thiacloprid in cultured human peripheral blood lymphocytes. *Environmental Toxicology*. 2014;**29**(6):631-641. DOI: 10.1002/tox.21790
- [78] Schwarzbacherová V, Wnuk M, Deregowska A, Holečková B, Lewinska A. In vitro exposure to thiacloprid-based insecticide formulation promotes oxidative stress, apoptosis and genetic instability in bovine lymphocytes. *Toxicology in Vitro*. 2019;**61**:104654. DOI: 10.1016/j.tiv.2019.104654
- [79] Fan LF, Zhao GH, Shi HJ, Liu MC, Li ZX. A highly selective electrochemical impedance spectroscopy-based aptasensor for sensitive detection of acetamiprid. *Biosensors and Bioelectronics*. 2013;**43**(1):12-18. DOI: 10.1016/j.bios.2012.11.033
- [80] Fogel MN, Schneider MI, Desneux N, Gonzalez B, Ronco AE. Impact of the neonicotinoid acetamiprid on immature stages of the predator *Eriopis connexa* (Coleoptera: Coccinellidae). *Ecotoxicology*. 2013;**22**(6):1063-1071. DOI: 10.1007/s10646-013-1094-5
- [81] National Center for Biotechnology Information. PubChem Compound Summary for CID 213021. Acetamiprid. 2021. Available from: <https://pubchem.ncbi.nlm.nih.gov/compound/Acetamiprid> [Accessed: 2021-07-15]
- [82] Patočka J, Ardila MC, Vázquez MV. Poisons of fogs of the family Dendrobatidae-inspiration for bioorganic chemistry. *Chemické Listy*. 2000;**94**:230-233. Available from: <http://chemicke-listy.cz/docs/full/archiv/2000-PDF/04-PDF/230-233.pdf> [Accessed: 2020-04-07]
- [83] Decourtye A, Devillers J. Ecotoxicity of Neonicotinoid Insecticides to Bees. In: Thany S.H. (eds) *Insect Nicotinic Acetylcholine Receptors*. *Advances in Experimental Medicine and Biology*, Springer, New York. 2010.**683**: 85-95. DOI: 10.1007/978-1-4419-6445-8\_8
- [84] Liu Z, Dai Y, Huang G, Gu Y, Ni J, Wei H, Yuan S. Soil microbial degradation of neonicotinoid insecticides imidacloprid, acetamiprid, thiacloprid and imidaclothiz and its effect on the persistence of bioefficacy against horsebean aphid *Aphis craccivora* Koch after soil application. *Pest Management Science*. 2011;**67**(10):1245-1252. DOI: 10.1002/ps.2174
- [85] Gupta S, Gajbhiye VT. Persistence of acetamiprid in soil. *Bulletin of Environmental Contamination and Toxicology*. 2007;**78**(5):349-352. DOI: 10.1007/s00128-007-9097-7
- [86] Obana H, Okihashi M, Akutsu K, Kitagawa Y, Hori S. Determination of neonicotinoid pesticide residues in vegetables and fruits with solid phase extraction and liquid chromatography mass spectrometry. *Journal of Agricultural and Food Chemistry*. 2003;**51**(9):2501-2505. DOI: 10.1021/jf0261102
- [87] Ferrer I, Thurman EM, Fernández-Alba AR. Quantitation and accurate mass analysis of pesticides in vegetables by LC/TOF-MS. *Analytical*

- Chemistry. 2005;77(9):2818-2825. DOI: 10.1021/ac048458x
- [88] Marín A, Martínez Vidal JL, Egea Gonzalez FJ, Garrido Frenich A, Glass CR, Sykes M. Assessment of potential (inhalation and dermal) and actual exposure to acetamiprid by greenhouse applicators using liquid chromatography-tandem mass spectrometry. *Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences*. 2004;804(2):269-275. DOI: 10.1016/j.jchromb.2004.01.022
- [89] Imamura T, Yanagawa Y, Nishikawa K, Matsumoto N, Sakamoto T. Two cases of acute poisoning with acetamiprid in humans. *Clinical Toxicology (Philadelphia, Pa.)*. 2010;48(8):851-853. DOI: 10.3109/15563650.2010.517207
- [90] Kocaman AY, Topaktaş M. In vitro evaluation of the genotoxicity of acetamiprid in human peripheral blood lymphocytes. *Environmental and Molecular Mutagenesis*. 2007;48(6):483-490. DOI: 10.1002/em.20309
- [91] Yamamoto, I. Nicotine to Nicotinoids: 1962 to 1997. In: *Nicotinoid insecticides and the nicotinic acetylcholine receptor*. Springer-Verlag, Tokyo. 1999. p. 3-27. DOI: 10.1007/978-4-431-67933-2\_1
- [92] USDA Forest Service. Imidacloprid: Human Health and Ecological Risk Assessment-Final Report. 2005. Available from: ForestServiceImidacloprid.pdf (umn.edu) [Accessed: 2021-04-22]
- [93] Elbert A, Becker B, Hartwig J, Erdelen C. Imidacloprid – a new systemic insecticide. *Pflanzenschutz-Nachrichten Bayer*. 1991;44(2):113-136
- [94] National Center for Biotechnology Information. PubChem Compound Summary for CID 86287518. Imidacloprid. 2021. Available from: <https://pubchem.ncbi.nlm.nih.gov/compound/Imidacloprid> [Accessed: 2021-07-17]
- [95] Mohamed F, Gawarammana I, Robertson TA, Roberts MS, Palangasinghe C, Zawahir S, Jayamanne S, Kandasamy J, Eddleston M, Buckley A, Dawson AH, Roberts DM. Acute human self-poisoning with imidacloprid compound: a neonicotinoid insecticide. *Plos One*. 2009;4(4):e5127. DOI: 10.1371/journal.pone.0005127
- [96] Smit CE, Posthumus Doodeman JAM, van Vlaardingen PLA, de Jong FMW. Ecotoxicity of imidacloprid to aquatic organisms: derivation of water quality standards for peak and long-term exposure. *Human and Ecological Risk Assessment: An International Journal*. 2015;21(6):1608-1630. DOI: 10.1080/10807039.2014.964071
- [97] Ewere EE, Reichelt-Brushett A, Benkendorff K. Impacts of neonicotinoids on molluscs: what we know and what we need to know. *Toxics*. 2021;9(2):21. DOI: 10.3390/toxics9020021
- [98] Hallmann CA, Foppen RPB, van Turnhout CAM, de Kroon H, Jongejans E. Declines in insectivorous birds are associated with high neonicotinoid concentrations. *Nature*. 2014;511:341-343. DOI: 10.1038/nature13531
- [99] Suchail S, Guez D, Belzunces LP. Discrepancy between acute and chronic toxicity induced by imidacloprid and its metabolites in *Apis mellifera*. *Environmental Toxicology and Chemistry*. 2009;20(11):2482-2486. DOI: 10.1002/etc.5620201113
- [100] Bonmatin JM, Moineau I, Charvet R, Colin ME, Fleche C, Bengsch ER. Behaviour of Imidacloprid



- in Fields. Toxicity for Honey Bees. In: Lichtfouse E, Schwarzbauer J, Robert D. (eds). *Environmental Chemistry*. Springer, Berlin, Heidelberg; 2005. p. 483-494. DOI: 10.1007/3-540-26531-7\_44
- [101] Mundhe SA, Birajdar SV, Chavan SS, Pawar NR. Imidacloprid poisoning: an emerging cause of potentially fatal poisoning. *Indian Journal of Critical Care Medicine*. 2017;**21**(11):786-788. DOI: 10.4103/ijccm.IJCCM\_152\_17
- [102] Roberts TR, Hutson DH. Imidacloprid. *Metabolic pathways of agrochemicals – Part 2: Insecticides and Fungicides*. The Royal Society of Chemistry. Cambridge, UK. 1999. p. 111-120
- [103] Nauen R, Tietjen K, Wagner K, Elbert A. Efficacy of plant metabolites of imidacloprid against *Myzus persicae* and *Aphis gossypii* (Homoptera: Aphididae). *Pesticide Science*. 1998;**52**(1):53-57. DOI: 10.1002/(SICI)1096-9063(199801)52:1<53::AID-PS621>3.0.CO;2-6
- [104] Turro NJ. *Modern molecular photochemistry*. Benjamin-Cummings. Menlo Park, CA. 1978. pp. 305
- [105] Eftink MR, Ghiron CA. Fluorescence quenching studies with proteins. *Analytical Biochemistry*. 1981;**114**(2):199-227. DOI: 10.1016/0003-2697(81)90474-7
- [106] Tian J, Liu J, He W, Hu Z, Yao X, Chen X. Probing the binding of scutellarin to human serum albumin by circular dichroism, fluorescence spectroscopy, FTIR, and molecular modeling method. *Biomacromolecules*. 2004;**5**(5):1956-1961. DOI: 10.1021/bm049668m
- [107] Huang Y, Zhang Z, Zhang D, Lv J. Flow-injection analysis chemiluminescence detection combined with microdialysis sampling for studying protein binding of drug. *Talanta*. 2001;**53**(4):835-841. DOI: 10.1016/S0039-9140(00)00569-5
- [108] Zhang J, Zhuang S, Tong C, Liu W. Probing the molecular interaction of triazole fungicides with human serum albumin by multispectroscopic techniques and molecular modeling. *Journal of Agricultural and Food Chemistry*. 2013;**61**(30):7203-7211. DOI: 10.1021/jf401095n
- [109] Ross PD, Subramanian S. Thermodynamics of protein association reactions: forces contributing to stability. *Biochemistry*. 1981;**20**(11):3096-3102. DOI: 10.1021/bi00514a017
- [110] Némethy G, Scheraga HA. The structure of water and hydrophobic bonding in proteins. III. The thermodynamic properties of hydrophobic bonds in proteins 1,2. *Journal of Physical Chemistry*. 1962;**66**(10):1773-1789. DOI: 10.1021/j100816a004
- [111] Pace CN. Measuring and increasing protein stability. *Trends in Biotechnology*. 1990;**8**:93-98. DOI: 10.1016/0167-7799(90)90146-O
- [112] Greenfield NJ. Using circular dichroism spectra to estimate protein secondary structure. *Nature Protocols*. 2006;**1**(6):2876-2890. DOI: 10.1038/nprot.2006.202
- [113] Mitsopoulou CA, Dagas CE, Makedonas C. Synthesis, characterization, DFT studies and DNA binding of mixed platinum(II) complexes containing quenuxaline and 1,2-dithiolate ligands. *Journal of Inorganic Biochemistry*. 2008;**102**(1):77-86. DOI: 10.1016/j.jinorgbio.2007.07.002
- [114] Zhang YL, Zhang X, Fei XC, Wang SL, Gao HW. Binding of bisphenol A and acrylamide to BSA and

- DNA: insights into the comparative interactions of harmful chemicals with functional biomacromolecules. *Journal of Hazardous Materials*. 2010;**182** (1-3):877-885. DOI: 10.1016/j.jhazmat.2010.06.131
- [115] Shahabadi N, Hadidi S. Spectroscopic studies on the interaction of calf thymus DNA with the drug levetiracetam. *Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy*. 2012;**96**:278-283. DOI: 10.1016/j.saa.2012.05.045
- [116] Zhang Y, Zhang G, Zhou X, Li Y. Determination of acetamiprid partial-intercalative binding to DNA by use of spectroscopic, chemometrics, and molecular docking techniques. *Analytical Bioanalytical Chemistry*. 2013;**405**:8871-8883. DOI: 10.1007/s00216-013-7294-2
- [117] Verebová V, Želonková K, Holečková B, Staničová J. The effect of neonicotinoid insecticide thiacloprid on the structure and stability of DNA. *Physiological Research*. 2019;**68**(Suppl.4):459-466. DOI: 10.33549/physiolres.934385
- [118] Sugio S, Kashima A, Mochizuki S, Noda M, Kobayashi K. Crystal structure of human serum albumin at 2.5 Å resolution. *Protein Engineering, Design and Selection*. 1999;**12**(6):439-446. DOI: 10.1093/protein/12.6.439
- [119] Lee P, Wu X. Modifications of human serum albumin and their binding effect. *Current Pharmaceutical Design*. 2015;**21**(14):1862-1865. DOI: 10.2174/1381612821666150302115025
- [120] Chen ZG, Zhang GM, Chen ZX, Gao WH. A resonance light-scattering off-on system for studies of the selective interaction between adriamycin and DNA. *Analytical and Bioanalytical Chemistry*. 2012;**402**(6):2163-2171. DOI: 10.1007/s00216-011-5672-1
- [121] Ahamdi F, Jamali N, Moradian R, Astinchap B. Binding studies of pyriproxyfen to DNA by multispectroscopic atomic force microscopy and molecular modeling methods. *DNA and Cell Biology*. 2012;**31**:259-268. DOI: 10.1089/dna.2011.1303
- [122] Ahmadi F, Jafari B, Rahimi-Nasrabadi M, Ghasemi S, Ghanbari K. Proposed model for in vitro interaction between fenitrothion and DNA, by using competitive fluorescence, <sup>31</sup>P NMR, <sup>1</sup>H NMR, FT-IR, CD and molecular modeling. *Toxicology in Vitro*. 2013;**27**(2):641-650. DOI: 10.1016/j.tiv.2012.11.004
- [123] Kumar CV, Turner RS, Asuncion EH. Groove binding of styrylcyanine dye to DNA double helix: the salt effect. *Journal of Photochemistry and Photobiology A: Chemistry*. 1993;**74**(2-3):231-238. DOI: 10.1016/1010-6030(93)80121-O
- [124] Lavery R, Pullman B. Molecular electrostatic potential on the surface envelopes of macromolecules: B-DNA. *International Journal of Quantum Chemistry*. 1981;**20**(1):259-272. DOI: 10.1002/qua.560200123
- [125] Ihmels H, Faulhaber K, Vedaldi D, Dall'Acqua F, Viola G. Intercalation of organic dye molecules into double-stranded DNA. Part 2: The annelated quinolinizinium ion as a structural motif in DNA intercalators. *Photochemistry and Photobiology*. 2005;**81**(5):1107-1115. DOI: 10.1562/2005-01-25-IR-427
- [126] Wang F, Huang W, Su L, Dong ZJ, Zhang S. Spectrofluorimetric study of the binding of codeine to nucleic acids. *Journal of Molecular Structure*. 2009;**927**(1-3):1-6. DOI: 10.1016/j.molstruc.2009.01.022
- [127] Sun YT, Zhang HQ, Bi SY, Zhou XF, Wang L, Yan YS. Studies on the arctiin and its interaction with DNA by spectral

methods. *Journal of Luminescence*. 2011; **131**(11):2299-2306. DOI: 10.1016/j.jlumin.2011.04.036

[128] Shahabadi N, Kashanian S, Ahmadipour Z. DNA binding and gel electrophoresis studies of a new silver(I) complex containing 1,9-dimethyl-1,10-phenanthroline ligands. *DNA and Cell Biology*. 2011;**30**(3):187-194. DOI: 10.1089/dna.2010.1104

[129] Temerk YM, Ibrahim MS, Kotb M, Schuhmann W. Interaction of antitumor flavonoids with dsDNA in the absence and presence of Cu(II). *Analytical and Bioanalytical Chemistry*. 2013;**405**:3839-3846. DOI: 10.1007/s00216-012-6675-2

[130] Sun N, Ding Y, Tao Z, You H, Hua X, Wang M. Development of an upconversion fluorescence DNA probe for the detection of acetamiprid by magnetic nanoparticles separation. *Food Chemistry*. 2018;**257**:289-294. DOI: 10.1016/j.foodchem.2018.02.148

[131] Ge W, Yan S, Wang J, Zhu L, Chen A, Wang J. Oxidative stress and DNA damage induced by imidacloprid in Zebrafish (*Danio rerio*). *Journal of Agricultural and Food Chemistry*. 2015;**63**(6):1856-1862. DOI: 10.1021/jf504895h

[132] Bebane PSA, Hunt BJ, Pegoraro M, Jones ARC, Marshall H, Rosato E, Mallon EB. The effects of the neonicotinoid imidacloprid in gene expression and DNA methylation in the buff-tailed bumblebee *Bombus terrestris*. *Proceeding of the Royal Society B*. 2019;**286**(1905):20190718. DOI: 10.1098/rspb.2019.0718

[133] Hou S, Jiang X, Du F, Cong H, Li Y, Jie N. Study on the interaction between nucleic acids and imidacloprid and its application. *Nucleosides, Nucleotides and Nucleic Acids*. 2009;**28**(11-12):989-997. DOI: 10.1080/15257770903362156

[134] Wang C, Chu Q, Chen C, Bo Z. Investigation of the mechanism of binding of thiacloprid to human serum albumin using spectroscopic techniques and molecular modeling methods. *Spectroscopy*. 2011;**25**:113-122. DOI: 10.3233/SPE-2011-0498

[135] Wang Y, Tang B, Zhang H, Zhou Q, Zhang G. Studies on the interaction between imidacloprid and human serum albumin: Spectroscopic approach. *Journal of Photochemistry and Photobiology B: Biology*. 2009;**94**:183-190. DOI: 0.1016/j.jphotobiol.2008.11.013

[136] Mikhailopulo KI, Serchenya TS, Kiseleva EP, Chernov YG, Tsvetkova TM, Kovganko NV, Sviridov OV. Interaction of molecules of the neonicotinoid imidacloprid and its structural analogs with human serum albumin. *Journal of Applied Spectroscopy*. 2008;**75**(6):857-863. DOI: 10.1007/s10812-009-9120-3

[137] Sjöholm I, Ekman B, Kober A, Ljungstedt-Pahlman I, Seiving B, Sjödin T. Binding of drugs to human serum albumin: XI. The specificity of three binding sites as studied with albumin immobilized in microparticles. *Molecular Pharmacology*. 1979;**16**(3):767-777

[138] Huang B, Zou GL, Yang YM. Studies on the interaction between adriamycin and bovine serum albumin. *Acta Chimica Sinica*. 2002;**60**(10):1867-1871

[139] Liu XF, Xia YM, Fang Y, Zou L, Liu LL. Interaction between natural pharmaceutical homologues of coumarin and bovine serum albumin. *Acta Chimica Sinica*. 2004;**64**(16):1484-1490

[140] Kang J, Liu Y, Xie MX, Li S, Jiang M, Wang YD. Interactions of human serum albumin with chlorogenic acid and ferulic acid. *Biochimica*

et Biophysica Acta (BBA) – General Subjects. 2004;**1674**(2):205-214. DOI: 10.1016/j.bbagen.2004.06.021

[141] Bao XY, Zhu ZW, Li NQ, Chen JG. Electrochemical studies of rutin interacting with hemoglobin and determination of hemoglobin. *Talanta*. 2001;**54**(4):591-596. DOI: 10.1016/S0039-9140(00)00667-6

[142] Chen CY, Zhao B, Wang ZW. Interaction of thiacloprid with bovine hemoglobin using spectroscopic and molecular modeling methods. *Spectroscopy*. 2010;**24**:559-566. DOI: 10.3233/SPE-2010-0477

[143] Sulkowska A, Równicka J, Bojko B, Sulkowski W. Interaction of anticancer drugs with human and bovine serum albumin. *Journal of Molecular Structure*. 2003;**651**:133-140. DOI: 10.1016/S0022-2860(02)00642-7

[144] Abou-Zied OK, Al-Shihi OIK. Characterization of subdomain IIA binding site of human serum albumin in its native, unfolded, and refolded states using small molecular probes. *Journal of the American Chemical Society*. 2008;**130**(32):10793-10801. DOI: 10.1021/ja8031289

[145] Ashoka S, Seetharamappa J, Kandagal PB, Shaikh SMT. Investigation of the interaction between trazodone hydrochloride and bovine serum albumin. *Journal of Luminiscence*. 2006;**121**(1):179-186. DOI: 10.1016/j.jlumin.2005.12.001

[146] Yan CN, Mei P, Guan ZJ, Liu Y. Studies on thermodynamics features of the interaction between imidacloprid and bovine serum albumin. *Chinese Journal of Chemistry*. 2007;**25**:1085-1089. DOI: 10.1002/cjoc.200790202

[147] Ranz RR. Introductory Chapter: Actuality and trends of beekeeping worldwide, Beekeeping - New Challenges. IntechOpen. 2020.

DOI: 10.5772/intechopen.86890.  
Available from: <https://www.intechopen.com/chapters/67687>  
[Accessed: 2021-08-21]

[148] Erickson BM. Neonicotinoid pesticides can stay in the US market, EPA says. *Chemical and Engineering News*. 2021; American Chemical Society. ISSN 0009-2347