

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

186,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



N-Heterocyclic Carbene Mediated Organocatalysis Reactions

*Yatheesh Narayana, Sandhya N.C., H.E. Dinesh,
Sridhar B. Thimmaiah, Kanchugarakoppal S. Rangappa
and Kempegowda Mantelingu*

Abstract

Arduengo et al., isolated the first ‘bottleable’ carbene, the first N-heterocyclic carbene (NHC) 1,3-di(adamantyl)imidazol-2-ylidene resulted to an explosion of experimental and theoretical studies of novel NHCs being synthesized and analyzed have huge practical significance. These compounds emerged as successful ligands for coordinating transition metals, the complexes with NHC show diverse applications in the field of catalysis and organic transformation, NHC as ligand to main group elements and their properties and applications. Here this chapter provides the concise overview of N-heterocycle carbene as an organocatalyst that provides different organic transformation on to a carbonyl group. The majority of the NHC catalyzed reactions are employed in the phenomenon of reversing the electrophilic character of carbonyl carbon to nucleophilic carbon (umpolung activity) on coordination suggests benzoin, Stetter and hydroacylation reactions. Also, non-umpolung activity of bis-electrophile α,β -unsaturated acylazoliums reaction with suitable bis-nucleophiles in the organic synthesis have been studied.

Keywords: NHC carbene, Breslow intermediate, benzoin reaction, Stetter reaction, hydroacylation reaction, annulation reaction, acylazoliums, α,β -unsaturated acylazolium

1. Introduction

The first isolable carbene stabilized with adjacent phosphorous and silicon are reported by Bertrand et al. [1]. Soon after in 1991 Arduengo et al., isolated a bottleable carbene in N-heterocyclic scaffold, said to be N-heterocyclic carbene (NHC) [2]. Substituted bulky groups in NHCs adjacent to the carbene carbon provide kinetically stabilized and sterically avoids the dimerization to corresponding olefins. Remarkable stability and simple synthetic protocols revealed enormous applications of NHCs on transition metal, main group elements and as organo catalysts highlights new area of research. From the last two decades NHC organo catalysis has shepherded to extensive applications in the carbon–carbon and carbon–heteroatom bond formation. NHCs as organocatalyst involving umpolung activity of the functional group with carbonyl carbon (majority reaction employ aldehydes as substrate) acts as a transient nucleophile rather than an electrophile. The obtained nucleophile acyl anion is commonly called as “Breslow intermediate” [3]. The purpose of this

chapter, to explore the NHC catalyzed transformations in organic chemistry involving in the benzoin reaction, Stetter reaction, α - β -unsaturated aldehydes in construction of heterocycles, β -functionalization of enals, hydroacylation of double bonds and triple bonds. NHCs are also known as non-umpolung mode transformations catalysis. Specially, the mode of reactivity in the generation of α - β -unsaturated acylazolium intermediates and are intercepted with various bis-nucleophiles for the enantioselective construction of various heterocyclic compounds [4–8].

2. Benzoin reaction

In 1958, Breslow proposed a mechanism in which the reaction precedes *via* an enaminol intermediate popularly known as Breslow intermediate, using thiazonium zwitterion nothing but the resonance structure of an NHC. Deprotonation of thiazonium salt (A) using base generates a nucleophilic thiazolidine on addition to aromatic aldehyde furnish the tetrahedral intermediate (C) followed by proton transfer in order to obtain enaminol (D). In case of benzoin condensation, nucleophilic attack of intermediate on to another equivalent of aldehyde leads to formation of 2-hydroxy ketone (E) with subsequent elimination of the thiazolidine (B) (NHC). Transformation of carbonyl compound as transient nucleophile with NHC example of umpolung reaction and Breslow intermediate can be thought of as acyl anion equivalent (Figure 1) [9].

2.1 Homo benzoin reaction

Homo benzoin reactions are less challenging due to chemoselectivity issue and oxidation of aldehyde to corresponding carboxylic acid that could be avoided by coupling exclusion of oxygen. An efficient benzoin reaction catalyzed by




Figure 1.
Proposed mechanism of benzoin reaction.

imidazonium carbene was shown by Xu and Xia in 2005 [10] (**Figure 2**). Iwamoto et al. shows NHC catalyzed benzoin reaction in aqueous media [11, 12]. Enantioselective benzoin reaction employing NHC catalyst have many illustrations by various groups [13] (**Figure 3**). The most efficient enantioselective benzoin reaction was reported by Connon et al. with >99% ee (**Figure 4**) [14].

2.2 Cross benzoin reaction




Figure 2.
 Efficient enantioselective homo benzoin reaction.




Figure 3.
 An efficient benzoin reaction catalyzed by imidazonium carbene.




Figure 4.
 The most efficient enantioselective benzoin reaction by Connon et al. with >99%.

In an NHC driven cross benzoin reaction, total four products are possible, a pair of homo benzoin and cross benzoin adducts each. Electronic or steric reasons one of the aldehydes may be significantly less reactive. Connon et al. found that cross benzoin reaction could be biased by synthetically useful amount for the synthesis of desirable cross coupling products (**Figure 5**) [15].

Glorious et al., demonstrated a highly selective cross benzoin reaction with broad substrate scope [16]. Young et al. showed reactivity controlled by the carbene catalyst. An intermolecular cross coupling of aromatic aldehyde for the formation of Breslow intermediate is followed by coupling with acetaldehyde using thiazonium carbene catalyst. In constant acyl anion generation from acetaldehyde preferred by triazolium carbene is followed by coupling with aromatic aldehyde (**Figure 6**) [17].




Figure 5.
Cross benzoin reaction by Connon et al.




Figure 6.
Reactivity controlled reaction by using different carbene catalyst.




Figure 7.
The first aza benzoin reaction by Enders et al.

2.3 Aza benzoin reaction

The coupling of Breslow intermediate to imines was reported by Murry et al. for the first time [18]. NHC addition to highly electrophilic N-BOC imines leads to the formation of corresponding aza-Breslow intermediate, but this can be reversible under suitable reaction condition to obtain a pure 2-amino ketone [19]. Enders et al., reported the first aza benzoin reaction with trifluoromethyl ketimines using achiral thiazolium precatalyst [20]. Where enals reacts with trifluoromethyl ketone, obtained excellent enantioselectivity using chiral triazolium catalyst, which also furnishes electronic diminished groups tolerance (**Figure 7**) [21].

3. The Stetter reaction

In 1976, Stetter developed the thiazolium catalyzed highly selective conjugate addition reaction of aromatic/aliphatic aldehydes with an array of Michael acceptors and in most cases these reactions proceed in an intermolecular fashion [22, 23]. The Stetter reaction can be catalyzed by broad range of thiazolium, triazolium and imidazolium carbene, mostly α,β -unsaturated ketones are used as Michael acceptor. The formation of 1,4-diketone, γ -ketonitriles and γ -ketoesters resulting from NHC catalyzed Stetter reaction which is not easy by conventional method [24–27]. A mechanism of benzoin reaction of similar fashion is observed in the Stetter reaction, the in-situ generated free carbene (**B**) of azolium salt (**A**) when treated with base reacts with aldehyde, generating the nucleophilic Breslow intermediate (**D**). This intermediate undergoes irreversible addition to the Michael acceptor generating intermediate (**F**), which on proton transfer and subsequent release of free carbene affords the desired Stetter product (**G**) (**Figure 8**) [28].




Figure 8.
 Proposed mechanism of Stetter reaction.

3.1 Intramolecular Stetter reaction

The first general intra molecular Stetter reaction was reported by Ciganek in 1995. By the next year, Enders et al. explains the first enantioselective intramolecular Stetter reaction. Later, the implementation of chiral NHC's resulted in asymmetric transformation leading to synthesis of enantioselective 1,4-bifunctional compounds. The amino indanol derived chiral triazolium salt and pentafluoroamyl-substituted were developed by Rovis et al. in 2002 for the most efficient catalyst for the enantioselective intramolecular reaction (**Figure 9**) [29–33].

3.2 Intermolecular Stetter reaction

The initial intermolecular reaction established by Endres et al. with chiral thiazolium catalyzed reaction of *n*-butanal with chalcones resulted in Stetter product with 40% ee [34, 35] (**Figure 10**).

Rovis et al. contributed in the asymmetric intermolecular Stetter reaction of glyoxamide derivatives as aldehyde component and alkylidene molecule as Michael acceptors [36, 37]. DiRocco and Rovis expanded the reaction of intermolecular




Figure 9.
Some examples of intramolecular Stetter reaction.




Figure 10.
 The first intermolecular Stetter reaction by Enders et al.




Figure 11.
 Intermolecular Stetter reaction with α,β -unsaturated aldehyde and β -nitrostyrenes.

Stetter reaction with α,β -unsaturated aldehyde as aldehyde component and β -nitrostyrenes using fluorinated triazolium salt (**Figure 11**) [38].

The synthesis of enantiomeric α -amino acid derivative was developed by Glorius et al. using *N*-allylamidoarylate as Michael accepter [39]. Biju co-workers reported the efficient enantioselective NHC catalyzed intermolecular Stetter reaction of aldehydes with α,β -unsaturated sulfones and vinyl phosphonates [40, 41].

3.3 Hydroacylation of enol ethers of double bonds and triple bonds

Acyl anion reaction has been extended followed by the addition to electron neutral as carbon–carbon multiple bonds. She and Pan explained for the first time using alkyl tosylates (**Figure 12**) and found the selectivity changes when the native substrate with a phenyl group [42]. The reaction resulted in the formation of benzofuranone and the mechanism involving the addition of the Breslow intermediate to the C–C double bond of the enol ether [43].

A very important reaction for the synthesis of chromanones from the intermolecular cyclization of 2-allyloxy benzaldehydes using thiazolium NHC-catalyst by Glorius et al. for the first time [44] (**Figure 13**). Biju et al. showed the coupling of cyclopropenes using achiral triazolium with aryl aldehyde coupling partners (**Figure 14**) [45].

3.4 Annulation reaction

The extended Breslow intermediate has explored very well especially for the synthesis of heterocyclic compounds. Bode and Glorius reported NHC catalyzed homoenolate reactivity leading to synthesis of useful molecule from simple enal cascade. The reaction involved in the formation of extended Breslow intermediate from enal with carbene followed by 1,2-addition to aryl aldehyde which was then cyclized to deliver the γ -lactone product (**Figure 15**).

Several groups explored NHC catalyzed to synthesize enantioselective γ -lactone [46–48], spirocyclic γ -lactones [49, 50]. [3 + 4] Annulation reaction between enals





Figure 12.
Proposed mechanism of hydroacylation reaction of enol ether.



Scheme 13

Figure 13.
The intermolecular cyclization of 2-allyloxy benzaldehydes using thiazolium NHC-catalyst.




Figure 14.
The coupling of cyclopropenes using achiral triazolium with aryl aldehyde.




Figure 15.
 Mechanism of homoenolate reactivity for the formation of γ -lactone product.

and o-quinonemethides (**Figure 16**) was reported by Ye et al. in 2013 to obtain dioxolane fused-quinone methides [51].

The [8 + 3] annulation of enals and tropone *via* conjugate addition followed by cyclization afforded a lactone (**Figure 17**), reported by Nair et al. [52].

A number of nitrogen-containing heterocycles generated by homoenolate has explored largely. Synthesis of enantioselective cyclic sulfonyl ketamine annulation [53], β -lactam formation [54], synthesis of pyrazolidinones [55], isoxazolidinone formation [56] and mainly nitroso coupling reactions [57] and others have many




Figure 16.
 [3 + 4] Annulation reaction between enals and o-quinonemethides.





Figure 17.
 [8 + 3] Annulation of enals and tropone.

**Figure 18.**

The first NHC generated homoenolate in the formation of cyclopentenones.


synthetic applications. The first NHC generated homoenolate further utilized in the formation of cyclopentenones was introduced by Nair et al. in 2006 [58, 59]. The homoenolate intermediate reacts with chalcones generates allyloxide, further cyclized 4-membered β -lactone which decarboxylate to provide cyclopentene (Figure 18).

4. α,β -Unsaturated acylazolium intermediate

An important strategy of non-umpolung transformation reaction proceeds through the α,β -unsaturated acylazolium conjugation addition of various bis-nucleophile, a wide variety of carbo cycles and heterocycles are synthesized. 1,2-addition followed by cyclization [4, 5, 60]. α,β -unsaturated acylazoliums can be generated from α,β -unsaturated aldehyde with external oxidants [61–66], ynals, 2-bromo enals [67–69], α,β -unsaturated esters [70] or acyl fluorides [71, 72], etc. (Figure 19).

The main application of acylazolium intermediate found in the biosynthesis of clavulanic acid (potent β -lactamase inhibitor) from conjugation addition of L-arginine to the α,β -unsaturated acylazolium, demonstrated by Merski and Townsend (Figure 20) [73, 74].

Lupton reported the Claisen type reaction of α,β -unsaturated enol esters to the corresponding acylazolium/enolate pair followed by rearranged to 2,3-dihydropyrazones (Figure 21) [71, 72].

**Figure 19.**

Generation of α,β -unsaturated acylazoliums.




Figure 20.
 Acylazolium intermediate found in the biosynthesis of clavulanic acid.




Figure 21.
 Claisen type reaction of α,β -unsaturated enol esters to 2,3-dihydropyranones.

The generated electrophilic acylazolium intermediates involved in various annulation and cycloaddition reaction with bis-nucleophiles. α,β -unsaturated acylazolium with various cyclic and acyclic bis-nucleophiles for the synthesis of dihydropyranones and dihydropyridines in a formal [3 + 3] annulation reaction (Figure 22), demonstrated by Biju et al. [68].

Cycloaddition reaction of α -bromo enals with 1,3-diketones by using different achiral NHC catalyst obtained the product with the same absolute configuration but different stereodirecting substituents (Figure 23) [67].

5. Conclusion

N-heterocyclic carbenes have had a broad scope in the field of organic chemistry, often tolerating the construction of complex molecules from simple starting materials. The various modes of NHC catalysis such as the generation of Breslow intermediates, homoenolates, α,β -unsaturated acylazoliums, NHC enolates can be engaged in the synthesis of various heterocycles and carbocycles. The use of enantiomerically pure carbene catalyst can result in asymmetric synthesis of the




Figure 22.
The synthesis of dihydropyranones in a formal $[3 + 3]$ annulation reaction.




Figure 23.
Cycloaddition reaction of α -bromoaldehydes with 1,3-diketones.

target molecules. This chapter has focused on the reactivity pathways which expand the variety reactions with suitable reaction companions beyond the old-fashioned aldehydes with enhanced catalytic conventions.

Conflict of interest

The authors declare no conflict of interest.

Author details

Yatheesh Narayana¹, Sandhya N.C.², H.E. Dinesh¹, Sridhar B. Thimmaiah³,
Kanchugarakoppal S. Rangappa^{4*} and Kempegowda Mantelingu^{1*}

¹ Department of Studies in Chemistry, University of Mysore, Mysore, Karnataka, India


² PG Department of Chemistry, St. Philomena's College, Mysore, Karnataka, India

³ Department of Studies in Chemistry, Maharani's Science College for Women, Bangalore, Karnataka, India

⁴ Vignana Bhavan, Institute of Excellence (IOE), University of Mysore, Mysore, Karnataka, India

*Address all correspondence to: rangappaks@gmail.com and kmlingu@gmail.com

IntechOpen

© 2021 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. 

References

- [1] Aigau A, Grutzmacher H, Baceiredo A, Bertrand G. Analogous α,α' -biscarbenoid triply bonded species: Synthesis of a stable λ^3 -phosphinocarbene- λ^5 -phosphaacetylene. *Journal of the American Chemical Society*. 1988;**110**: 6463-6466. DOI: 10.1021/ja00227a028
- [2] Arduengo AJ III, Harlow RL, Kline M. A stable crystalline carbene. *Journal of the American Chemical Society*. 1991;**113**:361-363. DOI: 10.1021/ja00001a054
- [3] Breslow R. On the mechanism of thiamine action. IV. Evidence from studies on model systems. *Journal of the American Chemical Society*. 1958; **80**(14):3719-3726. DOI: 10.1021/JA01547A064
- [4] Zhang C, Hooper JF, Lupton DW. N-Heterocyclic carbene catalysis via the α,β -unsaturated acyl azolium. *ACS Catalysis*. 2017;**7**:2583-2596. DOI: 10.1021/acscatal.6b03663
- [5] Mahatthananchai J, Bode JW. On the mechanism of N-heterocyclic carbene-catalyzed reactions involving acyl azoliums. *Accounts of Chemical Research*. 2014;**47**:696-707. DOI: 10.1021/ar400239v
- [6] Ryan SJ, Candish L, Lupton DW. Acyl anion free N-heterocyclic carbene organo catalysis. *Chemical Society Reviews*. 2013;**42**:4906-4917. DOI: 10.1039/C3CS35522E
- [7] De Sarkar S, Biswas A, Samanta RC, Studer A. Catalysis with N-heterocyclic carbenes under oxidative conditions. *Chemistry - A European Journal*. 2013; **19**:4664-4678. DOI: 10.1002/chem.201203707
- [8] Knappke CEI, Imami A, Jacobi von Wangelin A. Oxidative N-heterocyclic carbene catalysis. *ChemCatChem*. 2012; **4**:937-941. DOI: 10.1002/cctc.201200133
- [9] Bugaut X, Glorius F. Organocatalytic umpolung: N-heterocyclic carbenes and beyond. *Chemical Society Reviews*. 2012;**41**:3511-3522. DOI: 10.1039/C2CS15333E
- [10] Xu L-W, Gao Y, Yin J-J, Li L, Xia C-G. Efficient and mild benzoin condensation reaction catalyzed by simple 1-N-alkyl-3-methylimidazolium salts. *Tetrahedron Letters*. 2005;**46**:5317. DOI: 10.1016/j.tetlet.2005.06.015
- [11] Iwamoto K, Hamaya M, Hashimoto N, Kimura H, Suzuki Y, Sato M. Benzoin reaction in water as an aqueous medium catalyzed by benzimidazolium salt. *Tetrahedron Letters*. 2006;**47**:7175-7177. DOI: 10.1016/j.tetlet.2006.07.153
- [12] Iwamoto K, Kimura H, Oike M, Sato M. Methylene-bridged bis (benzimidazolium) salt as a highly efficient catalyst for the benzoin reaction in aqueous media. *Organic & Biomolecular Chemistry*. 2008;**6**: 912-915. DOI: 10.1039/B719430G
- [13] Bugaut X. Benzoin and aza-benzoin. In: Molander GA, Knochel P, editors. *Comprehensive Organic Synthesis*. 2nd ed. Vol. 1. Oxford, UK: Elsevier; 2014. pp. 424-470
- [14] Baragwanath L, Rose CA, Zeitler K, Connon SJ. Highly Enantioselective benzoin condensation reactions involving a bifunctional protic pentafluorophenyl-substituted triazolium precatalyst. *The Journal of Organic Chemistry*. 2009;**74**: 9214-9217. DOI: 10.1021/jo902018j
- [15] Rose CA, Gundala S, Connon SJ, Zeitler K. Chemoselective crossed acyloin condensations: Catalyst and substrate control. *Synthesis*. 2011;**2**: 190-198. DOI: 10.1055/s-0030-1258363
- [16] Kuhl N, Glorius F. Direct and efficient N-heterocyclic

carbene-catalyzed hydroxymethylation of aldehydes. *Chemical Communications*. 2011;**47**:573. DOI: 10.1039/C0CC02416C

[17] Jin MY, Kim SM, Han H, Ryu DH, Yang JW. Switching regioselectivity in crossed acyloin condensations between aromatic aldehydes and acetaldehyde by altering N-heterocyclic carbene catalysts. *Organic Letters*. 2011;**13**: 880-883. DOI: 10.1021/ol102937w

[18] Murry JA, Frantz DE, Soheili A, Tillyer R, Grabowski EJJ, Reider PJ. Synthesis of α -amido ketones via organic catalysis: Thiazolium-catalyzed cross-coupling of aldehydes with acylimines. *Journal of the American Chemical Society*. 2001;**123**:9696-9697. DOI: 10.1021/ja0165943

[19] DiRocco DA, Rovis T. Catalytic asymmetric cross-aza-benzoin reactions of aliphatic aldehydes with N-Boc-protected imines. *Angewandte Chemie, International Edition*. 2012;**51**: 5904-5906. DOI: 10.1002/anie.201202442

[20] Enders D, Henseler A, Lowins S. N-Heterocyclic carbene catalyzed nucleophilic acylation of trifluoromethyl ketimines. *Synthesis*. 2009;**24**:4125-4128. DOI: 10.1055/s-0029-1217070

[21] Sun L-H, Liang Z-Q, Jia W-Q, Ye S. Enantioselective N heterocyclic carbene catalyzed aza-benzoin reaction of enals with activated ketimines. *Angewandte Chemie, International Edition*. 2013;**52**: 5803-5806. DOI: 10.1002/ange.201301304

[22] Stetter H. Catalyzed addition of aldehydes to activated double bonds—A new synthetic approach. *Angewandte Chemie (International Ed. in English)*. 1976;**15**:639-647. DOI: 10.1002/anie.197606391

[23] Stetter H, Kuhlmann H. In: Paquette LA, editor. *Organic Reactions*.

Vol. 40. New York: Wiley & Sons; 1991. p. 407

[24] Read de Alaniz J, Rovis T. The catalytic asymmetric intramolecular Stetter reaction. *Synlett*. 2009(**8**): 1189-1207. DOI: 10.1055/s-029-1216654

[25] Rovis T. Development of chiral bicyclic triazolium salt organic catalysts: The importance of the N-aryl substituent. *Chemistry Letters*. 2008;**37**: 2. DOI: 10.1246/cl.2008.2

[26] Christmann M. New developments in the asymmetric Stetter reaction. *Angewandte Chemie, International Edition*. 2005;**44**:2632. DOI: 10.1002/anie.200500761

[27] Read de Alaniz J, Kerr MS, Moore JL, Rovis R. Scope of the asymmetric intramolecular Stetter reaction catalyzed by chiral nucleophilic triazolinyldene carbenes. *The Journal of Organic Chemistry*. 2008;**73**:2033. DOI: 10.1021/jo702313f

[28] Moore JL, Silvestri AP, Read de Alaniz J, DiRocco DA, Rovis T. Mechanistic Investigation of the enantioselective intramolecular Stetter reaction: Proton transfer is the first irreversible step. *Organic Letters*. 2011; **13**:1742-1745. DOI: 10.1021/ol200256a

[29] Kerr MS, Read de Alaniz J, Rovis T. A highly enantioselective catalytic intramolecular Stetter reaction. *Journal of the American Chemical Society*. 2002;**124**:10298-10299. DOI: 10.1021/ja027411v

[30] Kerr MS, Rovis T. Enantioselective synthesis of quaternary stereocenters via a catalytic asymmetric Stetter reaction. *Journal of the American Chemical Society*. 2004;**126**:8876-8877. DOI: 10.1021/ja047644h

[31] Kerr MS, Read de Alaniz J, Rovis T. An efficient synthesis of achiral and chiral 1,2,4-triazolium salts: Bench

stable precursors for N-heterocyclic carbenes. *The Journal of Organic Chemistry*. 2005;**70**:5725-5728. DOI: 10.1021/jo050645n

[32] Moore JL, Kerr MS, Rovis T. Enantioselective formation of quaternary stereocenters using the catalytic intramolecular Stetter reaction. *Tetrahedron*. 2006;**62**:11477-11482. DOI: 10.1016/j.tet.2006.06.042

[33] Cullen SC, Rovis T. Catalytic asymmetric Stetter reaction onto vinylphosphine oxides and vinylphosphonates. *Organic Letters*. 2008;**10**:3141-3144. DOI: 10.1021/ol801047k

[34] Enders D. *Stereoselective Synthesis*. Heidelberg: Springer-Verlag; 1993. pp. 63-90

[35] Enders D, Bockstiegel B, Dyker H, Jegelka U, Kipphardt H, Kownatka D, et al. *Dechema-Monographies*. Vol. 129. Weinheim: VCH; 1993. p. 209

[36] Liu Q, Rovis T. Enantio and diastereoselective intermolecular Stetter reaction of glyoxamide and alkylidene ketoamides. *Organic Letters*. 2009;**11**: 2856. DOI: 10.1021/ol901081a

[37] Liu Q, Perreault S, Rovis T. Catalytic asymmetric intermolecular Stetter reaction of glyoxamides with alkylidenemalonates. *Journal of the American Chemical Society*. 2008;**130**: 14066. DOI: 10.1021/ja805680z

[38] DiRocco DA, Rovis T. Catalytic asymmetric intermolecular Stetter reaction of enals with nitroalkenes: Enhancement of catalytic efficiency through bifunctional additives. *Journal of the American Chemical Society*. 2011; **133**:10402. DOI: 10.1021/ja203810b

[39] Jousseau T, Wurz NE, Glorius F. Highly enantioselective synthesis of α -amino acid derivatives by an NHC-catalyzed intermolecular Stetter

reaction. *Angewandte Chemie, International Edition*. 2011;**50**:1410. DOI: 10.1002/anie.201006548

[40] Bhunia A, Yetra SR, Bhojgude SS, Biju AT. Efficient synthesis of γ -keto sulfones by NHC-catalyzed intermolecular Stetter reaction. *Organic Letters*. 2012;**14**:2830. DOI: 10.1021/ol301045x

[41] Patra A, Bhunia A, Biju AT. Facile synthesis of γ -ketophosphonates by an intermolecular Stetter reaction onto vinylphosphonates. *Organic Letters*. 2014;**16**:4798. DOI: 10.1021/ol502262d

[42] He J, Zheng J, Liu X, She X, Pan X. N-Heterocyclic carbene catalyzed nucleophilic substitution reaction for construction of benzopyrones and benzofuranones. *Organic Letters*. 2006;**8**:4637-4640. DOI: 10.1021/ol061924f

[43] Zhang J, Xing C, Tiwari B, Chi YR. Catalytic activation of carbohydrates as formaldehyde equivalents for Stetter reaction with enones. *Journal of the American Chemical Society*. 2013;**135**: 8113. DOI: 10.1021/ja401511r

[44] Hirano K, Biju AT, Piel I, Glorius F. N-Heterocyclic carbene-catalyzed hydroacylation of unactivated double bonds. *Journal of the American Chemical Society*. 2009;**131**:14190. DOI: 10.1021/ja906361g

[45] Biju AT, Glorius F. Intermolecular N-heterocyclic carbene catalyzed hydroacylation of arynes. *Angewandte Chemie, International Edition*. 2010;**49**: 9761-9764. DOI: 10.1002/anie.201005490

[46] Sohn SS, Rosen EL, Bode JW. N-Heterocyclic carbene-catalyzed generation of homoenolates: γ -butyrolactones by direct annulations of enals and aldehydes. *Journal of the American Chemical Society*. 2004;**126**: 14370-14371. DOI: 10.1021/ja044714b

- [47] Burstein C, Glorius F. Organocatalyzed conjugate umpolung of α,β -unsaturated aldehydes for the synthesis of γ -butyrolactones. *Angewandte Chemie, International Edition*. 2004;**43**:6205-6208. DOI: 10.1002/anie.200461572
- [48] Burstein C, Tschan S, Xie X, Glorius F. N-Heterocyclic carbene-catalyzed conjugate umpolung for the synthesis of γ -butyrolactones. *Synthesis*. 2006:2418-2439. DOI: 10.1055/s-2006-942447
- [49] Nair V, Vellalath S, Poonoth M, Mohan R, Suresh E. N-Heterocyclic carbene catalyzed reaction of enals and 1,2-dicarbonyl compounds: Stereoselective synthesis of spiro γ -butyrolactones. *Organic Letters*. 2006;**8**: 507-509. DOI: 10.1021/ja0625677
- [50] Nair V, Vellalath S, Poonoth M, Suresh E, Viji S. N-Heterocyclic carbene catalyzed reaction of enals and diaryl-1,2diones via homoenolate: Synthesis of 4,5,5-trisubstituted γ -butyrolactones. *Synthesis* 2007, **No. 20**, 3195–3200. DOI: 10.1055/s-2007-990781
- [51] Lv H, Jia W-Q, Sun L-H, Ye S. N-Heterocyclic carbene catalyzed [4+3] annulation of enals and o-quinone methides: Highly enantioselective synthesis of benzo- ϵ -lactones. *Angewandte Chemie, International Edition*. 2013;**52**:8607-8610. DOI: 10.1002/ange.201303903
- [52] Nair V, Poonoth M, Vellalath S, Suresh E, Thirumalai R. An N-heterocyclic carbene-catalyzed [8+3] annulation of tropone and enals via homoenolate. *The Journal of Organic Chemistry*. 2006;**71**:8964-8965. DOI: 10.1021/jo0615706
- [53] Rommel M, Fukuzumi T, Bode JW. Cyclic ketimines as superior electrophiles for NHC-catalyzed homoenolate additions with broad scope and low catalyst loadings. *Journal of the American Chemical Society*. 2008;**130**: 17266-17267. DOI: 10.1021/ja807937m
- [54] He M, Bode JW. Enantioselective, NHC-catalyzed bicyclo- β -lactam formation via direct annulations of enals and unsaturated N-sulfonyl ketimines. *Journal of the American Chemical Society*. 2008;**130**:418-419. DOI: 10.1021/ja0778592
- [55] Chan A, Scheidt KA. Direct amination of homoenolates catalyzed by N-heterocyclic carbenes. *Journal of the American Chemical Society*. 2008;**130**: 2740-2741. DOI: 10.1021/ja711130p
- [56] Seayad J, Patra PK, Zhang Y, Ying JY. Organocatalytic synthesis of N-phenylisoxazolidin-5-ones and a one-pot synthesis of β -amino acid esters. *Organic Letters*. 2008;**10**:953-956. DOI: 10.1021/ol800003n
- [57] Ikota H, Ishida T, Tsukano C, Takemoto Y. Synthesis of 3,3-disubstituted indoline-2-thiones catalyzed by an N-heterocyclic carbene. *Chemical Communications*. 2014;**50**:8871-8874. DOI: 10.1039/C4CC04047C
- [58] Nair V, Vellalath S, Poonoth M, Suresh E. N-Heterocyclic carbene-catalyzed reaction of chalcones and enals via homoenolate: An efficient synthesis of 1,3,4-trisubstituted cyclopentenones. *Journal of the American Chemical Society*. 2006;**128**:8736-8737. DOI: 10.1021/ja0625677
- [59] Nair V, Babu BP, Vellalath S, Varghese V, Raveendran AE, Suresh E. Nucleophilic heterocyclic carbene catalyzed annulation of enals to chalcones in methanol: A stereoselective synthesis of highly functionalized cyclopentanes. *Organic Letters*. 2009;**11**: 2507-2510. DOI: 10.1021/ol900571x
- [60] Ryan SJ, Candish L, Lupton DW. Acyl anion free N-heterocyclic carbene organocatalysis. *Chemical Society*

Reviews. 2013;**42**:4906-4917. DOI: 10.1039/C3CS35522E

[61] De Sarkar S, Studer A. NHC-catalyzed Michael addition to α,β -unsaturated aldehydes by redox activation. *Angewandte Chemie, International Edition*. 2010;**49**: 9266-9269. DOI: 10.1002/anie.201004593

[62] Rong Z-Q, Jia M-Q, You S-L. Enantioselective N-heterocyclic carbene-catalyzed Michael addition to α,β -unsaturated aldehydes by redox oxidation. *Organic Letters*. 2011;**13**: 4080-4083. DOI: 10.1021/ol201595f10

[63] Wanner B, Mahatthananchai J, Bode JW. Enantioselective synthesis of dihydropyridinones via NHC-catalyzed aza-Claisen reaction. *Organic Letters*. 2011;**13**:5378-5381. DOI: 10.1021/ol202272t

[64] Kravina AG, Mahatthananchai J, Bode JW. Enantioselective, NHC-catalyzed annulations of trisubstituted enals and cyclic N-sulfonylimines via α,β -unsaturated acyl azoliums. *Angewandte Chemie, International Edition*. 2012;**51**:9433-9436. DOI: 10.1002/anie.201204145

[65] De Sarkar S, Grimme S, Studer A. NHC catalyzed oxidations of aldehydes to esters: Chemoselective acylation of alcohols in presence of amines. *Journal of the American Chemical Society*. 2010;**132**:1190-1191. DOI: 10.1021/ja910540j

[66] Maji B, Vedachalan S, Ge X, Cai S, Liu X-W. N-Heterocyclic carbene-mediated oxidative esterification of aldehydes: Ester formation and mechanistic studies. *The Journal of Organic Chemistry*. 2011;**76**:3016-3023. DOI: 10.1021/jo200275c

[67] Sun F-G, Sun L-H, Ye S. N-Heterocyclic carbene-catalyzed enantioselective annulation of

bromoenal and 1,3-dicarbonyl compounds. *Advanced Synthesis and Catalysis*. 2011;**353**:3134-3138. DOI: 10.1002/adsc.201100622

[68] Yetra SR, Bhunia A, Patra A, Mane MV, Vanka K, Biju AT. Enantioselective N-heterocyclic carbene-catalyzed annulations of 2-bromoaldehydes with 1,3-dicarbonyl compounds and enamines via chiral α,β -unsaturated acylazoliums. *Advanced Synthesis and Catalysis*. 2013;**355**: 1089-1097. DOI: 10.1002/adsc.201300219

[69] Yetra SR, Kaicharla T, Kunte SS, Gonnade RG, Biju AT. Asymmetric N-heterocyclic carbene (NHC)-catalyzed annulation of modified enals with enolizable aldehydes. *Organic Letters*. 2013;**15**:5202-5205. DOI: 10.1021/ol4026155

[70] Cheng J, Huang Z, Chi YR. NHC organocatalytic formal LUMO activation of α,β -unsaturated esters for reaction with enamides. *Angewandte Chemie, International Edition*. 2013;**52**: 8592-8596. DOI: 10.1002/anie.201303247

[71] Ryan SJ, Candish L, Lupton DW. N-Heterocyclic carbene-catalyzed generation of α,β -unsaturated acylimidazoliums: Synthesis of dihydropyranones by their reaction with enolates. *Journal of the American Chemical Society*. 2009;**131**: 14176-14177. DOI: 10.1021/ja905501z

[72] Candish L, Lupton DW. The total synthesis of (–)-7-deoxyloganin via N-heterocyclic carbene catalyzed rearrangement of α,β -unsaturated enol esters. *Organic Letters*. 2010;**12**: 4836-4839. DOI: 10.1021/ol101983h

[73] Merski M, Townsend CA. Observation of an acryloyl-thiamin diphosphate adduct in the first step of clavulanic acid biosynthesis. *Journal of the American Chemical Society*. 2007;

129:15750-15751. DOI: 10.1021/
ja076704r

[74] Khaleeli N, Li R, Townsend CA.
Origin of the β -lactam carbons in
clavulanic acid from an unusual
thiamine pyrophosphate-mediated
reaction. *Journal of the American
Chemical Society*. 1999;121:9223-9224