We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Clinical Magnetic Resonance Neuroimaging in Fibromyalgia

Nicolás Fayed* and Javier Garcia-Campayo**

*Department of Radiology, Quirón Hospital, Zaragoza, Spain**

*Somatoform Disorders and Fibromyalgia Unit, Miguel Servet Hospital, University of Zaragoza, Spain***

Key points

- Magnetic resonance spectroscopy (MRS) offers invaluable information about living tissues with special contribution to the diagnosis and prognosis of the central nervous system diseases.
- Diffusion tensor imaging (DTI) detects subtle degradation of white matter microstructure in fibromyalgia.
- Perfusion magnetic resonance imaging (MRI) offers higher spatial resolution than radionuclide techniques such as positron emission tomography and single-photon emission computed tomography.
- Voxel-based morphometry (VBM) is a recent methodology that can simultaneously visualize group differences or statistical effects on gray and white matter, throughout the whole brain.
- Functional magnetic resonance imaging (fMRI) studies regularly confirm that multiple brain regions are invoked to execute even ostensibly simple tasks.

1. Introduction

Imaging human internal organs with exact and non-invasive methods is very important in medicine for medical diagnosis, treatment and follow-up, as well as for clinical research. Today, one of the most important tools for this purpose is magnetic resonance imaging (MRI). MRI scanners are based on the nuclear magnetic resonance (NMR) phenomenon, which was detected independently by Bloch, Purcell et al. in 1946. The advent of neuroimaging techniques yielding neuropsychological data in addition to structural information was of particular interest for scientific and clinical research into fibromyalgia and psychiatric disorders. These are groups of conditions for which any structural changes evident in anatomical imaging sequences generally correlate poorly with clinical diagnostic categories, pointing to underlying pathophysiology and severity of disease. T1- and T2-dependent MR sequences, the mainstay of routine clinical neuroimaging, are frequently insensitive to the underlying pathological processes in these conditions. Focal or global atrophy due to associated neuronal loss is also frequently absent.

The remit of this section is an overview of some general issues concerning the application of spectroscopy, diffusion, perfusion, morphometry and functional techniques in fibromyalgia, and their clinical impact in the context of other available tests.

2. Magnetic Resonance Spectroscopy (MRS)

This technique enables us to study the chemical composition of living tissues. It is based on the chemical shift of atoms. The concentration of some metabolites is determined from spectra that may be acquired in several ways.

3. Voxel placement

Currently the spectra may be acquired with single-voxel (SV) or multivoxel techniques (MV). The SV technique is readily available on most scanners. Voxels must be positioned away from sources of susceptibility artifacts and lipids. For diffuse processes, a $2 \times 2 \times 2$ -cm (8 cm^3) voxel is routinely used. For local lesions, the SV can be reduced in volume. The SV technique offers the advantages of better spatial location, more homogeneity, better water suppression and speed. However, only one spectrum can be obtained per acquisition.

However, The MV technique makes it possible to obtain multiple spectra simultaneously per acquisition and to assess a greater area of the brain but with smaller spectral resolution. To date, the SV is still superior to MV on the grounds of reproducibility (Sauter et al., 1991; Hsu et al., 1999, 2001; Law et al., 2004). For both SV and MV, the MR scanner employs a process known as shimming to narrow peak linewidths within the spectra. For SV studies, improving field homogeneity is performed with basic, zero-ordered shimming on clinical MR scanners. For MV, the simultaneous production of uniform field homogeneity in multiple regions requires higher order shimming. To obtain high-quality spectra, blood products, air, fat, necrotic areas, cerebrospinal fluid, metal, calcification and bone should be avoided. In such areas differing magnetic susceptibility results in a non-homogenous field that hinders the production of diagnostic quality spectra.

4. MRS pulse sequence

Two different approaches are generally used for proton spectroscopy of the brain: a) single-voxel methods based on the stimulated echo acquisition mode (STEAM) and b) point resolved spectroscopy (PRESS) pulse sequences and spectroscopy imaging (SI), also known as chemical shift imaging (CSI). These latter studies are usually done in two dimensions, using a variety of different pulse sequences (spin-echo (SE), usually PRESS).

The basic principle underlying single-voxel localization techniques is to use three mutually orthogonal slice selective pulses and design the pulse sequence to collect only the echo signal from the point (voxel) in space where all three slices intersect. In STEAM, three 90° pulses are used and the stimulated echo is collected. All other signals (echoes) should be dephased by the large crusher gradient applied during the so-called mixing time. Crusher gradients are necessary for consistent formation of the stimulated echo and removal of unwanted coherences. In PRESS, the second and third pulses are refocusing (180°) pulses, and crusher gradients are applied around these pulses to select the desired SE signal arising

from all three RF pulses and dephasing unwanted coherences. STEAM and PRESS are generally similar but differ in a few key aspects.

With regard to the mode of acquisition, PRESS can be performed with short and long echo times (TEs) and there is complete recovery of signal. STEAM can be performed with very short TEs, but there is incomplete recovery of signal and a precise volume element (voxel) is formed. The PRESS mode is used more than STEAM because it increases the signal/noise ratio and is less sensitive to movement artifacts (Maheshwari et al. 2000). A short TE (20-40 ms) allows us to increase the signal/noise ratio and to visualize most metabolite peaks, with the inconvenience of some degree of peak overlapping. Time matters in clinical practice, so short TEs are preferable. In our experience with a 1.5T GE Signa Horizon-clinical scanner a TE of 30 ms and a TR of 2500 ms have proven valuable (Fayed et al., 2006). Recently, a TE-averaged PRESS technique has been yielding highly simplified spectra with better suppression of signals not pertaining to assessed metabolites, such as that of macromolecules. TE is increased from 35 ms to 355 ms in steps of 2.5 ms with two acquisitions per step (Hancu et al., 2005).

5. MR spectra quantification

The most commonly used spectroscopy is that originating from a Hydrogen nucleus (proton ^1H -MRS). This technique is based on the differences in resonance obtained from hydrogen nuclei depending on the surrounding atoms (chemical shift). Each metabolite being assessed discloses a different hydrogen resonance frequency and appears in a different site in the spectrum. The most frequently evaluated metabolites are N-acetyl-aspartate (NAA), myo-inositol (mI), choline (Ch), creatine (Cr) and glutamate+glutamine (Glx). The position of the metabolite signal is identified on the horizontal axis by its chemical shift, scaled in units referred to as parts per million (ppm). With the appropriate factors considered, such as the number of protons, relaxation times and so forth, a signal can be converted into a metabolite concentration by measuring the area under the curve. Because water is the main component of living beings and its concentration is much higher than that of metabolites, it becomes necessary to suppress the resonance signal from the hydrogen of water (Maheshwari et al. 2000 and Bonavita et al. 1999). A plot showing peak amplitudes and frequencies is obtained. Each spectrum shows peaks corresponding to the different metabolite values: Myo-inositol (mI), 3.56 and 4.06 ppm; Choline compounds (Ch), 3.23 ppm; Creatine (Cr), 3.03 and 3.94 ppm; γ N-acetyl-aspartate (NAA), 2.02; 2.5 and 2.6 ppm; glutamine and glutamate (Glx), 2.1-2.55 ppm and 3.8 ppm. Ratios between metabolites and creatine are also of great value as they counteract the systematic errors of measurements.

6. Fitting of model spectra

A more recent program, called a linear combination model ("LCModel") (Provencher, 1993), fits in vivo spectra as a linear superposition of high-resolution "basis" spectra that are acquired from model solutions of the metabolites present in the organ of interest. Advantages of LCModel are that all pre-processing steps, automatic phase correction as well as modelling of a smooth baseline are included. Standardized basis sets are available for the most common clinical MR machines (both 1.5 and 3 T).

7. Fibromyalgia

Fibromyalgia is a chronic and disabling musculoskeletal pain disorder of unknown, characterized by a history of widespread pain for at least three months and patients reporting of tenderness in at least 11 of 18 defined tender points when digitally palpated with about 4 kg per unit area of force. Other frequent accompanying symptoms are fatigue, sleep disturbance and depressed mood (Wolfe et al, 1990; Giesecke et al, 2003). With an estimated lifetime prevalence of 2% in community samples (Wolfe et al, 1995), it accounts for 15% of outpatient rheumatology visits and 5% of primary care visits (Wolfe 1989). The prognosis for symptomatic recovery is generally poor (Wolfe et al., 1997). Recent meta-analysis (Garcia Campayo et al., 2008) demonstrates that both pharmacological and psychological treatments show moderate effectiveness with a mean effect size of 0.49. The variables that related most with improvement in outcome were younger age of the patients and shorter duration of the disorder. On the contrary, gender and type of treatment (pharmacological or psychological) did not affect outcome (Garcia Campayo et al., 2008).

8. Usefulness of different neuroimaging techniques in Fibromyalgia

a) ¹H MRS in Fibromyalgia patients

Glutamate has been implicated as an important mediator in the neurotransmission, potentiation, and negative affect associated with pain, and it has been related to chronic pain sensitization (Dickenson et al., 2002). Experimental pain models for fibromyalgia have revealed elevated glutamate levels in the posterior insula. MRS spectroscopy studies have shown that dynamic changes in glutamate and glutamine levels in the insula were associated with improvements in clinical and experimental pain in fibromyalgia (Harris et al., 2008). Also, a recent MR spectroscopy study concludes with the involvement of the posterior insula in pain (Harris et al., 2009)

A previous MRS study by our group demonstrated the differences in metabolites between FM patients and controls. (A)

Our data suggest that Glx plays a role in this augmented pain processing in those individuals who have elevated Glx levels, which is entirely consistent with the literature and knowledge regarding FM. In our study there is a significant correlation in the posterior cingulate gyrus between Glu+Gln (Glx) and Glx/Cr with depression, pain measured in an objective way, such as by sphygmomanometer, and global function assessed with the Fibromyalgia Impact Questionnaire (FIQ). Since astrocytes participate in the uptake, metabolism, and recycling of glutamate, we hypothesize that an astrocyte deficit may account for the alterations in glutamate/GABA neurotransmission in depression. Factors such as stress, excess glucocorticoids, altered gene expression of neurotrophic factors and glial transporters, and changes in extracellular levels of neurotransmitters released by neurons may modify glial cell numbers and affect the neurophysiology of depression (Rajkowska et al., 2007). Other studies found that absolute concentrations of Glx, Glu, and creatine+phosphocreatine (Cr) were significantly higher in adult bipolar patients in all mood states compared to healthy controls (Yildiz-Yesiloglu et al. 2006).

Elevated Glx inside the astrocyte leads to increased cellular osmolarity in the brain. Within 30 minutes of glutamate administration, electron microscopy reveals massive acute swelling of neuronal cell bodies and dendrites. Consequently, water shifts from the extracellular fluid space to the intracellular fluid space resulting in edema of the astrocytes. Relevant clinical

manifestations are thought to be secondary to this edema (Häussinger et al., 2000). To compensate for the increased cellular osmolarity, myo-inositol shifts to the extracellular fluids space, leading to a reduction in its concentration inside the astrocyte. In consequence, Glx is an excitatory amino acid, and its increase indicates that the metabolic function of patients with fibromyalgia differs from controls. Elevation of Gln levels may result in permanent cerebral damage.

In addition, high correlations were found between some clinical variables and certain brain metabolites, for instance: ratios of myo-Inositol/Creatine and NAA + *N*-acetyl aspartyl glutamate (NAA+NAAG) in the left hippocampus with pain; myo-Inositol in the right hippocampus and myo-Inositol in the posterior cingulate gyrus with catastrophization.

A recent HMRS study, showed decreased NAA levels within the hippocampus of individuals with FM (Wood et al., 2009). In another study, a reduction in the absolute concentration of NAA in the right and left hippocampi was reported in a sample of 15 patients with fibromyalgia (Emad et al. 2008). Lower hippocampal NAA levels suggest neuronal or axonal metabolic dysfunction, or some combination of these processes. Neuronal loss in the hippocampus has not yet been studied in our series. This should be assessed in further studies, in order to gauge atrophic changes within the hippocampus. Some studies found that the persistence of elevated Ca²⁺ in hippocampal neurons exposed to glutamate correlated with the extent of neuronal death, and that a large rise in Ca²⁺ in cultured hippocampal neurons, following glutamate application, predicted cell death (Mattson et al., 1989). Hippocampal dysfunction may be partly responsible for some of the phenomena associated with FM. Blocking NMDA receptors in the hippocampal formation reduces nociceptive behaviours; this in turn supports the hypothesis that the hippocampal formation is involved in pain-related neural processing and expression of pain-related behaviours (McKenna et al. 2001).

Finally, our study confirms the reduction in myo-Inositol levels in both hippocampi and reduction in myo-Inositol/Cr ratios in the left hippocampus and sensitivomotrice area. In all cases, brain metabolites were lower in the patient group compared to controls. Changes in myo-Inositol have previously been described in patients suffering from bipolar disorder (Silverstone et al. 2005). Research suggests that lithium functions primarily by decreasing myo-inositol concentrations in bipolar patients (Harwood et al., 2005).

Patients suffering from clinical depression generally have decreased levels of inositol in their cerebrospinal fluid (Barkai et al., 1978). This deficiency could be explained by a reduced synthesis or more interestingly by an increased consumption of the compound (Lentner, 1986). We speculate that myo-Inositol might, via its conversion to glucuronic acid, be consumed in protective detoxification reactions of the brain and could be associated with depression.

b) Diffusion Tensor Imaging (DTI)

White matter structure

Neuroimaging reveals changes in the white matter (WM) structure in the human brain. White matter comprises half of the human brain and consists of bundles of myelinated axons connecting neurons in different brain regions (Fields, 2008). Grey matter is composed of neuronal cell bodies and dendrites concentrated in the outer layers of the cortex. Microstructural changes in white matter can be revealed by specialized MRI brain imaging techniques such as diffusion tensor imaging (DTI). This method analyses the of proton

diffusion in tissue, which is more restricted in white matter than in grey matter. The anisotropy increases with increased myelination, diameter and axon compaction.

DTI measurement

Water molecules in the brain are in constant Brownian motion, and although the movement of these protons affects conventional structural imaging, diffusion weighted imaging (DWI) and diffusion tensor imaging (DTI) allow quantification of this microscopic movement within each voxel. The main advantage of using DTI, rather than DWI, is that DTI reflects the underlying diffusion properties of the sample, independently of the orientation of the tissue with respect to the direction of measurements. DTI is thus a robust quantitative technique that is independent of how the subject has been oriented inside the scanner magnet and gradient coils. In regions with few or no constraints imposed by physical boundaries, such as CSF in the ventricles, water movement is random in every direction and is isotropic. In contrast to CSF, the path of a water molecule in a WM fibre is constrained by physical boundaries, such as the axon sheath, causing the movement along the long axis of a fibre to be greater than radial diffusion across the.

These data can be used to calculate the probable anatomy of white matter bundles in living brain, a process called tractography. Orientation is calculated from the eigenvectors defining proton diffusion in three dimensions in each voxel. Using algorithms, the principal eigenvalue vector is connected to the next voxel to trace the fibre structure and orientation in white matter tracts (Jones et al., 2002)

Diffusion tensor imaging (DTI) yields quantitative measures for tissue water mobility as a function of the direction of water motion and is probed by application of diffusion-sensitization gradients in multiple directions (Basser et al., 1996). Basser et al. describe the use of multivariate linear regression to calculate D from a non-diffusion-weighted image plus six or more diffusion-weighted measurements in a non-collinear direction. The diffusion weighting is obtained by simultaneously applying diffusion gradients along combinations of the three physical axes.

The appropriate mathematical combination of the directional diffusion-weighted images provides quantitative measures of water diffusion for each voxel via the apparent diffusion coefficient (ADC), as well as the degree of diffusion directionality, or anisotropy (Sundgren et al. 2004). Myelin is a major diffusion barrier for water, and gives white matter its high anisotropy. Demyelinating diseases are characterized by partial or total loss of myelin, with consequent loss of neuronal function.

Fibromyalgia

DTI allows in vivo mapping of the anatomic connections in the human brain; previous studies have identified and confirmed the existence of an anatomic circuitry for the functionally characterized top-down influences on pain processing via brainstem structures in humans (Hadjipavlou et al., 2006) Fractional anisotropy (FA) is a measure of the portion of the diffusion tensor from anisotropy.

Previous studies with diffusion tensor imaging in FM patients showed alterations in the right thalamus and significantly lower fractionated anisotropy in comparison with controls. A negative correlation was seen between the FA values in the right thalamus and clinical pain in the FM group (Sundgren et al., 2007). Other authors confirmed that DTI in the brain

of patients with FM appears to be more sensitive than volumetric imaging of voxel-based morphometry (VBM) and increased pain intensity scores were correlated with changes in DTI measurements in the right superior frontal gyrus. Increased fatigue was correlated with changes in the left superior frontal and left anterior cingulate gyrus, and self-perceived physical impairment was correlated with changes in the left postcentral gyrus. Higher intensity scores for stress symptoms were correlated negatively with diffusivity in the thalamus and FA in the left insular cortex (Lutz et al., 2008).

In a previous study by our group, DWI and DTI were not sensitive enough to detect changes in fibromyalgia patients. This may be explained because intraparenchymal injection of glutamate or other excitatory aminoacids cause neuronal cell bodies and dendrites to be predominantly affected, whereas axons and terminal boutons, originating from cell bodies outside of the affected region, remain largely intact (Coyle et al., 1981).

c) Perfusion MRI

Positron emission tomography (PET) (Jones et al., 1991) and single-photon emission computed tomography (SPECT) (Kwiatek et al., 2000) have been used to identify focal changes in regional cerebral blood flow (CBF) in patients with fibromyalgia. However, the low spatial resolution of PET and SPECT, and the ionizing radiation emitted from the nuclear medicine tracers are major concerns. MR perfusion techniques have also been developed and offer higher spatial resolution without the use of ionizing radiation (Rosen et al., 1990; Belliveau et al., 1990; Detre et al., 1992).

MR perfusion techniques are based on exogenous or endogenous tracers. In the method based on exogenous tracers, a paramagnetic agent such as gadolinium dimeglumine gadopentate (Gd-DTPA) is injected, and the resulting decrease and subsequent recovery of the MR signal is used to estimate perfusion (Ostergaard et al., 1996; Liu et al., 1999). In the method using endogenous tracers, the magnetization of the spins of arterial water are non-invasively labelled using radiofrequency (RF) pulses, and the regional accumulation of the label is measured in the tissues by comparison with an image acquired without labelling (Edelman et al., 1994; Kim, 1995; Kwong et al., 1995).

d) Voxel-based morphometry (VBM)

Tissue volumes in the CNS, and in particular changes in volume over time, are sensitive markers of a range of neurological disease states and disease progression. Measurement of brain volume requires segmentation of the brain from the rest of the tissues in the head and neck. Whilst this can be performed manually or in a semiautomated fashion (see for example Harris et al., 1999; Bokde et al., 2002), automated procedures are likely to be more reproducible and rapid. This is understandable as the size of the structures involved is usually relatively small, making the analysis less tedious than a manual segmentation of the whole brain. Additionally, many structures, such as the hippocampus, are difficult to segment in an automated fashion, but are relatively easily identified and manually or semiautomatically outlined, given appropriate software. Some progress has been made in automating segmentation procedures, with methods including the use of deformable shape models.

Tensor-based morphometry (TBM) is a relatively new image analysis technique that identifies regional structural differences in the brain, across groups or over time, from the

gradients of the deformation fields that warp images to a common anatomical template. The anatomical information is encoded in the spatial transformation. Therefore, accurate inter-subject non-rigid registration is an essential tool. With the new advent of recent and powerful non-rigid registration algorithms based on the large deformation paradigm, TBM is being increasingly used (Lepore et al., 2008; Chiang et al., 2007; Lee et al., 2007).

In FM, Valet et al. (2009) found significant grey-matter decreases in the prefrontal, cingulate, and insular cortex. These regions are known to be critically involved in the modulation of subjective pain experiences. Patients presented a decrease in grey matter volume in the prefrontal cortex, the amygdala, and the anterior cingulate cortex (ACC). The duration of pain or functional pain disability did not correlate with grey matter volumes. A trend of inverse correlation of grey matter volume reduction in the ACC with the duration of pain medication intake has been detected. Some authors (Burgmer et al., 2009) suggest that structural changes in the pain system are associated with FM. As disease factors do not correlate with reduced grey matter volume in areas of the cingulo-frontal cortex and the amygdala in patients, one possible interpretation is that volume reductions might be a precondition for central sensitization in fibromyalgia.

e) Functional MR imaging (fMRI)

Another more recent imaging method is functional MRI (fMRI) for the mapping of activation patterns in the brain. This is an important technique for better understanding of brain function. When a brain region is activated, new energy must be transported to this region, which leads to increased blood flow to that part of the brain. This can be imaged by repetitive MR scans and detected by appropriate signal processing methods.

Over the years, many have viewed Fibromyalgia syndrome (FMS) as a so-called "functional disorder" and patients have experienced a concomitant lack of interest and legitimacy from the medical profession. The symptoms have not been explained by peripheral mechanisms alone or by specific central nervous system mechanisms.

Functional MRI blood oxygenation level dependent activation studies on patients who have FM have demonstrated augmented sensitivity to painful pressure and the association of this augmentation with variables such as depression and catastrophizing, and have also been used to evaluate the symptoms of cognitive dysfunction.

The fMRI-analysis by Jensen et al. (2009), revealed no differences in activity in brain regions related with attention and affection or regions with sensory projections from the stimulated body area. However, when there is a primary lesion in the descending pain regulating system (the rostral anterior cingulate cortex), the patients failed to respond to pain provocation. The attenuated response to pain in these cases is the first demonstration of a specific brain region where the impairment of pain inhibition in FMS patients is expressed. These results validate previous reports of dysfunctional endogenous pain inhibition in FM, and advance the understanding of the central pathophysiologic mechanisms, providing a new direction for the development of successful treatments in FM.

Fibromyalgia is the prototypical functional chronic pain condition, and it affects 2-4% of individuals. Although the aetiology of this disorder remains largely unknown, emerging data suggest that FM arises through augmentation of central pain processing pathways. This hypothesis is largely based upon findings of previous functional neuroimaging studies showing that FM patients display augmented neuronal responses to both innocuous and

painful stimuli (Gracely et al., 2002; Cook et al., 2004), confirming the allodynia and hyperalgesia seen in this condition (Petzke et al., 2003).

Studies with functional neuroimaging support the hypothesis of central pain augmentation in FMS. Differences of activation in the fronto-cingulate cortex, the supplemental motor areas and the thalamus were found between both groups with distinct differences in BOLD-signals changes over the time course of pain stimulation, even during anticipation of pain. These results support the hypothesis that central mechanisms of pain processing in the medial pain system, favourable cognitive/affective factors even during the anticipation of pain, may play an important role for pain processing in patients with FMS (Burgmer et al., 2009).

9. Summary, conclusions and future directions

In this chapter, several techniques used for the diagnosis of fibromyalgia are discussed. At present, there are no other non-invasive techniques that can provide equivalent information and, as a consequence, MRS, DTI tractography and functional MRI are expected to be a powerful combined technique for researching brain anatomy and disease *in situ* in human beings.

Diffusion tensor imaging (DTI) in combination with magnetic resonance spectroscopy (MRS) and functional MRI (fMRI) may provide clinicians with information about ongoing pathological changes in FM. Future studies may investigate whether DTI can detect Many factors may influence the ability of DTI to discriminate patients with FM from healthy individuals. These factors include the field strength of the magnet, spatial resolution, signal-to-noise ratio, contrast-to-noise ratio and image artifacts. DTI and MRS are non-invasive and do not require the use of radioactive tracers, suggesting its potential safe application for longitudinal follow-up and repeated assessments.

Continued improvements in the design of imaging equipment and analysis algorithms are progressively improving the specificity of the biological parameters that can be calculated, allowing detailed quantitative characterization of microvascular structure in a wide range of pathological tissues, including fibromyalgia.

10. References

1. Barkai IA, Dunner DL, Gross HA, Mayo P, Fieve RR. Reduced myo-inositol levels in cerebrospinal fluid from patients with affective disorder. *Biol Psychiatry* 1978; 13: 65-72.
2. Basser PJ, Pierpaoli C. Microstructural and physiological features of tissues elucidated by quantitative-diffusion-tensor MRI. *J Magn Reson B* 1996; 111: 209-19.
3. Belliveau JW, Rosen BR, Kantor HL, Rzedzian RR, Kennedy DN, McKinstry RC, Vevea JM, Cohen MS, Pykett IL, Brady TJ. Functional cerebral imaging by susceptibility-contrast NMR. *Magn Reson Med* 1990; 14: 538-46.
4. Bloch F. Nuclear induction. *Phys Rev* 1946, 70:460-474.
5. Bonavita S, Di Salle F, Tedeschi G. Proton MRS in neurological disorders. *Eur J Radiol* 1999; 30: 125-31.
6. Burgmer M, Gaubitz M, Konrad C, Wrenger M, Hilgart S, Heuft G, Pfleiderer B. Decreased gray matter volumes in the cingulo-frontal cortex and the amygdala in patients with fibromyalgia. *Psychosom Med* 2009; 71: 566-73.

7. Burgmer M, Pogatzki-Zahn E, Gaubitz M, Wessoleck E, Heuft G, Pfeleiderer B. Altered brain activity during pain processing in fibromyalgia. *Neuroimage* 2009; 44: 502-8.
8. Chiang MC, Dutton RA, Hayashi KM, Lopez OL, Aizenstein HJ, Toga AW, Becker JT, Thompson PM: 3D pattern of brain atrophy in HIV/AIDS visualized using tensor-based morphometry. *NeuroImage* 2007; 34: 44-60.
9. Cook DB, Lange G, Ciccone DS, Liu WC, Steffener J, Natelson BH. Functional imaging of pain in patients with primary fibromyalgia. *J Rheumatol* 2004; 31: 364-78.
10. Coyle JT, Bird SJ, Evans RH, Gulley RL, Nadler JV, Nicjlas WJ, et al. Excitatory amino acid neurotoxins: selectively and mechanisms of action. *Neurosci Res Prog* 1981; 19: 1-427
11. Detre JA, Leigh JS, Williams DS, Koretsky AP. Perfusion imaging. *Magn Reson Med*. 1992; 23:37-45.
12. Edelman RR, Siewert B, Darby DG, Thangaraj V, Nobre AC, Mesulam MM, Warach S. Qualitative mapping of cerebral blood flow and functional localization with echo-planar MR imaging and signal targeting with alternating radio frequency. *Radiology* 1994; 192: 513-20.
13. Emad Y, Ragab Y, Zeinhom F, El-Khouly G, Abou-Zeid A, Rasker JJ. Hippocampus dysfunction may explain symptoms of fibromyalgia syndrome. A study with single-voxel magnetic resonance spectroscopy. *J Rheumatol* 2008; 35: 1371-7.
14. Fayed N, Morales H, Modrego PJ, Pina MA. Contrast/Noise ratio on conventional MRI and choline/creatine ratio on proton MRI spectroscopy accurately discriminate low-grade from high-grade cerebral gliomas. *Acad Radiol* 2006;13: 728-37.
15. Fayed N, Olmos S, Morales H, Modrego PJ. Physical Basis of Magnetic Resonance Spectroscopy and its Application to Central Nervous System Diseases. *Am J Applied Sci* 2006; 3:1836-1845.
16. Fields RD. White matter in learning, cognition and psychiatric disorders. *Trends Neurosci* 2008; 31: 317-376.
17. Hadjipavlou G, Duncley P, Behrens TE, Tracey I. Determining anatomical connectivities between cortical and brainstem pain processing regions in humans: a diffusion tensor imaging study in healthy controls. *Pain* 2006; 123:169-78.
18. Hancu I, Zimmerman EA, Sailasuta N, Hurd RE. 1H MR spectroscopy using TE averaged PRESS: a more sensitive technique to detect neurodegeneration associated with Alzheimer's disease. *Magn Reson Med* 2005; 53:777-82.
19. Harris RE, Sundgren PC, Pang Y, Hsu M, Petrou M, Kim SH et al. Dynamic levels of glutamate within the insula are associated with improvements in multiple pain domains in fibromyalgia. *Arthritis Rheum* 2008; 58:903-7.
20. Harris RE, Sundgren PC, Craig AD, Kirshenbaum E, Sen A, Napadow V et al. Elevated insular glutamate in fibromyalgia is associated with experimental pain. *Arthritis Rheum* 2009; 60: 3146-52.
21. Harwood AJ. Lithium and bipolar mood disorder: the inositol-depletion hypothesis revisited. *Mol Psychiatry* 2005; 10: 117-26.
22. Häussinger D, Kircheis G, Fischer R, Schliess F, vom Dahl S. Hepatic encephalopathy in chronic liver disease: a clinical manifestation of astrocyte swelling and low-grade cerebral edema? *J Hepatol* 2000; 32: 1035-8.
23. Hsu YY, Chang C, Chang CN, Chu NS, Lim KE, Hsu JC. Proton MR spectroscopy in patients with complex partial seizures: single-voxel spectroscopy versus chemical-shift imaging. *Am J Neuroradiol*. 1999; 20(4): 643-51.

24. Hsu YY, Chen MC, Lim KE, Chang C. Reproducibility of hippocampal single-Voxel proton MR spectroscopy and chemical shift imaging. *Am J Roentgenol* 2001; 176: 529-36.
25. Garcia-Campayo J, Magdalena J, Magallón R, Fernández E, Salas M, Andrés E. A meta-analysis of the efficacy of fibromyalgia treatment according to level of care. *Arthr Res Ther* 2008; 10: R81.
26. Gracely RH, Petzke F, Wolf JM, Clauw DJ. Functional magnetic resonance imaging evidence of augmented pain processing in fibromyalgia. *Arthritis Rheum* 2002; 46: 1333-43.
27. Giesecke T, Williams DA, Harris RE, Cupps TR, Tian X, Tian TX, Gracely RH, Clauw DJ. Subgrouping of fibromyalgia patients on the basis of pressure-pain threshold and psychological factors. *Arthritis Rheum* 2003; 48: 2916-22.
28. Jensen KB, Kosek E, Petzke F, Carville S, Fransson P, Marcus H, Williams SC, Choy E, Giesecke T, Mainguy Y, Gracely R, Ingvar M. Evidence of dysfunctional pain inhibition in Fibromyalgia reflected in rACC during provoked pain. *Pain* 2009; 144: 95-100.
29. Jones AK, Brown WD, Friston KJ, Qi LY, Frackowiak RS. Cortical and subcortical localization of response to pain in man using positron emission tomography. *Proc Biol Sci* 1991; 244: 39-44.
30. Jones DK, Williams SC, Gasston D, Horsfield MA, Simmons A, Howard R. Isotropic resolution diffusion tensor imaging with whole brain acquisition in a clinically acceptable time. *Hum Brain Mapp* 2002; 15: 216-30.
31. Kim SG. Quantification of relative cerebral blood flow change by flow-sensitive alternating inversion recovery (FAIR) technique: application to functional mapping. *Magn Reson Med* 1995; 34: 293-301.
32. Kwiatek R, Barnden L, Tedman R, Jarrett R, Chew J, Rowe C, Pile K. Regional cerebral blood flow in fibromyalgia: single-photon-emission computed tomography evidence of reduction in the pontine tegmentum and thalami. *Arthritis Rheum* 2000; 43: 2823-33.
33. Kwong KK, Chesler DA, Weisskoff RM, Donahue KM, Davis TL, Ostergaard L, Campbell TA, Rosen BR. MR perfusion studies with T1-weighted echo planar imaging. *Magn Reson Med* 1995; 34: 878-87.
34. Law M. MR spectroscopy of brain tumors. *Magn Reson Imaging* 2004; 15:291-313.
35. Lee AD, Leow AD, Lu A, Reiss AL, Hall S, Chiang MC, Toga AW, Thompson, PM:. 3D pattern of brain abnormalities in Fragile X syndrome visualized using tensor-based morphometry. *Neuroimage* 2007; 34: 924-938.
36. Lentner C. (ed.). Geigy Scientific Tables, Vol. 4, p. 211, Ciba-Geigy Ltd, Basel, 1986.
37. Lepore, N, Brun C, Chou YY, Chiang MC, Dutton R, Hayashi K, Luders E, Lopez O, Aizenstein H, Toga A, Becker J, Thompson P: Generalized Tensor-Based Morphometry of HIV/AIDS Using Multivariate Statistics on Deformation Tensors. *IEEE Transactions on Medical Imaging* 2008; 27: 129-141.
38. Liu HL, Pu Y, Liu Y, Nickerson L, Andrews T, Fox PT, Gao JH. Cerebral blood flow measurement by dynamic contrast MRI using singular value decomposition with an adaptive threshold. *Magn Reson Med* 1999; 42:167-72.
39. Lutz J, Jager L, de Quervain D, Krauseneck T, Padberg F, Wichnalek M et al. White and gray matter abnormalities in the brain of patients with fibromyalgia: a diffusion-tensor and volumetric imaging study. *Arthritis Rheum* 2008; 58: 3960-9.
40. Maheshwari SR, Fatterpekar GM, Castillo M, Mukherji SK. Proton MR spectroscopy of the brain. *Semin Ultrasound CT MR*. 2000; 21: 434-51.

41. Mattson MP, Murrain M, Guthrie PB, and Kater SB. Fibroblast growth factor and glutamate: opposing roles in the generation and degeneration of hippocampal neuroarchitecture. *J. Neurosci* 1989; 9: 3728-3740.
42. Ostergaard L, Sorensen AG, Kwong KK, Weisskoff RM, Gyldensted C, Rosen BR. High resolution measurement of cerebral blood flow using intravascular tracer bolus passages. Part II: Experimental comparison and preliminary results. *Magn Reson Med* 1996; 36:726-36.
43. Petzke F, Clauw DJ, Ambrose K, Khine A, Gracely RH. Increased pain sensitivity in fibromyalgia: effects of stimulus type and mode of presentation. *Pain* 2003; 105: 403-13.
44. Provencher SW. Estimation of metabolite concentrations from localized in vivo proton NMR spectra. *Magn Reson Med* 1993; 30:672-9.
45. Purcell EU, Torrey HC, Pound RV. Resonance absorption by nuclear magnetic moments in a solid. *Phys Rev* 1946, 69:37-38.
46. Rajkowska G, Miguel-Hidalgo JJ. Gliogenesis and glial pathology in depression. *CNS Neurol Disord Drug Targets* 2007; 6: 219-33.
47. Rosen BR, Belliveau JW, Vevea JM, Brady TJ. Perfusion imaging with NMR contrast agents. *Magn Reson Med* 1990; 14: 249-65.
48. Sauter R, Schneider K, Wicklow K, Kolem H. Localized ¹H MRS of the human brain: Single-voxel versus CSI techniques. *J. Magn Reson Imaging* 1991, 1:241.
49. Silverstone PH, McGrath BM, Kim H. Bipolar disorder and myo-inositol: a review of the magnetic resonance spectroscopy findings. *Bipolar Disord* 2005; 7: 1-10.
50. Sundgren PC, Dong Q, Gómez-Hassan D, Mukherji SK, Maly P, Welsh R. Diffusion tensor imaging of the brain: review of clinical applications. *Neuroradiology* 2004; 46: 339-50.
51. Sundgren PC, Petrou M, Harris RE, Fan X, Foerster B, Mehrotra N et al. Diffusion-weighted and diffusion tensor imaging in fibromyalgia patients: a prospective study of whole brain diffusivity, apparent diffusion coefficient, and fraction anisotropy in different regions of the brain and correlation with symptom severity. *Acad Radiol* 2007; 14: 839-46.
52. Valet M, Gündel H, Sprenger T, Sorg C, Mühlau M, Zimmer C, Henningsen P, Tölle TR. Patients with pain disorder show gray-matter loss in pain-processing structures: a voxel-based morphometric study. *Psychosom Med* 2009; 71: 49-56.
53. Wolfe F, Smythe HA, Yunus MB, Bennet RM, Bombardier C, Goldenberg DL, Tugwell P, Campbell SM. The American College of Rheumatology 1990 Criteria for the classification of fibromyalgia. Report of the Multicenter Criteria Committee. *Arthritis Rheum* 1990; 33: 160-172.
54. Wolfe F, Ross K, Anderson J, Russell IJ. The prevalence and characteristics of fibromyalgia in the general population. *Arthritis Rheum* 1995; 38: 19-28.
55. Wolfe F. Fibromyalgia, the clinical syndrome. *Rheum Dis Clin NA* 1989; 15: 1-18.
56. Wolfe F, Anderson J, Harkness D, Bennet RM, Caro XJ, Goldenberg DL, Russell IJ, Wood PB, Ledbetter CR, Glabus MF, Broadwell LK, Patterson JC. Hippocampal metabolite abnormalities in fibromyalgia: correlation with clinical features. *J Pain* 2009; 10: 47-52.
57. Yildiz-Yesiloglu A, Ankerst DP. Neurochemical alterations of the brain in bipolar disorder and their implications for pathophysiology: a systematic review of the in vivo proton magnetic resonance spectroscopy findings. *Prog Neuropsychopharmacol Biol Psychiatry* 2006; 30: 969-95.

58. Yunus MB. Health status and disease severity in fibromyalgia: results of a six-center longitudinal study. *Arthritis Rheum* 1997; 40: 1571-79.

11. Annex. Some examples of new neuroimagen techniques in fibromyalgia.

Fig. 1. Magnetic Resonance Spectroscopy
Typical in-vivo proton magnetic resonance spectrum depicts the localization of major peaks for *N*-acetylaspartate (2.02 ppm), creatine/phosphocreatine complex (3.02 ppm), choline (3.22 ppm), glutamine and glutamate (2.1 to 2.55 ppm); and myo-inositol (3.56 ppm)

Fig. 2. Diffusion Tensor Imaging
Example of color-encoded fiber orientation maps. Fibers that are predominantly oriented left-right are shown in red, anterior-posterior fibers are shown in green, and superior-inferior fibers are shown in blue.

Fig. 3. Perfusion study

Neuroimaging

Edited by Cristina Marta Del-Ben

ISBN 978-953-307-127-5

Hard cover, 136 pages

Publisher Sciyo

Published online 17, August, 2010

Published in print edition August, 2010

Neuroimaging has become a crucial technique for Neurosciences. Different structural, functional and neurochemical methods, developed in recent decades, have allowed a systematic investigation on the role of neural substrates involved in functions performed by the central nervous system, whether normal or pathological. This book includes contributions from the general area of the neuroimaging to the understanding of normal functions and abnormalities of the central nervous system.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Javier Garcia-Campayo and Nicolas Fayed (2010). Clinical Magnetic Resonance Neuroimaging in Fibromyalgia, Neuroimaging, Cristina Marta Del-Ben (Ed.), ISBN: 978-953-307-127-5, InTech, Available from: <http://www.intechopen.com/books/neuroimaging/clinical-magnetic-resonance-neuroimaging-in-fibromyalgia>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2010 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike-3.0 License](https://creativecommons.org/licenses/by-nc-sa/3.0/), which permits use, distribution and reproduction for non-commercial purposes, provided the original is properly cited and derivative works building on this content are distributed under the same license.

IntechOpen

IntechOpen