We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Defining Interactions of Herbicides with Other Agrochemicals Applied to Peanut

¹David L. Jordan, Professor

²Gurinderbir S. Chahal, Graduate Research Assistant

³Sarah H. Lancaster, Former Graduate Research Assistant

⁴Joshua B. Beam, Former Graduate Research Assistant

⁵Alan C. York, William Neal Reynolds Professor Emeritus

¹*Department of Crop Science, North Carolina State University,
Box 7620, Raleigh, NC 27695-7620.*

²*Department of Crop Science, North Carolina State University,
Box 7620, Raleigh, NC 27695-7620.*

³*Department of Crop Science, North Carolina State University,
Box 7620, Raleigh, NC 27695-7620.*

*Current position: Assistant Professor, Department of Plant
and Soil Sciences, Oklahoma State University, 368
Agricultural Hall, Stillwater, OK 74078-6028.*

⁴*Department of Crop Science, North Carolina State University
Box 7620, Raleigh, NC 27695-7620.*

*Current position: North Carolina Business Manager, Sunny
Ridge Farm Inc., 1900 5th Street Northwest,
Winter Haven, FL 33881-2106.*

⁵*Department of Crop Science, North Carolina State University
Box 7620, Raleigh, NC 27695-7620.
United States of America*

1. Introduction

Management strategies to protect peanut (*Arachis hypogaea* L.) from pest damage require multiple applications of fungicides, herbicides, and insecticides. Additionally, micronutrients and plant growth regulators are often applied to improve nutrient balance and to manage peanut growth and development. Over fifty active ingredients can be used to manage pests in peanut, often with more than one formulated product commercially available. Timing of application of pesticides, micronutrients, and plant growth regulators often coincide during the growing season, and co-application of these agrochemicals is desirable if pesticide, micronutrient, and plant growth regulator performance and peanut tolerance are not compromised. In addition to potential interactions related to physiological effects on plants and other organisms, application variables such as commercial formulation, adjuvant, water quality, and environmental stress can affect agrochemical compatibility.

Physical compatibility, in particular formation of precipitates in spray tanks and equipment, is a concern for farmers when co-applying agrochemicals. Defining potential interactions among these agrochemicals is important in developing appropriate weed management programs and implementing integrated pest management strategies for peanut.

Considerable research has been conducted during the last four decades to define interactions among agrochemicals (Barrett, 1993; Green, 1989; Green and Bailey, 1987; Hatzios and Penner, 1985; Putnam and Penner, 1974). Most of these reviews are focused on interactions of herbicides in mixture with other herbicides, fungicides, insecticides, nematicides, and adjuvants, in general, but not for a particular crop. Some of these reviews summarized the mechanisms responsible for the interactions of herbicides with other agrochemicals and the statistical methodology for characterization of agrochemical combinations (Barrett, 1993; Green, 1989; Hatzios and Penner, 1985; Jianhua *et al.*, 1995). Since these reviews were published, many of new agrochemicals have received registration for different crops and for other uses. Defining interactions of these new agrochemicals is important when developing pest management strategies for a cropping system. This chapter reviews some of the interactions discussed in the earlier work, but also elucidates the interactions and/or compatibility of herbicides with other agrochemicals used in peanut production systems.

2. Peanut production systems

Mechanized production systems utilize a wide range of agrochemicals to manage peanut growth and development and minimize the impact of pests on peanut yield and quality (Lynch and Mack, 1995; Sherwood *et al.*, 1995; Wilcut *et al.*, 1995). Pests that can potentially impact peanut are diverse (Table 1). Yield loss from weed interference or from damage caused by insects, diseases, and nematodes can be substantial in peanut if pest control strategies are not implemented in a timely manner.

Monocotyledonous weeds, including annual and perennial grasses and sedges, as well dicotyledonous weeds, are prevalent in peanut production systems in the United States (Webster, 2009; Wilcut *et al.*, 1995). Comprehensive herbicide programs, in combination with appropriate cultural practices, are employed to manage weeds and minimize interference and subsequent yield loss (Wilcut *et al.*, 1987a 1987b 1990 1995). Herbicides are often applied in mixtures either prior to planting (preplant incorporated in conventional tillage or preplant to emerged weeds in reduced tillage), immediately following planting (preemergence), or after peanut and weeds have emerged (postemergence) (Burke *et al.*, 2004; Clewis *et al.*, 2007; Richburg *et al.*, 1995 1996; Wilcut *et al.*, 1994a 1994b 1995).

Agrochemicals with efficacy against insects and plant parasitic nematodes are often applied in the seed furrow at planting and include organophosphate and carbamate insecticides (Brecke *et al.*, 1996; Drake *et al.*, 2009; Funderburk *et al.*, 1998; Minton *et al.*, 1990; Minton and Morgan, 1974; Riley *et al.*, 1997). In-furrow insecticides also reduce incidence of tomato spotted wilt of peanut (caused by tomato spotted wilt virus, a *Tospovirus* vectored by several species of thrips) (Brown *et al.*, 2003; Hurt *et al.*, 2003). Pyrethroid insecticides are often applied to peanut foliage to control beet armyworm, corn earworm, fall armyworm, potato leaf hopper, and two-spotted spider mites. Chlorpyrifos can be applied at pegging, 45 to 70 days after peanut emergence, to control lesser cornstalk borer (Mack *et al.*, 1989 1991) and southern corn rootworm (Brandenburg and Herbert, 1991; Chapin and Thomas, 1993).

Depending on environmental and edaphic conditions and a range of agronomic and pest management practices, application of insecticides may be needed throughout the growing season to protect peanut from pest damage.

Disease, caused by viruses, bacteria, or fungi, can reduce peanut yield considerably when not controlled (Sherwood *et al.*, 1995). Fungicides are applied routinely to peanut to control foliar-borne diseases, including early leaf spot, late leaf spot, and web blotch (Brenneman *et al.*, 1994; Culbreath *et al.*, 2008; Shew and Waliyar, 2005). Fungicides are also applied to control the soil-borne disease stem rot and Sclerotinia blight (Brenneman *et al.*, 1994; Culbreath *et al.*, 2008; Smith *et al.*, 1992). Although variation is noted among geographical regions, years, and environmental conditions, during a typical growing season fungicides are applied either singly or in combination beginning approximately 45 days after peanut emergence and continuing throughout the remainder of the growing season, which can approach 135 or more days (Sherwood *et al.*, 1995; Smith and Littrell, 1980). Fungicide programs to control early and late leaf spot and stem rot often include bi-weekly sprays during this period. Fungicides applied to control these diseases provide protection for a period of two weeks under most environmental conditions (Shew and Waliyar, 2005). The soil fumigant metam sodium is often applied to peanut to control *Cylindrocladium* black rot (Cline and Beute, 1986). A period of at least two weeks between fumigation and peanut planting is required to allow the fumigant to dissipate, making weed control prior to planting challenging under some environmental conditions, especially excessive rainfall, that allow weeds to emerge between fumigation and planting operations (Van Gundy and McKenry, 1977).

The micronutrients boron and manganese are applied routinely to optimize peanut growth and development and, in the case of boron, to ensure proper kernel development (Gascho and Davis, 1995; Harris and Brolman, 1966; Powell *et al.*, 1996). Because peanut is often grown on coarse-textured soils, boron can be deficient due to leaching. Single, and in some cases, multiple applications of boron-containing foliar solutions are applied 45 to 70 days after peanut emergence (Gascho and Davis, 1995). Manganese deficiency occurs frequently in peanut because of liming to achieve a target soil pH above 6.0. Correcting a manganese deficiency is achieved by foliar applications when visible symptoms become apparent, although some growers apply manganese irrespective of plant symptomology (Powell *et al.*, 1996).

Excessive vine growth of peanut can reduce row visibility at digging and vine inversion (Mitchem *et al.*, 1996). Prohexadione calcium is currently the only plant growth regulator applied to manage vine growth in order to facilitate efficient digging. Prohexadione calcium inhibits gibberellin biosynthesis in responsive plants (Grossman *et al.*, 1994) and is applied when 50% of vines from adjacent peanut rows have met, and an application is repeated 2 to 3 weeks later (Mitchem *et al.*, 1996). This timing of application is generally 70 to 90 days after peanut emergence (Mitchem *et al.*, 1996). In addition to prohexadione calcium, a wide range of products are available at the distributor level that contain micronutrient combinations, synthetic plant growth regulators, and other ingredients perceived to have value. Many of these products are not applied routinely to peanut.

3. Agrochemicals used in peanut

A diversity of pesticide active ingredients is available for peanut (Table 2) (Brandenburg, 2010; Jordan, 2010; Shew, 2010). Currently, 19 herbicide active ingredients, 16 insecticide

active ingredients, and 20 fungicide active ingredients representing the major modes of action can be applied during the peanut growing season. Three fumigants, two micronutrients, and one plant growth regulator are registered for use in peanut. Within herbicide, insecticide, fungicide, and fumigant categories, a range of formulated products are available for most active ingredients. These products are often manufactured and sold through distributor networks and are numerous. Additionally, spray adjuvants are recommended with some, but not all, agrochemicals to increase performance and compatibility.

Presence of biotic and abiotic stresses often occur simultaneously during the peanut growing season, and timing of application for many agrochemicals overlap (Table 3). Practitioners prefer limiting the number of trips across fields in order to increase efficiency of managing peanut. This approach is preferable because of convenience, savings in time, reduced application costs, and freeing labor for other operations. Additionally, applying multiple pesticides with different modes of action is an important resistance management strategy for pests (Brandenburg, 2010; Jordan, 2010; Shew, 2010). This approach is feasible as long as adverse interactions, primary increased crop injury or decreased pest control, do not occur. Defining interactions among agrochemicals is important in assisting growers and their advisors as they make decisions on co-application of these products.

4. Herbicide – Herbicide interactions

A considerable amount of research has been conducted to define interactions among herbicides used in peanut. Herbicides applied in combination either preplant incorporated or preemergence generally increase the spectrum of weed control or the length of residual weed control (Wilcut *et al.*, 1987b 1995). For example, pendimethalin is often applied in combination with alachlor, dimethenamid-*P*, metolachlor, or *S*-metolachlor to improve early season weed control (Bridges *et al.*, 1984; Wehtje *et al.*, 1988; Wilcut *et al.*, 1994b 1995; Wilcut and Swann, 1990). Alachlor, dimethenamid-*P*, metolachlor, or *S*-metolachlor can be applied with diclosulam, flumioxazin, or imazethapyr preemergence to enhance weed control with a single application (Clewis *et al.*, 2007; Grichar *et al.*, 1992, 1996, 2000, 2008; Scott *et al.*, 2001). Combinations of preplant incorporated or preemergence herbicides currently registered for use in peanut have not been shown to increase peanut injury over either herbicide component applied alone (Wilcut *et al.*, 1995). However, several herbicides that are no longer registered for peanut increased peanut injury when co-applied as compared to the herbicides applied alone (Wilcut *et al.*, 1995).

In reduced tillage systems, herbicides are needed to control winter weeds and summer annual weeds that have emerged prior to planting peanut. These herbicide applications include glyphosate, paraquat, or 2,4-D alone or in combinations with other herbicides. Combinations of glyphosate and 2,4-D broaden the spectrum of weed control compared with each herbicide applied alone (Flint and Barrett, 1989a). However, in some instances, 2,4-D can negatively affect efficacy of glyphosate, but this interaction is typically noted only on grass weeds (Flint and Barrett, 1989b). Efficacy of paraquat is generally not negatively affected by 2,4-DB (Wehtje *et al.*, 1992a). Glyphosate and paraquat can also be applied with herbicides that provide residual weed control. This approach is designed to control emerged weeds and provide residual weed control prior to and following planting (Wilcut *et al.*, 1995).

Paraquat is often applied at peanut emergence or up to 28 days after peanut emergence (Carley *et al.*, 2009; Wilcut *et al.*, 1990). Other non-residual herbicides such as bentazon or acifluorfen plus bentazon as well as residual herbicides such as alachlor, diclosulam, dimethenamid-*P*, imazethapyr, metolachlor, or *S*-metolachlor are applied postemergence to broaden the spectrum of control (Askew *et al.*, 1999; Bailey *et al.*, 1999; Grey *et al.*, 2002; Grichar and Colburn, 1996). Injury associated with paraquat can be reduced by co-application with bentazon (Jordan *et al.*, 2003b; Wehtje *et al.*, 1992b). However, the chloroacetamide herbicides alachlor, dimethenamid-*P*, metolachlor, or *S*-metolachlor applied with paraquat can increase peanut injury (Jordan *et al.*, 2003b). Diclosulam and imazethapyr did not affect injury potential from paraquat (Jordan *et al.*, 2003b). Weed control with these herbicide combinations generally increases depending on the weed species and size of the weed (Wilcut *et al.*, 1995). For example, bentazon and imazethapyr co-applied can increase control of emerged common cocklebur and yellow nutsedge, while control of annual grasses by paraquat can be reduced when paraquat is co-applied with bentazon (Wehtje *et al.*, 1992b; Wilcut *et al.*, 1994b). Residual control by chloroacetamide herbicides, diclosulam, and imazethapyr was not affected by paraquat applied alone or with bentazon (Grichar *et al.*, 2000; Wilcut *et al.*, 1995).

Co-application of postemergence herbicides with efficacy against dicotyledonous weeds and sedges generally increases control of weeds or broadens the spectrum of control compared with components of the mixture applied alone (Green, 1989; Hatzios and Penner, 1985; Jianhua *et al.*, 1995; Wilcut *et al.*, 1995). In contrast, efficacy of clethodim and sethoxydim, often referred to as graminicides, can be reduced when applied in mixture with herbicides that control dicotyledonous weeds and sedges (Culpepper *et al.*, 1998 1999; Grichar, 1991; Jordan and York, 1989; Minton *et al.*, 1989; Mueller *et al.*, 1989; Myers and Coble, 1992; Vidrine *et al.*, 1995). The interaction of bentazon and sethoxydim is one of the most notable examples of reduced graminicide efficacy caused by a herbicide that controls dicotyledonous plants and sedges (Rhodes and Coble, 1984a 1984b; Wanamarta and Penner, 1989; Wanamarta *et al.*, 1989). Annual and perennial grass control by sethoxydim is reduced by bentazon through reduced absorption of sethoxydim into grasses (Rhodes and Coble, 1984b; Wanamarta and Penner, 1989; Wanamarta *et al.*, 1989). The mechanism of reduced control is associated with physical interactions of the herbicides in the spray solution prior to reaching the target weed (Penner, 1989; Thelen *et al.*, 1995). Acifluorfen and imazethapyr also can reduce efficacy of clethodim and sethoxydim (Burke and Wilcut, 2003; Grichar, 1991; Lassiter and Coble, 1987; Myers and Coble, 1992). In contrast to reduced grass control when these herbicides are co-applied, control of dicotyledonous plants and sedges is not reduced by clethodim and sethoxydim (Dotray *et al.*, 1993; Holshouser and Coble, 1990; Isaacs *et al.*, 2003). Efficacy of clethodim can also be reduced by acifluorfen, acifluorfen plus bentazon, bentazon, imazethapyr, imazapic, lactofen, and 2,4-DB (Grichar *et al.*, 2002; Myers and Coble, 1992; York *et al.*, 1993). The magnitude of reduced efficacy can be minimized or eliminated by applying the herbicides sequentially, increasing the graminicide rate, or applying more efficacious adjuvants (Burke *et al.*, 2004; Jordan, 1995; Myers and Coble, 1992; Wanamarta and Penner, 1989; Wanamarta *et al.*, 1989). Grass species, plant size, and plant stress also can affect the magnitude of negative interactions (Green, 1989; Hatterman-Valenti *et al.*, 2006). York and Wilcut (1995) reported that bentazon reduced control of yellow and purple nutsedge by imazethapyr.

Chloroacetamide herbicides can be applied postemergence without injuring peanut (Grichar *et al.*, 1996, 2008; Jordan *et al.*, 2003b). While these herbicides provide residual control of grasses and some dicotyledonous and sedge weeds, they do not control weeds that have emerged (Foy and Witt, 1997; Grichar *et al.*, 2000; Richburg *et al.*, 1995). These herbicides can be applied with herbicides that have efficacy against emerged weeds. Dimethenamid-*P* and *S*-metolachlor did not reduce grass control by the graminicides clethodim or sethoxydim or the dicotyledonous and sedge herbicides acifluorfen, acifluorfen plus bentazon, or imazapic (Grichar *et al.*, 2000; Wilcut *et al.*, 1995). However, visible injury caused by acifluorfen increased when acifluorfen was applied with chloroacetamide herbicides (Jordan *et al.*, 2003b). Johnson *et al.* (1993) reported that injury from postemergence application of paraquat was not increased when following several chloroacetamide herbicides applied at planting, in contrast with injury observed when the herbicides were co-applied.

5. Herbicide – Insecticide interactions

Timing of application of herbicides and insecticides overlap during much of the growth cycle of peanut (Table 3). As with other crops, potential interactions between herbicides and insecticides applied in the seed furrow to control thrips and suppress plant parasitic nematodes can occur (Hauser *et al.*, 1976 1981). Acephate and aldicarb applied in the seed furrow at planting did not affect injury potential of peanut following postemergence application of acifluorfen plus bentazon or bentazon; however, the insecticide phorate applied in the seed furrow enhanced visible injury associated with bentazon, although this injury was generally transient (Swann and Herbert, 1999). Although interactions of nicosulfuron (Bailey and Kapusta, 1994; Morton *et al.*, 1994; Rahman and James, 1993) and pyriithiobac-sodium (Allen and Snipes, 1995) increased injury in corn (*Zea mays* L.) and cotton (*Gossypium hirsutum* L.), respectively. However, chlorpyrifos applied at planting did not affect peanut response to diclosulam, *S*-metolachlor, or flumioxazin applied preemergence or acifluorfen, acifluorfen plus bentazon, imazapic, or paraquat plus bentazon applied postemergence (Jordan *et al.*, 2008). Efficacy of graminicides can be affected by insecticides applied to peanut. Carbaryl and dimethoate applied postemergence in combination with sethoxydim reduced annual grass control; no adverse effect was noted when acephate was mixed with sethoxydim (Byrd and York, 1988). Pyrethroid insecticides did not affect efficacy of postemergence herbicides (Allen and Snipes, 1995).

6. Herbicide – Fungicide interactions

Similar to herbicides and insecticides, timing of application of postemergence herbicides and fungicides to control foliar and soil-borne diseases overlap considerably during the peanut growing season (Table 3). Fungicides are applied beginning approximately 45 days after peanut emergence and can be applied until a few weeks prior to digging and vine inversion. Efficacy of clethodim and sethoxydim can be reduced by co-application with copper-containing fungicides or azoxystrobin, chlorothalonil, and pyraclostrobin (Jordan *et al.*, 2003a; Lancaster *et al.*, 2005a 2008). Fluazinam and tebuconazole did not reduce grass control compared with graminicides applied alone (Jordan *et al.*, 2003a; Lancaster *et al.*, 2005a 2005b). Efficacy of herbicides that control dicotyledonous and sedge weeds is not generally

affected by fungicides (Jordan *et al.*, 2003a). As was noted for interactions of herbicides, weed species and size and plant stress can affect the magnitude of interactions between herbicides and fungicides (Jordan *et al.*, 2003a).

Although not used in peanut, efficacy of glyphosate was not affected by azoxystrobin, pyraclostrobin, or tebuconazole (Grichar and Prostko, 2009). Weed control by metribuzin, rimsulfuron, and thifensulfuron-methyl applied to tomato (*Lycopersicon esculentum* Mill.) was not affected by azoxystrobin or pyraclostrobin (Robinson and Nurse, 2008). However, pyraclostrobin increased tomato injury from thifensulfuron-methyl when co-applied (Robinson and Nurse, 2008). Chlorothalonil increased persistence of metolachlor in soil although cyproconazole, flutriafol, and tebuconazole did not affect dissipation of metolachlor (White *et al.*, 2009).

7. Herbicide – Micronutrient interactions

Boron and manganese are the primary micronutrients applied to peanut. Occasionally, these can affect herbicide performance. For example, efficacy of clethodim and imazethapyr was reduced by micronutrients for some, but not all, weeds evaluated (Jordan *et al.*, 2006; Lancaster *et al.*, 2005b).

8. Herbicide – Plant growth regulator interactions

Prohexadione calcium is the primary plant growth regulator available for use in peanut. Efficacy of the herbicides acifluorfen, acifluorfen plus bentazon, bentazon, imazethapyr, imazapic, lactofen, and 2,4-DB was not affected by prohexadione calcium (Beam *et al.*, 2002). However, other plant growth regulator products developed by agrochemical distributor chains are numerous and have not been evaluated sufficiently to make recommendations on compatibility with herbicides.

9. Co-application of multiple components

The previous discussion focused on co-applications that have only two components. However, there is considerable interest in compatibility of three or more pesticides, micronutrients, adjuvants, or plant growth regulators and their impact on pest control and crop management. With respect to weed control, efficacy of clethodim, sethoxydim, and 2,4-DB were compared when these herbicides were applied alone or with fungicides and insecticides (Jordan *et al.*, 2003a; Lancaster *et al.*, 2005a 2005b). Although results often supported previous findings with components from two groups of pesticides, no clear relationships were established with respect to combinations of three or more pesticides (Lancaster *et al.*, 2005a 2005b). More recently, research is being conducted to compare herbicide efficacy with mixtures containing various levels of fungicide, insecticide, micronutrient, or adjuvant (Chahal *et al.*, 2009a 2009b).

10. Herbicide effects on other agrochemicals

The focus of this review has been the impact of agrochemicals used in peanut on herbicide efficacy. However, defining the impact of herbicides on insect and disease control and

response of peanut to micronutrient and plant growth regulator applications is important. Preliminary research has shown that the herbicides clethodim and 2,4-DB do not affect performance of chlorothalonil, pyraclostrobin, or the prepackage combination of prothioconazole plus tebuconazole (Chahal *et al.*, 2009a 2009b). Paraquat and 2,4-DB did not affect chlorothalonil efficacy (Choate *et al.*, 1998). Katan and Eshel (1973) discussed possible mechanisms of interactions among herbicides and pathogens. With respect to peanut, Baysinger *et al.* (1999) reported that sporulation of early leaf spot was reduced by acifluorfen and lactofen. In-vitro, chlorothalonil efficacy against early blight (*Alternaria solani*) was reduced by metribuzin while susceptibility of *Pseudo cercospora herpotrichoides* to cyproconazole increased following exposure to dicamba, bromoxynil, or ioxynil (Kataria and Gisi, 1990; Levesque and Rahe, 1992).

The influence of the postemergence herbicides clethodim, imazapic, lactofen, and 2,4-DB on boron and manganese absorption into peanut leaves was evaluated, and results suggested that while herbicides could affect accumulation of boron and manganese in leaf tissue, the adjuvant associated with these herbicides may have been the primary factor in influencing absorption (Jordan *et al.*, 2006 2009a).

11. Application variables that can affect interactions

A wide range of application variables can affect interactions of herbicides with other agrochemicals. Adjuvant selection, herbicide rate, commercial formulation, active ingredient, length of time between applications of components, spray volume, water quality, weed species, and environmental conditions can affect interactions of agrochemicals. For example, the negative effect of bentazon was reduced by including ammonium sulfate and other more efficacious adjuvants with clethodim and sethoxydim (Jordan, 1995; Jordan and York, 1989; Penner, 1989; Wanamarta and Penner, 1989; York *et al.*, 1990). Applying a higher rate of the herbicide that may be adversely affected can compensate for interactions (Chernicky and Slife, 1986; Rhodes and Coble, 1984a 1984b). Differential response to clethodim was noted when applied with different formulations of chlorothalonil (Jordan *et al.*, 2003a). Increasing the interval between applications of components of the mixture can overcome negative interactions, especially herbicide-herbicide interactions (Green, 1989; Grichar and Boswell, 1987; Putnam and Ries, 1967). Applying graminicides in higher spray volumes can exacerbate the negative influence of herbicides and fungicides on weed control by graminicides (Buhler and Burnside, 1984; Jordan *et al.*, 2003a; Kells and Wanamarta, 1987). Water quality, in particular presence of cations that can form complexes in the spray solution, can influence the propensity of herbicides to perform poorly in pesticide combinations (Buhler and Burnside, 1983a 1983b; Hatzios and Penner, 1985; Sandberg *et al.*, 1978; Stahlman and Phillips, 1979; Thelen *et al.*, 1995; Whisenant and Bovey, 1993; Wills and McWhorter, 1985).

Environmental conditions that affect plant response to agrochemicals can influence the magnitude of interactions. Negative effects of interactions associated with efficacy of systemic herbicides, especially graminicides, are exacerbated when grasses are stressed and physiological processes that reduce absorption and translocation occur (Burke *et al.*, 2004; Burke and Wilcut, 2003; Green, 1989; Wanamarta and Penner, 1989; Wanamarta *et al.*, 1989).

12. Challenges of defining interactions and making recommendations

Major challenges in recommending practices associated with interactions of herbicides with other agrochemicals include the diversity of products available, the diversity of weeds present in the field that can vary in response to herbicides, differences in water quality within and across production regions, and adjuvant recommendations associated with herbicides versus other agrochemicals. Compounding these complex variables is the unpredictable response often observed due to environmental conditions. Additionally, new active ingredients are being marketed that have not been evaluated for possible interactions, and as patents expire, some of these active ingredients are formulated differently from the product receiving the initial registration. The extensive number of possible combinations, especially when multiple components are considered, is challenging from a research standpoint when considering the logistics of research trials needed to address possible interactions. Although attempts are made to establish research trials in a manner similar to practitioner operations, techniques associated with maintaining spray solutions and applying materials with small-plot equipment differ from commercial applications and adds to the challenges in research.

One reasonable criticism of the current approach to defining interactions, which is dictated by the number of combinations and the logistics of experimentation, is the lack of defining the impact of interactions on the larger context of the production system. For example, when a reduction in weed control by a fungicide occurs, how detrimental to peanut yield and economic value is this reduction when considering alternatives to prevent the interaction from occurring? Also, dose response curves are often used to define interactions of pesticides, and while these can be more informative than selection of a single rate of the components in the mixture, the number of treatment combinations required for this approach is often not feasible because of resource constraints.

13. Future research

A considerable knowledge base has been developed to define interactions of herbicides with other agrochemicals with respect to weed control in peanut. However, the effect of herbicides on performance of fungicides and insecticides is limited but no less important than defining impacts on herbicide efficacy. As new active ingredients and new formulations of active ingredients become available, additional research will be needed to define interactions among these agrochemicals. Although interactions of herbicide-herbicide combinations have been defined broadly and in some cases in detail, research elucidating the mechanism of reduced control associated with co-application of fungicides, insecticides, or plant growth regulator and micronutrients is limited (Duke *et al.*, 2007). Finally, determining the impact of interactions in the overall production system would be beneficial.

Pest	Latin bionomical and authority
Weeds^a	
Bermudagrass	<i>Cynodon dactylon</i> (L.) Pers.
Bristly starbur	<i>Acanthospermum hispidum</i> DC.
Broadleaf signalgrass	<i>Urochloa platyphylla</i> (Nash) R.D. Webster
Common cocklebur	<i>Xanthium strumarium</i> L.
Common lambsquarters	<i>Chenopodium album</i> L.
Common ragweed	<i>Ambrosia artemisiifolia</i> L.
Crabgrass spp.	<i>Digitaria</i> spp.
Crowfootgrass	<i>Dactyloctenium aegyptium</i> (L.) Willd.
Eclipta	<i>Eclipta prostrata</i> L.
Florida beggarweed	<i>Desmodium tortuosum</i> (Sw.) DC.
Florida pusley	<i>Richardia scabra</i> L.
Goosegrass	<i>Eleusine indica</i> (L.) Gaertn.
Hairy indigo	<i>Indigofera hirsuta</i> Harvey
Jimsonweed	<i>Datura stramonium</i> L.
Johnsongrass	<i>Sorghum halepense</i> (L.) Pers.
Morningglory spp.	<i>Ipomoea</i> spp.
Nutsedge spp.	<i>Cyperus</i> spp.
Palmer amaranth	<i>Amaranthus palmeri</i> S. Wats.
Pigweed spp.	<i>Amaranthus</i> spp.
Prickly sida	<i>Sida spinosa</i> L.
Sicklepod	<i>Senna obtusifolia</i> (L.) H.S. Irwin & Barneby
Spurge spp.	<i>Chamaesyce</i> spp.
Texas millet	<i>Urochloa texana</i> (Buckl.) R. Webster
Tropic croton	<i>Croton glandulosus</i> var. septentrionalis Muell.-Arg.
Insects^b	
Beet armyworm	<i>Spodoptera exigua</i> Hübner
Corn earworm	<i>Heliothis zea</i> Boddie
Lesser cornstalk borer	<i>Elasmopalpus lignosellus</i> Zeller

Pest	Latin bionomical and authority
Southern corn rootworm	<i>Diabrotica undecimpunctata</i> Howardi
Thrips	<i>Frankliniella</i> spp.
Diseases^c	
Aspergillus crown rot	<i>Aspergillus niger</i>
Botrytis	<i>Botrytis cinerea</i> Pers.
<i>Cylindrocladium</i> black rot	<i>Cylindrocladium parasiticum</i> Crous, Wingfield, and Alfenas
Early leaf spot	<i>Cercospora arachidicola</i> Hori
Late leaf spot	<i>Cercosporidium personatum</i> Berk. & Curtis
Pythium	<i>Pythium</i> spp.
Rhizoctonia limb rot	<i>Rhizoctonia solani</i> Kuhn
Sclerotinia blight	<i>Sclerotinia minor</i> Jagger
Spotted wilt	Tomato spotted wilt, caused by a <i>Tospovirus</i>
Stem rot	<i>Sclerotium rolfsii</i> Sacc.
Web blotch	<i>Phoma arachidicola</i> Marasas, Pauer, and Boerema
Plant parasitic nematodes^c	
Lesion nematode	<i>Pratylenchus brachyurus</i>
Northern root knot	<i>Meloidogyne hapla</i>
Peanut root knot	<i>Meloidogyne arenaria</i>
Ring	<i>Criconebella ornata</i>
Sting	<i>Belonolaimus longicaudatus</i>
Two-spotted spider mite^b	<i>Tetranychus urticae</i> Koch

^aWebster, T. M. 2009. Weed survey – southern states. Proc. South. Weed Sci. Soc. 62:509-524.

^bBrandenburg, R. L. 2010. Peanut insect management. Pages 85-102 in 2010 Peanut Information. North Carolina Cooperative Extension Service Publication AG-331.

^cShew, B. B. 2010. Peanut disease management. Pages 103-130 in 2010 Peanut Information. North Carolina Cooperative Extension Service Publication AG-331.

Table 1. Common and scientific names for major pests found in peanut in the United States.

Herbicides	Insecticides	Fungicides	Fumigants	Plant growth regulators
Acifluorfen	Acephate	Azoxystrobin	Dichloropropene plus chloropicrin	Prohexadione calcium
Alachlor	Aldicarb	Basic copper sulfate	Metam sodium	
Bentazon	<i>Bacillus thuringiensis</i>	Boscalid	1,3 dichloropropene	
Carfentrazone	Carbaryl	Chlorothalonil		
Clethodim	Chlorpyrifos	Dodine		
Diclosulam	Disulfoton	Fludioxomil		
Dimethenamid- <i>p</i>	Esfenvalerate	Fluoxastrobin		
Ethafluralin	Fenpropathrin	Flutolanil		
Flumioxazin	Gamma-cyhalothrin	Mancozeb		
Glyphosate	Indoxacarb	Mancozeb and copper hydroxide		
Imazapic	Lambda-cyhalothrin	Mefenoxam		
Imazethapyr	Malathion	Metconazole		
Lactofen	Methomyl	Pentachloronitrobenzene (PCNB)		
Metolachlor	Phorate	Propiconazole		
Paraquat	Propargite	Prothioconazole		
Pendimethalin	Spinosad	Pyraclostrobin		
Sethoxydim		Sulfur		
S-metolachlor		Tebuconazole		
Sulfentrazone		Thiophanate methyl		
2,4-DB		Trifloxystrobin		

^aBrandenburg, R. L. 2010. Peanut insect management. Pages 85-102 in 2010 Peanut Information. North Carolina Cooperative Extension Service Publication AG-331.

^bJordan, D. L. 2010. Weed management in peanuts. Pages 55-83 in 2010 Peanut Information. North Carolina Cooperative Extension Service Publication AG-331.

^cShew, B. B. 2010. Peanut disease management. Pages 103-130 in 2010 Peanut Information., Peanut Information 2010. North Carolina Cooperative Extension Service AG-331.

Table 2. Pesticide active ingredients registered for use in peanut in the United States during 2010.^{a,b,c}

	April	May	June	July	August	September
Weeds						
Dicotyledonous						
Monocotyledonous						
Insects						
Beet armyworm						
Corn earworm						
Lesser cornstalk borer						
Southern corn rootworm						
Tobacco thrips						
Diseases						
Aspergillus crown rot						
Botrytis						
Cylindrocladium black rot						
Cercospora spp.						
Pythium						
Rhizoctonia limb rot						
Sclerotinia blight						
Spotted wilt						
Stem rot						
Web blotch						
Plant parasitic nematodes						
Two-spotted spider mite						
Nutrient deficiency						
Boron						
Manganese						
Peanut vine management						
Prohexadione calcium						

Table 3. Biotic and abiotic stresses and approximate timing of management that can occur in peanut during the growing season in the United States.

14. References

Allen, R. L. and C. E. Snipes. 1995. Interactions of foliar insecticides applied with pyrithiobac. Weed Technol. 9:512-517.

- Askew, S. D., J. W. Wilcut, and J. R. Cranmer. 1999. Weed management in peanut (*Arachis hypogaea*) with flumioxazin preemergence. *Weed Technol.* 13:594-598.
- Bailey, J. A. and G. Kapusta. 1994. Soil insecticide and placement influence corn (*Zea mays*) tolerance to nicosulfuron. *Weed Technol.* 8:598-606.
- Bailey, W. A., J. W. Wilcut, D. L. Jordan, C. W. Swann, and V. B. Langston. 1999. Response of peanut (*Arachis hypogaea*) and selected weeds to diclosulam. *Weed Technol.* 13:771-776.
- Barret, M. 1993. Interactions of herbicides and other agrochemicals in plants: Interactions in mixtures with other herbicides and with safeners, fungicides, insecticides, and nematicides. Pages 113-132 in J. Altman (ed.) *Pesticide Interactions in Crop Production: Beneficial and Deleterious Effects*. CRC press Boca Raton, FL.
- Baysinger, J. A., H. A. Melouk, and D. S. Murray. 1999. Acifluorfen, acifluorfen plus 2,4-DB, and lactofen decreased the sporulation of early leaf spot lesions. *Peanut Sci.* 26:4-8.
- Beam, J. B., D. L. Jordan, A. C. York, J. E. Bailey, T. G. Isleib, and T. E. McKemie. 2002. Interaction of prohexadione calcium with agrichemicals applied to peanut (*Arachis hypogaea* L.). *Peanut Sci.* 29:29-35.
- Brandenburg, R. L. 2010. Peanut insect management. Pages 85-102 in 2010 Peanut Information. North Carolina Cooperative Extension Service Publication AG-331.
- Brandenburg, R. L. and D. A. Herbert, Jr. 1991. Effect of timing on prophylactic treatments for southern corn rootworm (Coleoptera:Chrysomelidae) in peanut. *J. Econ. Entomol.* 84: 1894-1898.
- Brecke, B. J., J. E. Funderburk, I. D. Teare, and D. W. Gorbet. 1996. Interactions of early-season herbicide injury, tobacco thrips injury, and cultivar on peanut. *Agron. J.* 88:14-18.
- Brenneman, T. B., H. R. Sumner, L. R. Chandler, J. M. Hammond, and A. K. Culbreath. 1994. Effect of application techniques on performance of propiconazole for peanut disease control. *Peanut Sci.* 21:134-138.
- Bridges, D. C., R. H. Walker, J. A. McGuire, and N. R. Martin. 1984. Efficiency of chemical and mechanical methods for controlling weeds in peanuts (*Arachis hypogaea*). *Weed Sci.* 32:584-591.
- Brown, S., J. Todd, A. Culbreath, J. Baldwin, J. Beasley, B. Kemerait, and E. Prostko. 2003. Minimizing spotted wilt of peanut. University of Georgia Cooperative Extension Service Bulletin 1165.
- Buhler, D. D. and O. C. Burnside. 1983a. Effect of spray components on glyphosate toxicity to annual grasses. *Weed Sci.* 31:124-130.
- Buhler, D. D. and O. C. Burnside. 1983b. Effect of water quality, carrier volume, and acid on glyphosate phytotoxicity. *Weed Sci.* 31:163-169.
- Buhler, D. D. and O. C. Burnside. 1984. Effect of application factors on the phytotoxicity of fluazifop-butyl, haloxyfop-methyl, and sethoxydim. *Weed Sci.* 32:574-583.
- Burke, I. C., A. J. Price, J. W. Wilcut, D. L. Jordan, A. S. Culpepper, and J. T. Ducar. 2004. Annual grass control in peanut (*Arachis hypogaea*) with clethodim and imazapic. *Weed Technol.* 18:88-92.
- Burke, I. C. and J. W. Wilcut. 2003. Physiological basis for antagonism of clethodim by imazapic on goosegrass (*Eleusine indica* (L.) Gaertn.). *Pestic. Biochem. Physiol.* 76:37-45.

- Byrd, J. D., Jr. and A. C. York. 1988. Interactions of carbaryl and dimethoate with sethoxydim. *Weed Technol.* 2:433-436.
- Carley, D. S., D. L. Jordan, R. L. Brandenburg, and L. C. Dharmasri. 2009. Factors influencing response of Virginia market type peanut (*Arachis hypogaea*) to paraquat under weed-free conditions. *Peanut Sci.* 36:180-189.
- Chahal, G. B. S., D. L. Jordan, J. Burton, S. B. Clewis, R. L. Brandenburg, and B. B. Shew. 2009a. Weed control in peanut with tank mixtures containing five components. *Proc. South. Weed Sci. Soc.* 62:124.
- Chahal, G. B. S., D. L. Jordan, B. B. Shew, R. L. Brandenburg, J. D. Burton, and D. Danehower. 2009b. Weed and disease control in peanut as influenced by co-application of agrichemicals. *Proc. Am. Peanut Res. and Education Soc.* 41:51.
- Chapin, J. W. and J. S. Thomas. 1993. Effects of chlorpyrifos on pod damage, disease incidence, and yield in two peanut fungicide programs. *Peanut Sci.* 20:102-106.
- Chernicky, J. P. and F. W. Slife. 1986. Effects of sublethal concentrations of bentazon, fluazifop, haloxyfop, and sethoxydim on corn (*Zea mays*). *Weed Sci.* 34:171-174.
- Choate, J., G. Wehtje, and K. L. Bowen. 1998. Interaction of paraquat-based weed control with chlorothalonil-based disease control in peanut. *J. Production Agriculture.* 11:191-195.
- Clewis, S. B., W. J. Everman, D. L. Jordan, and J. W. Wilcut. 2007. Weed management in North Carolina peanuts (*Arachis hypogaea*) with S-metolachlor, diclosulam, flumioxazin, and sulfentrazone systems. *Weed Technol.* 21:629-635.
- Cline, W. O. and M. K. Beute. 1986. Effect of metam sodium, peanut genotype and inoculum density on incidence of *Cylindrocladium* black rot. *Peanut Sci.* 13:41-45.
- Culbreath, A. K., R. C. Kemeraite, Jr., and T. B. Brenneman. 2008. Management of leaf spot diseases of peanut with prothioconazole applied alone or in combination with tebuconazole or trifloxystrobin. *Peanut Sci.* 35:149-158.
- Culpepper, A. S., A. C. York, and C. Brownie. 1999. Influence of bromoxynil on annual grass control by graminicides. *Weed Sci.* 47:123-128.
- Culpepper, A. S., A. C. York, K. M. Jennings, and R. B. Butts. 1998. Interaction of bromoxynil and postemergence graminicides in large crabgrass (*Digitaria sanguinalis*). *Weed Technol.* 12:554-559.
- Dotray, P. A., L. C. Marshall, W. B. Parker, D. L. Wyse, D. A. Somers, and B. G. Gengenbach. 1993. Herbicide tolerance and weed control in sethoxydim-tolerant corn (*Zea mays*). *Weed Sci.* 41:213-217.
- Drake, W. L., D. L. Jordan, R. L. Brandenburg, B. R. Lassiter, P. D. Johnson, and B. M. Royals. 2009. Peanut cultivar response to damage from tobacco thrips and paraquat. *Agron. J.* 101:1388-1393.
- Duke, S. O., D. E. Wedge, A. L. Cerdeira, and M. B. Matallo. 2007. Herbicide Effects on Plant Disease. *Pest Manage. Sci.* 18:36-40.
- Flint, J. L. and M. B. Barrett. 1989a. Effects of glyphosate combinations with 2,4-D or dicamba on field bindweed (*Convolvulus arvensis*). *Weed Sci.* 37:12-18.
- Flint, J. L. and M. B. Barrett. 1989b. Antagonism of glyphosate toxicity to johnsongrass (*Sorghum halepense*) by 2,4-D and dicamba. *Weed Sci.* 37:700-705.
- Foy, C. L. and H. L. Witt. 1997. SAN 582, alachlor, and metolachlor control triazine-resistant (TR) smooth pigweed (*Amaranthus hybridus*) in no-till corn (*Zea mays*). *Weed Technol.* 11:623-625.

- Funderburk, J. E., D. W. Gorbet, I. D. Tearre, and J. Stavisky. 1998. Thrips injury can reduce peanut yield and quality under conditions of multiple stress. *Agron. J.* 90:563-566.
- Gascho, G. J. and J. G. Davis. 1995. Soil fertility and plant nutrition. Pages 383-418 in H. E. Pattee and H. T. Stalker (eds.) *Advances in Peanut Science*. Am. Peanut Res. and Education Soc., Stillwater, OK.
- Green, M. J. 1989. Herbicide antagonism at the whole plant level. *Weed Technol.* 3:217-226.
- Green, J. M. and S. P. Bailey. 1987. Herbicide interactions with herbicides and other agricultural chemicals. Pages 37-61 in C. G. McWhorter and M. R. Gebhardt (eds.) *Methods of Applying Herbicides*, Weed Sci. Soc. Am., Champaign, IL.
- Grey, T. L., D. C. Bridges, E. F. Eastin, and G. E. MacDonald. 2002. Influence of flumioxazin rate and herbicide combinations on weed control in peanut (*Arachis hypogaea* L.). *Peanut Sci.* 29:24-29.
- Grichar, W. J. 1991. Sethoxydim and broadleaf herbicide interaction effects on annual grass control in peanuts (*Arachis hypogaea*). *Weed Technol.* 5:321-324.
- Grichar, W. J., B. A. Besler, K. D. Brewer, and T. A. Baughman. 2002. Grass control in peanut (*Arachis hypogaea*) with clethodim and selected broadleaf herbicide combinations. *Peanut Sci.* 29:85-88.
- Grichar, W. J. and T. E. Boswell. 1987. Herbicide combinations in peanut (*Arachis hypogaea*). *Weed Technol.* 1:290-293.
- Grichar, J. E. and A. E. Colburn. 1996. Flumioxazin for weed control in Texas peanut (*Arachis hypogaea* L.). *Peanut Sci.* 23:30-36.
- Grichar, W. J., A. E. Colburn, and P. A. Baumann. 1996. Yellow nutsedge (*Cyperus esculentus*) control in peanut (*Arachis hypogaea*) as influenced by method of metolachlor application. *Weed Technol.* 10:278-281.
- Grichar, W. J., P. A. Dotray, and T. A. Baughman. 2008. Yellow nutsedge (*Cyperus esculentus*) control and peanut tolerance to S-metolachlor and diclosulam combinations. *Weed Technol.* 22:442-447.
- Grichar, W. J., R. G. Lemon, D. C. Sestak, and K. D. Brewer. 2000. Comparison of metolachlor and dimethenamid for yellow nutsedge (*Cyperus esculentus* L.) control and peanut (*Arachis hypogaea* L.) injury. *Peanut Sci.* 27:26-30.
- Grichar, W. J., P. R. Nester, and A. E. Colburn. 1992. Nutsedge (*Cyperus* spp.) control in peanuts (*Arachis hypogaea*) with imazethapyr. *Weed Technol.* 6:396-400.
- Grichar, J. L. and E. P. Prostko. 2009. Effect of glyphosate and fungicide combinations on weed control in soybeans. *Crop Protection.* 28:619-622.
- Grossman, K., K. S. Koenig, and J. Kwiatkowski. 1994. Phytohormonal changes in intact shoots of wheat and oilseed rape treated with the acylcyclohexanedione growth retardant prohexadione calcium. *Physiol. Plant.* 90:139-143.
- Harris, H. C. and J. B. Brolman. 1966. Comparison of calcium and boron deficiencies of peanut. I. Physiological and yield differences. *Agron. J.* 58:575-578.
- Hatterman-Valenti, H. M., A. Pitty, and M. D. K. Owen. 2006. Effect of environment on giant foxtail (*Setaria faberi*) leaf wax and fluazifop-P absorption. *Weed Sci.* 54:607-614.
- Hatzios, K. K. and D. Penner. 1985. Interaction of herbicides with other agricultural chemicals in higher plants. *Rev. Weed Sci.* 1:1-64.

- Hauser, E. W., G. A. Buchanan, W. J. Ethredge, M. D. Jellum, and S. R. Ceci. 1976. Interactions among peanut cultivars, herbicide sequences, and a systemic insecticide. *Peanut Sci.* 3:56-62.
- Hauser, E. W., G. A. Buchanan, J. E. Harvey, W. L. Currey, D. W. Gorbet, and N. A. Minton. 1981. Pesticide interactions with peanut cultivars. *Peanut Sci.* 8:142-144.
- Holshouser, D. L. and H. D. Coble. 1990. Compatibility of sethoxydim with five postemergence broadleaf herbicides. *Weed Technol.* 4:128-133.
- Hurt, C., R. Brandenburg, D. Jordan, B. Shew, T. Isleib, M. Linker, A. Herbert, P. Phipps, C. Swann, and W. Mozingo. 2003. Managing tomato spotted wilt virus in peanuts in North Carolina and Virginia. North Carolina Cooperative Extension Service Publication AG-638.
- Isaacs, M. A., H. P. Wilson, and J. E. Toler. 2003. Combinations of sethoxydim with postemergence broadleaf herbicides in sethoxydim-resistant corn (*Zea mays*). *Weed Technol.* 17:224-228.
- Jianhua, Z., A. S. Hamill, and E. W. Susan. 1995. Antagonism and synergism between herbicides: Trends from previous studies. *Weed Technol.* 9:86-90.
- Johnson, W. C., Jr., D. L. Colvin, and B. G. Mullinix, Jr. 1993. Comparative response of three peanut cultivars to multiple herbicide applications. *Peanut Sci.* 20:17-20
- Jordan, D. L. 1995. Influence of adjuvants on the antagonism of graminicides by broadleaf herbicides. *Weed Technol.* 9:741-747.
- Jordan, D. L. 2010. Weed management in peanuts. Pages 55-83 in 2010 Peanut Information. North Carolina Cooperative Extension Service Publication AG-331.
- Jordan, D. L. 2010. Peanut production practices. Pages 25-53 in 2010 Peanut information. North Carolina Cooperative Extension Service Publication AG-331.
- Jordan, D. L., R. L. Brandenburg, P. D. Johnson, B. M. Royals, and B. Watson. 2008. Interactions of chlorpyrifos and herbicides applied to peanut (*Arachis hypogaea* L.). *Peanut Sci.* 35:32-37.
- Jordan, D. L., A. S. Culpepper, W. J. Grichar, J. T. Ducar, B. J. Brecke, and A. C. York. 2003a. Weed control with combinations of selected fungicides and herbicides applied postemergence to peanut (*Arachis hypogaea* L.). *Peanut Sci.* 30:1-7.
- Jordan, D. L., S. H. Lancaster, J. E. Lanier, P. D. Johnson, J. B. Beam, and A. C. York. 2009a. Influence of application variables on efficacy of manganese-containing fertilizers applied to peanut. *Proc. Am. Peanut Res. and Education Soc.* 41:82-83.
- Jordan, D. L., S. H. Lancaster, J. E. Lanier, P. D. Johnson, J. B. Beam, A. C. York, R. L. Brandenburg, F. R. Walls, S. Casteel, and C. Hudak. 2006. Influence of application variables on efficacy of boron-containing fertilizers applied to peanut (*Arachis hypogaea* L.). *Peanut Sci.* 33:104-111.
- Jordan, D. L., J. F. Spears, and J. W. Wilcut. 2003b. Tolerance of peanut (*Arachis hypogaea* L.) to herbicides applied postemergence. *Peanut Sci.* 30:8-13.
- Jordan, D. L. and A. C. York. 1989. Effects of ammonium fertilizers and BCH 81508 S on antagonism with sethoxydim plus bentazon mixtures. *Weed Technol.* 3:450-454.
- Katan, J. and Y. Eshel. 1973. Interactions between herbicides and plant pathogens. *Residue Rev.* 45:145-177.
- Kataria, H. R. and U. Gisi. 1990. Interactions of fungicide-herbicide combinations against plant pathogens and weeds. *Crop Protection.* 9:403-409.

- Kells, J. J. and G. Wanamarta. 1987. Effect of adjuvant and spray volume on quackgrass (*Agropyron repens*) control with selective postemergence herbicides. *Weed Technol.* 1:129-132.
- Lancaster, S. H., D. L. Jordan, and P. D. Johnson. 2008. Influence of graminicides formulation on compatibility with other pesticides. *Weed Technol.* 22:580-583.
- Lancaster, S. H., D. L. Jordan, A. C. York, I. C. Burke, F. T. Corbin, Y. S. Sheldon, J. W. Wilcut, and D. W. Monks. 2005a. Influence of selected fungicides on efficacy of clethodim and sethoxydim. *Weed Technol.* 19:397-403.
- Lancaster, S. H., D. L. Jordan, A. C. York, J. W. Wilcut, D. W. Monks, and R. L. Brandenburg. 2005b. Interactions of clethodim and sethoxydim with selected agrichemicals applied to peanut. *Weed Technol.* 19:456-461.
- Lassiter, R. B. and H. D. Coble. 1987. Carrier volume effects on the antagonism of sethoxydim by bentazon. *Weed Sci.* 35:541-546.
- Levesque, C. A. and J. E. Rahe. 1992. Herbicide interactions with fungal root pathogens, with special reference to glyphosate. *Annual Review Phytopathology.* 30:579-602.
- Lynch, R. E. and T. P. Mack. 1995. Biological and biotechnical advances for insect management in peanut. Pages 95-159 in H. E. Pattee and H. T. Stalker (eds.) *Advances in Peanut Science.* Am. Peanut Res. and Education Soc., Stillwater, OK.
- Mack, T. P., J. E. Funderburk, R. E. Lynch, E. G. Braxton, and C. B. Backman. 1989. Efficacy of chlorpyrifos in soil to lesser cornstalk borer (Lepidoptera: Pyralidae). *J. Econ. Entomol.* 82: 1224-1229.
- Mack, T. P., J. E. Funderburk, and M. G. Miller. 1991. Efficacy of selected granular insecticides in soil in 'Florunner' peanut fields to larvae of lesser cornstalk borer (Lepidoptera: Pyralidae). *J. Econ. Entomol.* 84: 1899-1904.
- Minton, B. W., M. E. Kurtz, and D. R. Shaw. 1989. Barnyardgrass (*Echinochloa crus-galli*) control with grass and broadleaf weed herbicide combinations. *Weed Sci.* 37:223-227.
- Minton, N. A., A. S. Csinos, and L. W. Morgan. 1990. Relationship between tillage, nematicide, fungicide, and insecticide treatments on pests and yield of peanut double-cropped with wheat. *Plant Disease* 74:1025-1029.
- Minton, N. A. and L. W. Morgan. 1974. Evaluation of systemic and nonsystemic pesticides for insect and nematode control in peanuts. *Peanut Sci.* 1:91-98.
- Mitchem, W. E., A. C. York, and R. B. Batts. 1996. Peanut response to prohexadione calcium, a new plant growth regulator. *Peanut Sci.* 23:1-9.
- Morton, C. A., R. G. Harvey, J. L. Wedberg, J. J. Kells, D. A. Landis, and W. E. Lueschen. 1994. Influence of corn rootworm insecticide on the response of field corn (*Zea mays*) to nicosulfuron. *Weed Technol.* 8:289-295.
- Mueller, T. C., W. W. Witt, and M. Barrett. 1989. Antagonism of johnsongrass (*Sorghum halepense*) control with fenoxaprop, haloxyfop, and sethoxydim by 2,4-D. *Weed Technol.* 3:86-89.
- Myers, P. F. and H. D. Coble. 1992. Antagonism of graminicide activity on annual grass species by imazethapyr. *Weed Technol.* 6:333-338.
- Penner, D. 1989. The impact of adjuvants on herbicide antagonism. *Weed Technol.* 3:227-231.
- Powell, N. L., C. W. Swann, and D. C. Martens. 1996. Foliar fertilization of Virginia-type peanut with MnEDTA- crop grade, pod yield, and value. *Peanut Sci.* 23:98-103.

- Putnam, A. R. and D. Penner. 1974. Pesticide interactions in higher plants. *Residue Rev.* 50:73-110.
- Putnam, A. R. and S. K. Ries. 1967. The synergistic action of herbicide combinations containing paraquat on *Agropyron repens* (L.) Beauv. *Weed Res.* 7:191-199.
- Rahman, A. and T. K. James. 1993. Enhanced activity of nicosulfuron in combination with soil-applied insecticides in corn (*Zea mays*). *Weed Technol.* 7:824-829.
- Rhodes, G. N., Jr. and H. D. Coble. 1984a. Influence of application variables on antagonism between sethoxydim and bentazon. *Weed Sci.* 32: 436-441.
- Rhodes, G. N., Jr. and H. D. Coble. 1984b. Influence of bentazon on absorption and translocation of sethoxydim in goosegrass. *Weed Sci.* 32:595-597.
- Richburg III, J. S., J. W. Wilcut, D. L. Colvin, and G. R. Wiley. 1996. Weed management in southeastern peanut (*Arachis hypogaea*) with AC 263,222. *Weed Technol.* 10:145-152.
- Richburg III, J. S., J. W. Wilcut, and E. F. Eastin. 1995. Weed management in peanut (*Arachis hypogaea*) with imazethapyr and metolachlor. *Weed Technol.* 9:807-812.
- Riley, D. G., G. K. Douce, and R. M. McPherson. 1997. Summary of losses from insect damage and costs of control in Georgia 1996. Georgia Agricultural Experiment Station, University of Georgia Special Publication No. 91.
- Robinson, D. E. and R. Nurse. 2008. Effect of herbicide-fungicide tank-mix combinations on weed control and tomato tolerance. *Acta Horticulturae (ISHS)* Pages 129-134.
- Sandberg, C. L., W. F. Meggitt, and D. Penner. 1978. Effect of diluent volume and calcium on glyphosate phytotoxicity. *Weed Sci.* 26:476-479.
- Scott, G. H., S. D. Askew, and J. W. Wilcut. 2001. Economic evaluation of diclosulam and flumioxazin systems in peanut (*Arachis hypogaea*). *Weed Technol.* 15:360-364.
- Sherwood, J. L., M. K. Beute, D. W. Dickson, V. J. Elliot, R. S. Nelson, C. H. Opperman, and B. B. Shew. 1995. Biological and biotechnical control in *Arachis* diseases. Pages 160-206 in H. E. Pattee and H. T. Stalker (eds.) *Advances in Peanut Science*. Am. Peanut Res. and Education Soc., Stillwater, OK.
- Shew, B. B. 2010. Peanut disease management. Pages 103-130 in 2010 Peanut Information. North Carolina Cooperative Extension Service Publication AG-331.
- Shew, B. B. and F. Waliyar. 2005. Peanut Diseases: Ecology and Control. *Encyclopedia of Pest Management* 1:1-5. (DOI: 10.1081/E-EPM-120041160).
- Smith, D. H. and R. H. Littrell. 1980. Management of peanut foliar diseases with fungicides. *Plant Disease.* 64:356-361.
- Smith, F. D., P. M. Phipps, and R. J. Stipes. 1992. Fluazinam: A new fungicide for control of Sclerotinia blight and other soilborne pathogens of peanut. *Peanut Sci.* 19:115-120.
- Stahlman, P. W. and W. M. Phillips. 1979. Effects of water quality and spray volume on glyphosate phytotoxicity. *Weed Sci.* 27:38-41.
- Swann, C. W. and D. A. Herbert, Jr. 1999. Influence of in-furrow thrips insecticides on response of peanut to bentazon and acifluorfen. *Proc. South. Weed Sci. Soc.* 52:67.
- Thelen, K. D., E. P. Jackson, and D. Penner. 1995. Characterizing the sethoxydim-bentazon interaction with proton nuclear magnetic resonance spectrometry. *Weed Sci.* 43:337-341.
- Van Gundy, S. D. and M. V. McKenry. 1977. Action of Nematicides. Pages 263-283 in J. G. Horsfall, J. G. And E. B. Cowling (eds.) *Plant Disease: An Advanced Treatise*, Vol. 1. Academic Press, NY.

- Vidrine, P. R., D. B. Reynolds, and D. C. Blouin. 1995. Grass control in soybean (*Glycine max*) with graminicides applied alone and in mixtures. *Weed Technol.* 9:68-72.
- Wanamarta, G. and D. Penner. 1989. Identification of efficacious adjuvants for sethoxydim and bentazon. *Weed Technol.* 3:60-66.
- Wanamarta, G., D. Penner, and J. J. Kells. 1989. The basis of bentazon antagonism on sethoxydim absorption and activity. *Weed Sci.* 37:400-404.
- Webster, T. M. 2009. Weed survey – southern states. *Proc. South. Weed Sci. Soc.* 62:509-524.
- Wehtje, G., J. W. Wilcut, T. V. Hicks, and J. McGuire. 1988. Relative tolerance of peanuts to alachlor and metolachlor. *Peanut Sci.* 15:53-56.
- Wehtje, G., J. W. Wilcut, and J. A. McGuire. 1992a. Paraquat behavior as influenced by 2, 4-DB in peanut (*Arachis hypogaea*) and selected weeds. *Peanut Sci.* 19:51-55.
- Wehtje, G., J. W. Wilcut, and J. A. McGuire. 1992b. Influence of bentazon on the phytotoxicity of paraquat to peanuts (*Arachis hypogaea*) and associated weeds. *Weed Sci.* 40:90-95.
- Whisenant, S. G. and R.W. Bovey. 1993. Water quality effects on stability and phytotoxicity of picloram and clopyralid. *J. Range Manage.* 46: 254-256.
- White, P. M., T. L. Porter, and A. K. Culbreath. 2009. Fungicide dissipation and impact on metolachlor-aerobic soil degradation and soil microbial dynamics. *Science of Total Environment.* 408:1393-1403. (doi:10.1016/j.scitotenv.20.9.11.012).
- Wilcut, J. W., J. S. Richburg III, E. F. Eastin, G. R. Wiley, F. R. Walls, Jr., and S. Newell. 1994a. Imazethapyr and paraquat systems for weed management in peanut (*Arachis hypogaea*). *Weed Sci.* 42:601-607.
- Wilcut, J. W. and C. W. Swann. 1990. Timing of paraquat applications for weed control in Virginia-type peanut (*Arachis hypogaea*). *Weed Sci.* 38:558-562.
- Wilcut, J. W., A. C. York, W. J. Grichar, and G. R. Wehtje. 1995. The biology and management of weeds in peanut (*Arachis hypogaea*). Pages 207-244 in H. E. Pattee and H. T. Stalker (eds.) *Advances in Peanut Science*. Am. Peanut Res. and Education Soc., Stillwater, OK.
- Wilcut, J. W., G. R. Wehtje, D. L. Colvin, and M. G. Patterson. 1987a. Economic assessment of herbicide systems for minimum-tillage peanut and conventional tillage peanut. *Peanut Sci.* 14:83-86.
- Wilcut, J. W., G. R. Wehtje, and T. V. Hicks. 1990. Evaluation of herbicide systems in minimum and conventional tillage peanut (*Arachis hypogaea*). *Weed Sci.* 38:243-248.
- Wilcut, J. W., G. R. Wehtje, and M. G. Patterson. 1987b. Economic assessment of weed control systems for peanuts (*Arachis hypogaea*). *Weed Sci.* 35:433-437.
- Wilcut, J. W., A. C. York, and G. R. Wehtje. 1994b. The control and interaction of weeds in peanut (*Arachis hypogaea*). *Rev. Weed Sci.* 6:177-205.
- Wills, G. D. and C. G. McWhorter. 1985. Effect of inorganic salts on the toxicity and translocation of glyphosate and MSMA in purple nutsedge (*Cyperus rotundus*). *Weed Sci.* 33:755-761.
- York, A. C., D. L. Jordan, and J. W. Wilcut. 1990. Effects of $(\text{NH}_4)_2\text{SO}_4$ and BCH 81508 S on efficacy of sethoxydim. *Weed Technol.* 4:76-80.
- York, A. C. and J. W. Wilcut. 1995. Interaction of bentazon and imazethapyr applied postemergence to nutsedge (*Cyperus* spp.). *Peanut Sci.* 22:150-154.
- York, A. C., J. W. Wilcut, and W. J. Grichar. 1993. Interactions of 2,4-DB with postemergence herbicides. *Peanut Sci.* 20:57-61.

Herbicides, Theory and Applications

Edited by Prof. Marcelo Larramendy

ISBN 978-953-307-975-2

Hard cover, 610 pages

Publisher InTech

Published online 08, January, 2011

Published in print edition January, 2011

The content selected in Herbicides, Theory and Applications is intended to provide researchers, producers and consumers of herbicides an overview of the latest scientific achievements. Although we are dealing with many diverse and different topics, we have tried to compile this "raw material" into three major sections in search of clarity and order - Weed Control and Crop Management, Analytical Techniques of Herbicide Detection and Herbicide Toxicity and Further Applications. The editors hope that this book will continue to meet the expectations and needs of all interested in the methodology of use of herbicides, weed control as well as problems related to its use, abuse and misuse.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

David L. Jordan, Gurinderbir S. Chahal, Sarah H. Lancaster, Joshua B. Beam, Alan C. York and William Neal Reynolds (2011). Defining Interactions of Herbicides with Other Crop Protection Products Applied to Peanut, Herbicides, Theory and Applications, Prof. Marcelo Larramendy (Ed.), ISBN: 978-953-307-975-2, InTech, Available from: <http://www.intechopen.com/books/herbicides-theory-and-applications/defining-interactions-of-herbicides-with-other-crop-protection-products-applied-to-peanut>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2011 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike-3.0 License](https://creativecommons.org/licenses/by-nc-sa/3.0/), which permits use, distribution and reproduction for non-commercial purposes, provided the original is properly cited and derivative works building on this content are distributed under the same license.

IntechOpen

IntechOpen