We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Synthesis and Evaluation of Pyrazine Derivatives with Herbicidal Activity

Martin Doležal¹ and Katarína Kráľová²

¹*Faculty of Pharmacy in Hradec Králové, Charles University in Prague*

²*Faculty of Natural Sciences, Comenius University in Bratislava*

¹*Czech Republic*

²*Slovak Republic*

1. Introduction

The pyrazine ring is a part of many polycyclic compounds of biological and/or industrial significance; examples are quinoxalines, phenazines, and bio-luminescent natural products pteridines, flavins and their derivatives. All these compounds are characterized by a low lying unoccupied π -molecular orbital and by the ability to act as bridging ligand. Due to these two properties 1,4-diazines, and especially their parent compound pyrazine, possess a characteristic reactivity. Pyrazine is a weak diacid base ($pK_1 = 0.57$; $pK_2 = -5.51$), weaker than pyridine, due to the induction effect of the second nitrogen (Bird, 1992). Its inherent bifunctionality and the low lying unoccupied molecular orbital permit pyrazine to form coordination polymers having unusual electrical and magnetic properties (Brown & Knaust, 2009). 1,4-Diazines may be employed to study inter- and intramolecular electron transfer in organic, inorganic and biochemical reactions. Autocondensation of α -aminocarbonyl compounds to the dihydropyrazine derivative, which is followed by oxidation on the final substituted pyrazine, or the condensation of α,β -dicarbonyl and α,β -diamino compounds forming during the fermentation of saccharides and peptides are the main routes of pyrazine ring building. Pyrazines are found mainly in processed food, where they are formed during dry heating processes *via* Maillard reactions (Maillard, 1912). They are also found naturally in many vegetables, insects, terrestrial vertebrates, and marine organisms, and they are produced by microorganisms during their primary or secondary metabolism (Adams et al., 2002; Beck et al., 2003; Wagner et al., 1999; Woolfson & Rothschild, 1990). The widespread occurrence of simple pyrazine molecules in nature, especially in the flavours of many food systems, their effectiveness at very low concentrations as well as the still increasing applications of synthetic pyrazines in the flavour and fragrance industry are responsible for the high interest in these compounds (Maga, 1992). Certain pyrazines, especially dihydropyrazines, are essential for all forms of life due their DNA strand-breakage activity and/or by their influencing of apoptosis (Yamaguchi, 2007). Synthetic pyrazine derivatives are also useful as drugs (antiviral, anticancer, antimycobacterial, etc.), fungicides, and herbicides (Doležal, 2006a). Furthermore, a simple pyrazine compound, 3-amino-6-chloro-pyrazine-6-carboxylic acid, has shown anti-auxin behaviour (Camper & McDonald, 1989). The importance of the pyrazine (1,4-diazine) ring for the biological

activity can be evaluated primarily according to the size of the studied molecules. In relatively small compounds, the pyrazine ring is necessary for biological action due to its resemblance (bioisosterism) to the naturally occurring compounds (e.g. nicotinamide, or pyrimidine nucleic bases). In bulky compounds the introduction of the pyrazine ring brings specific chemical and physicochemical properties for the molecule as a whole, such as basic and slightly aromatic character (Doležal, 2006a). A fully comprehensive study of the pyrazines including reactivity and synthesis is beyond the scope of this work but can be found in the literature (Brown, 2002; Joule & Mills, 2010).

Herbicides are generally considered as growth inhibitors, thus their different inhibitory responses have been studied in various culture systems. Plant tissue and cell cultures provide model systems for the study of various molecular, physiological, organism and genetic problems. These systems have been used in the study of herbicides and other xenobiotics (Linsmaier & Skoog, 1965).

2. Pyrazine herbicides

The most successful pyrazine derivative was diquat-dibromide (see Fig. 1, the structure I). This non-selective, contact herbicide has been used to control many submerged and floating aquatic macrophytes which interferes with the photosynthetic process, releasing strong oxidizers that rapidly disrupt and inactivate cells and cellular functions (at present banned in many EU countries). Severe oral diquat intoxication has been associated with cerebral haemorrhages and severe acute renal failure (Peiró et al., 2007). Also quinoxaline herbicides (containing the pyrazine fragment) are very useful herbicides. Among them propaquizafop (Fig. 1, II) and quizalofop-ethyl (Fig. 1, III) are the most important derivatives (Frater et al., 1987; Sakata et al., 1983).

Fig. 1. Structures of diquat-dibromide (I), propaquizafop (II) and quizalofop-ethyl (III).

2.1 Diquat

Diquat-dibromide (6,7-dihydrodipyrido[1,2-a:2',1'-c]pyrazinediium-dibromide; for the structure see Fig. 1, I) is a quaternary ammonium salt used as a non-selective contact herbicide and desiccant, absorbed by the foliage with some translocation in the xylem. It is used for preharvest desiccation of many crops, as a defoliant on hops, for general weed control on non crop land etc. (Ritter et al., 2000; Ivany, 2005). It is applied as an aquatic

herbicide in many countries since the late 1950s for control of emergent and submerged aquatic weeds (Ritter et al., 2000). According to Massachusetts Department of Agricultural Resources (2010) following weeds are controlled by diquat: i) submersed aquatics: *Ultricularia*, *Ceratophyllum demersum*, *Elodea* spp., *Najas* spp., *Myriophyllum* spp., *Hydrilla verticillata*, *Potamogeton* spp.; ii) floating aquatics: *Salvinia* spp., *Eichhornia crassipes*, *Pistia Stratiotes*, *Lemna* spp., *Hydrocotyle* spp.; iii) marginal weeds: *Typha* spp.; iv) algae: *Pithophora* spp., *Spyrogyra* spp. (filamentous algae). Diquat is stable in neutral and acidic solutions but unstable in alkaline medium. It breaks down by the UV radiation and the degradation increases with pH > 9 (Diaz et al., 2002). It is also biodegraded in water by microorganisms that uses this herbicide as a source of carbon or nitrogen (Petit et al., 1995).

Trade names for diquat-dibromide formulations included Desiquat[®], Midstream[®], Reglone[®], and Reglex[®]. Mixtures of diquat with another quaternary herbicide paraquat (1,1'-dimethyl-4,4'-bipyridinium-dichloride) were sold under trade names including Actor[®], Dukatalon[®], Opal[®], Pathclear[®] (also includes simazine and aminotriazole), Preeglox[®], Preglone[®], Seccatutto[®], Spray Seed[®], and Weedol[®] (Lock & Wilks, 2001).

Fig. 2. Scheme of the photosynthetic electron transport in photosystem I (PS I). (Figure taken from <http://www.bio.ic.ac.uk/research/barber/psIimages/PSI.jpg> with permission of Prof. Barber, Imperial College London).

The first paper dealing with the mode of action of diquat was published in 1960 by Mees who indicated that oxygen and light were essential for its herbicidal effect. Later Zweig et al. (1965) found that diquat caused a deviation of electron flow from photosystem (PS) I what resulted in an inhibition of NADP⁺ reduction and the production of a reduced diquat radical. In Fig. 2 is shown scheme of the photosynthetic electron transport (PET) in PS I.

In plants, the PS I complex catalyzes the oxidation of plastocyanin and the reduction of ferredoxin (F_d). From the primary donor, P700, electrons are transferred to the primary

acceptor, A_0 and then to plastoquinone (A_1) operating as a single electron acceptor. From A_1 electrons are transferred to a 4Fe-4S cluster (F_X) and subsequently to two 4Fe-4S clusters, F_A and F_B , located on the stromal side of the reaction center close to F_X . PS I produces a strong reductant that transfers electrons to F_d . Ferredoxin, one of the strongest soluble reductants found in cells, operates in the stromal aqueous phase of the chloroplast, transferring electrons from PS I to ferredoxin-NADP⁺ oxidoreductase. The final electron acceptor in the photosynthetic electron transport chain is NADP⁺, which is fully reduced by two electrons (and one proton) to form NADPH, a strong reductant which serves as a mobile electron carrier in the stromal aqueous phase of the chloroplast (Whitmarsch, 1998).

Due to deviation of electron flow from F_d , an inhibition of NADP⁺ reduction occurs and a reduced diquat radical is formed. Davenport (1963) found that in the presence of oxygen the reduced diquat free radical was reoxidized with the production of hydrogen peroxide. Thus, an one-electron reduction of diquat results in a cation free radical that reacts rapidly with molecular oxygen and generates reactive oxygen species such as the superoxide anion radical (Mason, 1990). Reactive oxygen species cause oxidative stress in the cell with consecutive damage of biological membranes. In herbicide classification diquat, similarly to paraquat, is classified as HRAC Group D herbicide causing PS I electron diversion (HRAC 2005). Injury to diquat-treated crop plants occurs in the form of spots of dead leaf tissue wherever spray droplets contact the leaves indicating that this herbicide belongs to membrane disruptors. The use of diquat for the control of aquatic weeds is widespread in the US (US Environmental Protection Agency, 1995) whereas it is forbidden in the EU (European Commission, 2001, 2002).

As mentioned above, diquat toxicity to both aquatic plants and animals originates from the formation of reactive oxygen species in both chloroplasts and mitochondria (Cedergreen et al., 2006; Sanchez et al., 2006). The field effects of diquat to natural strands of aquatic vegetation were studied by Peterson et al. (1997) and Campbell et al. (2000). The filamentous cyanobacteria were slightly less tolerant than the unicellular cyanobacteria and the most sensitive was genus *Anabena* (Peterson et al., 1997). Gorzerino et al. (2009) showed that diquat, used as the commercial preparation Reglone 2®, inhibited the growth of *Lemna minor* in indoor microcosms. According to findings of Campbell et al. (2000) diquat has a minimal ecological impact to benthic invertebrates and fish; on the other hand, aquatic plants in the vicinity of application to surface waters appear to be at risk (nevertheless this is expected, as diquat-dibromide kills aquatic plants). However, Koschnick et al. (2006) observed that the accession of *Landoltia* from Lake County (Florida) had developed resistance to diquat and the resistance mechanism was independent of photosynthetic electron transport.

2.2 Patented pyrazine herbicides

The control of unwanted vegetation by means of chemical agents, *i.e.* herbicides, is an important aspect of modern agriculture and land management's. While many chemicals that are useful for the control of unwanted vegetation are known, new compounds that are more effective generally, are more effective for specific plant species, are less damaging to desirable vegetation, are safer to man or the environment, are less expensive to use or have other advantageous attributes, are desirable (Benko, 1997). Many structural variations of pyrazine compounds with herbicidal properties can be found in the patent literature.

Several thiazolopyrazines exhibited pre-emergent herbicidal activity when applied as aqueous drenches to soil planted with seeds of certain plants. For example, application of 4000 ppm of compound IV (Fig. 3) resulted in emergence inhibition of crabgrass (50% of the

control) and barnyard grass (*Echinochloa crus-galli* (L.) P. Beauv.) (45% of the control). Due to the treatment with a dose of 2 lb per acre of compound V (Fig. 3), the emergence of cotton reached only 30% of the control (Tong, 1978).

Böhner & Meyer (1989a, 1989b, 1990) prepared a set of aminopyrazinones (Fig. 3, VI) and aminotriazinones and tested these compounds for their herbicidal action before emergence of the plants. It was found that application of 70.8 ppm of some compounds on the substrate vermiculite resulted in very potent inhibition of seed germination of *Nasturtium officinalis*, *Agrostis tenuis*, *Stellaria media* and *Digitaria sanguinalis*. Due to the treatment with compound where $R^1 = \text{CH}_3$, $R^2 = \text{OCH}_3$, $R^3 = \text{H}$, $R^7 = \text{H}$, $R^8 = \text{COOCH}_3$, $X = \text{O}$ plants have not germinated and completely died. After spraying of 21 days old spring barley (*Hordeum vulgare*) and spring rye (*Secale*) plants shoots with an active substance VI (up to 100 g per hectare) new additional growth of plants reached only 60-90% of the control. For grasses *Lolium perenne*, *Poa pratensis*, *Festuca ovina*, *Dactylis glomerata* and *Cynodon dactylon* sprayed with the same dose of an active substance (Fig. 3, VII) reduction in new additional growth in comparison with the untreated control (10-30% of control) was observed, too (Böhner & Meyer, 1989a, 1989b, 1990).

Benko et al. (1997) patented a series of *N*-aryl[1,2,4]triazolo[1,5-*a*]pyrazine-2-sulfonamides as good pre- and post-emergence selective herbicides with good growth regulating properties. Excellent pre-emergence activity against pigweed and morning glory and very good post-emergence herbicidal activity against morning glory and velvet leaf (*Abutilon theophrasti*) have been exhibited by the title compounds.

Dietsche (1977) patented as herbicides a group of substituted 6,7-dichloro-3,4-dihydro-2*H*-pyrazino(2,3-*b*)(1,4)oxazines showing hundred-percent inhibitory effectiveness when applied as pre- as well as post-emergence herbicides (4000 ppm) for pigweeds.

Shuto et al. (2000) patented as useful active ingredients of herbicides a series of pyrazin-2-one derivatives (Fig. 3, VIII, IX) where R^1 is hydrogen or alkyl, R^2 is haloalkyl, R^3 is optionally substituted alkyl, alkenyl or alkynyl and Q is optionally substituted phenyl. Some compounds showed superb effectiveness against *Abutilon theophrasti* and *Ipomoea hederacea* when applied as foliar or soil surface treatment on upland fields (2000 g/ha).

Griffin et al. (1990) patented alkylpyrazine compounds (Fig. 3, X) with plant growth regulating activity, where R^1 is $\text{C}_1\text{-C}_4$ alkyl optionally substituted with halogen or cyclopropyl, optionally substituted with $\text{C}_1\text{-C}_4$ alkyl; R^2 is $\text{C}_1\text{-C}_8$ alkyl, $\text{C}_2\text{-C}_8$ alkenyl, or $\text{C}_2\text{-C}_8$ alkynyl optionally substituted with halogen; $\text{C}_3\text{-C}_6$ cycloalkyl, $\text{C}_3\text{-C}_6$ cycloalkenyl, $\text{C}_3\text{-C}_6$ cycloalkylalkyl, $\text{C}_3\text{-C}_6$ cycloalkenylalkyl, phenylalkenyl or phenylalkynyl each optionally substituted on the ring group; R^3 is hydrogen or $\text{C}_1\text{-C}_4$ alkyl; R^4 is hydrogen, $\text{C}_1\text{-C}_4$ alkyl, halogen, alkylamino, cyano, or alkoxy; n is 0 or 1; and salts, ethers, acylates and metal complexes thereof. The treatment of plants with these compounds can lead to the leaves developing a darker green colour. In dicotyledonous plants such as soybean and cotton, there may be promotion of side shooting. The compounds may be useful in rendering plants resistant to stress since they can delay the emergence of plants grown from seeds, shorten stem height and delay flowering. Engel et al. (1999) patented herbicidal pyrazine derivatives (Fig. 3, XI) which are suitable very effectively control weeds and grass weeds mainly in crops such as wheat, rice, corn, soybean and cotton, without significantly damaging the crops. It could be stressed that this effect occurs in particular at low application rates. In addition, these compounds can also be used in crops which have been made substantially resistant to the action of herbicides by breeding and/or by the use of genetic engineering methods.

N-pyrazinyl-haloacetamides (Fig. 3, XII) where R is hydrogen, hydrocarbonyl, halogen, epoxy, hydroxy, alkoxy, mercapto, alkylsulfanyl, nitro, cyano or amino, R' is hydrogen or

hydrocarbonyl, X is halogen, m is integer from 1 to 4 and n is 0, 1 or 2 showed herbicidal activity. For example, spraying of the 2,2,2-trichloro-*N*-pyrazinyl acetamide on the soil resulted in 100% growth inhibition of wild oats (dosage 1.12 g m⁻²) and yellow foxtail or cultured rice (dosage 1.12 g m⁻²) (Fischer, 1988).

Novel pyrazine-sulfonylcarbamates and thiocarbamates (Fig. 3, XIII) (where Z is oxygen or sulfur and R is C₁-C₄ alkyl, phenyl or benzyl; whereas the pyrazine ring may be variously further substituted) have been found to be good selective herbicides and therefore they are suitable for use in crops of cultivated plants. Moreover, these compounds can damage problem weeds which till then have only been controlled with total herbicides (Böhner et al., 1987). By means of surface treatment it is possible to damage perennial weeds to their roots. Moreover, the compounds are effective when used in very low rates of application and they are able to potentiate the phytotoxic action of other herbicides against certain noxious plants and to reduce the toxicity of such herbicides to some cultivated plants. These compounds can be used also as plant growth regulators causing inhibition of vegetative plant growth what results in substantial increase of the yield of plants. Böhner et al. (1987) synthesized and patented also a set of novel pyrazinyl sulfonamides of the formula Q-SO₂-NH₂ where Q is substituted pyrazine group which could be useful in controlling weeds and are suitable for selectively influencing plant growth. The compounds can be used as pre- and post-emergence herbicides and as plant growth regulators for growth inhibition of cereals (e.g. *Hordeum vulgare* or summer rye (*Secale*)) and grasses (e.g. *Lolium perenne*, *Poa partensis*, *Festuca ovina*, *Cynodon dactylon*). Selective inhibition of the vegetative growth of many cultivated plants permits more plants to be grown per unit of crop area, resulting in significant increase in yield with the same fruit setting and in the same crop area.

Zondler et al. (1989) prepared a set of 2-arylmethyliminopyrazines (Fig. 3, XIV) and tested them for their pre-emergent and post-emergent herbicidal action, as well as for their plant growth regulating activity. Compounds with R⁵ = 4-Cl, R⁶ = 2-Cl, R⁷ = H and R¹ = SCH₃H₇(n) or SCH₂CH=CH₂ showed excellent pre-emergent effect (dose 4 kg/ha) against *Echinochloa crus-galli* and *Monocharia vag*. The last compound was active already at application rate of 500 g/ha. The 2-arylmethylimino-pyrazines were found to be also effective post-emergence herbicides and can be used for growth inhibition of tropical leguminous cover crops (e.g. *Centrosema plumieri* and *Centrosema pubescens*), growth regulation in soybeans and growth inhibition of cereals, too.

Cyanatothiomethylthiopyrazines have been found to be active as pesticides and find particular usage as fungicides, bactericides, nematocides and herbicides (Mixan et al., 1978). Arylsulfanylpurazine-2,3-dicarbonitriles have high herbicidal activity (Takematsu et al., 1984; Portnoy, 1978). Takematsu et al. (1981) patented 2,3-dicyanopyrazines (Fig. 3, XV) as compounds with high herbicidal activity as well as useful active ingredients of herbicides. The compounds have ability to inhibit the germination of weeds and/or wither their stems and leaves, and therefore exhibit an outstanding herbicidal effect as an active ingredient of pre-emergence and/or post-emergence herbicides in submerged soil treatment, foliar treatment of weeds, upland soil treatment, etc.

Compounds where A represents a phenyl group which may have 1 or 2 substituents selected from the class consisting of halogen atoms and lower alkyl groups containing 1 to 3 carbon atoms and B represents an ethylamino, *n*-propylamino, *n*- or *iso*-butylamino, 1-carboxyethylamino, 1-carboxy-*n*-propylamino, 1-carboxy-*iso*-butylamino, 1-carboxy-*n*-pentylamino or allylamino group have the property of selectively blanching (causing

chlorosis, *i.e.* inhibiting the formation of chlorophyll and/or the acceleration of its decomposition) of weeds without chlorosis of useful crops. Hence, these compounds are most suitable as high selective herbicides of chlorosis type.

Fig. 3. Structures of patented thiazolopyrazines (IV,V), aminopyrazinones (VI,VII), substituted pyrazin-2-ones (VIII,IX), arylalkylpyrazines (X, XI), *N*-pyrazinyl-haloacetamides (XII), pyrazine-sulfonylcarbamates and thiocarbamates (XIII), 2-arylmethyliminopyrazines (XIV), substituted 2,3-dicyanopyrazines (XV), pyridopyrazines (XVI), aryloxy pyrazines (XVII) and pyrimidinopyrazines (XVIII).

Takematsu et al. (1984) also patented a set of 2,3-dicyano-6-phenylpyrazine herbicides with outstanding herbicidal activities on paddy weeds in submerged soil treatment. Because they

are not phytotoxic to rice, they can effectively control weeds in paddies. The compounds exhibited herbicidal activity against important upland weeds such as *Digitaria adscendens*, *Polygonum persicaria*, *Galinsoga ciliata*, *Amaranthus viridis*, *Chenopodium album*, *Chenopodium ficifolium*, *Echinochloa crus-galli* (without damaging upland crops) as well as against a very broad range of other upland weeds including *Galium aparin*, *Rumex japonicus*, *Erigeron philadelphicus*, *Erigeron annuus*, and *Capsella bursapastoria*.

Cordingley et al. (2008) prepared herbicidal effective pyridopyrazines (Fig. 3, XVI) with R^1, R^2 independently = H, alkyl, halo, CN, aryl, etc.; R^3 = H, (halo)alkyl, alkenyl, etc.; R^4 = (un)substituted heteroaryl; and R^5 = OH or group metabolizable to OH) or a salt or *N*-oxide thereof. XVI applied post-emergence at 1000 g/ha completely controlled *Solanum nigrum* and *Amaranthus retroflexus*. Also substituted aryloxypyrazines (Fig. 3, XVII) possess interesting herbicidal effect (Niederman & Munro, 1994). For example, in tests against 8 plants, title compound XVII at 5 kg/ha (foliar spray) gave complete kill of *Echinochloa crus-galli* with no damage to rice. Test data include foliar, pre-emergence, and soil drench applications against the 8 plants for most compounds. Sato et al. (1993) patented pyrimidinopyrazines (Fig. 3, XVIII) (R^1 = H, halo, alkoxy, alkylamino, alkyl, haloalkyl; R^2 = Ph, substituted Ph, benzyl, pyridyl, thienyl, furyl; R^3 = SR^4 , OR^5 , NR^6R^7 ; R^4, R^5, R^6, R^7 = H, alkyl, alkenyl, alkynyl; NR^6R^7 may form 3-7 membered ring), useful as herbicides, were prepared and showed herbicidal activity against *Stellaria neglecta* at 0.63 kg/ha.

2.2.1 Structure-activity relationships in series of herbicidal 2,3-dicyanopyrazines

Nakamura et al. (1983) synthesized sixty six 2,3-dicyano-5-substituted pyrazines and measured their herbicidal activities against barnyard grass in pot tests to clarify the relationship between chemical structure and activity. The activity of 59 derivatives showed parabolic dependence on the hydrophobic substituent parameter at the 5-position of the pyrazine ring, indicating that the compounds should pass through a number of lipoidal-aqueous interfaces to reach a critical site for biological activity. It was found that the moiety of 2,3-dicyanopyrazine is essential for herbicidal activity, and the 5-substituent on the pyrazine ring plays an important role in determining the potency of this activity and that *para*-substituted phenyl derivatives show undesirable effects on the potency of the activity at the ultimate site of herbicidal action.

Nakamura et al. (1983a) also synthesized sixty eight 6-substituted 5-ethylamino and 5-propylamino-2,3-dicyanopyrazines and tested their herbicidal activities against barnyard grass using pot tests. In general, these compounds induced chlorosis against young shoots of barnyard grass and inhibited their growth. The most active compound was 2,3-dicyano-5-propylamino-6-(*m*-chlorophenyl)-pyrazine. The results indicated that the structure of the 5-ethylamino and 5-propylamino-2,3-dicyanopyrazine moieties is an important function for the herbicidal activity and that the potency of activity of these two series of compounds is determined by the hydrophobic and steric parameters of substituents at the 6-position of the pyrazine ring.

3. Design, synthesis and evaluation of the pyrazinecarboxamides with herbicidal activity

The structural diversity of organic herbicides continues to increase; therefore classification of herbicides should be based on their chemical structure. The chlorinated aryloxy acids dominated for long period, later were replaced by chemicals of many distinct chemical

classes, including triazines, amides (haloacetanilides), benzonitriles, carbamates, thiocarbamates, dinitroanilines, ureas, phenoxy acids, diphenyl ethers, pyridazinones, bipyridinium compounds, ureas and uracils, sulfonylureas, imidazolinones, halogenated carboxylic acids, and many other compounds. Carboxamide or anilide moieties are present in many used herbicides, *i.e.* alachlor, acetochlor, benoxacor, butachlor, diflufenican, dimethenamid, diphenamid, isoxaben, karsil, napropamide, pretilachlor, propyzamide, dicryl, diflufenican, flufenacet, mefenacet, mefluidide, metolachlor, naphthalan, picolinafen, propachlor, propanil, propham, solan (The Merck Index, 2006). Carboxamide or anilide herbicides are nonionic and moderately retained by soils. The sorption of several carboxamide herbicides has been investigated (Weber & Peter, 1982). The *N*-substituted phenyl heterocyclic carboxamides are an important class of herbicides as protoporphyrinogen-IX oxidase inhibitors with advantages such as high resistance to soil leaching, low toxicity to birds, fish, and mammals, and slow development of weed resistance (Hirai, 1999).

We have designed and prepared a series of 113 carboxamide herbicides derived from pyrazinecarboxylic acid and various substituted anilines. The final compounds XIX were prepared by the anilinolysis of substituted pyrazinoylchlorides (Doležal, 1999, 2000, 2002, 2006b, 2007, 2008a, 2008b). Their chemical structure, hydrophobic parameters ($\log P$ calculated by ACD/logP ver. 1.0, 1996), and photosynthesis-inhibiting activity, structure-activity relationship (SAR) were studied. We synthesized in preference: *i*) the compounds with the lipophilic and/or electron-withdrawing substituents on the benzene moiety (R^3), *ii*) the compounds with the hydrophilic and/or electron-donating groups on the benzene part of molecule (R^3), and finally *iii*) the compounds with the lipophilic alkyl (R^2), *i.e.* methyl ($-\text{CH}_3$) or *tert*-butyl ($-\text{C}(\text{CH}_3)_3$) and/or halogen (chlorine) substitution (R^1) on the pyrazine nucleus, for their synthesis and structure see Fig. 4 and Table 1.

Fig. 4. Synthesis and structure of substituted *N*-phenylpyrazine-2-carboxamides (XIX).

3.1 Inhibition of photosynthetic electron transport by substituted *N*-phenylpyrazine-2-carboxamides

3.1.1 Photosynthetic electron transport in photosystem II

Photosystem II uses light energy to drive two chemical reactions: the oxidation of water and the reduction of plastoquinone. Five of redox components of PS II are known to be involved in transferring electrons from H_2O to the plastoquinone pool: the water oxidizing manganese cluster (Mn_4), the amino acid tyrosine (Y_z), the reaction center chlorophyll (P680), pheophytin, and two plastoquinone molecules, Q_A and Q_B (Fig. 5). Tyrosine, P680, pheophytin (Pheo), Q_A , and Q_B are bound to two key polypeptides (D_1 and D_2) that form the reaction center core of PS II and also provide ligands for the (Mn_4) cluster (Whitmarsh,

1998). After primary charge separation between P680 (chlorophyll *a*) and pheophytin (Pheo), $P680^+/Pheo^-$ is formed. Then electron is subsequently transferred from pheophytin to a plastoquinone molecule Q_A (permanently bound to PS II) acting as a one-electron acceptor.

Fig. 5. Scheme of the photosynthetic electron transport in photosystem II (PS II). (Taken from Photosystem II in <http://www.bio.ic.ac.uk/research/barber/psIIimages/PSII.jpg> with permission of Prof. Barber, Imperial College London).

From Q_A^- the electron is transferred to another plastoquinone molecule Q_B (acting as a two-electron acceptor); two photochemical turnovers of the reaction centre are necessary for the full reduction and protonation of Q_B . Because Q_B is loosely bound at the Q_B -site, reduced plastoquinone then unbinds from the reaction centre and diffuses in the hydrophobic core of the membrane and Q_B -binding site will be occupied by an oxidized plastoquinone molecule (Whitmarsh, 1998). Several commercial herbicides inhibit Photosynthetic electron transport (PET) by binding at or near the Q_B -site, preventing access to plastoquinone (e.g. Oettmeier, 1992). Photosystem II is the only known protein complex that can oxidize water, which results in the release of O_2 into the atmosphere. Oxidation of water is driven by the oxidized primary electron donor, $P680^+$ which oxidizes a tyrosine on the D_1 protein (Y_z) and four Mn ions present in the water oxidizing complex undergo light-induced oxidation, too. Water oxidation requires two molecules of water and involves four sequential turnovers of the reaction centre whereby each photochemical reaction creates an oxidant that removes one electron. The net reaction results in the release of one O_2 molecule, the deposition of four protons into the inner water phase, and the transfer of four electrons to the Q_B -site (producing two reduced plastoquinone molecules) (Whitmarsh & Govindjee, 1999).

PET in chloroplasts can be estimated by electrochemical measurements of oxygen concentration using Clark electrode (PET through the whole photosynthetic apparatus is registered) or by spectrophotometric methods enabling the monitoring of PET through individual parts of photosynthetic apparatus. The site of action of PET inhibitors can be

more closely specified by the use of chlorophyll fluorescence (*e.g.* Joshi & Mohanty, 2004) or by electron paramagnetic resonance (EPR) (*e.g.* Doležal et al., 2001a).

3.1.2 Hill reaction activity of *N*-phenylpyrazine-2-carboxamides

The Hill reaction is formerly defined as the photoreduction of an electron acceptor by the hydrogens of water, with the evolution of oxygen. *In vivo*, or in the organism, the final electron acceptor is NADP⁺, in isolated chloroplasts an artificial electron acceptor that changes colour as it is reduced, is applied. We tested a large series of pyrazinecarboxamides (XIX) for their activity related to oxygen evolution rate (OER) using spinach chloroplasts and 2,6-dichlorophenol-indophenol (DCPIP) as an electron acceptor what intercepts the electrons before they transfer to cytochrome *b_f* complex. Because the site of DCPIP action is plastoquinone pool (PQ) on the acceptor side of PS II (Izawa, 1980) this method is suitable for PET monitoring through PS II. The PET-inhibiting activities of the studied compounds XIX (expressed as IC₅₀ values) are summarized in Table 1.

No.	R ¹	R ²	R ³	IC ₅₀	Ref.	No.	R ¹	R ²	R ³	IC ₅₀	Ref.
1	Cl	H	2-Br	334	a	58	Cl	tBu	2-Cl,5-OH	652	i
2	H	tBu	2-Br	171	a	59	H	CH ₃	3-Br	648	b
3	Cl	tBu	2-Br	315	a	60	H	CH ₃	3-C≡CH	668	b
4	Cl	H	3,5-Br-4-OH	995	a	61	Cl	tBu	3-C≡CH	385	b
5	H	tBu	3,5-Br-4-OH	404	a	62	Cl	tBu	3-C≡N	375	b
6	Cl	tBu	3,5-Br-4-OH	590	a	63	H	CH ₃	3-Cl	174	b
7	Cl	H	3-OCH ₃	500	a	64	H	CH ₃	3-NO ₂	402	b
8	H	tBu	3-OCH ₃	800	a	65	H	CH ₃	2-C≡N-4-NO ₂	550	b
9	Cl	tBu	3-OCH ₃	644	a	66	H	CH ₃	3-I-4-CH ₃	317	b
10	Cl	H	3,5-OCH ₃	533	a	67	H	CH ₃	2-COOH	75	b
11	H	tBu	3,5-OCH ₃	317	a	68	Cl	tBu	3-F	262	c
12	Cl	tBu	3,5-OCH ₃	435	a	69	H	tBu	3-OH-4-Cl	105	c
13	Cl	H	5-Br-2-OH	146	a	70	Cl	tBu	3-OH-4-Cl	44	c
14	H	tBu	5-Br-2-OH	80	a	71	Cl	tBu	2-Cl	43	c
15	Cl	tBu	5-Br-2-OH	42	a	72	H	tBu	2-Cl	371	c
16	Cl	H	3,4-Cl	105	a	73	H	H	2-Cl	47	c
17	H	tBu	3,4-Cl	1525	a	74	Cl	H	2-CH ₃	1072	e
18	Cl	tBu	3,4-Cl	130	a	75	H	tBu	2-CH ₃	440	e
19	Cl	H	3-F	565	d	76	Cl	tBu	2-CH ₃	244	e
20	Cl	H	2,4-F	539	d	77	Cl	H	3-CH ₃	486	e
21	Cl	H	4-Cl	486	d	78	H	tBu	3-CH ₃	148	e
22	Cl	H	4-CH(CH ₃) ₂	118	d	79	Cl	tBu	3-CH ₃	118	e
23	H	tBu	3-F	313	d	80	H	tBu	2-OCH ₃	286	e
24	H	tBu	2,4-F	371	d	81	Cl	tBu	2-OCH ₃	97	e
25	H	tBu	4-Cl	1502	d	82	Cl	H	3-Br	313	e
26	H	tBu	4-CH(CH ₃) ₂	110	d	83	H	tBu	3-Br	81	e
27	Cl	tBu	3-F	129	d	84	Cl	tBu	3-Br	107	e
28	Cl	tBu	2,4-F	106	d	85	Cl	H	3,5-CF ₃	26	e

No.	R ¹	R ²	R ³	IC ₅₀	Ref.	No.	R ¹	R ²	R ³	IC ₅₀	Ref.
29	Cl	tBu	4-Cl	43	d	86	H	tBu	3,5-CF ₃	114	e
30	Cl	tBu	4-CH(CH ₃) ₂	52	d	87	Cl	tBu	3,5-CF ₃	241	e
31	Cl	H	2-OH	66	f	88	Cl	H	2,6-CH ₃	649	e
32	Cl	H	3-OH	2288	f	89	H	tBu	2,6-CH ₃	229	e
33	Cl	H	4-OH	3322	f	90	Cl	tBu	2,6-CH ₃	242	e
34	Cl	H	2-OH-5-Cl	8	f	91	H	H	2-Cl-5-OH	722	g
35	H	tBu	2-OH	205	f	92	H	H	4-F	480	g
36	H	tBu	3-OH	431	f	93	H	H	2-CF ₃	376	g
37	H	tBu	4-OH	314	f	94	H	H	3-CF ₃	130	g
38	H	tBu	2-OH-5-Cl	465	f	95	H	H	4-CH ₃	1475	g
39	Cl	tBu	2-OH	435	f	96	Cl	H	2-Cl-5-OH	624	g
40	Cl	tBu	3-OH	262	f	97	Cl	H	4-F	384	g
41	Cl	tBu	4-OH	43	f	98	Cl	H	2-CF ₃	557	g
42	Cl	tBu	2-OH-5-Cl	105	f	99	Cl	H	3-CF ₃	229	g
43	Cl	H	4-Cl-3-CH ₃	595	h	100	Cl	H	4-CH ₃	1524	g
44	Cl	H	3-I-4-CH ₃	51	h	101	H	tBu	4-F	524	g
45	H	tBu	4-Cl-3-CH ₃	190	h	102	H	tBu	2-CF ₃	55	g
46	Cl	tBu	2-F	69	h	103	H	tBu	3-CF ₃	283	g
47	Cl	tBu	4-CF ₃	184	h	104	H	tBu	4-CH ₃	164	g
48	H	H	4-F	480	i	105	Cl	tBu	2-Cl-5-OH	625	g
49	Cl	H	4-F	384	i	106	Cl	tBu	4-F	103	g
50	H	tBu	4-F	524	i	107	Cl	tBu	2-CF ₃	205	g
51	Cl	tBu	4-F	103	i	108	Cl	tBu	3-CF ₃	173	g
52	H	H	3-Cl	290	i	109	Cl	tBu	4-CH ₃	73	g
53	Cl	H	3-Cl	262	i	110	Cl	H	2,4,6-CH ₃	495	j
54	H	tBu	3-Cl	47	i	111	H	tBu	2,4,6-CH ₃	434	j
55	H	tBu	3-Cl	103	i	112	Cl	tBu	2,4,6-CH ₃	195	j
56	H	H	2-Cl-5-OH	722	i	113	H	tBu	4-COCH ₃	664	j
57	Cl	H	2-Cl-5-OH	624	i	-	-	-	-	-	-

Table 1. IC₅₀ values (in μmol dm⁻³) related to PET inhibition in spinach chloroplasts by substituted pyrazinecarboxamides XIX (Ref. Doležal et al., 2006b^(a), 2008a^(b), 2001b^(c), 2000^(d), 2002^(e), 1999^(f), 2008b^(g), 2007^(h), 2004⁽ⁱ⁾, 2001a^(j)).

The compounds **1-18** inhibited PET in spinach chloroplasts; however the inhibitory activity of the majority of these compounds was relatively low. The IC₅₀ values varied in the range from 42 to 1589 μmol dm⁻³, the most efficient inhibitors was 5-*tert*-butyl-6-chloro-*N*-(5-bromo-2-hydroxyphenyl)-pyrazine-2-carboxamide (**15**, Table 1). The dependence of PET-inhibiting activity of compounds **1-18** on the lipophilicity of the compounds (log *P*) is shown in Fig. 6, A. Markedly lowered solubility of **4-6** as well as **17** due to insertion of two halogen atoms (Br or Cl) in R³ substituent resulted in decreased inhibitory activity of these compounds. Based on the dependence of PET-inhibiting activity on log *P* of the rest compounds, these can be divided into two groups. In both groups increase of compound activity with increasing lipophilicity can be observed. Thus, with the exception of compounds **14** and **15** (R² = 5-Br-2-OH) it can be assumed, that the introduction of lipophilic

R^1 (Cl) and R^2 (*tert*-butyl, tBu) substituents, respectively, can result in partial decrease of the aqueous solubility and so in reduced inhibitory activity.

In other set of studied compounds **19–30**, compound **25** exhibited very low activity due to its low aqueous solubility (Table 1). As shown (Fig. 6, B), the PET-inhibiting activity of other compounds from the set expressed as $\log (1/IC_{50})$ increased linearly with increasing compound lipophilicity ($\log P$). The most active compounds from the set were 5-*tert*-butyl-6-chloro-*N*-(4-chlorophenyl)-pyrazine-2-carboxamide (**29**, $IC_{50} = 43 \mu\text{mol dm}^{-3}$) and 5-*tert*-butyl-6-chloro-*N*-(4-isopropylphenyl)-pyrazine-2-carboxamide (**30**, $IC_{50} = 52 \mu\text{mol dm}^{-3}$).

The inhibitory activity of the compounds **31–42** (Table 1) was affected not only by the lipophilicity of the compounds but also by the value of Hammett's constants of R^3 substituents. Very low activity of compounds **32** and **33** was connected with their low aqueous solubility. The most active compounds from this set were 6-chloro-*N*-(5-chloro-2-hydroxyphenyl)-pyrazine-2-carboxamide (**34**, $IC_{50} = 8 \mu\text{mol dm}^{-3}$) and 5-*tert*-butyl-6-chloro-*N*-(4-hydroxyphenyl)-pyrazine-2-carboxamide (**41**, $IC_{50} = 43 \mu\text{mol dm}^{-3}$), the activity of rest compounds from the set varied between 66 (**31**) and $465 \mu\text{mol dm}^{-3}$ (**38**).

Fig. 6. The dependence of PET-inhibiting activity of compounds **1–18** (A) and compounds **19–30** (B) on the lipophilicity of the compounds ($\log P$).

It was found that from the aspect of inhibitory activity it is much more favourable when on the phenyl ring (R^3 substituent) halogen atom occurs in *meta* and methyl moiety in *para* position (**44**, $IC_{50} = 51 \mu\text{mol dm}^{-3}$) in comparison with compound **43** where $R^3 = 4\text{-Cl-3-CH}_3$ ($IC_{50} = 595 \mu\text{mol dm}^{-3}$). However, the inhibitory activity of the above mentioned compound **43** can be increased by introduction of *tert*-butyl substituent instead of H in R^2 (**45**, $IC_{50} = 190 \mu\text{mol dm}^{-3}$). The IC_{50} values related to PET-inhibiting activity of compounds **48–58** varied in the range from 47.0 (**54**) to $722 \mu\text{mol dm}^{-3}$ (**56**). The inhibitory activity of majority of these compounds was relatively low, the most efficient inhibitors were 5-*tert*-butyl-6-chloro-*N*-(4-fluorophenyl)-pyrazine-2-carboxamide (**51**), *N*-(2-chloro-5-hydroxyphenyl)-pyrazine-2-carboxamide (**55**, both $IC_{50} = 103.0 \mu\text{mol dm}^{-3}$), and especially 5-*tert*-butyl-6-chloro-*N*-(3-chlorophenyl)-pyrazine-2-carboxamide (**54**, $IC_{50} = 47.0 \mu\text{mol dm}^{-3}$). Their $\log P$ values calculated ranged between 3.28 and 4.18.

In the set of compounds **59–67** the PET-inhibiting activity of compounds **61**, **62**, **63**, **66** and **67** (Fig. 7, A) expressed as $\log (1/IC_{50})$ showed a linear decrease with increasing values of lipophilicity parameter ($\log P$). On the other hand, the biological activity of compounds **59**,

60, **64** and **65** was significantly lower and linear decrease of PET-inhibiting activity with increasing $\log P$ values was less sharp indicating that the biological activity of compounds **59-67** depended both on the compound lipophilicity as well as on Hammett's constants σ of the substituent R^2 . The most active PET inhibitor from this set was found to be 2-(5-methylpyrazine-2-carboxamido)-benzoic acid (**67**, $IC_{50} = 75.0 \mu\text{mol dm}^{-3}$) (Doležal et al., 2008a). From the set of compounds **68-73** the most active inhibitors with comparable inhibitory activity were compounds 5-*tert*-butyl-6-chloro-*N*-(3-chloro-4-hydroxyphenyl)-pyrazine-2-carboxamide (**70**, $IC_{50} = 44 \mu\text{mol dm}^{-3}$), 5-*tert*-butyl-6-chloro-*N*-(2-chlorophenyl)-pyrazine-2-carboxamide (**71**, $IC_{50} = 43 \mu\text{mol dm}^{-3}$) and *N*-(2-chlorophenyl)-pyrazine-2-carboxamide (**73**, $IC_{50} = 47 \mu\text{mol dm}^{-3}$).

Fig. 7. The dependence of PET-inhibiting activity of compounds **59-67** (A) and compounds **74-90** (B) on the lipophilicity of the compounds ($\log P$).

In the set of compounds **74-90** the IC_{50} values related to PET inhibition varied in the range from 26 (**85**) to 1072 $\mu\text{mol dm}^{-3}$ (**74**), see Table 1. In general, the inhibitory activity of these compounds depended on their lipophilicity showing a quasi-parabolic trend (Fig. 7, B). However, the studied compounds could be divided into two groups. The compounds with 2-CH₃ substituents on the phenyl ring (**74**, **75**, **76**, **88**, **89** and **90**, squares in Fig. 7, B) had lower biological activity than the other investigated compounds with comparable $\log P$ values. Consequently, it can be assumed that the methyl substituent in *ortho* position of the benzene ring is disadvantageous from the viewpoint of interactions with the photosynthetic apparatus. On the other hand, compound **85** (6-chloro-*N*-(3,5-trifluoro-methylphenyl)-pyrazine-2-carboxamide) exhibited higher inhibitory activity than expected.

The majority of compounds **91-109** inhibited PET in spinach chloroplasts; however their inhibitory activity was rather low. From the obtained results it can be concluded that the activity depended on the lipophilicity and also on the electron accepting or withdrawing power of R^3 substituent(s). The most effective inhibitor was compound **102** (5-*tert*-butyl-*N*-(2-trifluoromethylphenyl)-pyrazine-2-carboxamide, $IC_{50} = 55 \mu\text{mol dm}^{-3}$). Among the three most active compounds **102**, **109** and **106** the optimal values of lipophilicity ranges from $\log P = 4.02-4.41$. On the other hand, for the group of compounds **105**, **108** and **107** with the highest lipophilicity, the PET-inhibiting activity showed a decrease with increasing compound lipophilicity. The most effective inhibitor from the compounds with $R^3 = 2,4,6$ -

CH₃ was 5-*tert*-butyl-6-chloro-*N*-(2,4,6-methylphenyl)-pyrazine-2-carboxamide (**112**, IC₅₀ = 195 μmol dm⁻³) (Doležal et al., 2001a).

3.1.3 Determination of the site of inhibitory action of *N*-phenylpyrazine-2-carboxamides in the photosynthetic electron transport chain by electron paramagnetic resonance spectroscopy and chlorophyll *a* fluorescence measurements

The site of inhibitory action of some *N*-phenylpyrazine-2-carboxamides XIX in the photosynthetic electron transport chain was investigated using spinach (*Spinacia oleracea* L.) chloroplasts. For this purpose electron paramagnetic resonance spectroscopy (EPR) and measurement of chlorophyll *a* fluorescence were used.

Intact chloroplasts of algae and vascular plants exhibit EPR signals in the region of free radicals ($g = 2.00$), which are stable during several hours (Hoff, 1979) and could be registered at laboratory temperature by conventional continual wave EPR apparatus. These signals were denoted as signal I ($g = 2.0026$, $\Delta B_{pp} = 0.8$ mT) and signal II ($g = 2.0046$, $\Delta B_{pp} = 2$ mT) indicating their connection with photosystem (PS) I and PS II, respectively (Weaver, 1968). Signal II consists from two components, namely signal II_{slow} which is observable in the dark and signal II_{very fast} which occurs at irradiation of chloroplasts by visible light and represents intensity increase of signal II at irradiation of chloroplasts by the visible light. It was found that signal II_{slow} belongs to the intermediate D[•] and signal II_{very fast} belongs to the intermediate Z[•]. Intermediates Z[•] and D[•] are tyrosine radicals which are situated at 161st position in D₁ and D₂ proteins which are located on the donor side of PS II (Svensson et al., 1991). The EPR signal I is associated with cation radical of chlorophyll *a* dimmer situated in the core of PS I (Hoff, 1979).

Using EPR spectroscopy it has been found that the studied compounds XIX affect predominantly the intensity of EPR signal II, mainly the intensity of its constituent signal II_{slow}. As mentioned above, the signal II_{slow} is well observable in the dark (see Fig. 8, full line) and it belongs to the D[•] intermediate, *i.e.* tyrosine (Tyr_D or Y_D) radical which is located on the donor side of PS II in the 161st position in D₂ protein (Svensson et al., 1991; see Fig. 5). From Fig. 8 it is evident that the intensity of signal II_{slow} has been decreased by the studied compounds (see Fig. 8, B and C, full lines). That means that in the suspension of spinach chloroplasts the 5-*tert*-butyl-6-chloro-*N*-(3-fluorophenyl)-pyrazine-2-carboxamide (**68**) and 5-*tert*-butyl-*N*-(3-hydroxy-4-chlorophenyl)-pyrazine-2-carboxamide (**69**) interact with the D[•] intermediate. Due to this interaction of the studied anilides with this part of PS II, the photosynthetic electron transport from the oxygen evolving complex to the reaction centre of PS II is impaired. Consequently, the electron transport between PS II and PS I is inhibited as well and a pronounced increase of signal I intensity in the light can be observed (see Fig. 8, B and C, dashed lines). The signal I ($g = 2.0026$, $\Delta B_{pp} = 0.8$ mT) belongs to the cation radical of chlorophyll *a* dimmer in the reaction centre of PS I (Hoff, 1979).

Similar site of action in the photosynthetic apparatus of spinach chloroplasts was confirmed for 2-alkylsulfanylpuridine-4-carbothioamides (Kráľová et al., 1997) and substituted benzanilides and thiobenzanilides (Kráľová et al., 1999). From Fig. 8 it is evident that the decrease of signal II_{slow} is greater in the presence of compound **69** (Fig. 8, B) than in presence of compound **68** (Fig. 8, C). These results are in agreement with those obtained for OER inhibition in spinach chloroplasts (Table 1, IC₅₀ = 105 μmol dm⁻³ for **69** and 262 μmol dm⁻³ for **68**).

1,5-Diphenylcarbazine (DPC) is an artificial electron donor acting in $Z^{\bullet}/D^{\bullet}$ intermediate (Jegerschöld & Styring, 1991). By addition of DPC to chloroplasts inhibited by PET inhibitors the supply of electrons to P680 is secured. However, the complete restoration of the electron transport to PS I occurs only in the case that photosynthetic electron transport chain between $Z^{\bullet}/D^{\bullet}$ and plastoquinone is not damaged. After addition of DPC to chloroplasts inhibited by the studied anilides up to 70-80%, the OER in the suspension of spinach chloroplasts was not completely restored. It was restored only up to 55-75% of the untreated control sample what indicated that also some member of the photosynthetic electron transport chain between $Z^{\bullet}/D^{\bullet}$ intermediate and plastoquinone is partially damaged by the studied compounds in the light (dashed lines).

Fig. 8. EPR spectra of the untreated spinach chloroplasts (A) and in the presence of 0.05 mol dm^{-3} of 5-*tert*-butyl-*N*-(3-hydroxy-4-chlorophenyl)-pyrazine-2-carboxamide (**69**, B) and 5-*tert*-butyl-6-chloro-*N*-(3-fluorophenyl)-pyrazine-2-carboxamide (**68**, C) registered in the dark (full lines) and in the light (dashed lines). (Ref. Doležal et al., 2001a; reprinted with permission of editor).

The effects of *N*-phenylpyrazine-2-carboxamides XIX on the photosynthetic centres of spinach chloroplasts were investigated by studying chlorophyll *a* fluorescence. Fluorescence emission spectra of spinach chloroplasts were recorded on fluorescence spectrophotometer F-2000 (Hitachi, Japan) using excitation wavelength $\lambda_{\text{ex}} = 436$ nm for monitoring fluorescence of chlorophyll *a* and the samples were kept in the dark 10 min before measuring (Doležal et al., 2001a). When chloroplasts were irradiated with the light of $\lambda_{\text{ex}} = 436$ nm, an emission band with the maximum at $\lambda = 686$ nm was observed. This band belongs to the pigment-protein complexes present mainly in photosystem II (Govindjee, 1995). It was found that chloroplasts treated with the studied compounds exhibited quenching of the emission of Chl *a* molecules. Fig. 9 presents the dependence of F/F_{contr} in the suspension of spinach chloroplasts (F_{contr} – fluorescence intensity at $\lambda = 686$ nm in the control, F – fluorescence intensity at $\lambda = 686$ nm in the presence of the studied compound)

on the concentration of 5-*tert*-butyl-*N*-(3-hydroxy-4-chlorophenyl)-pyrazine-2-carboxamide (**69**), 5-*tert*-butyl-6-chloro-*N*-(3-hydroxy-4-chlorophenyl)-pyrazine-2-carboxamide (**70**), 5-*tert*-butyl-6-chloro-*N*-(2-chlorophenyl)-pyrazine-2-carboxamide (**71**), and 5-*tert*-butyl-(2-chlorophenyl)-pyrazine-2-carboxamide (**72**). The greater is the fluorescence quenching, the more efficient is the interaction of the inhibitor with pigment-protein complexes in photosystem II. For the investigated compounds the intensity of this interaction showed a decrease in the following order: **70** > **69** > **72** > **71** (Fig. 9).

Fig. 9. Dependence of the fluorescence quenching on the concentration of 5-*tert*-butyl-6-chloro-*N*-(3-hydroxy-4-chlorophenyl)-pyrazine-2-carboxamide (**70**, squares), 5-*tert*-butyl-*N*-(3-hydroxy-4-chlorophenyl)-pyrazine-2-carboxamide (**69**, circles), 5-*tert*-butyl-(2-chlorophenyl)-pyrazine-2-carboxamide (**72**, down triangles) and 5-*tert*-butyl-6-chloro-*N*-(2-chlorophenyl)-pyrazine-2-carboxamide (**71**, up triangles) (F_{contr} = fluorescence of the untreated suspension of spinach chloroplasts; F = fluorescence of anilide treated suspension of spinach chloroplasts; $\lambda = 686$ nm). (Ref. Doležal et al., 2001a; reprinted with permission of editor).

The most effective compounds (**70** and **71**) contained two Cl substituents in their molecules. The results of fluorescence study obtained for compounds **70**, **69** and **72** are in agreement with those obtained for OER evolution in spinach chloroplasts (Table 1; $\text{IC}_{50} = 44$ (**70**), 105 (**69**) and $371 \mu\text{mol dm}^{-3}$ (**72**)). However, the fluorescence of the chloroplast suspension was not affected by 5-*tert*-butyl-6-chloro-*N*-(2-chlorophenyl)-pyrazine-2-carboxamide (**71**) which can be considered as relatively effective inhibitor of OER ($\text{IC}_{50} = 43 \mu\text{mol dm}^{-3}$). This can be explained with the decreased aqueous solubility of this compound. Whereas in the OER experiments the investigated compounds were dissolved in dimethyl sulfoxide, in fluorescence experiments ethanolic solutions were used and after evaporation of the solvent the compound was dissolved directly in the aqueous chloroplast suspension. Consequently, it can be assumed that the fluorescence was not affected due to insolubility of compound **71**.

in this suspension. The quenching of the fluorescence intensity at $\lambda = 686$ nm produced by the studied compounds suggested PS II as the site of action of the studied compounds.

3.1.4 Inhibition of oxygen evolution rate in suspensions of *Chlorella vulgaris* by *N*-phenylpyrazine-2-carboxamides

The inhibition of oxygen evolution rate (OER) in the suspension of *Chlorella vulgaris* was investigated with two model inhibitors (compounds **69** and **72**). The dependences of OER (expressed as the percentage of the untreated control sample) on the concentrations of compounds 5-*tert*-butyl-*N*-(3-hydroxy-4-chlorophenyl)-pyrazine-2-carboxamide (**69**) and 5-*tert*-butyl-*N*-(2-chlorophenyl)-pyrazine-2-carboxamide (**72**) are shown in Fig. 10. It is evident that both investigated compounds inhibited OER in the suspension of *Chlorella vulgaris* algae. Compound **69** was more effective inhibitor than compound **72** what is reflected in the corresponding IC_{50} values ($99 \mu\text{mol dm}^{-3}$ for **69** and $329 \mu\text{mol dm}^{-3}$ for **72**). These results are in good agreement with those obtained for inhibition of OER in spinach chloroplasts (Table 1). The introduction of hydroxyl moiety in compound **69** enhanced its photosynthesis-inhibiting activity with respect to that of compound **72** approximately threefold.

Fig. 10. Dependence of OER in the suspension of *Chlorella vulgaris* (expressed as the percentage of the control) on the concentration of 5-*tert*-butyl-*N*-(2-chlorophenyl)-pyrazine-2-carboxamide (**72**, triangles) and 5-*tert*-butyl-*N*-(3-hydroxy-4-chlorophenyl)-pyrazine-2-carboxamide (**69**, circles). (Ref. Doležal et al., 2001a; reprinted with permission of editor).

3.1.5 Reduction of chlorophyll content in *Chlorella vulgaris* by *N*-phenylpyrazine-2-carboxamides

Toxic effects of environmental pollutants on algae which are essential components of aquatic ecosystems can directly affect the structure and function of ecosystem (Campanella et al., 2000). Herbicides can alter species composition of an algal community what could result in modified structure and function of aquatic communities. Ma et al. (2000) examined the effects of 40 herbicides (belonging to 18 different chemical classes with nine different modes of action) on the green alga *Raphidocelis subcapitata* (formerly named *Selenastrum capricornutum*) and found that the highest acute toxicity exhibited herbicides acting as photosynthesis inhibitors. Photosynthetic pigments have often been used as biomarkers of exposure to different classes of herbicides in autotrophic plants including algae (Blaise, 1993;

Sandmann, 1993). The inhibitory effectiveness of some substituted pyrazinecarboxamides related to reduction of chlorophyll content in *Chlorella vulgaris* expressed by IC₅₀ values is summarized in Table 2. The dependence of log (1/IC₅₀) on the compound lipophilicity (log *P*) showed a quasi-parabolic course (Fig. 11).

No.	R ¹	R ²	R ³	IC ₅₀	Ref.
43	Cl	H	4-Cl-3-CH ₃	80	h
44	Cl	H	3-I-4-CH ₃	44	h
45	H	tBu	4-Cl-3-CH ₃	89	h
79	Cl	tBu	3-CH ₃	63	e
84	Cl	tBu	3-Br	67	e
85	Cl	H	3,5-CF ₃	125	e
86	H	tBu	3,5-CF ₃	208	e
87	Cl	tBu	3,5-CF ₃	356	e
88	Cl	H	2,6-CH ₃	79	e
95	H	H	4-CH ₃	71	b
97	Cl	H	4-F	32	b
100	Cl	H	4-CH ₃	37	b
102	H	tBu	2-CF ₃	33	b

Table 2. IC₅₀ values (in μmol dm⁻³) related to reduction of chlorophyll content in *Chlorella vulgaris* of some substituted pyrazinecarboxamides XIX. Cultivation conditions: 7 days; photoperiod 16 h light/8 h dark; photosynthetic active radiation 80 μmol m⁻² s⁻¹; pH = 7.2; Chl content in the suspensions at the beginning of the cultivation was 0.01 mg dm⁻³. (Ref. Doležal et al., 2007^(h), 2002^(e), 2008a^(b)).

Fig. 11. The dependence of antialgal activity expressed as log (1/IC₅₀) on the lipophilicity (log *P*) of some substituted pyrazinecarboxamides XIX.

However, differences in IC₅₀ values of compounds with comparable lipophilicity indicate that the biological activity is affected beside of lipophilicity also by the electronic properties

of R³ substituent(s). Because of too low aqueous solubility of many compounds from the tested set of pyrazinecarboxamides XIX the compounds fall out during experiment (7 days) and the corresponding IC₅₀ values could be determined only for limited number of compounds.

4. Photosynthesis-inhibiting pyrazine analogues of chalcones

Chalcones and related compounds "chalconoids" are aromatic ketones containing two aromatic rings linked with three carbon chain. The presence of an unsaturated double bound is typical for chalcones. Hence, chalcones are 1,3-diarylprop-2-ones. They show antibacterial, antifungal, antitumor and anti-inflammatory properties (Dimmock et al., 1999). The aim of our project was the isosteric replacement of a phenyl moiety in chalcones with the pyrazine ring to form some pyrazine analogues of chalcones ("diazachalcones"). Several series (thirty two compounds) of ring substituted (*E*)-3-phenyl-1-(pyrazin-2-yl)-prop-2-en-1-ones XX (Fig. 12) were prepared in our laboratories by means of modified Claisen-Schmidt condensation of acetylpyrazines with aromatic aldehydes (Opletalová et al., 2002, Opletalová et al., 2006, Chlupáčová et al., 2005).

Fig. 12. Pyrazine analogues of chalcones XX (R¹ = H, alkyl; R² = OH, NO₂, Cl).

Ring substituted (*E*)-3-phenyl-1-(pyrazin-2-yl)-prop-2-en-1-ones XX were tested for their activity related to OER inhibition in spinach chloroplasts and *Chlorella vulgaris* as well as reduction of chlorophyll content in statically cultured suspensions of freshwater alga *Chlorella vulgaris*. The corresponding IC₅₀ values are summarized in Tables 3 and 4.

No.	R ¹	R ²	OER inhibition/IC ₅₀	
			<i>S. oleracea</i>	<i>C. vulgaris</i>
114	tBu	2-OH	167	78
115	isoBu	2-OH	144	63
116	nBu	2-OH	184	147
117	nPro	2-OH	187	100
118	tBu	4-OH	315	279
119	isoBu	4-OH	235	232
120	nBu	4-OH	306	265
121	nPro	4-OH	399	514

Table 3. IC₅₀ values (in μmol dm⁻³) related to OER inhibition in spinach chloroplasts and *Chlorella vulgaris* by diazachalcones XX. (Ref. Opletalová et al., 2002).

The inhibition of OER in spinach chloroplasts by substituted diazachalcones XX (**114-121**) (Fig. 12) has been investigated spectrophotometrically, using DCPIP as an electron acceptor (Král'ová et al., 1992). For the study of OER inhibition in the algal suspensions a Clark type electrode has been used. The IC₅₀ values of compounds **114-121** related to OER inhibition varied in the range of 144-399 µmol dm⁻³ for spinach chloroplasts) and 63-514 µmol dm⁻³ for algal suspension of *Chlorella vulgaris* (Table 3). 2-Hydroxy substituted derivatives were found to be more effective inhibitors of photosynthesis than the 4-hydroxy substituted ones. The inhibitory activity of 2-hydroxy substituted derivatives was affected also by the branching of R¹ substituent: OER inhibition in photosynthesizing organisms by the isomers with branched alkyl chain (*tert*-butyl, *isobutyl*) was more pronounced than by the isomer with unbranched alkyl substituent (*n*-butyl) (Opletalová et al., 2002).

No.	R ¹	R ²	IC ₅₀		Ref.
			PET inhibition <i>S. oleracea</i>	Chl. content reduction <i>C. vulgaris</i>	
122	H	2-NO ₂	ND	70.6	k
123	tBu	2-NO ₂	325.0	ND	k
124	isoBu	2-NO ₂	ND	118.0	k
125	nBu	2-NO ₂	393.0	585.0	k
126	nPro	2-NO ₂	ND	123.0	k
127	H	3-NO ₂	658.0	19.6	k
128	tBu	3-NO ₂	461.0	ND	k
129	isoBu	3-NO ₂	340.0	62.8	k
130	nBu	3-NO ₂	236.0	ND	k
131	nPro	3-NO ₂	ND	18.6	k
132	H	4-NO ₂	ND	44.9	k
133	tBu	4-NO ₂	ND	ND	k
134	isoBu	4-NO ₂	ND	ND	k
135	nBu	4-NO ₂	706.0	ND	k
136	nPro	4-NO ₂	ND	238.3	k
137	H	3-OH	877.0	32.5	l
138	tBu	3-OH	105.0	238.3	l
139	isoBu	3-OH	256.0	65.5	l
140	nBu	3-OH	ND	95.9	l
141	nPro	3-OH	ND	69.9	l
142	H	4-Cl	ND	24.5	l
143	tBu	4-Cl	181.0	ND	l
144	isoBu	4-Cl	246.0	ND	l
145	nBu	4-Cl	374.0	ND	l

Table 4. IC₅₀ values (in µmol dm⁻³) related to PET inhibition in spinach chloroplasts and IC₅₀ values (in µmol dm⁻³) related to reduction of chlorophyll content in statically cultivated *Chlorella vulgaris* determined for diazachalcones XX. (Ref. Opletalová et al., 2006^(k), Chlupáčová et al., 2005^(l)), ND - not determined.

The effects of substituted diazachalcones XX (**114-121**) on the photosynthetic centres of chloroplasts were investigated by studying chlorophyll *a* fluorescence. The decreased intensity of the emission band at 686 nm, belonging to the pigment-protein complexes in photosystem (PS) II, suggested PS II as the site of action of the studied compounds (Kráľová et al., 1998).

Using EPR spectroscopy it has been found that in spinach chloroplasts the intensity of EPR signal II, mainly the intensity of its constituent signal II_{slow}, showed a decrease by the studied compounds **114-121**. Consequently it can be concluded that the studied compounds, similarly to *N*-phenylpyrazine-2-carboxamides (Doležal et al., 2001a), interact with D[•] intermediate, *i.e.* with the tyrosine radical in 161st position (Tyr_D; Y_D) which is located in D₂ protein on the donor side of PS II (Fig. 5). Due to interaction of the studied compounds with D[•] intermediate PET from the oxygen evolving complex to the core of PS II is impaired. A pronounced increase of EPR signal I intensity in the light belonging to the cation-radical of chlorophyll *a* dimmer in the core of PS I indicated that the electron transport between PS II and PS I is inhibited as well. However, addition of DPC to chloroplasts inhibited by the studied compounds completely restored the reduction of DCPIP indicating that the core of PS II (P680) and a part of the electron transport chain - at least up to plastoquinone - remained intact. These results are in accordance with those obtained with 2-alkylsulfanylpiperidine-4-carbothioamides (Kráľová et al., 1997). Similar study with anilides of 2-alkylpyridine-4-carboxylic acids has shown that also the core of PS II was partially impaired by these inhibitors of photosynthetic electron transport (Kráľová et al., 1998a). On the other hand, after addition of DPC to chloroplasts inhibited by the studied *N*-phenylpyrazine-2-carboxamides **68** and **69** up to 70-80%, the OER in the suspension of spinach chloroplasts was restored only up to 55-75% of the untreated control sample indicating that also some member of the photosynthetic electron transport chain between Z[•]/D[•] and plastoquinone was partially damaged by the these compounds (Doležal et al., 2001a).

In general, in the series of diazachalcones **122-145** the most effective reduction of chlorophyll content in the suspensions of *C. vulgaris* showed compounds with R¹ = H (Table 4): **127** (R² = 3-NO₂; IC₅₀ = 19.6 μmol dm⁻³), **142** (R² = 4-Cl; IC₅₀ = 24.5 μmol dm⁻³), **137** (R² = 3-OH; IC₅₀ = 32.5 μmol dm⁻³), **132** (R² = 4-NO₂; IC₅₀ = 44.9 μmol dm⁻³) and **122** (R² = 2-NO₂; IC₅₀ = 70.6 μmol dm⁻³). However, the highest anti-algal activity from this series showed compound **131** (R¹ = CH₃CH₂CH₂, R² = 3-NO₂; IC₅₀ = 18.6 μmol dm⁻³). On the other hand, the most effective inhibitors of PET in spinach chloroplasts were found to be two compounds with R¹ = C(CH₃)₃, namely **138** (R² = 3-OH; IC₅₀ = 105 μmol dm⁻³) and **143** (R² = 4-Cl; IC₅₀ = 181 μmol dm⁻³) whereby IC₅₀ values for several compounds could not be determined due to too low solubility of these compounds.

5. Conclusion

Pyrazines are a class of compounds that occur almost ubiquitously in nature. The worldwide distribution of pyrazines in plants, insects, terrestrial vertebrates, marine organisms, fungi and bacteria, their specific properties, including their using as drugs, fungicides and herbicides invite reasonable attention. Our review brings the basic information about some commercially produced pyrazine herbicides including their mechanism of action as well as survey of patented herbicidal pyrazine derivatives. Special attention was paid to the original compounds from series of 113 substituted *N*-

phenylpyrazine-2-carboxamides XIX and 32 diazachalcones XX prepared and evaluated in our laboratories. In first series, pyrazinecarboxamides XIX connected via -CONH- bridge with substituted anilines can form centrosymmetric dimer pairs with the peptidic carboxamido group of some peptides, needed for binding to the receptor site, possibly by formation of hydrogen bonds. All compounds were tested as potential inhibitors of the photosynthetic electron transport in spinach chloroplasts. Based on the obtained results it could be assumed that the biological activity of the studied substituted pyrazinecarboxamides did not depend exclusively on the compound lipophilicity but it was also affected by electron accepting or withdrawing power of the substituents on the aromatic benzene ring. The site of action of some substituted *N*-phenylpyrazine-2-carboxamides XIX in the photosynthetic apparatus of spinach chloroplasts was studied using fluorescence and EPR spectroscopy. It was found that the studied compounds cause quenching of the chlorophyll *a* fluorescence at 685 nm belonging mainly to the pigment–protein complexes in photosystem (PS) II. The extent of the fluorescence quenching correlated with the effectiveness of the compounds concerning inhibition of oxygen evolution rate (OER) in spinach chloroplasts. Using EPR spectroscopy it was confirmed that the title compounds interact with the intermediate $D^{\bullet}$ (Tyr_D), *i.e.* with the tyrosine radical, which is situated on the donor side of PS II at the 161th position of D₂ protein. It was found that the studied compounds inhibit OER not only in the suspension of spinach chloroplasts but also in the suspensions of *Chlorella vulgaris*. Introducing of Cl substituents into aromatic ring as well as pyrazine moiety of the studied molecules enhanced the effectiveness of OER–inhibiting activity. Some *N*-phenylpyrazine-2-carboxamides XIX reduced chlorophyll content in *Chlorella vulgaris* whereby their biological activity was affected beside of lipophilicity also by the electronic properties of R³ substituent(s). The most effective inhibitor from the series XIX was 6-chloro-*N*-(5-chloro-2-hydroxyphenyl)-pyrazine-2-carboxamide (**34**, IC₅₀ = 8 μmol dm⁻³; Doležal, 1999).

The studied pyrazine analogues of chalcones, diazachalcones XX also reduced the rate of oxygen evolution in spinach chloroplasts and *C. vulgaris*, whereby the inhibitory activity of *ortho*-hydroxyl substituted derivatives XX was greater than that of *para*-hydroxyl substituted ones. The lowest IC₅₀ values were found with compounds having a branched alkyl group on the pyrazine ring. The photosynthesis-inhibiting activity of nitro derivatives was lower than that of the corresponding hydroxylated analogs. In general, in the series of diazachalcones with R² = 2-NO₂; 3-NO₂; 4-NO₂; 3-OH and 4-Cl, the most effective reduction of chlorophyll content in the suspensions of *C. vulgaris* showed compounds with R¹ = H. It was confirmed that studied diazachalcones interact with $D^{\bullet}$ intermediate, *i.e.* with the tyrosine radical in 161st position (Tyr_D) which is located in D₂ protein on the donor side of PS II and that they do not damage the core of PS II (P680) and a part of the electron transport chain - at least up to plastoquinone.

6. Acknowledgments

We dedicate the present article to Assoc. Prof. Jiří Hartl, former leader of our research team. Thanks to all our colleague and numerous graduate and undergraduate students who participated on the synthesis of biologically active pyrazines. The work was partly supported by the Ministry of Education of the Czech Republic (MSM002160822) and Sanofi Aventis Pharma Slovakia.

7. References

- ACD/logP, ver. 1.0 (Advanced Chemistry Development Inc., Toronto, 1996)
- Adams, T. B.; Doull, J.; Feron, V. J.; Goodman, J. I.; Marnett, L. J.; Munro, I. C.; Newberne, P. M.; Portoghese, P. S.; Smith, R. L.; Waddell, W. J. & Wagner, B. M. (2002). The FEMA GRAS assessment of pyrazine derivatives used as flavor ingredients. *Food Chem. Toxicol.*, Vol. 40, No 4, 429-451, ISSN 0278-6915
- Blaise, C. R. (1993). Practical laboratory applications with micro-algae for hazard assessment of aquatic contaminants. In: Richardson, M. (ed.), *Ecotoxicology Monitoring*, pp. 83-107, VCH, ISBN 3-527-28560-1, Weinheim
- Beck, H. C.; Hansen, A. M. & Lauritsen, F. R. (2003). Novel pyrazine metabolites found in polymyxin biosynthesis by *Paenibacillus polymyxa*. *FEMS Microbiol. Lett.*, Vol. 220, 67-73, ISSN 0378-1097
- Benko, Z.; Jachetta, J. J.; Costales, M. J. & Arndt K. E. (1997). N-aryl[1,2,4]triazolo[1,5-a]pyrazine-2-sulfonamide herbicides. US Patent 5602075, February 11, 1997
- Bird, C. W. (1992). Heteroaromaticity, 5, a unified aromaticity index. *Tetrahedron*, Vol. 48, No. 2, 335-340, ISSN 0040-4020
- Böhner, B.; Fory, W.; Schurter, R. & Pissiotas, G. (1987). Pyrazine-sulfonylcarbamates and thiocarbamates. US Patent 4684730, August 4, 1987
- Böhner, B.; Fory, W.; Schurter, R. & Pissiotas, G. (1987a). Pyrazinyl sulfonamides useful as intermediates for herbicides. US Patent 4707551, November 17, 1987
- Böhner, B. & Meyer, W. (1989a). Aminopyrazinones. US Patent 4871853, October 3, 1989
- Böhner, B. & Meyer, W. (1989b). Aminopyrazinones and aminotriazinones. US Patent 4795486, January 3, 1989
- Böhner, B. & Meyer, W. (1990). Aminopyrazinones and aminotriazinones. US Patent 4920223, April 24, 1990
- Brown, D. J. (2002). *The Pyrazines, Supplement I. The Chemistry of Heterocyclic Compounds*. John Wiley & Sons, Inc., ISBN 9780471403821, New York, 576 p.
- Brown III, C. J. & Knaust, J. M. (2009). 2,5-Dimethylpyrazine 1,4-dioxide. *Acta Cryst. Sect. E - Struct. Reports Online*, Vol. 65 (December), No. 12, o3052-sup-9, ISSN 1600-5368
- Campanella, L.; Cubadda, F. Sammartino, M. P. & Saoncella, A. (2000). An algal biosensor for the monitoring of water toxicity in estuarine environments. *Water Res.*, Vol. 25, No. 1, 69-76, ISBN 0043-1354
- Campbell, K. R.; Bartell, S. M. & Shaw, J. L. (2000). Characterizing aquatic ecological risks from pesticides using a diquat dibromide case study. II. Approaches using quotients and distributions. *Environ. Toxicol. Chem.*, Vol. 19, No. 3, 760-774, ISSN 0730-7268
- Camper, N. D. & McDonald, S. K. (1989). Tissue and cell cultures as model system in herbicide research. *Rev. Weed Sci.*, Vol. 4, No. 2, 169-190, ISSN 1042-4148
- Cedergreen, N.; Kamper, A. & Streibig, J. C. (2006). Is prochloraz a potent synergist across aquatic species? A study on bacteria, daphnia, algae and higher plants. *Aquat. Toxicol.*, Vol. 78, No. 3, 243-252, ISSN 0166-445X
- Chlupáčová, M.; Opletalová, V.; Kuneš J. & Kráľová, K. (2005). Synthesis of 3-alkyl- and 3-arylsulfanyl-1,3,-diphenylpropane-1-ones and their effects on two

- photosynthesizing organisms. *Folia Pharm. Univ. Carol.*, Vol. 31-32, 109-115, ISSN 1210-9495
- Cordingley, M. R.; Turnbull, M. D.; Carter, N. B. & Crowley, P. J. (2009). *Herbicide pyridopyrazines and their preparation and use in controlling plants*. WO 2009090402, January 16, 2009
- Davenport, H. E. (1963). The mechanism of cyclic phosphorylation by illuminated chloroplasts. *Proc. R. Soc. B*, Vol. 157, March 26, pp. 332-345, ISSN 1471-2954
- Diaz, T. G.; Merás, I. D. & Franco, M. F. (2002). Stopped flow kinetic-spectrophotometric determination of Diquat in waters. *Water Res.*, Vol. 36, No. 3, 783-787, ISSN 0043-1354
- Dietsche, T. J. (1977). 6,7-Dichloro-3,4-dihydro-2H-pyrazino(2,3-b)(1,4) oxazines. US Patent 49029657, June 14, 1977
- Dimmock, J. R.; Elias, D. W.; Beazely, M. A. & Kandepu, N. M. (1999). Bioactivities of chalcones. *Curr. Med. Chem.*, Vol. 6, No. 12, 1125-1149, ISSN 0929-8673
- Doležal, M.; Hartl, J.; Miletín, M.; Macháček, M. & Kráľová, K. (1999). Synthesis and photosynthesis-inhibiting activity of some anilides of substituted pyrazine-2-carboxylic acids. *Chem. Pap.*, Vol. 53, No. 2, 126-130, ISSN 0366-6352
- Doležal, M.; Vičík, R.; Miletín, M. & Kráľová, K. (2000). Synthesis and antimycobacterial, antifungal, and photosynthesis-inhibiting evaluation of some anilides of substituted pyrazine-2-carboxylic acids. *Chem. Pap.*, Vol. 54, No. 4, 245-249, ISSN 0366-6352
- Doležal, M.; Kráľová, K.; Šeršeň, F. & Miletín, M. (2001a). The site of action of pyrazine-2-carboxylic acids in the photosynthetic apparatus. *Folia Pharm. Univ. Carol.*, Vol. 26, 13-20, ISSN 1210-9495
- Doležal, M., Miletín, M.; Hejský, R.; Kráľová, K. & Kuneš, J. (2001b). Synthesis of some amides of substituted pyrazine-2-carboxylic acids and their photosynthesis-inhibiting activity. *5th International Electronic Conference on Synthetic Organic Chemistry (ECSOC-5)*, C009, Seijas, J. A. (Ed.), ISBN 3-906980-06-5, CD-ROM edition, September 1-30, 2001, MDPI, Basel
- Doležal, M.; Miletín, M.; Kuneš, J. & Kráľová, K. (2002). Substituted amides of pyrazine-2-carboxylic acids: Synthesis and biological activity. *Molecules*, Vol. 7, No. 7, 363 - 373, ISSN 1420-3049
- Doležal, M.; Palek, L.; Röslerová, Z.; Kuneš, J.; Buchta, V. & Kráľová, K. (2004). Substituted pyrazinecarboxamides: Preparation and biological activity. *8th International Electronic Conference on Synthetic Organic Chemistry (ECSOC-8)*, C001, Seijas, J. A. & Tato, M. P. V. (Ed.), ISBN 3-906980-15-4, CD-ROM edition, November 1-30, 2004, MDPI, Basel
- Doležal, M. (2006a). Biological active pyrazines of natural and synthetic origin. *Chem. Listy* Vol. 100, No. 11, 959-966, ISSN 0009-2770
- Doležal, M.; Palek, L.; Vinšová, J.; Buchta, V.; Jampílek, J. & Kráľová, K. (2006b). Substituted pyrazinecarboxamides: Synthesis and biological evaluation. *Molecules*, Vol. 11, No. 4, 242-256, ISSN 1420-3049
- Doležal, M.; Tůmová, L.; Kešetovičová, D.; Tůma, J. & Kráľová, K. (2007). Substituted N-phenylpyrazine-2-carboxamides, their synthesis and evaluation as herbicides and abiotic elicitors. *Molecules*, Vol. 12, No. 12, 2589-2598, ISSN 1420-3049

- Doležal, M.; Čmedlová, P.; Palek, L.; Vinšová, J.; Kuneš, J.; Buchta, V.; Jampílek, J. & Kráľová, K. (2008a). Synthesis and antimycobacterial evaluation of substituted pyrazinecarboxamides. *Eur. J. Med. Chem.*, Vol. 43, No. 5, 1105-1113, ISSN 0223-5234
- Doležal, M.; Lešňanská, K.; Peško, M.; Jampílek, J. & Kráľová, K. (2008b). Pyrazinecarboxamides, their synthesis and evaluation as potential herbicides. *12th International Electronic Conference on Synthetic Organic Chemistry (ECSOC-12)*, C0004, Seijas, J. A. & Tato, M. P. V. (Ed.), ISBN 3-906980-19-7, CD-ROM edition, November 1-30, 2008, MDPI, Basel
- Engel, S.; Nubling, C.; Kardorff, U.; Kast, J.; von Deyn, W.; Plath, P.; Walter, H.; Westphalen, K. & Gerber, M. (1999). *Herbicidal pyrazine derivatives*. US Patent 5939359, August 17, 1999
- European Commission, (2001). Commission Directive 2001/21/EC of 5 March 2001 amending Annex I to Council Directive 91/414/EEC concerning the placing of plant protection products on the market to include amatol, diquat, predate and thiabendazole as active substances. *Off. J. Eur. Communities L69*, 17-21, May 10, 2001.
- European Commission, (2002). Opinion of the scientific committee on plants on an additional question from the commission concerning the evaluation of diquat in the context of Council Directive 91/414/EEC, *Scientific Committee on Plants SCP/DIQUAT-bis/002-Final 14 January 2002*. European Commission, Belgium, 12pp.
- Fischer, G. C. (1987). *Biologically active N-pyrazinyl-haloacetamides*. US Patent 4742167, May 3, 1988
- Frater, G.; Suchy, M.; Wenger, J. & Winternitz, P. (1987). *[(Isopropylideneamino)oxy]-ethyl-2-[[6-chloroquinoxalinyloxy]phenoxy]propionate postemergent herbicide*. US Patent 4687849, August 18, 1987
- Gorzerino, C.; Quemeneur, A.; Hillenweck, A.; Baradat, M.; Delous, G.; Ollitrault, M.; Azam, D.; Caquet, T. & Lagadic, L. (2009). Effects of diquat and fomesafen applied alone and in combination with a nonylphenol polyethoxylate adjuvant on *Lemna minor* in aquatic indoor microcosms. *Ecotox. Environ. Saf.*, Vol. 72, No. 3, 802-810, ISSN 0147-6513
- Govindjee (1995). Sixty-three years since Kautsky: Chlorophyll *a* fluorescence. *Aust. J. Plant Physiol.*, Vol. 22, No. 4, 131-160, ISSN 0310-7841
- Griffin, D.; Rice, M. & Elliott R. (1990). *Pyrazine compound useful as plant growth regulators*. US Patent 4975112, December 4, 1990
- Hirai, K. (1999). Structural evolution and synthesis of diphenylethers, cyclic imides and related compounds. In: *Peroxidizing Herbicides*. Böger, P.; Wakabayashi, K., Eds.; pp. 15-71, Springer, 1999; ISBN 3540645500, Berlin
- HRAC: *Classification of Herbicides According to Mode of Action*, 2005 <http://www.hracglobal.com/Publications/ClassificationofHerbicideModeofAction/tabid/222/Default.aspx> (23 June 2010)
- Hoff, A. J. (1979). Application of ESR in photosynthesis. *Phys. Rep.*, Vol. 54, No. 2, 75-200, ISSN 0370-1573
- Ivany, J. A. (2005). Response of three potato (*Solanum tuberosum*) cultivars to pyrafluen-ethyl used as desiccant in Canada. *Crop Protect.*, Vol. 24, No. 9, 836-841, ISSN 0261-2194

- Izawa, S. (1980). Acceptors and donors for chloroplast electron transport. *Methods Enzymol.*, Vol. 69, Part C, pp. 413-434. Edited by Anthony San Pietro, Academic Press, ISBN 0121819698, New York – London
- Jegerschöld, C. & Styring, S. (1991). Fast oxygen-independent degradation of the D1 reaction center protein in photosystem-II. *FEBS Lett.*, Vol. 280, No. 1, 87-90, ISSN 0014-5793
- Joshi, M. K. & Mohanty, P. (2004). Chlorophyll *a* fluorescence as a probe of heavy metal ion toxicity in plants. In: *Chlorophyll a fluorescence: A signature of photosynthesis. Advances in Photosynthesis and Respiration.* ed. Papageorgiou, G. C. & Govindjee, Vol. 19, pp. 637-661, Springer, ISBN 1-4020-3217-X, Dordrecht
- Joule, J. A. & Mills, K. (2010). *Heterocyclic chemistry.* John Wiley and Sons, ISBN 9781405133005, pp. 249-288, Chichester
- Linsmaier, E. M. & Skoog, F. (1965). Organic growth factor requirements of tobacco tissue cultures. *Physiol. Plant*, Vol. 18, No. 1, 100-127, ISSN 0031-9317
- Lock, E. A. & Wilks, M. F. (2001). *Diquat.* In: *Handbook of Pesticide Toxicology, Volume 2. Agents*, Krieger R. (ed.), 1605-1622, Academic Press, San Diego, USA, ISBN 0124262600
- Koschnick, T. J.; Haller, W. T. & Glasgow, L. (2006). Documentation of landoltia (*Landoltia punctata*) resistance to diquat. *Weed Sci.*, Vol. 54, No. 4, 615-619, ISSN 0043-1745
- Kráľová, K.; Šeršeň, F. & Sidóová, E. (1992). Photosynthesis inhibition produced by 2-alkylthio-6-R-benzothiazoles. *Chem. Pap.*, Vol. 46, No. 5, 348-350, ISSN 0366-6352
- Kráľová, K.; Šeršeň, F.; Klimešová, V. & Waisser, K. (1997). Effect of 2-alkylthio-4-pyridinecarbothioamides on photosynthetic electron transport in spinach chloroplasts. *Collect. Czech. Chem. Commun.*, Vol. 62, No. 3, 516-520, ISSN 0010-0765
- Kráľová, K.; Šeršeň, F. & Opletalová, V. (1998a). Inhibition of oxygen evolution in spinach chloroplasts and *Chlorella vulgaris* by 2',5'-diazachalcones. *Folia Pharm. Univ. Carol.*, Vol. 23 Suppl., pp. 71-72, ISSN 1210-9495
- Kráľová, K.; Šeršeň, F.; Miletín, M. & Hartl, J. (1998b). Inhibition of photosynthetic electron transport by some anilides of 2-alkylpyridine-4-carboxylic acids in spinach chloroplasts. *Chem. Pap.*, Vol. 52, No. 1, 52-55, ISSN 0366-6352
- Kráľová, K., Šeršeň, F., Kubicová, L., & Waisser, K. (1999). Inhibitory effects of substituted benzanilides on photosynthetic electron transport in spinach chloroplasts. *Chem. Pap.*, Vol. 53, No. 5, 328-331, ISSN 0366-6352
- Ma, J. Y.; Wang, S. F.; Wang, P. W.; Ma, L. J.; Chen, X. L. & Xu, R. F. (2006). Toxicity assessment of 40 herbicides to the green alga *Raphidocelis subcapitata*. *Ecotox. Environ. Saf.*, Vol. 63, No. 3, 456-462, ISBN 0147-6513
- Maga, J. A. (1992). Pyrazine update. *Food Rev. Inter.*, Vol. 8, No. 4, 479-558, ISSN 8755-9129
- Maillard, L. C. (1912). Action des acides aminés sur les sucres: formation des mélanoidines par voie méthodique. *Compt. Rend.*, Vol. 154, 66-68, ISSN 1631-0748
- Mason, R. P. (1990). Redox cycling of radical anion metabolites of toxic chemicals and drugs and the Marcus theory of electron transfer. *Environ. Health Perspect.*, Vol. 87, No. 7, 237-243, ISSN 0091-6765
- Massachusetts Department of Agricultural Resources, Aquatic Vegetation Management. Appendix III – Diquat, 19 pp., <http://www.mass.gov/agr/pesticides/aquatic/docs/diquat.pdf> (June 23, 2010)

- Mees, G. C. (1960). Experiments on the herbicidal action of 1,1'-ethylene-2,2'-dipyridylium dibromide. *Ann. Appl. Biol.*, Vol. 48, No. 3, pp. 601-612, ISSN 0003-4746
- The Merck Index*, 14th Edition, 2006 by Merck & Co., Inc., Whitehouse Station, ISBN 0-911910-00-X, NJ, USA
- Mixan, C. E. & Goralski, C. T. (1978). *Cyanatothiomethylthio pyrazines*. US Patent 4101546, July 18, 1978
- Nakamura, A.; Ataka, T.; Segawa, H.; Takeuchi, Y. & Takematsu, T. (1983). Studies on herbicidal 2,3-dicyanopyrazines. 1. Structure-activity relationship of herbicidal 2,3-dicyano-5-substituted pyrazines. *Agric. Biol. Chem.*, Vol. 47, No. 7, pp. 1555-1560, ISSN 0002-1369
- Nakamura, A.; Ataka, T.; Segawa, H.; Takeuchi, Y. & Takematsu, T. (1983a). Studies on herbicidal 2,3-dicyanopyrazines. 2. Structure-activity relationships of herbicidal 5-ethylamino- and 5-propylamino-2,3-dicyanopyrazines. *Agric. Biol. Chem.*, Vol. 47, No. 7, pp. 1561-1567, ISSN 0002-1369
- Niedermann, H. P. & Munro, D. (1994). *Herbicidal pyrazines and pyrimidines*. GB 2285045, December 12, 1994
- Oettmeier, W. (1992). Herbicides of Photosystem II. In: *Topics in photosynthesis*, Vol. 11: The Photosystems: Structure, Function and Molecular Biology (Barber, J. ed.) pp. 349-408, Elsevier Science Publishers B. V., ISBN 0444894403, Amsterdam, Netherlands.
- Opletalová, V.; Hartl, J.; Patel, A.; Palát, Jr. K. & Buchta, V. (2002). Ring substituted 3-phenyl-1-(2-pyrazinyl)-2-propen-1-ones as potential photosynthesis-inhibiting, antifungal and antimycobacterial agents. *Farmaco*, Vol. 57, No. 2, pp. 135-144, ISSN 0014-827X
- Opletalová, V.; Pour, M.; Kuneš, J.; Buchta, V.; Silva, L.; Kráľová, K.; Chlupáčová, M.; Meltrová, D.; Peterka, M. & Posledníková, M. (2006). Synthesis and biological evaluation of (E)-3-(nitrophenyl)-1-(pyrazin-2-yl)prop-2-en-1-ones. *Collect. Czech. Chem. Commun.*, Vol. 71, No. 1, pp. 44-58, ISSN 0010-0765
- Peiró, A. M.; Zapater, P.; Alenda, C.; Ramírez, A.; Gutiérrez, A.; Pérez-Mateo, M. & Such, J. (2007). Hepatotoxicity related to paraquat and diquat absorption through intact skin. *Dig. Dis. Sci.*, Vol. 52, No. 11, pp. 3282-3284, ISSN 0163-2116
- Peterson, H. G.; Boutin, C.; Freemark, K. E. & Martin, P. A. (1997). Toxicity of hexazinone and diquat to green algae, diatoms, cyanobacteria and duckweed. *Aquat. Toxicol.*, Vol. 39, No. 2, pp. 111-134, ISSN 0166-445X
- Petit, V.; Cabridenc, R.; Swannell, R. P. J. & Sokhi, R. S. (1995). Review of strategies for modelling the environmental fate of pesticides discharged into riverine systems. *Environ. Int.*, Vol. 21, No. 2, pp. 167-176, ISSN 0160-4120
- Portnoy, R. C. (du Pont de Nemours, E. I., and Co.) (1978). *Arylthiodicyanopyrazines as plant disease control agents*. US Patent 4113724, September 12, 1978
- Ritter, A. M.; Shaw, J. L.; Williams, W. M. & Travis, K. Z. (2000). Characterizing aquatic ecological risks from pesticides using a diquat bromide case study. I. Probabilistic exposure estimates. *Environ. Toxicol. Chem.*, Vol. 19, No. 3, 749-759, ISSN 0730-7268
- Sakata, G.; Makino, K.; Kawamura, Y.; Ura, Y.; Ikai, T. & Kawamura, Y. (1983). NCI-96683, a new selective herbicide for annual and perennial grass weed control in broad leaf crops. *10th Int. Congr. Plant Prot., Proc. Conf.*, Vol. 1, pp. 315-23. ISBN 978-

- 0901436795, Brighton, England, 20-25 November, 1983, British Crop Protection Council
- Sanchez, W.; Palluel, O.; Lagadic, L.; Ait-Aissa, S. & Porcher, J. M. (2006). Biochemical effects of nonylphenol polyethoxylate adjuvant, Diquat herbicide and their mixture on the three-spined stickleback (*Gasterosteus aculeatus* L.). *Mar. Environ. Res.*, Vol. 62, Suppl. S, pp. S29-S33, ISSN 0141-1136
- Sandmann, G. (1992). Spectral determination of carotenoid precursors in *Scenedesmus* cells treated with bleaching herbicides. In: Böger, P. & Sandmann, G. (eds.), *Target Assays for Modern Herbicides and Related Compounds*, pp. 3-8. Lewis Publishers, ISBN 0-87371-539-X, Boca Raton, Florida
- Sato, J.; Kondo, Y.; Kudo, Y.; Nawamaki, T.; Watanabe, S.; Ishikawa, K. & Ito, Y. (1993). *Preparation of pyrazine derivatives as herbicides*. JP 06192252, September 21, 1993
- Shuto, A.; Hoshi, H. & Sanemitsu, Y. (2000). *Pyrazin-2-one derivatives, their use, and intermediates for their production*. US Patent 6015774, January 18, 2000
- Svensson, B.; Vass, I. & Styring, S. (1991). Sequence analysis of the D₁ and D₂ reaction center proteins of photosystem II. *Z. Naturforsch.*, Vol. 46c, No. 9-10, 765-776, ISBN 0939-5075
- Takematsu, T.; Segawa, H., Takamoro, M.; Ataka, T.; Chatani, M. & Nakamura, A. (1981). *2,3-Dicyanopyrazines*. US Patent 4259489, March 31, 1981
- Takematsu, T.; Segawa, H.; Miura, T.; Ataka, T.; Chatani, M. & Nakamura, A. (1984). *2,3-Dicyano 6-phenyl pyrazine herbicides*. US Patent 4460403, July 17, 1984
- Tong, Y. C. (1978). *Microbicidal/fungicidal thiazolopyrazines*. US Patent 4075207, February 21, 1978
- US Environmental Protection Agency, 1995. *Reregistration Eligibility Decision (RED) Diquat Dibromide*, EPA738-R-95-016. Office of Prevention Pesticides and Toxic Substances, Washington, DC, 303 pp., October 17, 2007, <http://www.epa.gov/oppsrrd1/REDs/0288.pdf> (June 23, 2010)
- Wagner, R.; Czerny, M.; Bielohradsky, J. & Grosch, W. (1999). Structure-odour-activity relationships of alkylpyrazines. *Z. Lebensm. Unters. Forsch. A*, Vol. 208, No. 5-6, 308-316, ISSN 1431-4630
- Weaver, E. C. (1968). EPR studies of free radicals in photosynthetic systems. *Annu. Rev. Plant. Physiol.*, Vol. 19 (June), 283-294, ISSN 0066-4294
- Weber, J. B. & Peter, C. J. (1982). Adsorption, bioactivity and evaluation of soil tests for alachlor, acetochlor and metolachlor. *Weed Sci.*, Vol. 30, No. 1, 14-20, ISSN 0043-1745
- Whitmarsh, J. (1998). Electron Transport and Energy Transduction. In: *Photosynthesis: A Comprehensive Treatise*, Raghavendra A. S., (ed.), pp. 87-110, Cambridge University Press, ISBN 0-521-57000-X, Cambridge
- Whitmarsh, J. & Govindjee (1999). The Photosynthetic Process. In: *Concepts in Photobiology: Photosynthesis and Photomorphogenesis*, G. S. Singhal, G. Renger, S. K. Sopory, K-D. Irrgang and Govindjee (eds.), pp. 11-51, Narosa Publishing House, ISBN 81-7319-151-4, New Delhi, India
- Woolfson, A. & Rothschild, M. (1990). Speculating about pyrazines. *Proc. R. Soc. Lond. B*, Vol. 242, 113-119, ISSN 0962-8452

- Yamaguchi, T. (2007). The relationship between the chemical structures of dihydropyrazine derivatives and DNA strand-breakage activity. *Chem. Pharm. Bull.*, Vol. 55, No. 4, 532-536, ISSN 0009-2363
- Zondler, H.; Eckhardt, W. & Nyfeler R. (1989). *2-Arylmethyliminopyrazines, pyrazines compositions and method of combating undesirable plant growth and of regulating plant growth*. US Patent 4849009, July 18, 1989
- Zweig, G.; Shavit, N. & Avron, M. (1965). Diquat (1,1'-ethylene-2,2'-dipyridylium dibromide) in photoreactions of isolated chloroplasts. *Biochim. Biophys. Acta*, Vol. 109, No. 2, 332-346, ISSN 0006-3002

Herbicides, Theory and Applications

Edited by Prof. Marcelo Larramendy

ISBN 978-953-307-975-2

Hard cover, 610 pages

Publisher InTech

Published online 08, January, 2011

Published in print edition January, 2011

The content selected in Herbicides, Theory and Applications is intended to provide researchers, producers and consumers of herbicides an overview of the latest scientific achievements. Although we are dealing with many diverse and different topics, we have tried to compile this "raw material" into three major sections in search of clarity and order - Weed Control and Crop Management, Analytical Techniques of Herbicide Detection and Herbicide Toxicity and Further Applications. The editors hope that this book will continue to meet the expectations and needs of all interested in the methodology of use of herbicides, weed control as well as problems related to its use, abuse and misuse.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Martin Doležal and Katarína Kráľova (2011). Synthesis and Evaluation of Pyrazine Derivatives with Herbicidal Activity, Herbicides, Theory and Applications, Prof. Marcelo Larramendy (Ed.), ISBN: 978-953-307-975-2, InTech, Available from: <http://www.intechopen.com/books/herbicides-theory-and-applications/synthesis-and-evaluation-of-pyrazine-derivatives-with-herbicidal-activity>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2011 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike-3.0 License](https://creativecommons.org/licenses/by-nc-sa/3.0/), which permits use, distribution and reproduction for non-commercial purposes, provided the original is properly cited and derivative works building on this content are distributed under the same license.

IntechOpen

IntechOpen