We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Damage and Replication Stress Responses

Haiying Wang¹, Ping Shen² and Wei-Guo Zhu¹

¹*Key laboratory of Carcinogenesis and Translational Research (Ministry of Education),
Department of Biochemistry and Molecular Biology,
Peking University Health Science Center,*

²*Department of Applied linguistics, Peking University Health Science Center,
China*

1. Introduction

The character of DNA replication is high fidelity. Precise and complete DNA replication is critical for the maintenance of genetic stability. Failures in these processes are major sources of genomic instability and will lead to cancer or other diseases. A wide variety of factors, such as DNA replication errors, spontaneous chemical reactions, reactive metabolic products, exogenous environmental agents or some anticancer therapeutics e.g. 5'-aza-2'-deoxycytidine (5-Aza-CdR), can cause DNA damage (Wang et al, 2008; Zhu et al, 2004; Chai et al, 2008; Hoeijmakers, 2009). It is estimated that DNA damage occurs at a rate of 1,000 to 1,000,000 molecular lesions per cell per day (Lodish, 2004). In order to deal with problems under which the genome is vulnerable to injury or replication stress, eukaryotic cells elaborate a genomic maintenance apparatus, which is termed the DNA damage response (DDR) and replication stress response, including various checkpoint, signal-transduction and effectors system, which monitor problems and trigger a comprehensive cellular response pathway to prevent genome integrity. The extent of DNA damage depends on the type of environment to which it is exposed (Hoeijmakers, 2001). So, organisms must be capable of recognizing and dealing with each type of damage. It is not surprising that there are various different types of DNA damage response and repair systems.

ATM and ATR are at the top of the DNA damage pathways. Although a cross-talk exists between the ATM and ATR pathways, ATM primarily seems to be involved in the detection of DNA double-strand breaks via Mre11/Rad50/Nbs1 complex (MRN), ATR is critical for cellular responses to a variety of DNA damage and stalled replication forks (Hefferin & Tomkinson, 2005). When these protein kinases activated, they eventually phosphorylate and modulate the downstream effectors (e.g., Chk1 and Chk2) and multiple additional substrates that initiate the cellular responses.

Dynamic changes in protein post-translational modifications play a significant role in most cellular signalling pathways. More and more proteins were found in a variety of post-translational modifications in response to DNA damage and genotoxic stress, such as phosphorylation, acetylation, sumoylation, methylation and ubiquitylation. Recent studies indicate that a crosstalk between multiple protein modifications exists, which collaboratively regulates signal transduction of DNA damage and genetic stresses.

Actually, the DNA damage and replication stress response consists of multiple interconnected pathways, which impact the cell cycle, DNA replication, DNA repair, transcriptional regulation, chromatin remodelling, metabolic and other cellular biological processes (Rouse & Jackson, 2002; Zhou & Elledge, 2000). In this chapter, we focus on recent findings of DNA damage response signalling pathways.

2. The DNA damage response pathways

The ability of cells to respond to DNA damage and replication stress response is critical for cellular survival. The evidence indicates that DNA damage and replication stress response are a cascade signal transductional process, which consists of multiple interconnected pathways through which sense damage or replication stress, transduce the damage signals, and trigger cellular responses, including cell cycle arrest, DNA repair or apoptosis (Shiloh, 2003; Bakkenist & Kastan, 2004; McGowan & Russell, 2004). In mammalian cells, PI3K family members, ATM and ATM-Rad3-related (ATR) are central to the entire DNA damage response (Elledge, 1996). All types of DNA lesions induce responses to these two main signalling pathways. Next, we will summarize the model of cells dealing with DNA damage and replication stress through these two pathways.

2.1 ATM dependent cellular response to DNA double strand breaks pathway

The DNA double strand breaks (DSBs) are the most dangerous damage type for the organisms because they are prone to cause genomic rearrangements, cancer predisposition, and cell death if not repaired correctly (Wyman & Kanaar, 2006). Many endogenous and exogenous factors may induce DSBs, such as IR, UV, reactive oxygen species (ROS) or topoisomerases inhibitors (Tanaka, 2006; Tanaka, 2007). Cellular responses to DSBs, include complex signal-transduction, cell-cycle-checkpoint and repair pathways, play a pivotal role in maintenance of the genome integrity. It is accepted that ATM is a central component of the DSB signalling cascade (Khanna & Jackson, 2001; Shiloh, 2001; Abraham, 2001).

2.1.1 ATM

ATM is the gene product mutated in ataxia telangiectasia discovered in 1995 and characterized by progressive cerebellar ataxia, immune deficiencies, radiation sensitivity, and an increased risk of cancer (Lavin & Shiloh, 1997; Savitsky et al, 1995). ATM is a serine-threonine kinase which belongs to the phosphatidylinositol-3-kinase (PI3K) like protein kinases (PIKK) family. In normal condition, ATM exists in an inactive form of dimer or multimer. Following DSBs, ATM was dissociated into an active monomer through autophosphorylation (Bakkenist & Kastan, 2003). Upon activation, ATM is recruited to DNA breaks where it initiates phosphorylation of several substrates such as p53, Mdm2, BRCA1, Chk2 and Nbs1 to initiate cell cycle arrest, DNA repair, and apoptosis (Lukas et al., 2003; Shiloh Y, 2006).

2.1.2 Mechanism for the activation of ATM

Many progresses have been made on understanding how DSBs activates ATM. Several investigations suggest that the Mre11-Rad50-Nbs1 (MRN) complex is involved in ATM activation and recruitment to the sites of DSBs (Uziel et al, 2003; Cerosaletti & Concannon, 2004), because attenuated activation and no recruitment of ATM to DSBs upon damage were found in Mre11- and Nbs1- deficient cell lines. Earlier studies have shown that MRN lies

downstream of the ATM mediated DNA damage signalling pathway because ATM can phosphorylate the components of the MRN complex in response to IR (Lim & Ki, 2000; Wu & Ranganathan, 2000; Zhao & Weng, 2000). However, further analyses demonstrate that the MRN complex is more like an upper actor of ATM pathway (Uziel et al, 2003 ; Difilippantonio et al, 2005; Carson et al, 2003). Because the MRN complex was reported to play a role in early detection of DSBs which initiates the localization of ATM to DSBs (Lee & Paull, 2004, 2005). It is now established that ATM at DSBs is a spatio-temporal dynamics mechanism. At first, change in chromatin structure caused by DNA DSB partially activates ATM (Berkovich et al, 2007). Activated ATM rapidly phosphorylates H2AX on its C-terminus, and γ -H2AX subsequently recruits MDC1 (mediator of DNA damage checkpoint protein 1) to bind to it and acts as a scaffold, in turn, recruits MRN at the flanking chromatin of DSBs (Burma et al, 2001; Stucki & Jackson, 2006; Lou et al, 2006; Stucki et al, 2005), which promotes accumulation of ATM to sites of DSBs, where it is fully activated (Lavin, 2008). In addition, MDC1 also mediates the interaction between ATM and γ -H2AX, which contributes to the extended phosphorylation of H2AX and the maintenance of the DSB response (Huen & Chen J, 2008). Autophosphorylation has been proposed as the other mechanism for ATM activation. Three phosphatases, PP2A, PP5 and WIP1, have been reported to be involved in the control of ATM activation. Autophosphorylation on Ser367, Ser1893, Ser1981 and a new site S2996 are present on activated ATM through dissociation of the inactive dimeric ATM to an active monomeric form (Bakkenist & Kastan, 2003; Kozlov et al, 2010). Recent reports indicate that there is a Nbs1-independent ATM activation pathway which regulates ATM activity through its effect on ATM autophosphorylation (Kanu & Behrens, 2007; Sun et al., 2005, Gupta et al., 2005; Richard et al, 2008). Interestingly, notwithstanding the difference on the importance of ATM autophosphorylation in humans and in mice, this is certainly the case in human cells that autophosphorylation of ATM at serine 1981 is required for the interaction of ATM with MDC1, which stabilizes ATM at DSBs and thereby promotes a full-scale response to DNA damage (Sairei et al, 2009). Once activated, ATM directly or indirectly phosphorylates approximately 30 substrates, such as Chk2, p53, BRCA1, RPAp34, H2AX, SMC1, HDMX, FANCD2, Rad17, Artemis or Nbs1, which are involved in cell cycle checkpoint control, apoptotic responses and DNA repair.

2.2 “two-man rule” of ATR in response to DNA damage

Like ATM, ATR (ATM- and Rad3-related) is a nuclear Ser/Thr kinase which belongs to the PIKK family (Bentley et al, 1996). ATR forms a stable heterodimer with its interacting partner ATRIP which can be activated by DNA damage (Cliby et al, 1998; Wright et al, 1998). Compared with the ATM, ATR can respond to a broader spectrum of genotoxic stimuli including DNA replication inhibitors (such as hydroxyurea), UV radiation, ionizing radiation, and agents that induce DNA interstrand cross-links and generate single-stranded DNA (Wright et al, 1998; Yang et al, 2003; Costanzo & Gautier, 2003; Wang et al, 2008). Once the break occurs, ATR is recruited by ATR-IP to the sites of DNA damage and interacts with RPA to initiate the response (Zou & Elledge, 2003b; Cortez et al, 2001; Wang et al, 2008).

2.2.1 Mechanisms of ATR activation

ATR is involved in many different types of DNA damage responses. The common feature is that ATR is activated by single strand DNA (ssDNA), which is a common intermediate structure that can be formed at sites of DNA damage and replication stress, or induced by

most cancer chemotherapies (Costanzo & Gautier, 2003; Zou & Elledge, 2003b). A study shows that both ssDNA and a 5' junction are sufficient to activate ATR signalling (MacDougall et al, 2007). ATR activation requires assembly of a protein complex on ssDNA, which begins with ATR-ATRIP complex loading on the RPA-coated ssDNA (Stokes et al, 2002; MacDougall et al, 2007; Byun et al, 2005). Earlier works have shown that RPA binds to ssDNA and then recruits ATR-ATRIP by interacting with ATRIP (Cortez et al, 2001; Ball et al, 2007). However, the recruitment to ssDNA is not sufficient for ATR activation (Ball et al., 2005; Namiki & Zou, 2006; Yoshioka et al., 2006), it requires additional ATR regulator, Rad9-Rad1-Hus1 (9-1-1) complex, a heterotrimeric ring-shaped structure like PCNA (Parrilla-Castellar et al., 2004). The 9-1-1 complex recognizes a DNA end that is adjacent to a stretch of RPA-coated ssDNA through working with RFC-RAD17 (Ellison & Stillman, 2003; Zou et al., 2003a; Bermudez et al., 2003). Current models for ATR activation suggest that the 9-1-1 mediated recruitment of TopBP1 to the ATR-ATRIP complex, and the ATR-activating domain of TopBP1 activates the kinase activity of ATR (Harper & Elledge, 2007; Cimprich & Cortez, 2008; Shiotani & Zou, 2009; Yan & Michael, 2009). In addition to be an activator of ATR, TopBP1 is also a substrate of ATR. The phosphorylation of TopBP1 on residue S1131 by ATM enhanced the interaction of it with ATR-ATRIP, which suggests that TopBP1 promotes a feed-forward signalling loop to amplify ATR-mediated signals (Yoo et al., 2007). Thus, sustained co-localization of the ATR-ATRIP and 9-1-1-TopBP1 complexes at the DNA damage site may increase their local concentration so that ATR activation is stimulated continually by TopBP1. However, recruitment of the ATR-ATRIP and 9-1-1-TopBP1 complexes to sites of DNA damage or stalled replication forks is independent events (Bonilla et al, 2008; Kondo et al, 2001; Melo et al, 2001; Zou et al, 2002; You et al, 2002; Lee et al, 2003). Therefore, there is the two-man rule in TopBP1-dependent regulation of ATR activity, by which ATRIP and 9-1-1 together control the TopBP1 to initiate ATR signalling.

As described above, activation of ATM involves its autophosphorylation, which helps it convert an inactive dimer form into an active monomers form. Some phosphorylation sites on ATR and ATRIP have been found (Cimprich & Cortez, 2008), unlike ATM, as yet, none of these identified modifications has been reported to contribute to ATR activation and the oligomerization status of ATR-ATRIP.

2.3 Interplay between ATM and ATR pathway

It was previously thought that ATM and ATR had overlapping but distinct roles in response to DNA damage. However, a current study demonstrate a high degree of cross-talk and connectivity. For instance, ATM and ATR collaborate in the IR-induced G2/M checkpoint, but incomplete DNA replication in mammalian cells can prevent M phase entry independent of ATR (Brown & Baltimore, 2003). Recently, Trenz et al indicate that both ATM and ATR promote Mre11-dependent restart of collapsed replication forks and prevent accumulation of DNA DSBs (Trenz et al, 2006). Another study showed that ATR is activated rapidly by IR, and both ATM and Mre11 enhance ATR signalling (Myers & Cortez, 2006). The new data demonstrate that ATR is required for the response to either replication stress or IR without any role for ATM (Paul et al. 2004).

3. DNA damage response pathways and cell cycle checkpoints

The maintenance of genome stability is critical to the survival and propagation of all cellular organisms. The cell cycle is required for cell growth and cell division into two daughter

cells. Cell cycle checkpoints are regulatory pathways that control the cell cycle events in the right order. DNA is vulnerable to diverse types of injury throughout the cell cycle. In response to DNA damage, checkpoint surveillance mechanisms initiate a cascade of events which coordinate cell cycle arrest and facilitate DNA repair pathways. These checkpoints include the G1/S, intra-S and G2/M of the cell cycle and are controlled by the ATM/Chk2 and ATR/Chk1 pathways. We will discuss the progresses of different signalling pathways involved in different checkpoints.

3.1 G1 Checkpoint

G1 checkpoint is the first checkpoint making the key decision of whether the cell should divide or arrest, which prevents the damaged DNA from being replication (Bartek & Lukas, 2001). The major player in the G1 checkpoint is the p53 protein. In normal cells, p53 is maintained at low levels due to interaction with MDM2, which targets p53 for degradation in the cytoplasm (Alarcon-Vargas & Ronai, 2002). In response to distinct or partially overlapping types of DNA damage, p53 is activated by ATM or ATR through phosphorylating different Ser/Thr residues directly and indirectly (Matsuoka et al., 2000; Maya et al., 2001; Shieh et al., 2000). The phosphorylation of Ser15 appears important in enhancing p53 transcriptional transactivation activity (Dumaz & Meek, 1999; Wang et al, 2008). The result of p53 activation is the up-regulation of various target genes (such as MDM2, GADD45a, and p21^{Waf1/Cip1}), some of which are involved in the DNA damage response. p21^{Waf1/Cip1} elicits G1 arrest through suppressing Cyclin E/Cdk2 kinase activity (Bartek & Lukas, 2001). In other p53 target genes, such as Gadd45 and BIG2, also lead to G1 arrest. p53 lead to G1 checkpoint arrest in multiple pathways, now, p53 is reported to contribute to maintain G1 checkpoint control via activating microRNAs directly.

3.2 S-phase Checkpoint

The S-phase checkpoint monitors cell cycle process and lowers the rate of DNA replication after DNA damage. ATM plays a primary role in contributing to S-phase checkpoints although it overlapping with the ATR dependent pathway in maintenance of the S-phase checkpoint. In response to ionizing radiation, ATM phosphorylates Nbs1 and Chk2 and triggers two parallel cascades of the DNA damage responses to activate the S-phase checkpoint. One is the ATM-Chk2-Cdc25A pathway; the other is the ATM dependent NBS1/BRCA1/SMC1 pathway, though the mechanism of this pathway is not well understood (Falck et al., 2002). On the other hand, the ATR-Chk1 pathway is also involved in the S-phase checkpoint arrest auxiliary in response to IR. Furthermore, the ATR-Chk1 pathway plays a dominant role in directing S-phase checkpoint arrest in response to UV damage and replication errors (Abraham, 2001).

3.3 G2 Checkpoint

The G2 cell cycle checkpoint is an important control point which functions to prevent damaged DNA from being segregated into daughter cells. This checkpoint activation depends on the maintenance of Cdc2 phosphorylation on T14 and Y15 (Rhind et al., 1997). ATM and ATR both indirectly modulate the phosphorylation status of these sites in response to DNA damage. Different from other checkpoints, ATR mainly controls the response to UV damage and replication blocks. The response to IR is also mediated primarily by ATR while ATM plays a supporting role (Graves et al, 2000). Upon DNA

damage, ATR and ATM phosphorylate their downstream kinases Chk1 and Chk2, respectively, and then phosphorylate the phosphatase Cdc25C on Ser216 (Peng et al., 1997). The phosphorylated Cdc25C binds with 14-3-3 protein and is sequestered in the cytoplasm, which prevents Cdc25C from dephosphorylating Cdc2 in the nucleus and the cells remain arrested in the G2 phase (Lopez-Girona et al., 1999; Peng et al., 1997).

p53 also plays a role in the G2/M checkpoint (Passalaris et al., 1999). Activated p53 in response to DNA damage results in G2/M checkpoint arrest through induction of GADD45 (Zhan et al., 1994). In addition, p53-dependent transcriptional repression of cdc2 and cyclin B may also contribute to the G2/M checkpoint (Passalaris et al., 1999).

4. DNA damage response and protein post-modifications

Post-translational modifications play a vital role in harmonizing cellular response to DNA damage. More and more proteins were found occurring in a variety of post-translational modifications including phosphorylation, acetylation, methylation and ubiquitylation in response to DNA damage or genotoxic stress. Recent research suggests that a crosstalk exists between multiple protein modifications. Here, we will summarize recent findings of protein post-translational modifications in coordinating the DNA damage response signalling cascade.

4.1 Protein phosphorylation modification in response to DNA damage

Signal transduction is predominantly mediated by a cascade of protein phosphorylation and dephosphorylation reactions, which is of prime importance for the organisms to sense the external and internal stimuli and generate the appropriate responses. Protein phosphorylation plays the same role in cellular DNA damage response. As indicated above, in responding to DSB signalling, ATM undergoes autophosphorylation, which seems to be instrumental in the monomerization and activation of ATM. It seems that DNA lesions activate various protein kinases, such as ATM and ATR, which transduce the damage signalling by directly phosphorylating or mediating the phosphorylation and activation of numerous substrates involved in the DNA repair machinery, the cell cycle checkpoints and apoptosis (Abraham, 2001; Osborn et al., 2002). So far, more than 700 proteins have been identified to be phosphorylated in response to DNA damage (Matsuoka et al., 2007). A signalling cascade is initiated starting with the phosphorylation of H2AX (γ -H2AX). γ -H2AX is a chromatin-based signal that regulates the assembly of DNA damage response proteins at the break sites and induction of DNA repairs (Lavin, 2008; Cook et al., 2009). So, the H2AX phosphorylation level is not only important as a marker of the DNA damage response, but also involve in DNA repair processes (van Attikum & Gasser, 2009).

The effector p53 stands at the cross-roads of cellular responses to various stresses (Appella & Anderson, 2001; Bode & Dong, 2004). DNA damage leads to specific phosphorylation modifications of p53 protein. Up to date, at least 20 phosphorylation sites have been detected in the p53 molecule in human cells following DNA damage (Bode & Dong, 2004). Some of which were phosphorylated by ATM in response to irradiation and chemotherapeutic drugs (Banin et al., 1998; Canman et al., 1998), whereas some are phosphorylated by ATR when cells are treated with UV or anti-cancer drugs (Appella & Anderson, 2001; Wang, 2008). Phosphorylation of p53 usually modulates its stability and sequence-specific DNA binding activity (Bode & Dong, 2004). Two major phospho-binding modules, the BRCA1 C-terminal repeat (BRCT) and the forkhead-associated (FHA) domain,

which are present in many proteins are involved in the cellular response to DNA damage, and facilitated protein-protein interactions in the recruitment and activation of damage signalling (Yu et al, 2003; Hofmann & Bucher, 1995; Li et al, 2002).

4.2 Protein acetylation in response to DNA damage

Phosphorylation is not the only post-translational modification in cellular response to DNA damage. For instance, following DSB, ATM is activated in the vicinity of the break and is recruited to the break site by the MRN complex where it is fully activated, facilitated by not only autophosphorylation but also acetylation of ATM (Bakkenist & Kastan, 2003; Sun et al, 2007). The study showed that after DNA damage, CK2 phosphorylates and releases HP1 β from chromatin which recruits a ATM-Tip60 complex to MRN at the break site. This promotes interaction between Tip60 acetyltransferase and the unbound histone H3 K9me3 leading to acetylation and activation of ATM (Sun et al, 2009). A single site at Lys3016 is acetylated by Tip60 acetyltransferase. This mutation inhibits the monomerization and up-regulation of ATM activation by DNA damage, further prevents ATM-dependent phosphorylation of p53 and checkpoint kinase-2 (Chk2) (Sun et al, 2005, 2007).

As the recruitment of the Tip60 acetyltransferase, the deacetylase enzymes HDAC1, HDAC2, HDAC4, SIRT1, and SIRT6 also have been observed at DSB sites in mammalian cells (Kao et al. 2003; Oberdoerffer et al. 2008; Kaidi et al. 2010; Miller et al. 2010). For instance, the MRN complex serves as a sensor for the detection of DSBs and is involved in the S phase checkpoint (Paull & Lee, 2005; van den Bosch et al, 2003). The acetylation level of NBS1 was recently reported to be tightly regulated by deacetylase SIRT1 (Yuan et al., 2007). Moreover, SIRT6-dependent deacetylation of the CtIP in response to DSBs stimulates the RPA and RAD51 foci, thus promoting ATR signalling and DSB repair (Kaidi et al. 2010). Furthermore, histone acetylation can regulate the dynamics of DDR factors in the vicinity of DNA breaks.

p53 acetylation also plays important roles in response to various types of DNA damage (Gu & Roeder, 1997; Lill et al, 1997; Nag et al, 2007). Transcription factors with histone acetyltransferase activity, p300/CBP, p300/CBP-associated factor (PCAF), and Tip60 are reported to be mainly responsible for the p53 acetylation (Liu et al., 1999; Sykes et al., 2006; Tang et al., 2006, 2008). p53 acetylation can increase its sequence-specific DNA binding capacity (Gu & Roeder, 1997; Zhao et al, 2006; Luo et al, 2004) or enhance its stabilization by inhibiting ubiquitination of p53 mediated by MDM2 (Li et al, 2002; Ito et al, 2002). Recently our studies indicate that histone deacetylase inhibitors and other chemical agents also induce p53 acetylation through the DNA damage response pathway (Zhao et al, 2006; Wang et al, 2008). Novel discoveries further confirm that p53 acetylation is an indispensable event for mediating the p53 response (Kruse & Gu, 2009). However, the regulatory mechanisms involving in this posttranslational modification are still largely unknown.

4.3 Protein ubiquitylation in response to DNA damage

Ubiquitylation is the process by which the 76-amino-acid polypeptide ubiquitin is attached to the target protein singly (monoubiquitylation) or in the form of polyubiquitin chains (polyubiquitylation) via the covalent bond. This is an enzyme cascade reaction which is involved by ubiquitin E1, E2 and E3 ligase proteins (Pickart, 2001). A growing number of evidences have shown that ubiquitylation and deubiquitylation are important regulatory mechanisms in response to DNA damage and genotoxic stresses. Assembly of DNA damage response proteins at the break site is catalyzed by the E3 ubiquitin ligases.

Consistent with these actions, several ubiquitin ligases have been shown to accumulate at sites of DNA breaks in mammalian cells, including BRCA1, RNF8, RNF168, RAD18, HERC2, and PRC1 (Polycomb-repressive complex 1) (Scully et al. 1997; Kolas et al. 2007; Doil et al. 2009; Huang et al. 2009; Stewart et al. 2009; Bekker-Jensen & Mailand, 2010; Chou et al., 2010; Lavin, 2008). For example, phosphorylation of MDC1 at ATM consensus sites promotes interaction with RNF8, the E3 ubiquitin ligase, which in turn ubiquitylates H2A, leading to the accumulation of 53BP1, BRCA1, and other proteins at the site of damage (Kolas et al, 2007; Mailand et al, 2007). Ubiquitylated H2A serves as an interacting partner for RNF168 that further propagates the ubiquitylation of H2A and other unknown targets at the double-strand break site (Doil et al, 2009; Stewart, 2009). In addition, FANCD2 (Fanconi anemia complementation group D2) is monoubiquitinated during the S phase (Taniguchi et al. 2002) and in response to various DNA damaging agents, which is required for its localization to DNA damage foci (Garcia-Higuera et al. 2001). It is demonstrated that ATR is required for efficient FANCD2 monoubiquitination and foci assembly in response to various genotoxic stresses, including IR and MMC. Another example is p53, which is kept at low level in unstressed cells through Mdm2-mediated polyubiquitination, which leads to nuclear export of p53 and subsequent proteasomal degradation. DNA damage attenuates polyubiquitination of p53, thereby stabilizes and activates p53 as a transcription factor, up-regulating expression of numerous proteins involved in cell cycle control, apoptosis and senescence (Toledo & Wahl, 2006; Bode & Dong, 2004).

Besides ubiquitylation, deubiquitylation has also been documented as an opposite way to regulate protein stability in response to genotoxic stress. A number of de-ubiquitylating enzymes (DUBs) were identified at double-strand breaks including USP3, USP28 and BRCC36. BRCC36 antagonizes RNF8-dependent ubiquitylation to maintain steady state levels required for appropriate signalling (Sobhian et al, 2007; Chen et al, 2006; Shao et al, 2009). USP3 is a chromatin-associated DUB that also antagonizes RNF8-mediated ubiquitylation (Nicassio et al, 2007). USP28 is a major regulator of DNA damage-induced apoptosis. It was shown that USP28 stabilizes CHK2, 53BP1 and a number of other DNA damage responsive proteins upon irradiation (Wu-Baer et al, 2003).

4.4 Crosstalk between post-translational modifications in response to DNA damage

Recent researches suggest that a crosstalk exists among multiple protein modifications, which collaboratively to regulate signal transduction of DNA damage and genetic stresses. p53, is subjected to multiple posttranslational modifications in response to genotoxic stress, which results in the accumulation of p53 and triggers its transcriptional activities. The damage-induced phosphorylation of p53 seems to be a signal for subsequent acetylation, because phosphorylation enhances its association with the CBP/p300 and PCAF to induce p53 acetylation in response to DNA damage, which results in p53 acetylation and further stabilized (Wang, 2008). Recent reports revealed that the Set8/Pr-Set7 methyltransferase suppresses p53 function in response to DNA damage (Shi et al. 2007), and lysine methylation of p53 by Set7/9 methyltransferase is important for its subsequent acetylation, which results in stabilization of the p53 protein (Ivanov et al., 2007). We also demonstrate that Set7/9 interacts with Sirt1 and induces a decrease in binding of Sirt1 to p53, and this relatively enhances p53 transactivity (Liu et al, 2011).

Apart from the above mentioned, H2AX, a variant form of H2A, is known to be acetylated by Tip60 acetyltransferase following DNA damage. Acetylated H2AX is required for its subsequent ubiquitylation via the ubiquitin-conjugating enzyme UBC13 (Ikura et al., 2007).

They suggested that acetylation-dependent ubiquitination by the Tip60-UBC13 complex leads to the release of H2AX from damaged chromatin, which enhanced histone dynamics and in turn stimulates a DNA damage response.

5. Conclusion

In summary, instability of genome is a constant problem of organisms. The coordination of DNA damage response (DDR) processes is required to maintain cellular viability and prevent diseases. The ATM and ATR protein kinases are master regulators of the DNA damage response. To further understand the molecular mechanisms through which the DDR operates, elucidate the genetic interactions between different DDR pathways and between DDR pathways and other cellular pathways, will be helpful for therapeutic strategies to treat many human disease.

6. Acknowledgment

This article was supported by National Natural Science Foundation of China 90919030 and 31070691 to Wei-Guo Zhu and 31071117 to Haiying Wang, and Beijing Natural Science Foundation 7112081(Study on the mechanism of HDAC inhibitor depsipeptide in cancer treatment) to Haiying Wang.

7. Abbreviations

The abbreviations used are: ATM, ataxia-telangiectasia mutated; ATR, ATM and Rad3-related; Chk, Checkpoint kinase; IR, ionizing radiation; PIKK, phosphoinositide-3 kinase-related kinases; ATR-IP, ATR-interacting protein; ROS, reactive oxygen species; DSB, double strand break; 5-Aza-CdR, 5'-aza-2'-deoxycytidine; DDR, DNA damage response; MRN, Mre11/Rad50/Nbs1; ssDNA, single strand DNA; 9-1-1, Rad9-Rad1-Hus1; FHA, forkhead-associated; BRCT, BRCA1 C-terminal; PRC1, Polycomb-repressive complex 1; DUBs, deubiquitylating enzymes.

8. References

- Abraham RT (2001) Cell cycle checkpoint signaling through the ATM and ATR kinases. *Genes Dev* 15(17):2177-2196.
- Alarcon-Vargas D & Ronai Z (2002) p53-Mdm2--the affair that never ends. *Carcinogenesis* 23(4):541-547.
- Appella E & Anderson CW (2001) Post-translational modifications and activation of p53 by genotoxic stresses. *Eur J Biochem* 268(10):2764-2772.
- Bakkenist CJ & Kastan MB (2003) DNA damage activates ATM through intermolecular autophosphorylation and dimer dissociation. *Nature* 421(6922):499-506.
- Bakkenist CJ & Kastan MB (2004) Initiating cellular stress responses. *Cell* 118(1):9-17.
- Ball HL, Ehrhardt MR, Mordes DA, Glick GG, Chazin WJ & Cortez D. (2007) Function of a conserved checkpoint recruitment domain in ATRIP proteins. *Mol Cell Biol* 27(9):3367-3377.

- Ball HL, Myers JS, & Cortez D (2005) ATRIP binding to replication protein A-single-stranded DNA promotes ATR-ATRIP localization but is dispensable for Chk1 phosphorylation. *Mol Biol Cell* 16(5):2372-2381.
- Banin S, Moyal L, Shieh S, Taya Y, Anderson C W, Chessa L, Smorodinsky N I, Prives C, Reiss Y, Shiloh Y, & Ziv Y (1998) Enhanced phosphorylation of p53 by ATM in response to DNA damage. *Science* 281(5383):1674-1677.
- Bartek J & Lukas J (2001) Mammalian G1- and S-phase checkpoints in response to DNA damage. *Curr Opin Cell Biol* 13(6):738-747.
- Bekker-Jensen S & Mailand N (2010) Assembly and function of DNA double-strand break repair foci in mammalian cells. *DNA Repair* 9(12):1219-1228.
- Bentley NJ, Holtzman DA, Flaggs G, Keegan KS, DeMaggio A, Ford JC, Hoekstra M & Carr AM. (1996) The *Schizosaccharomyces pombe* rad3 checkpoint gene. *Embo J* 15(23):6641-6651.
- Berkovich E, Monnat RJ, Jr., & Kastan MB (2007) Roles of ATM and NBS1 in chromatin structure modulation and DNA double-strand break repair. *Nat Cell Biol* 9(6):683-690.
- Bermudez VP, Lindsey-Boltz LA, Cesare AJ, Maniwa Y, Griffith JD, Hurwitz J & Sancar A. (2003) Loading of the human 9-1-1 checkpoint complex onto DNA by the checkpoint clamp loader hRad17-replication factor C complex in vitro. *Proc Natl Acad Sci U S A* 100(4):1633-1638.
- Bode AM DZ (2004) Post-translational modification of p53 in tumorigenesis. *Nat Rev Cancer* 4(10):793-805.
- Bonilla CY, Melo JA, & Toczyski DP (2008) Colocalization of sensors is sufficient to activate the DNA damage checkpoint in the absence of damage. *Mol Cell* 30(3):267-276.
- Brown EJ & Baltimore D (2003) Essential and dispensable roles of ATR in cell cycle arrest and genome maintenance. *Genes Dev* 17(5):615-628.
- Burma S, Chen BP, Murphy M, Kurimasa A, & Chen DJ (2001) ATM phosphorylates histone H2AX in response to DNA double-strand breaks. *J Biol Chem* 276(45):42462-42467.
- Byun TS, Pacek M, Yee MC, Walter JC, & Cimprich KA (2005) Functional uncoupling of MCM helicase and DNA polymerase activities activates the ATR-dependent checkpoint. *Genes Dev* 19(9):1040-1052.
- Canman CE, Lim DS, Cimprich KA, Taya Y, Tamai K, Sakaguchi K, Appella E, Kastan M B & Siliciano, JD. (1998) Activation of the ATM kinase by ionizing radiation and phosphorylation of p53. *Science* 281(5383):1677-1679.
- Carson CT, Schwartz RA, Stracker TH, Lilley CE, Lee DV & Weitzman MD. (2003) The Mre11 complex is required for ATM activation and the G2/M checkpoint. *Embo J* 22(24):6610-6620.
- Cerosaletti K & Concannon P (2004) Independent roles for nibrin and Mre11-Rad50 in the activation and function of Atm. *J Biol Chem* 279(37):38813-38819.
- Chai G, Li L, Zhou W, Wu L, Zhao Y, Wang D, Lu S, Yu Y, Wang H, McNutt MA, Hu YG, Chen Y, Yang Y, Wu X, Otterson GA & Zhu WG. (2008) HDAC inhibitors act with 5-aza-2'-deoxycytidine to inhibit cell proliferation by suppressing removal of incorporated abases in lung cancer cells. *PLoS One* 3(6):e2445.
- Chen X, Arciero CA, Wang C, Broccoli D, & Godwin AK (2006) BRCC36 is essential for ionizing radiation-induced BRCA1 phosphorylation and nuclear foci formation. *Cancer Res* 66(10):5039-5046.

- Chou DM, Adamson B, Dephore NE, Tan X, Nottke AC, Hurov KE, Gygi SP, Colaiacovo MP & Elledge SJ. (2010) A chromatin localization screen reveals poly (ADP ribose)-regulated recruitment of the repressive polycomb and NuRD complexes to sites of DNA damage. *Proc Natl Acad Sci U S A* 107(43):18475-18480.
- Cimprich KA & Cortez D (2008) ATR: an essential regulator of genome integrity. *Nat Rev Mol Cell Biol* 9(8):616-627.
- Cliby WA, Roberts CJ, Cimprich KA, Stringer CM, Lamb JR, Schreiber SL & Friend SH. (1998) Overexpression of a kinase-inactive ATR protein causes sensitivity to DNA-damaging agents and defects in cell cycle checkpoints. *Embo J* 17(1):159-169.
- Cook PJ, Ju BG, Telese F, Wang X, Glass CK & Rosenfeld MG.. (2009) Tyrosine dephosphorylation of H2AX modulates apoptosis and survival decisions. *Nature* 458(7238):591-596.
- Cortez D, Guntuku S, Qin J, & Elledge SJ (2001) ATR and ATRIP: partners in checkpoint signaling. *Science* 294(5547):1713-1716.
- Costanzo V & Gautier J (2003) Single-strand DNA gaps trigger an ATR- and Cdc7-dependent checkpoint. *Cell Cycle* 2(1):17.
- Difilippantonio S, Celeste A, Fernandez-Capetillo O, Chen HT, Reina San Martin B, Van Laethem F, Yang YP, Petukhova GV, Eckhaus M, Feigenbaum L, Manova K, Kruhlak M, Camerini-Otero RD, Sharan S, Nussenzweig M & Nussenzweig A. (2005) Role of Nbs1 in the activation of the Atm kinase revealed in humanized mouse models. *Nat Cell Biol* 7(7):675-685.
- Doil C, Mailand N, Bekker-Jensen S, Menard P, Larsen DH, Pepperkok R, Ellenberg J, Panier S, Durocher D, Bartek J, Lukas J & Lukas C. (2009) RNF168 binds and amplifies ubiquitin conjugates on damaged chromosomes to allow accumulation of repair proteins. *Cell* 136(3):435-446.
- Dumaz N & Meek DW (1999) Serine15 phosphorylation stimulates p53 transactivation but does not directly influence interaction with HDM2. *Embo J* 18(24):7002-7010.
- Elledge SJ (1996) Cell cycle checkpoints: preventing an identity crisis. *Science* 274(5293):1664-1672.
- Ellison V & Stillman B (2003) Biochemical characterization of DNA damage checkpoint complexes: clamp loader and clamp complexes with specificity for 5' recessed DNA. *PLoS Biol* 1(2):E33.
- Falck J, Petrini JH, Williams BR, Lukas J, & Bartek J (2002) The DNA damage-dependent intra-S phase checkpoint is regulated by parallel pathways. *Nat Genet* 30(3):290-294.
- Garcia-Higuera I, Taniguchi T, Ganesan S, Meyn MS, Timmers C, Hejna J, Grompe M & D'Andrea AD. (2001) Interaction of the Fanconi anemia proteins and BRCA1 in a common pathway. *Mol Cell* 7(2):249-262.
- Graves PR, Yu L, Schwarz JK, Gales J, Sausville EA, O'Connor PM & Piwnicka-Worms H. (2000) The Chk1 protein kinase and the Cdc25C regulatory pathways are targets of the anticancer agent UCN-01. *J Biol Chem* 275(8):5600-5605.
- Gu W & Roeder RG (1997) Activation of p53 sequence-specific DNA binding by acetylation of the p53 C-terminal domain. *Cell* 90(4):595-606.
- Gupta A, Sharma GG, Young CS, Agarwal M, Smith ER, Paull TT, Lucchesi JC, Khanna KK, Ludwig T & Pandita TK. (2005) Involvement of human MOF in ATM function. *Mol Cell Biol* 25(12):5292-5305.

- Harper JW & Elledge SJ (2007) The DNA damage response: ten years after. *Mol Cell* 28(5):739-745.
- Hefferin ML & Tomkinson AE (2005) Mechanism of DNA double-strand break repair by non-homologous end joining. *DNA Repair (Amst)* 4(6):639-648.
- Hoeijmakers JH (2009) DNA damage, aging, and cancer. *N Engl J Med* 361(15):1475-1485.
- Hoeijmakers JH (2001) Genome maintenance mechanisms for preventing cancer. *Nature* 411(6835):366-374.
- Hofmann K & Bucher P (1995) The FHA domain: a putative nuclear signalling domain found in protein kinases and transcription factors. *Trends Biochem Sci* 20(9):347-349.
- Huang J, Huen MS, Kim H, Leung CC, Glover JN, Yu X & Chen J. (2009) RAD18 transmits DNA damage signalling to elicit homologous recombination repair. *Nat Cell Biol* 11(5):592-603.
- Huen MS & Chen J (2008) The DNA damage response pathways: at the crossroad of protein modifications. *Cell Res* 18(1):8-16.
- Ikura T, Tashiro S, Kakino A, Shima H, Jacob N, Amunugama R, Yoder K, Izumi S, Kuraoka I, Tanaka K, Kimura H, Ikura M, Nishikubo S, Ito T, Muto A, Miyagawa K, Takeda S, Fishel R, Igarashi K & Kamiya K. (2007) DNA damage-dependent acetylation and ubiquitination of H2AX enhances chromatin dynamics. *Mol Cell Biol* 27(20):7028-7040.
- Ito A KY, Lai CH, Kovacs JJ, Higashimoto Y, Appella E, Yao TP (2002) MDM2-HDAC1-mediated deacetylation of p53 is required for its degradation. *EMBO J.* 21(22):6236-6245.
- Ivanov GS, Ivanova T, Kurash J, Ivanov A, Chuikov S, Gizatullin F, Herrera-Medina EM, Rauscher F 3rd, Reinberg D & Barlev NA. (2007) Methylation-acetylation interplay activates p53 in response to DNA damage. *Mol Cell Biol* 27(19):6756-6769.
- Kaidi A, Weinert BT, Choudhary C, & Jackson SP (2010) Human SIRT6 promotes DNA end resection through CtIP deacetylation. *Science* 329(5997):1348-1353.
- Kanu N & Behrens A (2007) ATMIN defines an NBS1-independent pathway of ATM signalling. *Embo J* 26(12):2933-2941.
- Kao GD, McKenna WG, Guenther MG, Muschel RJ, Lazar MA & Yen TJ. (2003) Histone deacetylase 4 interacts with 53BP1 to mediate the DNA damage response. *J Cell Biol* 160(7):1017-1027.
- Khanna KK & Jackson SP (2001) DNA double-strand breaks: signaling, repair and the cancer connection. *Nat Genet* 27(3):247-254.
- Kolas NK, Chapman JR, Nakada S, Ylanko J, Chahwan R, Sweeney FD, Panier S, Mendez M, Wildenhain J, Thomson TM, Pelletier L, Jackson SP & Durocher D. (2007) Orchestration of the DNA-damage response by the RNF8 ubiquitin ligase. *Science* 318(5856):1637-1640.
- Kondo T, Wakayama T, Naiki T, Matsumoto K, & Sugimoto K (2001) Recruitment of Mec1 and Ddc1 checkpoint proteins to double-strand breaks through distinct mechanisms. *Science* 294(5543):867-870.
- Kozlov SV, Graham ME, Jakob B, Tobias F, Kijas AW, Tanuji M, Chen P, Robinson PJ, Taucher-Scholz G, Suzuki K, So S, Chen D & Lavin MF. (2011) Autophosphorylation and ATM activation: additional sites add to the complexity. *J Biol Chem* 286(11):9107-9119.
- Kruse JP & Gu W (2009) Modes of p53 regulation. *Cell* 137(4):609-622.

- Kumagai A, Lee J, Yoo HY, & Dunphy WG (2006) TopBP1 activates the ATR-ATRIP complex. *Cell* 124(5):943-955.
- Lavin MF & Shiloh Y (1997) The genetic defect in ataxia-telangiectasia. *Annu Rev Immunol* 15:177-202.
- Lavin MF (2008) Ataxia-telangiectasia: from a rare disorder to a paradigm for cell signalling and cancer. *Nat Rev Mol Cell Biol* 9(10):759-769.
- Lee J, Kumagai A, & Dunphy WG (2003) Claspin, a Chk1-regulatory protein, monitors DNA replication on chromatin independently of RPA, ATR, and Rad17. *Mol Cell* 11(2):329-340.
- Lee JH & Paull TT (2004) Direct activation of the ATM protein kinase by the Mre11/Rad50/Nbs1 complex. *Science* 304(5667):93-96.
- Lee JH & Paull TT (2005) ATM activation by DNA double-strand breaks through the Mre11-Rad50-Nbs1 complex. *Science* 308(5721):551-554.
- Li J, Williams BL, Haire LF, Goldberg M, Wilker E, Durocher D, Yaffe MB, Jackson SP & Smerdon SJ. (2002) Structural and functional versatility of the FHA domain in DNA-damage signaling by the tumor suppressor kinase Chk2. *Mol Cell* 9(5):1045-1054.
- Li M LJ, Brooks CL, Gu W (2002) Acetylation of p53 inhibits its ubiquitination by Mdm2. *J Biol Chem*. 277(52):50607-50611.
- Lill NL GS, Ginsberg D, DeCaprio J, Livingston DM. (1997) Binding and modulation of p53 by p300/CBP coactivators. *Nature*. 387(6635):823-827.
- Lim DS & Kim ST. (2000) ATM phosphorylates p95/nbs1 in an S-phase checkpoint pathway. *Nature* 404(6778):613-617.
- Liu L SD, Trievel RC, Zhang HB, Marmorstein R, Halazonetis TD, Berger SL (1999) p53 sites acetylated in vitro by PCAF and p300 are acetylated in vivo in response to DNA damage. *Mol Cell Biol*. 19(2):1202-1209.
- Liu X, Wang D, Zhao Y, Tu B, Zheng Z, Wang L, Wang H, Gu W, Roeder RG & Zhu WG. (2011) Methyltransferase Set7/9 regulates p53 activity by interacting with Sirtuin 1 (SIRT1). *Proc Natl Acad Sci U S A* 108(5):1925-1930.
- Lodish H, Berk A, Matsudaira P, Kaiser CA, Krieger M, Scott MP, Zipursky SL & Darnell J.(2004). *Molecular Biology of the Cell*, p963. WH Freeman: New York, NY. 5th ed.
- Lopez-Girona A, Furnari B, Mondesert O, & Russell P (1999) Nuclear localization of Cdc25 is regulated by DNA damage and a 14-3-3 protein. *Nature* 397(6715):172-175.
- Lou Z, Minter-Dykhouse K, Franco S, Gostissa M, Rivera MA, Celeste A, Manis JP, van Deursen J, Nussenzweig A, Paull TT, Alt FW & Chen J. (2006) MDC1 maintains genomic stability by participating in the amplification of ATM-dependent DNA damage signals. *Mol Cell* 21(2):187-200.
- Lukas C, Falck J, Bartkova J, Bartek J, & Lukas J (2003) Distinct spatiotemporal dynamics of mammalian checkpoint regulators induced by DNA damage. *Nat Cell Biol* 5(3):255-260.
- Luo J, Li, M., Tang, Y., Laszkowska, M., Roeder, R. G. & Gu, W. (2004) Acetylation of p53 augments its site-specific DNA binding both in vitro and in vivo. *Proc Natl Acad Sci U S A* 101(8):2259-2264.
- MacDougall CA, Byun TS, Van C, Yee MC, & Cimprich KA (2007) The structural determinants of checkpoint activation. *Genes Dev* 21(8):898-903.

- Mailand N, Bekker-Jensen S, Faustrup H, Melander F, Bartek J, Lukas C & Lukas J. (2007) RNF8 ubiquitylates histones at DNA double-strand breaks and promotes assembly of repair proteins. *Cell* 131(5):887-900.
- Matsuoka S, Rotman G, Ogawa A, Shiloh Y, Tamai K & Elledge SJ. (2000) Ataxia telangiectasia-mutated phosphorylates Chk2 in vivo and in vitro. *Proc Natl Acad Sci U S A* 97(19):10389-10394.
- Matsuoka S, Ballif BA, Smogorzewska A, McDonald ER 3rd, Hurov KE, Luo J, Bakalarski CE, Zhao Z, Solimini N, Lerenthal Y, Shiloh Y, Gygi SP & Elledge SJ. (2007) ATM and ATR substrate analysis reveals extensive protein networks responsive to DNA damage. *Science* 316(5828):1160-1166.
- Maya R, Balass M, Kim ST, Shkedy D, Leal JF, Shifman O, Moas M, Buschmann T, Ronai Z, Shiloh Y, Kastan MB, Katzir E & Oren M. (2001) ATM-dependent phosphorylation of Mdm2 on serine 395: role in p53 activation by DNA damage. *Genes Dev* 15(9):1067-1077.
- McGowan CH & Russell P (2004) The DNA damage response: sensing and signaling. *Curr Opin Cell Biol* 16(6):629-633.
- Melo JA, Cohen J, & Toczyski DP (2001) Two checkpoint complexes are independently recruited to sites of DNA damage in vivo. *Genes Dev* 15(21):2809-2821.
- Miller KM, Tjeertes JV, Coates J, Legube G, Polo SE, Britton S & Jackson SP. (2010) Human HDAC1 and HDAC2 function in the DNA-damage response to promote DNA nonhomologous end-joining. *Nat Struct Mol Biol* 17(9):1144-1151.
- Myers JS & Cortez D (2006) Rapid activation of ATR by ionizing radiation requires ATM and Mre11. *J Biol Chem* 281(14):9346-9350.
- Nag A, Germaniuk-Kurowska A, Dimri M, Sassack M, Gurumurthy CB, Gao Q, Dimri G, Band H, & Band V (2007) An essential role of human ADA3 in p53 acetylation. *J Biol Chem*. 282,8812-8820.
- Nakanishi K, *et al.* (2002) Interaction of FANCD2 and NBS1 in the DNA damage response. *Nat Cell Biol* 4(12):913-920.
- Namiki Y & Zou L (2006) ATRIP associates with replication protein A-coated ssDNA through multiple interactions. *Proc Natl Acad Sci U S A* 103(3):580-585.
- Nicassio F, Taniguchi T, Ranganathan V, New HV, Moreau LA, Stotsky M, Mathew CG, Kastan MB, Weaver DT & D'Andrea AD. (2007) Human USP3 is a chromatin modifier required for S phase progression and genome stability. *Curr Biol* 17(22):1972-1977.
- Oberdoerffer P, Michan S, McVay M, Mostoslavsky R, Vann J, Park SK, Hartlerode A, Stegmuller J, Hafner A, Loerch P, Wright SM, Mills KD, Bonni A, Yankner BA, Scully R, Prolla TA, Alt FW & Sinclair DA. (2008) SIRT1 redistribution on chromatin promotes genomic stability but alters gene expression during aging. *Cell* 135(5):907-918.
- Osborn AJ, Elledge SJ, & Zou L (2002) Checking on the fork: the DNA-replication stress-response pathway. *Trends Cell Biol* 12(11):509-516.
- Pandita TK, Lieberman HB, Lim DS, Dhar S, Zheng W, Taya Yand Kastan MB. (2000) Ionizing radiation activates the ATM kinase throughout the cell cycle. *Oncogene* 19(11):1386-1391
- Parrilla-Castellar ER, Arlander SJ, & Karnitz L (2004) Dial 9-1-1 for DNA damage: the Rad9-Hus1-Rad1 (9-1-1) clamp complex. *DNA Repair (Amst)* 3(8-9):1009-1014.

- Passalaris TM, Benanti JA, Gewin L, Kiyono T, & Galloway DA (1999) The G(2) checkpoint is maintained by redundant pathways. *Mol Cell Biol* 19(9):5872-5881.
- Paul R, Andreassen, Alan D. D'Andrea & Toshiyasu Taniguchi.. (2004) Cell cycle-dependent dynamic localization of a bacterial response regulator with a novel di-guanylate cyclase output domain. *Genes Dev* 18(6):715-727.
- Paull TT & Lee JH (2005) The Mre11/Rad50/Nbs1 complex and its role as a DNA double-strand break sensor for ATM. *Cell Cycle* 4(6):737-740.
- Peng CY, Graves PR, Thoma RS, Wu Z, Shaw AS & Piwnica-Worms H. (1997) Mitotic and G2 checkpoint control: regulation of 14-3-3 protein binding by phosphorylation of Cdc25C on serine-216. *Science* 277(5331):1501-1505.
- Pickart CM (2001) Mechanisms underlying ubiquitination. *Annu Rev Biochem* 70:503-533.
- Rhind N, Furnari B, & Russell P (1997) Cdc2 tyrosine phosphorylation is required for the DNA damage checkpoint in fission yeast. *Genes Dev* 11(4):504-511.
- Richard DJ, Bolderson E, Cubeddu L, Wadsworth RI, Savage K, Sharma GG, Nicolette ML, Tsvetanov S, McIlwraith MJ, Pandita RK, Takeda S, Hay RT, Gautier J, West SC, Paull TT, Pandita TK, White MF & Khanna KK. (2008) Single-stranded DNA-binding protein hSSB1 is critical for genomic stability. *Nature* 453(7195):677-681.
- So S, Davis AJ, & Chen DJ (2009) Autophosphorylation at serine 1981 stabilizes ATM at DNA damage sites. *J Cell Biol* 187(7):977-990.
- Savitsky K, Bar-Shira A, Gilad S, Rotman G, Ziv Y, Vanagaite L, Tagle DA, Smith S, Uziel T, Sfez S, Ashkenazi M, Pecker I, Frydman M, Harnik R, Patanjali SR, Simmons A, Clines GA, Sartiel A, Gatti RA, Chessa L, Sanal O, Lavin MF, Jaspers NG, Taylor AM, Arlett CF, Miki T, Weissman SM, Lovett M, Collins FS & Shiloh Y. (1995) A single ataxia telangiectasia gene with a product similar to PI-3 kinase. *Science* 268(5218):1749-1753.
- Scully R, Chen J, Ochs RL, Keegan K, Hoekstra M, Feunteun J & Livingston DM. (1997) Dynamic changes of BRCA1 subnuclear location and phosphorylation state are initiated by DNA damage. *Cell* 90(3):425-435.
- Shao G, Lilli DR, Patterson-Fortin J, Coleman KA, Morrissey DE & Greenberg RA. (2009) TheRap80-BRCC36 de-ubiquitinating enzyme complex antagonizes RNF8-Ubc13-dependent ubiquitination events at DNA double strand breaks. *Proc Natl Acad Sci U S A* 106(9):3166-3171.
- Shi X, Kachirskaja I, Yamaguchi H, West LE, Wen H, Wang EW, Dutta S, Appella E & Gozani O.(2007) Modulation of p53 function by SET8-mediated methylation at lysine 382. *Mol Cell* 27(4):636-646.
- Shieh SY, Ahn J, Tamai K, Taya Y, & Prives C (2000) The human homologs of checkpoint kinases Chk1 and Cds1 (Chk2) phosphorylate p53 at multiple DNA damage-inducible sites. *Genes Dev* 14(3):289-300.
- Shiloh Y (2001) ATM and ATR: networking cellular responses to DNA damage. *Curr Opin Genet Dev* 11(1):71-77.
- Shiloh Y (2003) ATM and related protein kinases: safeguarding genome integrity. *Nat Rev Cancer* 3(3):155-168.
- Shiloh Y (2006) The ATM-mediated DNA-damage response: taking shape. *Trends Biochem Sci* 31(7):402-410.
- Shiotani B & Zou L (2009) ATR signaling at a glance. *J Cell Sci* 122(Pt 3):301-304.

- Sobhian B, Shao G, Lilli DR, Culhane AC, Moreau LA, Xia B, Livingston DM & Greenberg RA. (2007) RAP80 targets BRCA1 to specific ubiquitin structures at DNA damage sites. *Science* 316(5828):1198-1202.
- Stewart GS (2009) Solving the RIDDLE of 53BP1 recruitment to sites of damage. *Cell Cycle* 8(10):1532-1538.
- Stokes MP, Van Hatten R, Lindsay HD, & Michael WM (2002) DNA replication is required for the checkpoint response to damaged DNA in *Xenopus* egg extracts. *J Cell Biol* 158(5):863-872.
- Stucki M, Clapperton JA, Mohammad D, Yaffe MB, Smerdon SJ & Jackson SP. (2005) MDC1 directly binds phosphorylated histone H2AX to regulate cellular responses to DNA double-strand breaks. *Cell* 123(7):1213-1226.
- Stucki M & Jackson SP (2006) gammaH2AX and MDC1: anchoring the DNA-damage-response machinery to broken chromosomes. *DNA Repair (Amst)* 5(5):534-543.
- Sun Y, Jiang X, Chen S, Fernandes N, & Price BD (2005) A role for the Tip60 histone acetyltransferase in the acetylation and activation of ATM. *Proc Natl Acad Sci U S A* 102(37):13182-13187.
- Sun Y, Xu Y, Roy K, & Price BD (2007) DNA damage-induced acetylation of lysine 3016 of ATM activates ATM kinase activity. *Mol Cell Biol* 27(24):8502-8509.
- Sun Y, Jiang X, Xu Y, Ayrappetov MK, Moreau LA, Whetstone JR & Price BD. (2009) Histone H3 methylation links DNA damage detection to activation of the tumour suppressor Tip60. *Nat Cell Biol* 11(11):1376-1382.
- Sykes SM, Mellert HS, Holbert MA, Li K, Marmorstein R, Lane WS & McMahon SB. (2006) Acetylation of the p53 DNA-binding domain regulates apoptosis induction. *Mol Cell* 24(6):841-851.
- Tanaka T, Halicka HD, Huang X, Traganos F, & Darzynkiewicz Z (2006) Constitutive histone H2AX phosphorylation and ATM activation, the reporters of DNA damage by endogenous oxidants. *Cell Cycle* 5(17):1940-1945.
- Tanaka T, Huang X, Halicka HD, Zhao H, Traganos F, Albino AP, Dai W & Darzynkiewicz Z. (2007) Cytometry of ATM activation and histone H2AX phosphorylation to estimate extent of DNA damage induced by exogenous agents. *Cytometry A* 71(9):648-661.
- Tang Y, Luo J, Zhang W, & Gu W (2006) Tip60-dependent acetylation of p53 modulates the decision between cell-cycle arrest and apoptosis. *Mol Cell* 24(6):827-839.
- Tang Y, Zhao W, Chen Y, Zhao Y, & Gu W (2008) Acetylation is indispensable for p53 activation. *Cell* 133(4):612-626.
- Taniguchi T, Garcia-Higuera I, Andreassen PR, Gregory RC, Grompe M & D'Andrea AD. (2002) S-phase-specific interaction of the Fanconi anemia protein, FANCD2, with BRCA1 and RAD51. *Blood* 100(7):2414-2420.
- Tibbetts RS, Brumbaugh KM, Williams JM, Sarkaria JN, Cliby WA, Shieh SY, Taya Y, Prives C & Abraham RT. (1999) A role for ATR in the DNA damage-induced phosphorylation of p53. *Genes Dev* 13(2):152-157.
- Toledo F & Wahl GM (2006) Regulating the p53 pathway: in vitro hypotheses, in vivo veritas. *Nat Rev Cancer* 6(12):909-923.
- Trenz K, Smith E, Smith S, & Costanzo V (2006) ATM and ATR promote Mre11 dependent restart of collapsed replication forks and prevent accumulation of DNA breaks. *Embo J* 25(8):1764-1774.

- Uziel T, Lerenthal Y, Moyal L, Andegeko Y, Mittelman L & Shiloh Y. (2003) Requirement of the MRN complex for ATM activation by DNA damage. *Embo J* 22(20):5612-5621.
- van Attikum H & Gasser SM (2009) Crosstalk between histone modifications during the DNA damage response. *Trends Cell Biol* 19(5):207-217.
- van den Bosch M, Bree RT, & Lowndes NF (2003) The MRN complex: coordinating and mediating the response to broken chromosomes. *EMBO Rep* 4(9):844-849.
- Wang H, Zhao Y, Li L, McNutt MA, Wu L, Lu S, Yu Y, Zhou W, Feng J, Chai G, Yang Y & Zhu WG.(2008) An ATM- and Rad3-related (ATR) signaling pathway and a phosphorylation-acetylation cascade are involved in activation of P53/P21^{Waf1/Cip1} in response to 5-aza-2'-deoxycytidine treatment. *J Biol Chem* 283(5):2564-2574.
- Wright JA, Keegan KS, Herendeen DR, Bentley NJ, Carr AM, Hoekstra MF & Concannon P. (1998) Protein kinase mutants of human ATR increase sensitivity to UV and ionizing radiation and abrogate cell cycle checkpoint control. *Proc Natl Acad Sci U S A* 95(13):7445-7450.
- Wu X & Ranganathan V. (2000) ATM phosphorylation of Nijmegen breakage syndrome protein is required in a DNA damage response. *Nature* 405(6785):477-482.
- Wu-Baer F, Lagazon K, Yuan W, & Baer R (2003) The BRCA1/BARD1 heterodimer assembles polyubiquitin chains through an unconventional linkage involving lysine residue K6 of ubiquitin. *J Biol Chem* 278(37):34743-34746.
- Wyman C & Kanaar R (2006) DNA double-strand break repair: all's well that ends well. *Annu Rev Genet* 40:363-383.
- Yan S & Michael WM (2009) TopBP1 and DNA polymerase alpha-mediated recruitment of the 9-1-1 complex to stalled replication forks: implications for a replication restart-based mechanism for ATR checkpoint activation. *Cell Cycle* 8(18):2877-2884.
- Yang J, Yu Y, Hamrick HE, & Duerksen-Hughes PJ (2003) ATM, ATR and DNA-PK: initiators of the cellular genotoxic stress responses. *Carcinogenesis* 24(10):1571-1580.
- Yoo HY, Kumagai A, Shevchenko A, & Dunphy WG (2007) Ataxia-telangiectasia mutated (ATM)-dependent activation of ATR occurs through phosphorylation of TopBP1 by ATM. *J Biol Chem* 282(24):17501-17506.
- Yoshioka K, Yoshioka Y, & Hsieh P (2006) ATR kinase activation mediated by MutSalpha and MutLalpha in response to cytotoxic O6-methylguanine adducts. *Mol Cell* 22(4):501-510.
- You Z, Kong L, & Newport J (2002) The role of single-stranded DNA and polymerase alpha in establishing the ATR, Hus1 DNA replication checkpoint. *J Biol Chem* 277(30):27088-27093.
- Yu X, Chini CC, He M, Mer G, & Chen J (2003) The BRCT domain is a phospho-protein binding domain. *Science* 302(5645):639-642.
- Yuan Z, Zhang X, Sengupta N, Lane WS, & Seto E (2007) SIRT1 regulates the function of the Nijmegen breakage syndrome protein. *Mol Cell* 27(1):149-162.
- Zhan Q, Bae I, Kastan MB, & Fornace AJ, Jr. (1994) The p53-dependent gamma-ray response of GADD45. *Cancer Res* 54(10):2755-2760.
- Zhao S & Weng YC. (2000) Functional link between ataxia-telangiectasia and Nijmegen breakage syndrome gene products. *Nature* 405(6785):473-477.
- Zhao Y, Lu S, Wu L, Chai G, Wang H, Chen Y, Sun J, Yu Y, Zhou W, Zheng Q, Wu M, Otterson GA & Zhu WG. (2006) Acetylation of p53 at lysine 373/382 by the histone

- deacetylase inhibitor depsipeptide induces expression of p21(Waf1/Cip1). *Mol Cell Biol* 26(7):2782-2790.
- Zhou BB & Elledge SJ (2000) The DNA damage response: putting checkpoints in perspective. *Nature* 408(6811):433-439.
- Zhu WG, HT, Ke Y, Wang P, Lu S, Duan W, Dai Z, Tong T, Villalona-Calero MA, Plass C & Otterson GA (2004) 5-aza-2'-deoxycytidine activates the p53/p21Waf1/Cip1 pathway to inhibit cell proliferation. *J Biol Chem*. 279(15):15161-15166.
- Zou L, Cortez D, & Elledge SJ (2002) Regulation of ATR substrate selection by Rad17-dependent loading of Rad9 complexes onto chromatin. *Genes Dev* 16(2):198-208.
- Zou L, Liu D, & Elledge SJ (2003a) Replication protein A-mediated recruitment and activation of Rad17 complexes. *Proc Natl Acad Sci U S A* 100(24):13827-13832.
- Zou L & Elledge SJ (2003b) Sensing DNA damage through ATRIP recognition of RPA-ssDNA complexes. *Science* 300(5625):1542-1548.

IntechOpen

DNA Replication-Current Advances

Edited by Dr Herve Seligmann

ISBN 978-953-307-593-8

Hard cover, 694 pages

Publisher InTech

Published online 01, August, 2011

Published in print edition August, 2011

The study of DNA advanced human knowledge in a way comparable to the major theories in physics, surpassed only by discoveries such as fire or the number zero. However, it also created conceptual shortcuts, beliefs and misunderstandings that obscure the natural phenomena, hindering its better understanding. The deep conviction that no human knowledge is perfect, but only perfectible, should function as a fair safeguard against scientific dogmatism and enable open discussion. With this aim, this book will offer to its readers 30 chapters on current trends in the field of DNA replication. As several contributions in this book show, the study of DNA will continue for a while to be a leading front of scientific activities.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Haiying Wang, Ping Shen and Wei-Guo Zhu (2011). Damage and Replication Stress Responses, DNA Replication-Current Advances, Dr Herve Seligmann (Ed.), ISBN: 978-953-307-593-8, InTech, Available from: <http://www.intechopen.com/books/dna-replication-current-advances/damage-and-replication-stress-responses>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2011 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike-3.0 License](https://creativecommons.org/licenses/by-nc-sa/3.0/), which permits use, distribution and reproduction for non-commercial purposes, provided the original is properly cited and derivative works building on this content are distributed under the same license.

IntechOpen

IntechOpen