We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

A Diachronic Study of Diabetic Nephropathy in Two Autochthonous Lines of Rats to Understand Diabetic Chronic Complications

Juan Carlos Picena, Silvana Marisa Montenegro, Alberto Enrique D'Ottavio, María Cristina Tarrés and Stella Maris Martínez
National University of Rosario
Argentina

1. Introduction

The worldwide epidemic of Type 2 Diabetes Mellitus, a result of recent deep changes in human lifestyle like sedentary and overly rich nutrition, has dramatic consequences in terms of morbidity, mortality and health care costs (Zimmet *et al.*, 2001). Type 2 Diabetes Mellitus is a complex and multifactorial metabolic disorder which includes distinct nosological entities whose common patterns are hyperglycaemia, resistance to insulin and progressive chronic complications (American Diabetic Association, 2009).

A significant proportion of patients will develop a clinically relevant diabetic nephropathy with glomerulopathy, hyalinization of afferent and efferent arterioles, tubular and interstitial lesions (Fioretto *et al.*, 2008). Nephropathy is a very serious complication in both Type 1 and Type 2 Diabetes Mellitus because patients with diabetic kidney disease are at a higher risk of mortality, mostly from cardiovascular complications, than diabetic patients without diabetic nephropathy (Dronavalli & Barkis, 2008).

In the Western World the incidence of end-stage renal failure associated to the burden of diabetic nephropathy has increased in recent years and represents a failure of current disease control measure with serious public health implications (Stewart *et al.*, 2006; Villar & Zaoui, 2010). In Latin America, where incidence of Type 2 Diabetes Mellitus is on the rise and has reached epidemic proportions, people usually have poor disease control with the consequence of very high rates of diabetic nephropathy (Caballero & Tenzer, 2007). A study performed in the Asian Pacific region has also shown that diabetic nephropathy was the most common cause of End-Stage Renal Disease in 9 of the 12 countries surveyed and 6 of the 12 countries had greater than 35% of their dialysis patients age 60 years and older (Lee, 2003). In United States, between 20% and 40% of patients ultimately develop diabetic nephropathy, the most common cause of End-Stage Renal Disease requiring dialysis (Dronavalli & Barkis, 2008). Since 2000, the adjusted rate of prevalent End-Stage Renal Disease cases in the United States population ages 65-74, has increased 25 % with an enormous economic impact on the healthcare systems (United States Renal Data System, 2010). In sum, worldwide incidence of diabetic nephropathy, a frequently silent and unrecognized disease, has increased about a 50% between 1998 and 2008 and is now the leading cause of chronic kidney disease and End-Stage Renal Disease (United States Renal

Data System, 2008). These epidemiological findings along with the several pathways which have been implicated in the progression of diabetic nephropathy, such as systemic and glomerular hypertension, metabolic derangements (hyperglycaemia, hyperlipidaemia, hyperinsulinaemia), and oxidative stress among others, indicate the complexity and the magnitude of diabetic nephropathy as a health concern.

Streptozotocin-induced pancreatic injury as well as uninephrectomy have been used for creating both rat and mouse models of Type 1 Diabetes Mellitus which develop renal injury with similarities to human diabetic nephropathy (Tesch & Allen., 2007). Nephropathy is also a complication in rat models of spontaneous Type 2 Diabetes Mellitus as the obese Zucker (fatty) rat, the Goto Kakizaki rat (Janssen *et al.*, 1999) and the hypertensive/NIH-corpulent rat (Nangaku *et al.*, 2005) and in mice models as the BTBR Ob/Ob mutant mice (Hudkins *et al.*, 2010). Either spontaneous or induced, rodent models of diabetes have been extensively used in diabetes research because of their usefulness for understanding the pathogenesis, the complications and the genetic or environmental influences that increase the risks of suffering Type 2 Diabetes Mellitus (Srinivasan & Ramarao, 2007). These models, which depict many clinical features or related phenotypes of the disease, differ among them not only in its metabolic alterations but in their etiopathogenia (Breyer *et al.* 2005; Chatzigeorgiou *et al.*, 2009). Although their use has allowed important advances toward the better understanding of human diabetic nephropathy, it has been recently pointed out that a lack of reliable animal models of diabetic complications still persists (Brosius *et al.*, 2009).

The eSS rat is a spontaneously non-obese diabetic model obtained in Rosario, Argentina, by genetic manipulation, described in detail by Martinez *et al.* (1993). eSS rats evidence a progressive impaired glucose tolerance from 70 days of age onwards with hyperglycaemia and glycosuria worsening with ageing, being these signs much more striking in males (Martínez *et al.*, 1988). Males exhibit insulin resistance and consequent insulin hypersecretion till 12 months of age but the hormone secretion decreases as diabetes progresses in the sequence of events from insulin resistance to overt diabetes (Martínez *et al.*, 1988). This finding is coincident with conspicuous changes in the pancreas of 18 month-old eSS males, such as disruption of islet architecture and areas of pronounced fibrosis (Tarrés *et al.*, 1992; Picena *et al.*, 2007).

During the second year of life, eSS males are known to develop glomerulosclerosis and tubular nephrosis similar to those found in long-term human diabetes (Tarrés *et al.*, 1992; Martínez *et al.*, 1993). Older eSS males also develop bilateral cataracts but not retinopathy (Tarrés *et al.*, 1990; Picena *et al.*, 2005). Hence, the eSS rat was proposed as a model of non-insulin dependent diabetes characterized by complex glucose and lipid disorders (Martínez *et al.*, 1993; Montanaro *et al.*, 2003; Daniele *et al.*, 2010).

eSMT rats (IIMe/Fm eSMT) derived from crossing eSS with β rats, both parent strains belonging to the IIM stock (Calderari *et al.*, 1991). β (IIMb/Fm β) is a line of moderately and spontaneous obese, fertile and hypertriglyceridemic rats (Calderari *et al.*, 1987; Hisano *et al.*, 1994). Upon crossing the F1 to obtain F2, the new colony eSMT was maintained as a closed one, where the full genetic history of each breeding rat is preserved (Tarrés *et al.*, 2000). Young eSMT animals are more robust than eSS and develop higher levels of fasting hyperglycaemia, glycosuria, glucose intolerance, blood triglycerides and total cholesterol (Tarrés *et al.*, 2000; Montenegro *et al.*, 2005). eSMT males of 6 and 9 months show islets with altered shapes and fibrosis, as well as sporadic images of apoptosis and at 12 months of age, islets are reduced in number and size, resembling the histoarchitecture of eSS males during

their second year of life (Tarrés *et al.*, 2000). The sexual dimorphism is less evident in eSMT than in eSS diabetic syndrome (Picena, 2007).

The aim of this chapter is to present a wide and diachronic study of renal histopathology in eSS and eSMT rats, two strains which develop hereditary type 2 diabetic syndromes with different expression of the metabolic disorders, in order to characterize morphological changes. The arrival to conclusions could improve our knowledge and understanding of diabetic nephropathy.

2. Material and methods

The study was conducted in highly inbred 139 eSS (males= 71; females= 68) and 122 eSMT (males= 50; females= 72) rats from 3 to 26-30 months of age. Eumetabolic Wistar male rats of 12 (n= 15) and 21 month-old (n=12) were used as controls.

Since the expected life span of the eSS and the eSMT rats in our laboratory ranges from 24-30 months, the morphological studies of diabetic nephropathy covered their entire life span. From 3 months of age onwards, groups of rats from eSS and eSMT strains were euthanized. Previously, glycaemia at 120 min after 10% glucose overload (200 mg/100 g body weight) via stomach tube was assessed. Blood samples were obtained by tail vein puncture. Plasma samples were analysed for glucose by the glucose oxidase enzymatic method using a commercial kit (Wiener Laboratories, Argentina).

Body and kidney weights were registered in every case. Sections of renal tissue were fixed in 10% neutral-buffered formalin for 24 hours, embedded in paraffin, cut into sections of 4 μ m thick, stained with hematoxylin-eosin (HE) and Periodic Acid-Schiff (PAS), and examined by light microscopy in a blinded fashion. In each histological specimen, capsular and glomerular diameters of 10 superficial and 10 juxtamedullary nephrons, cut at the vascular pole, were measured through a calibrated Shimadzu ® linear scale, placed in the eyepiece of a light microscopy. Correlations between age and body weights, and age and kidney weights were also performed.

Glomeruli were classified as small when the diameter was lower than 80 μ m. Mesangial expansion, defined as the thickening mesangium by the deposit of an acidophilic, amorphous, PAS positive matrix substance and by the increase of cellular components, was considered diffuse when the glomerulus was wholly affected, and segmental if it was partially affected. Depending on its severity, it was graded as mild (ME +) or severe (SE ++). Fifty glomeruli per kidney were randomly selected in each animal. Percentages (%) of glomeruli displaying the same degree of injury (small glomeruli, ME+ and SE ++) were obtained per animal. Correlations between age and percentage of small glomeruli, and age and mildly and severely affected glomeruli were calculated. Nodular increases in mesangial matrix as well as the presence of sclerotic or atubular glomeruli and capsular drop were also sought for. Tubulointerstitial and vascular injuries were considered on the basis of tubular dilation and atrophy, interstitial infiltration and fibrosis as well as of intimal thickening and arteriolar hyalinosis, respectively.

eSS rats were maintained in the breeding facilities of our School of Medicine, Rosario University, Rosario, Argentina. Wistar rats came from the animal breeding facilities in the School of Biochemical Science, Rosario University. Breeding conditions were the same for all the animals, including temperature regulation (24°C) and light-darkness cycles as well as the artificial air exchange. In all the cases, the individuals had remained housed since they were 21 days old, in hanging collective cages. All animals were fed on a complete commercial

diet, special for laboratory rats, and water was *ad libitum*. These experimental conditions were maintained until the animals were euthanized.

All data are reported as the mean ± standard deviation. Statistical processing was partly carried out with the software SPSS 15, using the parametric and non-parametric methods referred below in the presentation of results.

All experimental procedures presented in this study were previously approved by the Bioethics Commission of School of Medicine, which assures adherence to the standards by the Guide for the Care and Use of Laboratory Animals.

3. Results

Body weights data in the experimental eSS and eSMT animals are seen in Table 1.

Age (months)	Line eSS		Line eSMT		Anova (effect: p)
	Males	Females	Males	Females	
3	266±21 (n=6)	137±14 (n=9)	300±43 (n=11)	207±13 (n=9)	Line:0.015 Sex:0.000 Interaction:0.000
6	342±26 (n=19)	245±43 (n=7)	368±55 (n=5)	284±31 (n=11)	Line:0.016 Sex:0.000 Interaction:0.481
9	362±17 (n=4)	232±14 (n=10)	397±61 (n=12)	308±40 (n=6)	Line: 0.004 Sex: 0.000 Interaction: 0.154
12	389±23 (n=14)	252±16 (n= 6)	419±61 (n=6)	338±34 (n=11)	Line:0.002 Sex:0.000 Interaction:0.215
15	399±72 (n=9)	337±15 (n=6)	431±91 (n=5)	370±42 (n=7)	Line:0.051 Sex:0.054 Interaction:0.889
18	402±51 (n=14)	345±36 (n=12)	416±47 (n=8)	350±35 (n=13)	Line:0.440 Sex:0.000 Interaction:0.031
21	357±40 (n=11)	338±26 (n=12)	378±36 (n=8)	348±58 (n=12)	Line:0.063 Sex:0.001 Interaction:0.055
24	282±25 (n=5)	254±52 (n=11)	297±61 (n=4)	319±57 (n=15)	Line:0.087 Sex:0.099 Interaction:0.034
≥ 26		242±26 (n=7)		313±37 (n=4)	Line:0.001

Values expressed as mean ± standard deviation.

Table 1. Body weights (g) in male and female eSS and eSMT rats from 3 to ≥26 months of age.

There was a significant line effect on body weight being eSMT heavier than eSS. A sex-dependent effect was also demonstrated remaining the males heavier than the females.

Table 1 also indicates that body weight increased in male and in female eSS rats till 18 months of age whilst eSMT rats underwent a similar evolution up to 15 months of age. Body weight was significantly correlated with age in both lines (eSS: $r=0.171$; $p=0.050$; eSMT: $r=0.218$; $p=0.016$). The low values obtained could be owed to the marked declination of body weight in the older animals.

As shown in Table 2, kidney weights increased up to 18 months of age in eSS males and till 15 months of age in eSMT males being always greater in eSMT rats than in eSS rats and in males than in females.

Age (months)	Line eSS		Line eSMT		Anova (effect: p)
	Males	Females	Males	Females	
3	2.30±0.27	1.25±0.25	2.40±0.22	1.94±0.24	Line:0.005 Sex:0.000 Interaction:0.050
6	2.79±0.32	2.03±0.15	2.84±0.45	2.40±0.19	Line:0.007 Sex:0.000 Interaction:0.022
9	3.55±0.38	2.26±0.16	3.70±0.31	2.55±0.38	Line:0.041 Sex:0.000 Interaction:0.041
12	3.15±0.63	2.63±0.10	4.02±0.94	2.97±0.29	Line: 0.004 Sex: 0.000 Interaction:0.001
15	3.85±0.52	2.75±0.26	4.68±0.36	3.22±0.28	Line:0.952 Sex:0.000 Interaction:0.027
18	4.06±0.57	2.87±0.45	4.39±0.55	3.20±0.44	Line:0.000 Sex:0.000 Interaction:0.026
21	3.68 ± 1.14	2.82±0.35	3.99±0.49	3.17±0.21	Line:0.546 Sex:0.014 Interaction:0.221
24	3.21 ± 1.13	2.78±0.43	3.88±0.46	3.12±0.37	Line:0.546 Sex:0.014 Interaction:0.032
≥ 26		2.67±0.27		3.01±0.10	Line:0.001

Values expressed as mean ± standard deviation

Table 2. Kidney weights (g) in male and female eSS and eSMT rats from 3 to ≥26 months of age

Direct and significant correlations were demonstrated between kidney weight and age in eSS and eSMT lines (eSS: $r=0.324$; $p=0.000$; eSMT: $r=0.498$; $p=0.000$).

Line and sex effects were shown on glucose tolerance being glycaemia after 120min glucose overload (G120) higher in eSMT rats than in eSS rats and in males than in females. A noticeable worsening of G120 with ageing was also evident (Table 3). eSS males reached the G120 threshold for diagnosis of overt diabetes ($G120\geq200$ mg/dl) from 6 months of age

onwards and eSMT males, from 3 months of age. Females displayed similar values later (eSS from 12 months of age and eSMT from 9 months of age). Wistar rats did not show any significant age-dependent changes remaining normoglycemic until 21 months of age ($G120: 109 \pm 10 \text{ mg/dl}$).

Age (months)	Line eSS		Line eSMT		Anova (effect: p)
	Males	Females	Males	Females	
3	168±37	124±08	201±75	132±20	Line:0.000 Sex:0.012 Interaction:0.330
6	237±44	179±39	264±41	187±28	Line:0.018 Sex:0.034 Interaction:0.342
9	254±43	188±27	303±34	207±35	Line:0.016 Sex:0.002 Interaction:0.431
12	299±64	201±32	324±43	223±42	Line:0.004 Sex:0.000 Interaction:0.154
15	301±55	222±25	356±44	270±33	Line:0.051 Sex:0.054 Interaction:0.889
18	309±49	226±46	429±27	289±26	Line:0.044 Sex:0.000 Interaction:0.330
21	314±42	235±38	433±52	299±35	Line:0.063 Sex:0.001 Interaction:0.752
24	317±35	240±32	438±64	301±27	Line:0.028 Sex:0.029 Interaction:0.349
≥ 26		242±22		309±46	Line:0.001

Values expressed as mean ± standard deviation

Table 3. Glycaemia after 120min glucose overload (mg/dl) in male and female eSS and eSMT rats from 3 to ≥26 months of age.

Present results indicate that at 3 month of age, when animals are still quite normoglycemic with the exception of eSMT males, kidneys did not show lesions visible under the light microscope and that renal injury became apparent from 6-month-old onward. The most frequent glomerular lesions were ME+ and SE++ diffuse mesangial expansion. Direct and significant correlations were evidenced between the percentage of diffuse mesangial expansion and the age (ME+: eSS $\rho = 0.873$; $p = 0.000$. eSMT $\rho = 0.689$; $p = 0.000$ and ME++: eSS $\rho = 0.832$; $p = 0.000$. eSMT $\rho = 0.863$; $p = 0.000$). Nevertheless, an increasing number of small glomeruli ($> 80 \mu\text{m}$) were also verified in ageing eSS and eSMT individuals (eSS $\rho = 0.774$; $p = 0.000$; eSMT $\rho = 0.641$; $p = 0.000$). In 12 month-old eSS rats, the glomerular diameter was higher in females (σ : $110 \pm 8 \mu\text{m}$ vs. ϕ : 120 ± 6 ; $p < 0.01$) whilst in eSMT strain these

values were not different between sexes (♂ : $102 \pm 9 \mu\text{m}$ vs. ♀ : 106 ± 11 ; $p > 0.05$). eSMT females showed similar values of glomerular diameter to those registered in eSS males (eSS ♂ : $110 \pm 8 \mu\text{m}$ eSMT vs. eSMT ♀ : 106 ± 11 ; $p > 0.05$) . Significant correlations between the percentages of small glomeruli, glomeruli with ME+ and with SE++ and age are shown in Figure 1.

Fig. 1. Correlations between age and percentage of small glomeruli (eSS $\rho = 0.774$; $p = 0.000$; eSMT $\rho = 0.641$; $p = 0.000$), ME+ (eSS $\rho = 0.873$; $p = 0.000$. eSMT $\rho = 0.689$; $p = 0.000$), and ME++ (eSS $\rho = 0.832$; $p = 0.000$. eSMT $\rho = 0.863$; $p = 0.000$) in male (●) and female (●) eSS and eSMT rats.

In both lines, glomerular damage began with thickening of the basement membrane in the peripheral sector of the glomeruli, followed by segmentary or diffuse mesangial expansion due to the deposit of an acidophilic, amorphous, PAS-positive substance in addition to increased cellularity (Fig 2 A and 2 B). Progressively, eSMT y eSS males showed a rising number of small glomeruli (Fig 2 C). While glomerular histology by light microscopy was rather normal in eSS rats younger than 12 months of age, a mild PAS positive thickening of glomerular basement membrane was commonly detected in older males. Mild or severe mesangial expansion was usual in 21 month-old eSS males as well as basement membrane thickening. In eSMT rats, mild thickening of the basement membrane was detected from 6 month-old males onwards, and worsened in older animals.

Fig. 2. (A) Mild diffuse mesangial thickening of the glomerular basement membrane (6-month-old eSMT male); (B) Segmentary mesangial expansion with an eccentric area of tuft-to-capsule adhesion (18-month-old eSMT male); (C) Left: small glomerulus (18-month-old eSS male); (D) Glomerulus totally occluded by fibrohyalinosis (21-month-old eSMT male); (E) Glomerulus with severe mesangial expansion and areas of tuft-to-capsule adhesion. Lymphocytic infiltrates near the vascular pole and intertubular interstitium. Atrophic tubules are also observed (24-month-old eSMT male); (F) Arrow: Kimmelstiel-Wilson-like nodular lesion (18-month-old eSMT male). HE and Periodic acid-Schiff (PAS) stain. 300X, except F (400X).

Prominent diffuse or segmentary mesangial expansion and, occasionally, areas of tuft-to-capsule adhesion were evident in the 12- and 18-month-old eSMT males. Severe injury was evidenced by many sclerotic glomeruli totally occluded (Fig 2 D) and a smaller number of atubular glomeruli with tuft adhesions in the older eSMT males (Fig 2 E). Prominent thickening of the peripheral glomerular basement membrane resulted in lobules with an abnormal architecture and a noticeable image of stiffness was observed in 60% of the eSMT rats and in 40% of the eSS after 21 months of age. Glomeruli with capsular drops or fibrin cap were not observed either in eSS or eSMT rats. A Kimmelstiel-Wilson-like nodular lesion was detected only in a 18-month-old eSMT male (Fig 2 F). Sporadically, periglomerular fibrosis was observed in eSMT and eSS females.

Fig. 3. (A) Intense lymphocytic infiltrate in renal pelvis (18-month-old eSMT female) (40X); (B) Mild tubular lesions (12-month-old eSS female) (300X); (C) Moderate tubular lesions characterized by disgregation of epithelial cells (6-month-old eSMT male) (300X); (D) Group of mildly dilated tubules with lumina containing protein casts (12-month-old eSS male) (300X). (E) Panoramic view of the renal medulla showing different tubular alterations, marked dilatation with protein content (15-month-old eSMT male) (100X); (F) Area of subcapsular retraction due to fibrosis. Parenchymal atrophy and lymphocytic infiltrates (24-month-old eSS male) (40X). HE and Periodic acid-Schiff (PAS) stain.

Areas of arteriolar hyalinosis and thickening of smooth muscle were observed in eSMT males from 9 months, in eSS males and eSMT females from 12 months and in eSS females from 21 month-old.

A few focal interstitial and pyelic mononuclear cell aggregates (Fig 3 A), concomitant with a mild tubular disruption, were apparent in 6-month-old eSS and eSMT males (Fig 3 B and 3 C). Groups of atrophic tubules with their lumina filled by hyaline casts were already noticeable in some 6-month-old eSS and eSMT males, progressing with ageing (Fig 3 D and 3 E). Prominent interstitial fibrosis and tubular atrophy and dilatation with vacuolization and desquamation of tubular cells, were observed in 100% of eSS and eSMT males at 18-24 months of age and older; these features were also detected in some 12-month-old eSS and eSMT females, affecting 80% of 24-month-old individuals. In older animals, subcapsular chronic inflammatory infiltrates, tubular atrophy and fibrosis lead to navel-like retractions of the kidney surface (Fig 3 F).

Renal damage, characterized by diffuse glomerulosclerosis and conspicuous tubular nephrosis, was registered in older eSMT and eSS males. Young eSS females showed the less severe kidney injury.

No changes were observed in glomerular, tubular or interstitial structures of Wistar controls.

4. Discussion and conclusions

Most animal models develop mild renal lesions in early phase diabetes, but not advanced lesions in late phase and even in the best diabetic models, aging is a critical feature in the development of lesions typical of the human DN (Nakagawa, 2009). The relative long-life-span of the eSS and eSMT rats, exceeding their metabolic disturbances, allowed the development of many of the characteristic lesions of diabetic nephropathy. This way, diabetic nephropathy worsened in both lines and sexes during the second year of life. Notwithstanding, results registered in 21-month-old control Wistar males demonstrated that ageing alone was unable to induce either a diabetic syndrome or diabetic renal lesions. It would also be argued that a particularly inherited liability of eSMT line might be an important single factor leading to a more severe renal injury. If so, findings point out that diabetic nephropathy developed as the result of complex interactions between the background genome response, body weight, sex and long-standing metabolic disorders.

Obesity is a very important factor in the beginning and rate of progression of metabolic disturbances and renal chronic complications of type 2 diabetes mellitus, in humans and in experimental models (Price *et al.*, 2002; Nangaku *et al.*, 2005). Even though eSMT strain is not considered an obese model, these rats are more robust than eSS and this relative overweight has been considered a key factor in the generation of their earlier and more severe hyperglycaemia, hyperlipidaemia and hyperinsulinaemia (Tarrés *et al.*, 2000; Montenegro *et al.*, 2005; Picena *et al.*, 2007). It is important to remark that eSS females, with the lowest body weight, had in turn the minor phenotypic diabetic expression. eSMT females, heavier than eSS females, displayed a more severe diabetic status and showed some glomerular alterations, as reduction in glomerulus's diameter, similar to those found in eSS males.

The degree and duration of hyperglycaemia constitute relevant risk factors regarding the occurrence and severity of renal impairment in people with both Type 1 (Alaveras *et al.*, 1997) and Type 2 Diabetes Mellitus (Ohkubo *et al.*, 1995) as well as in induced diabetic

animal models (Kern & Engelman, 1990; Petersen *et al.*, 1988). This way, the greater glucose metabolic disorder in eSS and eSMT males was clearly part of their more severe renal lesions. Several studies have pointed toward an association between sex hormones and the risk of Type 2 Diabetes Mellitus in men and women (Ding *et al.*, 2006). A study in the OLETF rats provided a strong support for this idea by showing that the incidence of Type 2 Diabetes Mellitus was 100% in males and 0% in females (Shi *et al.*, 1994). The higher incidence and severity of nephropathy in Type 2 diabetic male patients and the opposite in women indicate that sex hormones impact in the development and progression of diabetic renal disease by mechanisms yet poorly understood (Maric, 2009). The influence of sex hormones over the eSS diabetic syndrome was studied in rats gonadectomized at 90 days of age (Tarrés *et al.*, 1997). Spayed animals showed higher body weight, early and impaired intolerance to glucose and a decreased number of large pancreatic islets. Despite their lower biomass, castrated males did not evidence impairment in their intolerance to glucose, changes in insulinemia or remarkable modifications in endocrine pancreas histology. Further studies are needed for elucidating the complex interactions between genotype, body weight and sex in the onset of diabetes and the severity of chronic complications in the eSS and eSMT lines.

Transforming growth factor-beta 1 (TGF-beta1) appears to be an important downstream mediator for the development of renal hypertrophy and the accumulation of mesangial extracellular matrix components (Ziyadeh & Wolf, 2008). In the OLETF rats, plasma TGF-beta 1 and vascular TGF-beta type II receptors exist to a greater extent in pre- and early stages of diabetes mellitus when compared with non diabetic rats (Hosomi *et al.*, 2002). Streptozotocin-induced type 2 diabetic rats fed a high fat diet developed a higher proportion of sclerosed glomeruli than type 1 diabetic rats and the expression of CTGF and TGF-beta was significantly increased, despite their lower blood glucose levels and proteinuria (Danda *et al.*, 2005). The connective tissue growth factor (CTGF) is also overexpressed and involving in the pathogenesis of diabetic nephropathy; results from type 2 diabetic ob/ob mice demonstrated that albuminuria strongly correlated with urinary CTGF excretion (Roestenberg *et al.*, 2006). The activated renin-angiotensin system (RAS) has been implicated in the acceleration of diabetic renal disease in type 2 diabetic OLETF rats, as indicated by the long-lasting renoprotective effects of temporary angiotensin II blockage during the prediabetic stage, independent of its effects on glucose metabolism (Nagai *et al.*, 2005). The observation that in the ZDF obese rats reactive oxygen species-associated angiotensinogen enhancement plays an important role in renal damage suggests that reactive oxygen species are partly involved in intrarenal angiotensinogen augmentation, leading to the development of diabetic nephropathy (Kobori *et al.*, 2007).

In summary, mechanisms underlying the development of diabetic nephropathy are extremely complex persisting unclear the nature of the relationship between diabetes and late-onset renal disease. Remain to be clarified whether genetic and environmental factors determining early stages of these disorders are independent from those controlling their progression (Brosius *et al.* 2009; Nobrega, 2009).

The morphological changes in glomeruli of both strains were characterized by basement membrane thickening and diffuse mesangial expansion and, particularly in older eSMT males, by a diffuse or segmental glomerular sclerosis, being exceptional the nodular sclerosis of Kimmelstiel-Wilson. Moreover, it also shows that, even when the principal lesions affected the glomeruli according to the severity of glomerulopathy, the disease extended to the tubulointerstitial compartment. That morphological pattern has been

observed in Type 1 diabetic patients (Najafian *et al.*, 2003) and in poorly controlled long-term Type 2 Diabetes Mellitus, ascribed in the Category II of Fioretto *et al.*'s classification (2008). However, they also suffered from retinopathy, a nonexistent complication in eSS and eSMT rats.

In congruence with Tervaert *et al.* (2010), most of eSS and eSMT animals less than 12-month-old could be classified in Class I (barely evident mesangial lesions) whilst rats exceeding 12 months of age could belong to Class II (diffuse mesangial glomerulosclerosis with moderate to severe expansion of mesangium). It is noteworthy to outline that although the naturally slower progress of the metabolic derangement in eSS males with respect to the eSMT, almost all males over 18 months of age in both lines could be classified as Class IV (advanced glomerulosclerosis). Even though, lesions compatible with Class III (nodular sclerosis or presence of one Kimmelstiel-Wilson nodule at least) were detected in only one 18-month-old eSMT male rat. This single finding makes it exceptional and fits with other spontaneous diabetic murine models (Yuzawa *et al.*, 2008).

It has been reported that nodular glomerulosclerosis is present in diabetic patients with more progressed clinical and pathological features (Jang & Park, 2009). Nevertheless, other authors have sustained that diffuse and nodular sclerosis could be two discrete patterns of glomerulopathy caused by different pathophysiological mechanisms as Kimmelstiel-Wilson lesion correlates with diabetic retinopathy but most of Type 2 diabetic patients without retinopathy develop diffuse mesangial sclerosis (Schwartz *et al.*, 1998). Interestingly, no lesions of retinopathy have ever been detected in either eSS or in eSMT rats (Tarrés *et al.*, 1990; Tarrés *et al.*, 2000; Picena *et al.*, 2005). Moreover, although retinopathy and microalbuminuria are positively associated in Type 1 diabetic patients, the substantial discordances between glomerulopathy and retinopathy reported in Type 2 Diabetes Mellitus (Kanauchi *et al.*, 1998; Boelter *et al.*, 2006) highlights that kidney and retina are distinct organs with diverse underlying pathophysiologic mechanisms (Blum *et al.*, 2011). Further studies on the eSS and eSMT lines rats could be useful for studying the aforesaid differences.

Renal enlargement, particularly perceptible in eSMT males via the increase in kidney weights and diffuse mesangial expansion, evokes early stages of diabetic nephropathy. Notwithstanding, the areas of tubular atrophy detected in some eSS and particularly in the eSMT males at ~6 month, suggest a more advanced phase of diabetic nephropathy. Moreover, progression to macroalbuminuria and uremia already demonstrated in eSS male rats from 6 to 12 months of age in eSS males (Daniele *et al.*, 2000), may be correlated with the increasing loss of glomerular filtration rate accompanying each nephron sclerosis during the progression of diabetic nephropathy. While hyperglycaemia-induced osmotic polyuria has been implied as a relevant role in the onset and evolution of tubulointerstitial injury by causing an early tubular cell injury in the dilated collecting ducts (Wang *et al.*, 2008), careful attention should be paid in the persistent glycosuria detected in eSS and eSMT rats (Martínez *et al.*, 1988, Montenegro *et al.*, 2005). Moreover, the differences found between diabetic rats and Wistar controls as well as between both diabetic lines in superficial and juxtamedullary glomeruli glucidic residues and in convoluted proximal tubules ones could be indicative of alterations in the filtration and in the intracellular tubular processes (Frontini *et al.*, 2008).

As suggested by Murata *et al.*, 2002 in the Otsuka Long-Evans Tokushima Fatty (OLEFT) rats, we guess that the phenomenon of apoptosis could be operating at glomerular level in eSS and eSMT males which displayed a noticeable and increasing number of smaller

glomeruli as they get older. Furthermore, apoptosis induced by chronic hyperglycaemia has been related by Picena *et al.*, (2007) with the dramatic reduction of the area of the islets of Langerhans described in older eSS males by Gomez Dumm *et al.*, (1989). In diabetic nephropathy, the degree of proteinuria correlates with the progression of glomerulosclerosis and tubulointerstitial fibrosis (Wolf & Ziyadeh, 2007). The early albuminuria from 24-hs collections, the loss of glomerular filtration rate in 6-month-old eSS males and the progression to macroalbuminuria and uremia from 6 to 12 months of age (Daniele *et al.*, 2000), could be due to an increasing number of glomeruli affected associated to the progressive worsening of glucose and lipid disorders, particularly the rising values of hyperglycaemia, triglycerides, non-esterified fatty acids, total cholesterol, and low-density lipoprotein cholesterol with high- density lipoprotein-cholesterol reduced, demonstrated in this strain at 12 month of age (Daniele *et al.*, 2010). Previous studies in humans and animals have also suggested that altered lipid metabolism causes glomerular injury and promotes deterioration of glomerular function (Diamond & Karnovsky, 1987; Kasiske *et al.*, 1988; Maric *et al.*, 2004).

In humans, oxidative stress is linked to multiple chronic diabetic complications (Giacco & Brownlee, 2010). It is increasingly evident that changes in cellular function resulting in oxidative stress play a key role in the development and progression of diabetic nephropathy (Forbes *et al.*, 2008). Moreover, oxidized low density lipoprotein particles and non-esterified fatty acids could damage the mesangial cells and the tubulointerstitial tissue through different pathogenetic mechanisms (Nosadini & Tonolo, 2011). Interestingly, in 12 month-old eSS rats increased basal glycaemia and fructosamine values correlate with those of lipid peroxidation substances and inversely with total antioxidant capacity (Daniele *et al.*, 2007).

In agreement with the data in diabetic patients and in other experimental diabetic models, and taking into account that hyperlipidaemia and oxidative stress have been syndicated as having major roles in the pathogenesis of the diabetic nephropathy (Kume *et al.*, 2008), we suggest that early hyperinsulinemia and lipid disorders, already demonstrated in eSS rats (Montanaro *et al.*, 2003. Daniele *et al.*, 2010) and in eSMT rats (Tarrés MC *et al.*, 2000) as well as the oxidative stress previously verified in eSS male rats (Daniele *et al.*, 2007), are key players in the development of the diabetic nephropathy in eSS and eSMT rats. Though new experiments could be undertaken in eSS and eSMT rats to brighten the genesis and evolution of their nephropathy, we conclude that these two spontaneously diabetic rats closely resemble the advanced human diabetic nephropathy and, simultaneously, have opened valuable opportunities for enhance our understanding about the interactions between glucose and lipid disorders and the pathways towards diabetic nephropathy, a relevant cause of morbidity and mortality in the diabetic population and the leading cause of end-stage renal failure in the Western World.

5. Acknowledgments

The authors are grateful to the School of Medicine, National University of Rosario, for financial support and to Wiener Laboratories for the their invaluable contribution of commercial kits

6. References

- Alaveras, AE; Thomas, SM; Sagriotis, A *et al.* (1997). Diabetic Nephropathy: Diagnosis, Prevention, and Treatment: Evaluation of Patients With Diabetic Nephropathy. *Nephrol Dial transplant* 12, Suppl. 2, 71-74, ISSN 0931-0509

- American Diabetic Association (ADA) (2009). Diagnosis and Classification of Diabetes Mellitus. *Diabetes Care* 32, Suppl.1, S62-S67, ISSN 0149-5992
- Blum, M; Saemann, A & Wolf, G (2011). The eye, the kidney and microcirculation Editorial Comments. *Nephrol Dial Transplant* 26, 1, 4-6, ISSN 0931-0509
- Boelter, MC; Gross; JL; Canani LH *et al.* (2006). Proliferative diabetic nephropathy with microalbuminuria in patients with type 2 diabetes. *Braz J Med Biol Res* 39, 8, 1033-1039, ISSN 0100-879X
- Breyer, MD; Bottinger, EP; Brosius, FC *et al.* (2005). Mouse models of diabetic nephropathy. *J Am Soc Nephrol* 16, 1, 27-45, ISSN 1046-6673
- Brosius, FC; Alpers, CE; Bottinger, EP *et al.* (2009) Mouse models of diabetic nephropathy. *J Am Soc Nephrol* 20, 12, 2503-2512, ISSN 1046-6673
- Caballero, AE & Tenzer, P (2007). Building cultural competency for improved diabetes care: Latino Americans and diabetes. *J Fam Pract* 56, 9, S29-S37, ISSN 0094-3509
- Calderari, SA; Gonzalez, AC & Gayol, MD (1987). Spontaneous hypertriglyceridemic obesity and hyperglycaemia in an inbred line of rats. *Int J Obes* 11, 6, 571-579, ISSN 0307-0565
- Calderari, SA; Font, MT; Garroq, O *et al.* (1991). The inbred IIM/Fm stock. *Rat News Let* 25, 1, 28-29. ISSN 0309-1848
- Chatzigeorgiou, A; Halapas, A; Kalafatakis, K *et al.* (2009). The use of animal models in the study of diabetes mellitus. *In Vivo* 23, 2, 245-258, ISSN 0258-851X
- Danda, RS; Habiba, NM; Rincon-Choles, H; Bhandari, BK *et al.* (2005). Kidney involvement in a nongenetic rat model of type 2 diabetes. *Kidney International* 68, 6, 2562-2571 ISSN 0085-2538
- Daniele, S; Arriaga, S; Martínez, SM *et al.* (2000). Onset and evolution of nephropathy in rats with spontaneous diabetes mellitus. *J Physiol Biochem* 56, 1, 45-54, ISSN 1138-7548
- Daniele, SM; Picena, JC; Montenegro, SM *et al.* (2007). Hígado graso no alcohólico en ratas machos de una línea con diabetes genética. *Electron J Biomed* 7, 3, 29-39, ISSN 1697-090X
- Daniele, SM; Montenegro, SM; Tarrés, MC *et al.* (2010). The eSS rat, a nonobese model of disordered glucose and lipid metabolism and fatty liver. *Diabetol Metabol Syndr* 2, 3, 1-5, ISSN 1758-5996
- Diamond, J & Karnovsky, M. (1987) Exacerbation of chronic aminonucleoside nephrosis by dietary cholesterol supplementation. *Kidney Int* 32, 5, 671-678, ISSN 0085-2538
- Ding, EL; Song, Y; Malik, VS *et al.* (2006). Sex differences of endogenous sex hormones and risk of type 2 diabetes: a systematic review and metaanalysis. *JAMA* 295, 11, 1288-1299, ISSN 0098748
- Dronavalli, S & Barkis GL (2008). Mechanistic Insights into Diuretic-Induced Insulin Resistance. *Hypertension* 52, 6, 1009-1011, ISSN 0194911X
- Fioretto, P; Caramori, ML & Mauer, M (2008). The kidney in diabetes: dynamic pathways of injury and repair. The Camillo Golgi Lecture. *Diabetologia* 51, 8, 1347-1355, ISSN 0012-186X
- Forbes, JM; Coughlan, MT & Cooper, ME (2008). Oxidative stress as a major culprit in kidney disease in diabetes. *Diabetes* 57, 6, 1446-1154, ISSN 0012-1797

- Frontini, AV; Hisano, N & D'Ottavio, AE (2008). La densidad óptica aplicada al estudio lectinohistoquímico de radicales glucídicos néfricos en ratas diabéticas tipo 2. *Electron J Biomed* 6, 1, 36-44, ISSN 1697-090X
- Giacco, F & Brownlee, M (2010). Oxidative stress and diabetic complications. *Circ Res* 29, 107, 1058-1070, ISSN 0009-7330
- Gomez Dumm, CL; Semino, MC & Gagliardino, JJ (1990). Sequential morphological changes in pancreatic islets of spontaneously diabetic rats. *Pancreas* 5, 5, 533-539, ISSN 0885-3177
- Hisano, N; D'Ottavio, AE & Calderari, SA (1994). Female reproductive profile in a fertile genetically obese line of rats. *J Exp Zool* 270, 5, 486-490, ISSN 1932-5223
- Hosomi, N; Noma, T; Ohyama H & *et al.* (2002). Vascular proliferation and transforming growth factor-beta expression in pre- and early stage of diabetes mellitus in Otsuka Long-Evans Tokushima fatty rats. *Atherosclerosis* 162, 1, 69-76 ISSN 0021-9150
- Hudkins, KL; Pichaiwong, W, *et al.* (2010). Ob/Ob mutant mice model progressive diabetic nephropathy. *J Am Soc Nephrol* 21, 9, 1533-1542, ISSN 1046-6673
- Jang, SH & Park, MH (2009). The morphologic patterns of diabetic nephropathy in Koreans. *Korean J Pathol* 43, 1, 36-42, ISSN 1738-1843
- Janssen, U; Phillips, AO & Floege, J (1999). Rodent models of nephropathy associated with type II diabetes. *J Nephrol* 12, 3, 159-172, ISSN 1121-8428
- Kanauchi, M; Kawano, T; Uyama H *et al.* (1998). Discordance between retinopathy and nephropathy in type 2 diabetes. *Nephron* 80, 2, 171-174, ISSN 0028-2766
- Kasiske, B; Cleary, M; O'Donnell, M *et al.* (1988). Mechanisms of glomerular injury in the obese Zucker rat. *Kidney Int* 33, 3, 667-672, ISSN 0085-2538
- Kern, TS & Engelman, RL (1990). Arrest of glomerulopathy in diabetic dogs by improved glycaemic control. *Diabetologia* 33, 522-525, ISSN: 0012-186X
- Kobori, H; Ozawa, Y; Suzaki, Y; Prieto-Carrasquero, MC (2006). Intratubular angiotensinogen in hypertension and kidney Diseases. *Am J Hypertens* 19, 5, 541-550 ISSN 0895-7061
- Kume, S; Uzu, T; Isshiki TK *et al* (2008). Peroxisome proliferator-activated receptors in diabetic nephropathy (Review Article). *PPAR Res* 4, 7, 1-11, ISSN 1687- 4765
- Lee, G (2003). End - stage renal disease in the Asian - Pacific region *Semin Nephrol* 23, 1, 107-114, ISSN 0270-9295
- Maric, C; Sandberg, K & Hinojosa-Laborde, C (2004). Glomerulosclerosis and tubulointerstitial fibrosis are attenuated with 17- β estradiol in the aging Dahl salt-sensitive rat. *J Am Soc Nephrol* 15, 6, 1546-1556, ISSN 1046-6673
- Maric, C (2009). Sex, diabetes and the kidneys. *Am J Physiol-Renal Physiol* 296, 4, F680-F688, ISSN 0363-6127
- Martínez, SM; Tarrés, MC; Montenegro, SM *et al* (1988). Spontaneous diabetes in eSS rats. *Acta diabetol lat* 25, 4, 303-313, ISSN 0001-5563
- Martínez, SM; Tarrés, MC; Picena, JC *et al* (1993). eSS rat, an animal model for the study of spontaneous non-insulin-dependent diabetes, In: *Lessons from Animal Diabetes IV*. E Shafrir, Ed. Smith-Gordon, 75-90, ISBN 1854630075, London, UK

- Montanaro, M; Rimoldi, OJ; Igal, RA *et al.* (2003). Hepatic $\Delta 9$, $\Delta 6$ and $\Delta 5$ Desaturations in the Non-Insulin Dependent Diabetic Mellitus eSS Rats. *Lipids* 38, 8, 827-832, ISSN 0024-4201
- Montenegro, SM; Tarrés, MC; Picena, JC *et al.* (2005). Conducta alimentaria y perfil glucémico en dos modelos murinos de diabetes genética: eSS y eSMT. *Biomédica* 25, 4, 441-450, ISSN 0120-4157
- Murata, I; Takemura, G; Asano, A *et al.* (2002). Apoptotic cell loss following cell proliferation in renal glomeruli of Otsuka Long-Evans Tokushima Fatty rats, a model of type 2 diabetes. *Am J Nephrol* 22, 5-6, 587-595, ISSN 0250-8095
- Nagai, Y; Yao, L; Kobori, H *et al.*, (2005). Temporary Angiotensin II blockade at the prediabetic stage attenuates the development of renal injury in Type 2 diabetic rats. *J Am Soc Nephrol* 16, 3, 703-711 ISSN 1046-6673
- Najafian, B; Kim, Y; Crosson, JT *et al.* (2003). Atubular glomeruli and glomerulotubular junction abnormalities in diabetic nephropathy. *J Am Soc Nephrol* 14, 4, 908-917, ISSN 1046-6673
- Nakagawa, T (2009). A new mouse model resembling human diabetic nephropathy: uncoupling of VEGF with eNOS as a novel pathogenic mechanism. *Clin Nephrol* 71, 2, 103-9, ISSN 0301-0430
- Nangaku, M; Izuhara, Y; Usuda, N *et al.* (2005). In a type 2 diabetic nephropathy rat model, the improvement of obesity by a low calorie diet reduces oxidative/carbonyl stress and prevents diabetic nephropathy. *Nephrol Dial. Transplant* 20, 12, 2661-2669, ISSN 0931- 0509
- Nobrega, MA; Solberg Woods, LS; Fleming, S; Jacob, HJ (2009). Distinct genetic regulation of progression of diabetes and renal disease in the Goto-Kakizaki rat. *Physiol Genomics* 39: 38-46 ISSN 1531-2267
- Nosadini, R & Tonolo, G. (2011). Role of oxidized low density lipoproteins and free fatty acids in the pathogenesis of glomerulopathy and tubulointerstitial lesions in type 2 diabetes. *Nutr Metab Cardiovasc Dis* 21, 2, 79-85, ISSN 0939-4753
- Ohkubo, Y; Kishikawa, H; Araki, E *et al.* (1995). Intensive insulin therapy prevents the progression of diabetic microvascular complications in Japanese patients with non-insulin-dependent diabetes mellitus: a randomized prospective 6-year study. *Diabetes Res Clin Pract* 28, 2, 103-117, ISSN 0168-8227
- Petersen, J; Ross, J & Rabkin, R (1988). Effect of insulin therapy on established diabetic nephropathy in rats. *Diabetes* 37, 10, 1346-1350, ISSN 0012-1797
- Picena, JC; Daniele, SM; Tarrés, MC *et al.* (2005). An animal model fits for studying divergences among diabetic microvascular complications. *Electron J Biomed* 3, 2, 39-43, ISSN 1697-090X
- Picena, JC; Montenegro, SM; Tarrés, MC *et al.* (2007). Modificaciones dinámicas en los islotes de Langerhans de dos líneas de ratas espontáneamente diabéticas: eSS y eSMT. *Medicina (B Aires)* 67, 4, 331-340, ISSN 0025-7680
- Price, DA; Lansang, MC; Osey, SY *et al.* (2002). Type 2 diabetes, obesity and the renal response to blocking the renin system with irbesartan. *Diabetic Med* 19, 10, 858-861, ISSN 0742-3071

- Roestenberg, P; van Nieuwenhoven, FA; Joles, JA; Trischberger, C *et al*, (2006). Temporal expression profile and distribution pattern indicate a role of connective tissue growth factor (CTGF/CCN-2) in diabetic nephropathy in mice. *Am J Physiol Renal Physiol* 290, 6, F1344-1354 ISSN 1522-1466
- Schwartz, MM; Lewis, EJ; Leonard-Martin, T *et al*. (1998). Renal pathology patterns in type II diabetes mellitus: relationship with retinopathy. The Collaborative Study Group. *Nephrol Dial. Transplant.* 13, 10, 2547-2552, ISSN 0931- 0509
- Shi, K; Mizuno, A; Sano, T *et al*. (1994). Sexual difference in the incidence of diabetes mellitus in Otsuka-Long-Evans-Tokushima-Fatty rats: effects of castration and sex hormone replacement on its incidence. *Metabolism* 43, 10, 1214-1220, ISSN 0026-0495
- SPSS (2006). IBM SPSS Statistics, 15.0.1 version, SPSS Inc, New York, USA
- Srinivasan, K & Ramarao, P (2007). Animal models in type 2 diabetes research: an overview. *Indian J Med Res* 125, 3, 451-472, ISSN 0971-5916
- Stewart, JH; McCredie, MR; Williams, SM *et al*. (2006). The enigma of hypertensive ESRD: observations on incidence and trends in 18 European, Canadian, and Asian-Pacific populations, 1998 to 2002. *Am J Kidney Dis* 48, 2, 183-191, ISSN 0272-6386
- Tarrés, MC; Martínez, SM; Montenegro, SM *et al*. (1990). Relación entre dieta, biomasa y expresión de la diabetes en la rata eSS. *Medicina (Buenos Aires)* 50, 3, 235-243, ISSN 0025-7680
- Tarrés, MC; Martínez, SM; Montenegro, SM *et al*. (1992). The eSS rat: A new model of non-insulin-dependent human diabetes. *Am J Pathol* 141, 3, 761-763, ISSN 0002-9440
- Tarrés, MC; Martínez, SM; Montenegro, SM *et al*. (1997). Influence of gonadectomy in eSS diabetic rats. *J Physiol Biochem* 53, 2, 211-216, ISSN 1138-7548
- Tarrés, MC; Montenegro, S; Martínez, SM *et al*. (2000). The eSMT rat: a murine model of type 2 human diabetes. Chapter: Animal models in the study of human diseases, In: *Animal Research and Welfare. A Partnership. Proceedings of the International Joint Meeting between 12th ICLAS General Assembly & Conference and the 7th Felasa Symposium*. Laboratory Animals Ltd., 93-94, ISBN: 0-901334-13-8. London, UK
- Tervaert, TW; Mooyaart, AL; Amann, K *et al*. (2010) Pathologic classification of diabetic nephropathy. *J Am Soc Nephrol* 21, 4, 556-563, ISSN 1046-6673
- Tesch, GH & Allen, TJ (2007). Rodent models of streptozotocin-induced diabetic nephropathy. *Nephrology (Carlton)* 12, 3, 261-266, ISSN1320-5358
- U.S Renal Data System (USRDS) (2008). Annual Data Report: Atlas of Chronic Kidney Disease and End-Stage Renal Disease in the United States, National Institutes of Health, National Institute of Diabetes and Digestive and Kidney Diseases. Bethesda, MD. 2, Précis, 29-33
- U.S Renal Data System (USRDS) (2010). Annual Data Report: Atlas of Chronic Kidney Disease and End-Stage Renal Disease in the United States, National Institutes of Health, National Institute of Diabetes and Digestive and Kidney Diseases, Bethesda, MD. 2, Précis, 209-222
- Villar, E & Zaoui, P (2010). Diabetes and chronic kidney disease: Lessons from renal epidemiology. *Nephrol Ther* 6, 7, 585-590, ISSN 1769-7255
- Wang, S; Mitu, GM & Hirschberg, R (2008). Osmotic polyuria: an overlooked mechanism in diabetic nephropathy. *Nephrol Dial Transplant* 23, 7, 2167-2172, ISSN 0931- 0509

- Wolf, G & Ziyadeh, FN (2007). Cellular and molecular mechanisms of proteinuria in diabetic nephropathy. *Nephron Physiol* 106, 2, 26-31, ISSN 1660-2137
- Yuzawa, Y; Niki, I, Kosugi, T *et al.* (2008). Overexpression of Calmodulin in Pancreatic β Cells Induces Diabetic Nephropathy. *J Am Soc Nephrol* 19, 9. 1701-1711, ISSN 1046-6673
- Ziyadeh, FN & Wolf, G (2008). Pathogenesis of the podocytopathy and proteinuria in diabetic glomerulopathy. *Curr Diabetes Rev* 4, 1, 39-45 ISSN 1573-3998
- Zimmet, P; Alberti, KGM & Shaw, J (2001). Global and societal implications of the diabetic epidemic. *Nature* 414, 12, 782-787, ISSN 0028-0836

Recent Advances in the Pathogenesis, Prevention and Management of Type 2 Diabetes and its Complications

Edited by Prof. Mark Zimring

ISBN 978-953-307-597-6

Hard cover, 442 pages

Publisher InTech

Published online 29, August, 2011

Published in print edition August, 2011

Type 2 diabetes “œmellitus” affects nearly 120 million persons worldwide- and according to the World Health Organization this number is expected to double by the year 2030. Owing to a rapidly increasing disease prevalence, the medical, social and economic burdens associated with the microvascular and macrovascular complications of type 2 diabetes are likely to increase dramatically in the coming decades. In this volume, leading contributors to the field review the pathogenesis, treatment and management of type 2 diabetes and its complications. They provide invaluable insight and share their discoveries about potentially important new techniques for the diagnosis, treatment and prevention of diabetic complications.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Juan Carlos Picena, Silvana Marisa Montenegro, Alberto Enrique D’Ottavio, María Cristina Tarrés and Stella Maris Martínez (2011). A Diachronic Study of Diabetic Nephropathy in Two Autochthonous Lines of Rats to Understand Diabetic Chronic Complications, *Recent Advances in the Pathogenesis, Prevention and Management of Type 2 Diabetes and its Complications*, Prof. Mark Zimring (Ed.), ISBN: 978-953-307-597-6, InTech, Available from: <http://www.intechopen.com/books/recent-advances-in-the-pathogenesis-prevention-and-management-of-type-2-diabetes-and-its-complications/a-diachronic-study-of-diabetic-nephropathy-in-two-autochthonous-lines-of-rats-to-understand-diabetic>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2011 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike-3.0 License](https://creativecommons.org/licenses/by-nc-sa/3.0/), which permits use, distribution and reproduction for non-commercial purposes, provided the original is properly cited and derivative works building on this content are distributed under the same license.

IntechOpen

IntechOpen