We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Cardiovascular Diseases in Kidney Transplantation

Roberto Marcén and Sara Jiménez

*Department of Nephrology. Ramón y Cajal Hospital,
University of Alcalá de Henares, Madrid
Spain*

1. Introduction

Cardiovascular diseases that include atherosclerotic diseases; coronary artery disease, cerebrovascular disease and peripheral vascular disease, and cardiac functional diseases; congestive heart failure, left ventricular hypertrophy and arrhythmias, are very common in the general population and are the first cause of mortality (Wilson & Culleton, 1998, Culleton & Wilson, 1998). Moreover, patients with chronic renal failure have an increased risk of cardiovascular disease compared with the general population, and after stratification by age and gender, cardiovascular mortality is 10 to 20 times more frequent independently of treatment; predialysis, dialysis or after kidney transplantation (Foley et al, 1998). Kidney transplantation is the best therapy of end-stage renal disease by reducing cardiovascular mortality (McDonald & Russ, 2002, Ojo et al, 2000, Wolfe et al, 1999). But even after transplantation, a recipient of 25 to 35 years of age has a 10 times higher risk of cardiovascular mortality than an individual of similar sex and gender without renal failure (Foley et al, 1998). Among the cardiovascular diseases, atherosclerotic diseases are the most frequently studied and are associated with patient outcome. Ischaemic heart disease was the cause of 53% of total mortality in a study performed in Scandinavia 15 years ago (Lindholm et al, 1995) and these findings were still prevalent in a later report (Aakhus et al, 2004). However, several authors have recently emphasized the importance of functional cardiopathies such as congestive heart failure on patient outcome (Rigatto et al, 2002).

The high incidence of cardiovascular events and mortality in renal transplant recipients has been attributed to the increased presence of traditional (Ojo, 2006) and nontraditional risk factors (Ducloux et al, 2004, De Mattos et al, 2006). As traditional risk factors do not fully explain the high cardiovascular risk it has been postulated that some of these risk factors, age, diabetes and smoking, could have a higher deleterious impact in transplant recipients than in the general population (Kasiske et al, 2000a). Other authors consider that transplant related (De Mattos et al, 2006) and nontraditional or emergent factors, hyperhomocysteinemia and inflammation, could play a predominant role in the appearance of cardiovascular events (Ducloux et al, 2004). Progressive chronic graft dysfunction and death with functioning graft are the most important causes of graft loss (Matas et al, 2002, Collins et al, 2008) and cardiovascular diseases are the leading cause of mortality in renal transplant recipients dying with a functioning graft (Pilmore et al, 2010). Consequently decreasing

cardiovascular mortality could improve patient and graft outcomes (Howard et al, 2002, Marcén et al, 2001, Morales, 2008, Vanrenterghem et al, 2008).

The present chapter reviews the cardiovascular diseases which affect renal transplant recipients and their impact on patient mortality, the risk factors associated with these complications and the therapeutical strategies to improve patient and graft outcomes.

2. Cardiovascular diseases

Renal transplant recipients are not healthy individuals. They have a past history of chronic renal failure and dialysis therapy, both having negative impact on cardiovascular risk and they present variable chronic renal failure stages. All cardiovascular diseases; atherosclerotic and functional cardiopathies can affect transplant recipients (Table 1).

Atherosclerotic diseases	Functional cardiopathies
Coronary artery diseases	Congestive heart failure
Cerebrovascular diseases	Left ventricular hypertrophy
Peripheral vascular diseases	Arrhythmias

Table 1. Most common cardiovascular diseases affecting transplant recipients

As in the general population, cardiovascular diseases are important causes of morbidity and mortality in renal transplant recipients. Heart diseases and cerebrovascular diseases accounted for about 35% mortality in our Unit (Figure 1).

Fig. 1. Causes of mortality in 224 renal transplant recipients with functioning graft (Ramón y Cajal Hospital)

2.1 Atherosclerotic diseases

2.1.1 Coronary artery disease

Coronary artery disease is the most frequent atherosclerotic disease in the general population and a leading cause of cardiovascular mortality. It is generally admitted that coronary artery disease is between 2 and 6 times more frequent in renal transplant recipients than in the general population (Aarkhus et al, 1999, Kasiske, 1988, Lentine et al, 2005a, Marcén et al, 2006, Massy, 2001). A very high number of events occurring during the first weeks after transplantation have been attributed to surgical stress, immunosuppression and silent disease while the patient was on dialysis (Kasiske et al, 2000, 2005, Lentine et al, 2005, Marcén et al, 2006). In a retrospective study from the USA, the prevalence of coronary artery disease, defined as acute myocardial infarction, coronary revascularization procedures, or death due to coronary disease, was 23% at 15 years of follow-up and it was the cause of 18.7% of mortality (Kasiske et al, 1996). In a multicenter, retrospective study performed in Spain, the prevalence of coronary artery disease was 6.8% at 5 years (Marcén et al, 2006) and very similar findings have been reported from France (Doucloux et al, 2004), two low risk countries. Data from the United Network for Organ Sharing (UNOS) and United States Renal Data System (USRDS) registry showed that the prevalence of acute myocardial infarction was between 6% to 11% at 3 years a 17% lower than that observed on waiting list patients (Kasiske et al, 2006, Lentine et al, 2005). Moreover, myocardial infarction has a poor prognosis and is the cause of high mortality (Herzog et al, 1998, 2000, Morales, 2008).

The diagnostic criteria and therapeutical measures must be those used in the general population and in patients with chronic renal failure (Murphy et al, 1998). As a high percentage of events occur in the first weeks or months after the procedure, efforts have to be made to prevent and treat coronary artery disease before transplantation. Several algorithms have been proposed in which patients are classified according to cardiovascular risk (Lentine et al, 2008b, Wang & Kasiske, 2010), but we have not yet the ideal diagnostic procedure for asymptomatic patients.

2.1.2 Cerebrovascular diseases

Cerebrovascular diseases include transient cerebral ischaemia, when focal neurologic symptoms resolve in 24 hours, and persistent neurological deficits documented by computed tomography or nuclear magnetic resonance. Patients with chronic renal disease have more severe atherosclerotic lesions in carotic arteries than the general population (Kennedy et al 2001). This could explain the 5 to 10 times higher risk of having ischaemic or haemorrhagic events when compared with the general population (Seliger et al, 2002, 2003a,b). In renal transplant recipients the annual incidence was from 0.5% to 2.3% (Abedini et al, 2009a, Aull-Watschinger et al, 2008, Cosio et al, 2005, Lentine et al, 2008a, Oliveras et al, 2003), and ischaemic events predominated in a proportion of 2-3:1 compared with haemorrhagic events (Aull-Watschinger et al, 2008, Oliveras et al, 2003). In a long-term study, by actuarial analysis, 15% of patients who survived with a functioning graft for 15 years experienced a major cerebrovascular event (Kasiske et al, 1996). The evolution of cerebrovascular diseases is poor and a mortality rate around of 50% three months after the event has been reported (Oliveras et al, 2003). In retrospective studies, cerebrovascular diseases were the cause of 5% to 8% of total mortality (Aull-Watschinger et al, 2008, Howard et al, 2002). Only the Assessment of Lescol in Renal Transplantation (ALERT) study has prospectively investigated these complications, the incidence was 8.8% during the 6.7 year

follow-up period, and 21% of the cerebrovascular events, 48% of haemorrhagic and 8% of ischaemic, were fatal and accounted for 9.9% of total mortality (Abedini et al, 2009a).

2.1.3 Peripheral vascular disease

The incidence and clinical outcome of this complication have been seldom studied in renal transplant recipients. The diagnostic criteria include: intermittent claudication, ulceration with long-vessel disease on flow studies, amputations, by-pass or percutaneous angioplasty. The frequency reported was variable. In an old study, in which only amputation or revascularization procedures were included, the cumulative incidence was 15% at 15 years (Kasiske et al, 1996). In other studies, the incidence increased with the length of follow-up, from 2.1% at 1 year to 5.9% at 10 years (Sung et al, 2000). Data from the UNOS registry have shown a cumulative incidence of 20% in diabetic and 5% in nondiabetic patients at 3 years (Snyder et al, 2006). As the disease develops along years it is difficult to distinguish risk factors due to transplantation from those present before. The need of amputation is low about 2 to 3% (Sung et al, 2000). The disease by itself is not a cause of mortality but patients suffering from it have an increased risk of death with a functioning graft (Snyder et al, 2006, Sung et al, 2000).

3. Functional heart diseases

3.1 Congestive heart failure

Congestive heart disease is defined as dysnea plus at least two of the following characteristics; increased yugular venous pressure, basal lung rales, lung hypertension in radiography or pulmonary edema (Harnett et al, 1995). In dialysis patients, congestive heart failure is 36 times more frequent than in the general population and it is a mortality risk factor (Collins 2002, Stack & Bloemberg, 2001). Congestive heart disease has been studied less than coronary artery disease in the renal transplant population and it is associated with coronary artery disease in 30% of cases (Rigatto, 2003a,b). Its annual incidence was 3-5 times that of the general population, reached a cumulative incidence of 18.3% at 3 years and was associated with poor graft function (Abbott et al, 2003b, Lentine et al, 2005, Rigatto, 2003a,b). It has a high impact on mortality, similar to that of coronary artery disease (Lentine et al, 2005).

3.2 Left ventricular hypertrophy

There are two types of left ventricular hypertrophy; concentric ventricular hypertrophy and dilatation with or without hypertrophy. The first one is associated with volume overload and the second with aortic insufficiency or severe anemia. Both types are more frequent in patients in renal failure than in the general population, reaching 20-50% in patients with chronic renal failure (Levin et al, 1996, Tucker et al, 1997) and up to 70% in those on dialysis (Foley et al, 1995, McGregor et al, 1998). Several prospective studies have shown that left ventricular hypertrophy improved during the first two years after transplantation but it was still present in about 40% of renal transplant recipients (Rigatto et al, 2000, Teruel et al, 1987). Factors related with no improvement were: age, left ventricular morphology, duration and severity of hypertension and time averaged pulse pressure (Rigatto et al, 2000). Moreover, renal transplant also improved ventricular function in most patients even in those with severe impairment (Parfrey et al 1995, Wali et al, 2005). However these findings have been recently questioned when cardiac structure was assessed by magnetic resonance

(Patel et al, 2008). Parameters of ventricular hypertrophy or impaired cardiac function were associated with increased risk of cardiovascular events and cardiovascular mortality in renal transplant recipients (Aull-Watschinger et al, 2008, McGregor et al, 2000). In a nonrandomized study, conversión from CNI to sirolimus regressed left ventricular mass thickness regardless of blood pressure changes, thus suggesting non-hemodynamic-effect mechanisms on the left ventricular mass (Paoletti et al, 2008).

3.3 Arrhythmias

Atrial fibrillation is the most common cardiac rythm disorder in the general population and in patients on dialysis (Harnett et al 1995, Zebe 2001). Data from renal transplant recipients, despite being a high risk population due to the pre-transplant history and the high prevalence of risk factors related to this complication such as hypertension and obesity, have only been recently reported. Registry studies from the USA have shown a cumulative prevalence around 7 % at 3 years (Abbott et al, 2003a, Lentine et al, 2006). Risk factors for postransplantation atrial fibrillation include older age, male gender, renal failure for hypertension, and coronary artery disease. As in the general population atrial fibrillation was associated with an increased cardiovascular mortality, up to 3 times higher than patients without the disease (Abbott et al, 2003, Lentine et al, 2006).

4. Cardiovascular risk factors

Three types of cardiovascular risk factors are generally identified in transplant recipients (Table 2). 1) Traditional risk factors are those which in the general population are associated with cardiovascular diseases, and their treatment decreases the incidence of these complications. They include; older age, hypertension, hypercholesterolemia, diabetes mellitus, tobacco smoking and obesity. Their incidence is mostly increased in advanced CKD stages (Ansell et al, 2007, Karthikeyan et al, 2004, Marcén et al, 2005). 2) Risk factors associated with the transplant; anaemia, graft dysfunction and related complications, proteinuria, and immunosuppression. Finally,3), non-traditional or emergent factors such as hyperhomocysteinemia and chronic inflammation.

Traditional risk factors	Transplant related factors	Nontraditional or emergent factors
Age Sex Hypertension Dislipidemia Diabetes Smoking Obesity	Anaemia Graft dysfunction Proteinuria Immunosuppression	Hyperhomocysteinemia Inflammation

Table 2. Cardiovascular risk factors

4.1 Traditional risk factors

4.1.1 Age and sex

Older age is a nonmodifiable cardiovascular risk factor in the general population. In renal transplant recipients it was associated with an increased risk for cardiovascular

atherosclerotic diseases; ischemic heart disease, cerebrovascular disease, and peripheral vascular disease (Kasiske et al, 1996, Kasiske et al, 2006, Marcén et al, 2006, Oliveras et al, 2003, Rigatto et al, 2002, Snyder et al, 2006), and also for functional heart diseases; congestive heart failure, left ventricular hypertrophy and arrhythmias (Abbott et al, 2003, Lentine et al 2006, Rigatto et al, 2000, Rigatto et al, 2002). Male gender is a risk factor for ischemic heart disease and peripheral vascular disease (Kasiske, 1988, Rigatto et al, 2002, Kasiske et al, 2006, Marcén et al, 2006, Snyder et al, 2006) and female gender for cerebrovascular disease and congestive heart failure (Abbott et al, 2003, Lentine et al, 2008a).

4.1.2 Hypertension

It is a common complication in renal transplant recipients. Its prevalence varies between 70 and 90%. There are several causes and mechanisms of high blood pressure and many patients have several of them. A previous history of hypertension, artery graft stenosis, the own recipient kidneys, overweight, chronic graft dysfunction and immunosuppressive agents such as calcineurin inhibitors (CNIs), cyclosporine and tacrolimus, are the conditions generally associated with post-transplant hypertension (Koomans & Ligteneberg, 2001, Zhang et al, 2003). Among CNIs, cyclosporine seems to increase blood pressure more than tacrolimus (Ligteneberg et al, 2001). It has been associated with an increased risk of ischemic heart disease, congestive heart failure, left ventricular hypertrophy (Rigatto 2002) and with mortality (Kasiske et al 2004, Fernández-Fresnedo et al, 2005).

One characteristic of post-transplant hypertension is the lack of control despite treatment. In a study of 1295 patients, only 12.4% had a normal blood pressure one year after grafting and more than 95% of them were on antihypertensive therapy (Kasiske et al, 2004). Others found in their series a higher number of patients with normal blood pressure without therapy (26.0%) but they also reported that 32.0% of patients had uncontrolled blood pressure while they were on treatment (Tutone et al, 2005). Cross-sectional studies have shown that between 60 to 100% of patients according to the stage of graft failure had a blood pressure above 130/80 mm Hg and most of them were on antihypertensive therapy (Karthikeyan et al, 2004, Marcén et al, 2009a).

There are not specific blood pressure levels for renal transplant recipients and the reference values are those of the general population. As the renal transplant recipients are considered a high risk population for cardiovascular diseases, a blood pressure of 130/80 mm Hg has been recommended. Treatment includes changing life style, reducing the diet sodium intake, physical activity, low consumption of alcohol and antihypertensive agents (Choubanian et al, 2003). There are no specific antihypertensive agents to treat post-transplant hypertension and all agents can be used. The prescription has to be done taking into account the characteristic of each patient (KDIGO, 2009, Park & Luan, 2005). Most patients need to be treated with several antihypertensive agents. Studies in which angiotensin converting enzyme inhibitors or angiotensin receptor blockers have been compared with other antihypertensive agents have shown controversial results, and there are no studies in which the superiority of one agent over the others on patient survival has been definitively established (Opelz et al, 2006, Hiremath et al, 2007). There are not randomized, prospective studies that have demonstrated the beneficial effects of controlling blood pressure in renal transplant recipients, but it has been assumed that they would be similar to those obtained in the general population. However, retrospective registry studies have shown that decreasing blood pressure even several years after hypertension appearance was associated with a better patient outcome (Opelz et al, 2005).

4.1.3 Dyslipidemia

Lipid disorders are more common in renal transplant recipients than in the general population and include high levels of cholesterol and triglycerides. Also frequent are high levels of LDL-cholesterol, lipoprotein (a) and apolipoprotein B, while HDL-cholesterol can be high, normal or low. Hypercholesterolemia, total serum cholesterol above 200 mg/dl and LDL-cholesterol above 100 mg/dl, have been observed in up to 90 % of patients (Aakhus et al, 1996, Hricik et al, 1994). Several factors have been associated with hyperlipidemia; genetic predisposition, body weight gain, graft dysfunction, proteinuria, diabetes, immunosuppressive and antihypertensive agents (Massy & Kasiske, 1996).

Among immunosuppressive agents, corticosteroids, CNIs and mammalian target of rapamycin inhibitors (mTORs), sirolimus and everolimus, are those most frequently associated with hyperlipidemia. The mechanisms of corticosteroid-induced hyperlipidemia are through promoting insulin resistance and hyperinsulinism, reduction of lipoprotein lipase activity, overproduction of triglycerides and secretion of VLDL-cholesterol (Hricik et al, 1994). CNIs inhibit bile acid synthesis and binding of the drugs to the LDL-cholesterol receptor with reduction of its activity. Also a decrease in lipoprotein lipase activity and impairment of LDL-cholesterol catabolism may be involved (Moore et al, 2001). These effects seem to be more prominent with cyclosporine than with tacrolimus (Ligtenberg et al, 2001, Moore et al, 2001, Vincenti et al, 2002). mTORs are the agents with stronger hyperlipidemic effect (Kasiske et al, 2008), which is more accentuated in those patients also treated with cyclosporine than in those treated with tacrolimus (Ciancio et al, 2004). The pathogenesis of mTOR dyslipidemia is unclear. A reduced catabolism of apo B100 could be the cause of increased triglycerides and cholesterol and decreased lipoprotein lipase activity and increased free fatty acid levels may be contributing factors. Their effects are dose dependent and rapidly reversible (Kasiske et al, 2008, Webster et al, 2006).

As in the general population, hypercholesterolemia and low HDL-cholesterol levels are associated with ischemic heart disease (Kasiske, 1988 Kasiske et al, 1996, Kasiske et al, 2006, Marcén et al, 2006, Rigatto et al, 2002). The treatment of this complication may follow the recommendations given to the general population and confirmed by the transplant guidelines (KDIGO, 2009). It is important to begin with a rich diet of monosaturated fats, but diet therapy does not control hyperlipidemia and lipid-lowering agents have to be added. 3-Hydroxy-3methylglutaryl coenzyme A reductase inhibitors (statins) are the elective pharmacologic agents in hypercholesterolemic patients. Fluvastatin, pravastatin and atorvastatin seem to have a more favourable safety profile over simvastatin and lovastatin. In patients with hypertriglyceridemia, gemfibrozil is the pharmacological agent of choice. Some observational studies have shown an association between statin therapy and better patient outcome (Cosio et al, 2002a, Wiesbauer et al, 2008). However, the Assessment of Lescol in Renal Transplantation (ALERT) study did not show differences in the primary compound end point, despite a reduction of 32% in the LDL-cholesterol blood levels at 5 years follow-up, between recipients treated with fluvastatin compared with those on placebo (Holdaas et al, 2003). A later evaluation of the study showed the benefits of the treatment but only when statin therapy started in the first two years after transplantation and in low-risk recipients (Holdaas et al 2005, Jardine et al 2004). In case of statin intolerance or hyperlipidemia of difficult control, ezetimibe that blocks the cholesterol absorption in the brush border, alone or combined with statins, is an efficient and safe alternative (Buchanan et al 2006, Langone & Chuang, 2006).

4.1.4 Diabetes

In the renal transplant recipients, two types of diabetes mellitus can be distinguished, diabetes mellitus as the cause of end-stage renal disease and new onset diabetes mellitus (NODAT). The prevalence of diabetes mellitus as the cause of renal failure is variable among countries. In the USA, more than 20% of the patients on the waiting list or transplanted have diabetes mellitus as the cause of renal failure (Collins et al, 2008). The incidence of NODAT varies between 2% to 50% in the first posttransplantation year according to the criteria used in its definition (Montori et al, 2002). When the American Diabetes Association criteria were used, the incidence of NODAT at 12 months was 13% and of glucose intolerance of 33% in a study performed at the Mayo Clinic (Cosio et al, 2005). Similar findings have been observed in a prospective study from Spain (Marcén et al, 2006). As the term NODAT does not include states of impaired fasting glucose and impaired glucose tolerance which pose a cardiovascular threat similar to overt diabetes, the term transplant associated hyperglycemia (TAH) has been proposed (Crutchlow & Bloom, 2008). The most common risk factors associated with NODAT or TAH include: race, blacks or hispanics, older age, obesity, family history, hepatitis C virus infection and some immunosuppressive agents such as corticosteroids, CNIs (tacrolimus) and mTORs (Crutchlow & Bloom, 2007, Montori et al, 2002).

The effects of immunosuppressive agents on glucose metabolism have been widely reviewed (Heisel et al, 2004, Miller, 2002, Morales & Dominguez, 2006). Both CNIs, cyclosporine and tacrolimus, cause NODAT by inducing insulin resistance or by impaired insulin secretion (Hornum et al, 2010). Early trials designed to compare the efficacy and security of cyclosporine and tacrolimus, and registry data showed a higher incidence of NODAT in patients treated with tacrolimus (Kasiske et al, 2003, Mayer et al, 1997, Vincenti et al, 2002) and more recent studies have confirmed these findings (Vincenti et al 2007). Also mTORs are associated with an increased risk of NODAT (Johnston et al, 2008). These agents induce hyperglucemia by impairing insulin-mediated suppression of hepatic glucose production, by ectopic triglyceride deposition leading to insulin resistance, or by direct β cell toxicity (Crutchlow & Bloom, 2007).

Single centre and registry studies have shown the association of NODAT with acute myocardial infarction, cerebrovascular events and mortality (Cosio et al 2002b, 2005, 2008, Kasiske et al 2003, Lentine et al, 2005, 2008). The treatment has the objective of preventing the symptoms due to uncontrolled hyperglucemia and the microvascular complications as the transplant recipients develop identical complications as the nontransplanted diabetic patients (Burroughs et al, 2007). The guidelines of the American Diabetes Association and the Joint Nacional Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure for patients with type 2 diabetes have been recommended (Haffner SM 2003).

4.1.5 Tobacco

The effects of tobacco on the health in the general population are well known. It is a risk factor of cardiovascular diseases, malignancies and respiratory diseases (Bartecchi et al, 1994). About 25% of the renal transplant population are active smokers after transplantation (Cosio et al 1999, Kasiske & Klinger, 2000, Zitt et al, 2007). In transplant recipients, tobacco was associated with cardiovascular diseases and mortality (Kasiske & Klinger, 2000). It has been reported that the negative impact of tobacco on health disappeared after five years, and some authors emphasize that efforts have to be made to convince the patients about the

benefits of avoiding smoking. There are few data about the influence of transplant on toxic habits, but some studies suggest that transplantation constituted a strong reason to give up smoking (Banas et al, 2008).

4.1.6 Obesity

Obesity is a growing health problem in the general population. Epidemiological studies have shown its association with a higher morbidity and mortality due to cardiovascular diseases (Allison et al, 1999). Transplant recipients have a tendency to gain body weight mostly in the first year after grafting. In a study performed in our Unit, the mean body weight gain in the first year was 5 kg or 8.7 % of the body weight at the time of transplantation and the percentage of obese patients increased nearly two fold, from 6.5% to 11.7% (Marcén et al, 2007). Inappropriate dietary habits, decreased physical activity, and increased appetite as a result of well-being and corticosteroid therapy are among the causes of overweight and obesity after transplantation.

Both body weight gain and obesity (body mass index $>30 \text{ kg/m}^2$) are associated with an increased risk for NODAT, hypertension, hyperlipidemia and metabolic syndrome which are cardiovascular risk factors (Armstrong et al, 2005, El-Agroudy et al 2004). The effects of weight gain and obesity on graft and patient outcome are controversial. Some authors do not find that the complication has any impact on patient mortality (Chang et al 2007, Marcén et al, 2007, Massarweh et al, 2005). However, other studies and registry data have shown their negative impact on patient survival due to an increased cardiovascular and infectious mortality (Aalten et al, 2006, El-Agroudy et al 2004, Meier-Kriesche et al, 2002). In our opinion, controlling weight gain and weight reduction in patients with marked obesity seems to be a goal to improve well-being and outcomes in renal transplant recipients.

4.2 Risk factors associated with transplantation

4.2.1 Anaemia

Post-transplant anaemia is a common complication that has only been recently studied and considered. Its prevalence depends greatly on the definition criteria and the time post-transplant. Nowadays, there is a trend toward the use of the World Health Organization criteria which define anaemia as serum haemoglobin less than 12 g/dl in women and less than 13 g/dl in men. The prevalence of anaemia is about 90% during the first posttransplant weeks and decreases to 25% to 35% at 12 months and remains stable or slightly increases thereafter (Kamar et al, 2008, Vanrenterghem et al, 2003, Yorgin et al, 2002). In cross-sectional studies the prevalence of anaemia reached more than one third of patients and it was severe, serum haemoglobin below 11 g/dl, in about 10% (Karthikeyan et al, 2004, Marcen et al, 2009a, Molnar et al, 2007, Vanrenterghem et al, 2003). The origin of anaemia is multifactorial and graft function is the most important factor (Shah et al, 2006, Vanrenterghem et al, 2003, Yorgin et al, 2002). However, it does not completely explain post-transplant anaemia, as renal transplant recipients have more severe anaemia for each chronic renal disease stage when compared with nontransplantation subjects (Chadban et al, 2007). Several immunosuppressive agents such as azathioprine, mycophenolate mofetil (MMF), enteric coated mycophenolic acid (EC-MPA) and mTORs may cause anemia due to bone marrow toxicity or to disorders on iron homeostasis (Augustine et al, 2004, Fishbane et al, 2009, Vanrenterghem et al, 2003, Wang et al, 2004). The combination of MMF or EC-MPA with mTORs is specially toxic for the bone marrow (Hricik, 2003, Rigatto, 2006). Other

medications as angiotensin converting enzyme inhibitors and angiotensin receptor blockers are known to cause anaemia and should be cautiously used.

Recent studies have shown an association between anaemia and graft survival, cardiovascular diseases and mortality. Post-transplant anaemia seems to be a risk factor of congestive heart failure and of left ventricular hypertrophy but not of ischaemic heart disease (Borrow et al, 2008, Rigatto et al, 2002, 2003b). In addition, anaemia has been associated with increased mortality in some studies (Chhabra et al, 2008, Imoagene-Oyediji et al, 2006, Kamar et al, 2008, Molnar et al, 2007) but not in others (Winkelmayer et al, 2006). The treatment of anaemia must follow the recommendations given for patients with chronic kidney disease in the Kidney Disease Outcomes Quality Initiative (KDOQI) guidelines (KDIGO, 2009). Iron supplementation and erythropoiesis stimulant agents should be administered to maintain serum haemoglobin between 11 and 12 g/dl.

4.2.2 Graft dysfunction

Renal function measured by the estimated glomerular filtration rate (eGFR) is a cardiovascular risk factor in the general population (Go et al, 2004). An important number of renal transplant recipients have different stages of renal failure (Figure 2), and at least two thirds have chronic renal failure defined by an eGFR below 60 ml/min/1.73m² (Ansell et al, 2007, Karthikeyan et al, 2004, Marcén et al, 2005). As cardiovascular diseases are the leading cause of renal transplant recipient mortality, it seems logical to think in the existence of some links between premature cardiovascular death and poor graft function. Registry and prospective studies have demonstrated a correlation between serum creatinine and cardiovascular events or cardiovascular mortality (Fellstrom et al, 2005, Meier-Kriesche et al, 2003, Pilmore et al, 2010, Soveri et al, 2006). However, for other authors the increased cardiovascular risk of patients with poor graft function is mostly due to the effects of hypertension and anaemia than to graft failure itself (Rigatto et al, 2002). Moreover, uncontrolled hyperparathyroidism mostly in recipients with poor graft function may be a risk factor for progression of coronary artery calcification (Mazzaferro et al, 2009). The treatment of renal dysfunction includes the control of hypertension and dyslipidemias (Arias et al, 2005, Opelz et al, 2005, Wiesbauer et al, 2008) as well as the use of non-nephrotoxic immunosuppressive agents as MMF, EC-MPA and belatacept and dose reduction or withdrawal of CNIs.

The prevalence of proteinuria in the renal transplant recipients is between 7.5 and 45% (Knoll, 2009). It is a risk factor of progressive renal function loss and of cardiovascular disease in nontransplantation patients. Retrospective studies have reported that proteinuria is an important predictor of cardiovascular events and mortality in renal transplant recipients (Fernandez-Fresnedo et al, 2002, Roodnat 2001). It is important to note that proteinuria is frequently associated with graft dysfunction, hypertension and obesity and the effects of proteinuria on cardiovascular events could be mediated by these conditions or viceversa. Treatment of proteinuria includes control of hypertension, maintaining blood pressure levels below 120/80 mm Hg, of dyslipidemias and of overweight, and avoiding immunosuppressive agents associated with proteinuria as mTORs (Amer & Cosio, 2009). Angiotensin converting enzyme inhibitors and angiotensin receptor blockers are the elective agents of treatment of hypertension in patients with proteinuria because of their antiproteinuric effects. However, these agents can deteriorate graft function and increase the severity of anaemia, both cardiovascular risk factors as well. Moreover, there are no definitive studies which support the effectiveness of this treatment.

Fig. 2. Distribution of chronic kidney disease stages at 12 months in 447 renal transplant recipients (Ramón y Cajal Hospital)

4.2.3 Immunosuppression

The effects of immunosuppressive agents in cardiovascular risk factors have been previously reviewed (Miller, 2002, Marcén, 2009b, Morales & Dominguez-Gil, 2006). In addition, the impact of these agents on each cardiovascular risk factor has already been described. Corticosteroids and CNIs are the agents with worst cardiovascular risk profile, increasing the incidence and severity of traditional risk factors such as hypertension, hyperlipidemia, diabetes and obesity. CNIs also are nephrotoxic and cause graft function impairment. mTORs increase the risk of anaemia, hyperlipidemia, diabetes and proteinuria (Webster et al, 2006, Schena et al, 2009). MMF and EC-MPA may increase the risk of anaemia but have no impact in the other cardiovascular risk factors. Combinations of CNIs with MMF and EC-MPA or with mTOR allow corticosteroid withdrawal without harm. On the other hand, MMF and EC-MPA and mTORs alone or in combination can be used for CNI dose reduction or withdrawal. All these strategies have been shown to be effective in improving cardiovascular risk but there are no prospective studies in which this improvement is followed by a lower incidence of cardiovascular events or mortality (Schena et al, 2009). Among the new immunosuppressive agents, belatacept improved short-term cardiovascular risk profile and graft function when compared with cyclosporine (Vincenti et al, 2010, Durrbach et al 2010). However, phase II trials with tasocitinib have shown no improvement in hyperlipidemia and blood pressure when compared with patients who received tacrolimus (Busque et al, 2009).

4.3 Emergent risk factors

4.3.1 Hyperhomocysteinemia

Homocystein is a sulfur amino acid produced in all cells, and hyperhomocysteinemia has been considered a cardiovascular risk factor in the general population. Hyperhomocysteinemia is present in around 60% to 70% of renal transplant recipients. Homocystein levels are related with graft function, folic acid levels, serum albumin, age,

and treatment with CNIs (Bostom et al, 1999, Ducloux et al, 2000, Friedman et al, 2002). The impact of homocystein levels on cardiovascular events is controversial. Some authors found an association between the homocystein levels and cardiovascular diseases (Ducloux et al, 2000, 2004) and a prospective study has shown that homocystein levels above 12 $\mu\text{mol/l}$ were associated with 2.44 times increased mortality (Winkelmayer et al, 2005). Other authors have not found this association (Hagen et al, 2001). The treatment consists in the administration of folic acid supplements (5 mg/day) even with normal folic acid levels (Fernandez-Miranda et al, 2000). The efficacy of this therapy in the prevention of cardiovascular events has been examined in the Folic Acid for Vascular Outcome Reduction in Transplantation (FAVORIT). This trial should contribute to answer this question (Bostom et al, 2009).

4.3.2 Inflammation

C-reactive protein (CRP) is an acute phase marker of inflammation. It is produced by the liver under control of several cytokines. In the general population, C-reactive protein is associated with obesity and poor renal function (Stuveling et al, 2003). It is also a negative predictor of acute myocardial infarction, cerebrovascular disease and cardiovascular mortality (Ridker et al, 1997). In renal transplant recipients, its levels are associated with graft function, waist circumference and smoking (Van Ree et al, 2005). The association between graft function and C-reactive protein levels could be explained by the situation of chronic low-grade inflammation, by being a marker of graft-mediated immune response or by a decreased renal excretion. As in the general population, it has been considered a risk factor of cardiovascular disease and mortality (Ducloux et al 2004, Winkelmayer et al 2004). Data from the ALERT study have confirmed the previous findings, baseline levels of CRP as well as of IL-6, another inflammation marker, were independently associated with major cardiovascular events and all-cause mortality (Abedini et al, 2009b). Results from a retrospective study have shown an association between MMF therapy and less inflammation than other immunosuppressive agents (Wong et al, 2007).

5. Therapeutical strategies

The management of each particular cardiovascular disease in renal transplant recipients should be similar to that used in the general population. In addition, clinical trials have demonstrated that cardiovascular events and cardiovascular mortality have been reduced by controlling blood glucose, lipid levels and blood pressure in the general population. As interventional studies are lacking in the transplant population, it seems reasonable to extrapolate these findings to transplant recipients. However, transplant recipients present differences from the general population, one of them is the high incidence of graft dysfunction. Preserving graft function has to be a goal in the management of transplant recipients and this could be partly accomplished by controlling the traditional cardiovascular risk factors and by a prudent use of immunosuppressive agents. CNIs minimization or withdrawal may be individually considered. Additional interventions such as treatment of anaemia with erythropoiesis stimulating agents could help in the prevention of cardiovascular diseases but the optimal haemoglobin threshold has to be determined. The benefits of lowering homocysteine levels have not been proved. In addition, long-term interventional studies should be performed in order to improve graft and patient outcomes (Table 3).

Control of cardiovascular risk factors	Preserving graft function	Treating emergent factors
Blood pressure Dislipidemias Diabetes Smoking Obesity Anaemia	Minimization/avoidance of CNIs Control of proteinuria	Folic acid supplements Aspirin

Table 3. Prevention and treatment of cardiovascular diseases after transplantation.

6. Summary

Cardiovascular diseases are common after transplantation. Coronary vascular disease, cerebrovascular disease and congestive heart failure are the diseases most commonly associated with mortality. The increased incidence of cardiovascular events could be partly explained by the high prevalence of traditional risk factors which are not adequately controlled and by the presence of renal dysfunction. Pretransplant evaluation of candidates, control of traditional risk factors and preservation of graft function should be the measures taken to improve patient outcome. The control of traditional risk factors has been effective in the reduction of cardiovascular events in the general population and there are no reasons to believe that it does not work in the transplant population. In addition, adequate control of traditional risk factors could preserve progression of graft failure. A prudent use of immunosuppressive agents could also help to improve the cardiovascular risk profile and graft function. The benefits of additional interventions need to be proved.

7. References

Aakhus, S., Dahl, K. & Widerøe, TE. (1986). Hyperlipidaemia in renal transplant patients. *J Intern Med.* Vol.239, No.5, (May 1996), pp.407-415. ISSN: 0954-6820.

Aakhus, S., Dahl, K. & Wideroe, TE. (1986). Cardiovascular morbidity and risk factors in renal transplant patients. *Nephrol Dial Transplant.* Vol. 14, No.3, (March 1999) pp.648-654. ISSN: 0931-0509.

Aakhus, S., Dahl, K. & Wideroe, TE. (1987). Cardiovascular disease in stable renal transplant patients in Norway: morbidity and mortality during a 5-yr follow-up. *Clin Transplant.* Vol. 18, No.5, (October 2004), pp.596-604. ISSN: 0902-0063.

Aalten, J., Christiaans, MA., de Fijter, H., Hené, R., van der Heijde, JH., Roodnat. J., Surachno, J. & Hoitsma, A. (1988). The influence of obesity on short- and long-term graft and patient survival after renal transplantation. *Transplant Int.* Vol.19, No.11 (November 2006), PP. 901-907. ISSN.0934-0874.

Abbott, KC., Reynols, JC., Taylor, AJ. & Agodoa, LY. (2001). Hospitalized atrial fibrillation after renal transplantation in the United States. *Am J Transplant* . Vol. 3, No.4, (June 2003a), pp.471-476. ISSN: 1600-6135.

Abbott, KC., Yuan, CM., Taylor, AJ., Cruess, DF. & Agodoa, LY. (1990). Early renal insufficiency and hospitalized heart disease after renal transplantation in the era of modern immunosuppression. *J Am Soc Nephrol* Vol.14, No.9, (November 2003b), pp.2358-2365, ISSN: 1046-6673.

- Abedini, S., Holme, I., Fellstrom, B., Jardine, A., Cole, E., Maes, B. & Holdaas, H. on behalf of the ALERT study group. (1963). Cerebrovascular events in renal transplant recipients. *Transplantation*. Vol.87, No.1, (January 2009a), pp.112-7. ISSN: 0041-1337.
- Abedini, S., Holme, I., März, W., Weihrauch, G., Fellström, B., Jardine, A., Cole, E., Maes, B., Neumayer, H-H., Gronhagen-Riska, C., Ambühl, P. & Holdaas, H on behalf of the ALERT study group. (2006). Inflammation in renal transplantation. *Clin J Am Soc Nephrol* Vol.4, No.7, (July 2009b), pp. 1246-1254. ISSN: 1555-9041.
- Allison, DB., Fontaine, KR., Manson, JE., Stevens, J. & Vantallie, TB. (1883). Annual deaths attributable to obesity in the United States. *JAMA* Vol.282, No.16, (October 1999), pp.1530-1538, ISSN. 0098-7484.
- Amer, H. & Cosio, FG. (1990). Significance and management of proteinuria in kidney transplant recipients. *J Am Soc Nephrol*. Vol.20, No.12, (December 2009), pp. 2490-2492, ISSN: 1046-6673.
- Ansell, D., Udayarej, UP., Steenkamp, R. & Dudley, CRK. (2001). Chronic renal failure in kidney transplant recipients. Do they receive optimum care?: data from the UK renal registry. *Am J Transplant* Vol.7, No.5, (May 2007), pp.1167-1176. ISSN:1600-6135.
- Arias, M., Fernández-Fresnedo, G., Rodrigo, E., Ruiz, JC., González-Cotorruelo, J. & Gómez-Alamillo, C. (1972). Non-Immunologic intervention in chronic allograft nephropathy. *Kidney Int* Vol.99 (December 2005), pp.S118-S123, ISSN: 0085-2538.
- Armstrong, KA., Campbell, SB., Hawley, CM., Nicol, DL., Johnson, DW. & Isbel, NM. (2001). Obesity is associated with worsening cardiovascular risk factor profiles and proteinuria progression in renal transplant recipients *Am J Transplant* . Vol.5, No.11, (November 2005), pp.2710-2718, ISSN.1600-6135.
- Augustine, JJ., Knauss, TC., Schulak, JA., Bodziak, KA., Siegel, C. & Hricik, DE. (2001). Comparative effects of sirolimus and mycophenolate mofetil on erythropoiesis in kidney transplant patients. *Am J Transplant*. Vol.4, No.12, (December 2004), pp.2001-2006, ISSN.1600-6135.
- Aull-Watschinger, S., Konstantin, H., Demetriou, D., Schillinger, M., Habicht, A., Hörl, WH. & Watschinger, B. (1986). Pre-transplant predictors of cerebrovascular events after kidney transplantation. *Nephrol Dial Transplant* Vol. 23, No. 4 (April 2008), pp. 1429-1435, ISSN: 0931-0509.
- Banas, MC., Banas, B., Wolf, J., Hoffman, U., Krüger, B., Böger, CA., Orth, SR. & Krämer, BK. (1986). Smoking behaviour of patients before and after renal transplantation. *Nephrol Dial Transplant* Vol.23, No.4, (April 2008), pp.1442-1446, ISSN.0931-0509.
- Bartecchi, CE., MacKenzie, TD. & Schrier, RW. (1812). The human cost of tobacco use (first of two parts). *New Engl J Med*. Vol.330, No.13 (March 1994), pp.907-912, ISSN. 0028-4793.
- Borrows, R., Loucaidou, M., Chusney, G., Borrows, S., Tromp, JV., Cairns, T., Griffith, M., Hakim, N., McLean, A., Palmer, A., Papalois, V. & Taube, D. (1986). Anaemia and congestive heart failure early post-renal transplantation. *Nephrol Dial Transplant*. Vol.23, No.5, (May 2008), pp.3:1728-1734, ISSN.0931-0509.
- Bostom, AG., Gohh, RY., Beaulieu, AJ., Han, H., Jacques, PF., Selhub, J., Dworkin, L. & Rosenberg, IH. (1963). Determinants of fasting plasma total homocysteine levels among chronic stable renal transplant recipients. *Transplantation*. Vol.68, No.2, (July 1999), pp.257-261, ISSN: 0041-1337.
- Bostom, AG., Carpenter, MA., Hunsicker, L., Jacques, PF., Kusek, JW., Levey, AS., McKenney, JL., Mercier, RY., Pfeffer, MA. & Selhub, J. (1981). Baseline

- characteristics of participants in the folic acid for vascular outcome reduction in transplantation (FAVORIT) trial. *Am J Kidney Dis.* Vol.53, No.1, (January 2009), pp.121-128, ISSN: 0272-6386.
- Buchanan, C., Smith, L., Corbett, J., Nelson, E. & Shihab F. (2001). A retrospective analysis of ezetimibe treatment in renal transplant recipients. *Am J Transplant* Vol.6, No.4, (April 2006), pp.770-774, ISSN: 1600-6135.
- Burroughs, TE., Swindle, J., Takemoto, S., Lentine, KL., Machnicki, G., Irish, WD., Brennan, DC. & Schnitzler, MA. (1963). Diabetic complications associated with new-onset diabetes mellitus in renal transplant recipients. *Transplantation* Vol.83, No.8 (April 2007), pp.1027-1034, ISSN: 0041-1337.
- Busque, S., Leventhal, J., Brennan, DC., Steinberg, S., Klintmalm, G., Shah, T., Mulgaonkar, S., Bromberg, JS., Vincenti, F., Harharan, S., Slakey, D., Peddi, VR., Fisher, RA., Lawendy, N., Wang, C. & Chan, G. (2001). Calcineurin-inhibitor-free immunosuppression based on the JAK inhibitor CP-690,550: a pilot study in the novo kidney allograft recipients. *Am J Transplant.* Vol.9, No.8 (August 2009), pp.1936-1945, ISSN: 1600-6135.
- Chadban, SJ., Baines, L., Polkinghorne, K., Jefferys, A., Dogra, S., Kanganas, C., Irish, A., Eris, J. & Walker, R. (1981). Anemia after kidney transplantation is not completely explained by reduced kidney function. *Am J Kidney Dis.* Vol.49, No.2, (February 2007), pp.301-309, ISSN: 0272-6386.
- Chhabra, D., Grafals, M., Skaro, AI., Parker, M. & Gallon, L. (2006). Impact of anemia after renal transplantation on patient and graft survival and on rate of acute rejection. *Clin J Am Soc Nephrol.* Vol.3, No.4, (July 2008), pp.1168-1174, ISSN: 1555- 9041.
- Chang, SH., Coates, PT. & McDonald, SP. (1963). Effects of body mass index at transplant on outcomes of kidney transplantation. *Transplantation.* Vol.84, No.8, (October 2007), pp.981-987, ISSN: 0041-1337.
- Chobanian, AV., Bakris, GL., Black, HR., Cushman, WC., Green, LA., Izzo, JL. Jones, DW., Materson, BJ., Oparil, S., Wright, JT. & Roccella, EJ. (1983). Seventh report of the joint national committee on prevention, detection, evaluation, and treatment of high blood pressure. The JNC 7 report. *JAMA.* Vol. 289, No.19, (December 2003), pp.2560-2572, ISSN: 0098-7484.
- Ciancio, G., Burke, GW., Gaynor, JJ., Mattiazzi, A., Roth, D., Kupin, W., Nicolas, M., Ruiz, P., Rosen, A. & Miller, J. (1963). A randomized long-term trial of tacrolimus/sirolimus versus tacrolimus/mycophenolate mofetil, versus cyclosporine (NEORAL)/sirolimus in renal transplantation II. Survival, function, and protocol compliance at 1 year. *Transplantation* Vol.77, No.2, (January 2004), pp.252-258, ISSN: 0041-1337.
- Collins AJ. (1972). Impact of congestive heart failure and other cardiac diseases on patients outcomes. *Kidney Int. Suppl* 81 (October 2002), pp.S3-7, ISSN: 0085-2538.
- Collins, AJ., Foley, RN., Herzog, C., Chavers, BM., Gilbertson, D., Ishani, A., Kasiske, BL., Liu, J., Mau, LW., McBean, M., Murray, A., St Peter, W., Guo, H., Li, Q., Li, S., Li, S., Peng, Y., Qiu, Y., Roberts, T., Skeans, M., Snyder, J., Solid, C., Wang, C., Weinhandl, E., Zaun, D., Arko, C., Chen, SC., Dalleska, F., Daniels, F., Dunning, S., Ebben, J., Frazier, E., Hanzlik, C., Johnson, R., Sheets, D., Wang, X., Forrest, B., Constantini, E., Everson, S., Eggers, PW. & Agodoa L. (2008). (1981). Excerpts from the United States Renal Data System 2007 annual data report. *Am J Kidney Dis.* Vol. 51, 1 Suppl 1, (January 2008), pp.S1-320, ISSN: 0272-6386.

- Cosio, FG., Falkenhaim, MF., Pesavento, TE., Yim, S., Alamir, A., Henry, ML. & Ferguson, RM. (1987). Patient survival after renal transplantation.II. The impact of smoking . *Clin Transplant* Vol.13, No.4, (August 1999), pp.336-341. ISSN. 0902-0063.
- Cosio, FG., Pesavento, TE., Pelletier, RP., Henry, M., Ferguson, RM., Kim, S. & Lemeshow, S. (1981). Patient survival after renal transplantation III: the effects of statins, *Am J Kidney Dis* Vol.40, No.3, (September 2002a), pp.638-643. ISSN. 0272-6386.
- Cosio, FG., Pesavento, TE., Kim, S., Osei, K., Henry, M. & Ferguson, RM. (1972). Patient survival after renal transplantation:IV. Impact of post-transplant diabetes. *Kidney Int.* Vol.62, No.4 (October 2002b), pp. 1440-1446. ISSN. 0085-2538.
- Cosio, FG., Kudva, Y., van der Velde, M., Larson, TS., Textor, SC., Griffin, MD. & Stegall, MD. (1972) New onset hyperglycemia and diabetes are associated with increased cardiovascular risk after kidney transplantation. *Kidney Int* . Vol.67, No.6, (June 2005), pp. 2415-2421. ISSN: 0085-2538.
- Cosio, FG., Hickson, LJ., Griffin, MD., Stegall, MD. & Kudva, Y. (2001). Patient survival and cardiovascular risk after kidney transplantation: the challenge of diabetes. *Am J Transplant* Vol.8, No.3, (March 2008), pp.593-599, ISSN. 1600-6135.
- Crutchlow MF & Bloom RD. (2006). Transplant-associated hyperglycemia:a new look at an old problem. *Clin J Am Soc Nephrol* Vol.2, No.2, (March 2007), pp.343-355, ISSN: 1555-9041
- Culleton, BF. & Wilson, PWF. (1990). Cardiovascular disease: risk factors, secular trends, and therapeutic guidelines. *J Am Soc Nephrol.* Vol.9, 12 suppl, (December 1998), pp:S5-S15, ISSN: 1046-6673.
- De Mattos, AM., Prather, J., Olyaei, AJ., Shibagaki, Y., Keith, DS., Mori, M., Norman, DJ. et Becker, T. (1972). Cardiovascular events following renal transplantation: role of traditional and transplant-specific risk factors. *Kidney Int.* Vol. 70, No. 4, (August 2006), pp.757-764, ISSN: 0085-2538.
- Ducloux, D., Motte, G., Challier, B., Gibey, R. & Chalopin, JM. (1990). Serum total homocysteine and cardiovascular occurrence in chronic stable renal transplant recipients: a prospective study. *J Am Soc Nephrol.* Vol.11, No.1, (January 2000), pp.134- 137, ISSN: 1046-6673.
- Ducloux, D., Kazory, A. & Chalopin, JM. (1972). Predicting coronary heart disease in renal transplant recipients: a prospective study. *Kidney Int* . Vol. 66, No.1, (July 2004), pp.441-447, ISSN: 0085-2538.
- Durrbach, A., Pestana, JM., Pearson, T., Vincenti, F., García, CD., Campistol, J., Rial, MdeC., Florman, S., Block, A., Di Russo, G., Xing, J., Garg, P. & Grinyo, J. (2001). A phase III study of belatacept versus cyclosporine in kidney transplants from extended criteria donors (BENEFIT-EXT study). *Am J Transplant.* Vol.10, No.3, (March 2010), pp.547-557, ISSN: 1600- 6135.
- El-Agroudy, AE., Wafa, EW., Greith, OE., El-Dein, ABS. & Ghoneim, MA. (1963). Weight gain after renal transplantation is a risk factor for patient and graft outcome. *Transplantation.* Vol15, No.77, (May 2004), pp.1381-1385. ISSN.0041-1337.
- Fellström, B., Jardine, AG., Soveri, I., Cole, E., Neumayer, HH., Maes, B., Gimpelewicz, C. & Holdaas, H. (1963). Renal dysfunction as a risk factor for mortality and cardiovascular disease in renal transplantation: experience from the Assessment of Lescol in Renal Transplant trial. *Transplantation* Vol.79; No.7, (August 2005), pp. 1160-1163, ISSN: 0041-1337.
- Fernández-Fresnedo, G., Escallada, R., Rodrigo, E., De Francisco, AL., Cotorruelo, JG., Sanz De Castro, S., Zubimendi, JA., Ruiz, JC. & Arias, M. (1963). The risk of

- cardiovascular disease associated with proteinuria in renal transplant patients. *Transplantation*. Vol.76, No.8, (April 2002), pp.1345-1348, ISSN: 0041-1337.
- Fernández-Fresnedo, G., Escallada, R., Martín de Francisco, AL., Ruiz, JC., Rodrigo, E., Sanz de Castro, S., González-Cotorruelo, J & Arias, M. (2001). Association between pulse pressure and cardiovascular disease in renal transplant patients. *Am J Transplant* Vol.5, No.2, (February, 2005), pp.394-398, ISSN: 1600-6135.
- Fernandez-Miranda, C., Gómez, P., Díaz-Rubio, P., Estenoz, J., Carillo, JL., Andrés, A. & Morales, JM. (1987). Plasma homocysteine levels in renal transplanted patients on cyclosporine or tacrolimus therapy: effect of treatment with folic acid. *Clin Transplant*. Vol.14, No.2, (April 2000), pp.110-114, ISSN: 0902-0063.
- Fishbane, S., Cohen, DJ., Coyne, DW., Djamali, A., Singh, AK. & Wish, JB. (1972). Posttransplant anemia: the role of sirolimus. *Kidney Int* . Vol.76, No.4, (August 2009), pp.376-382, ISSN:0085-2538.
- Foley, RN., Parfrey, PS., Harnett, JD., Kent, GM., Murray, DC. & Barré, PE. (1990). The prognostic importance of left ventricular geometry in uremic cardiomyopathy. *J Am Soc Nephrol*. Vol.5, No.12, (June 1995), pp. 2024-2031, ISSN: 1046-6673.
- Foley, RN. Parfrey, PS. & Sarnak, MJ (1981). Clinical epidemiology of cardiovascular disease in chronic renal disease. *Am J Kidney Dis*. Vol.32, No.5, suppl.3, (November 1998), pp.S112-S119, ISSN: 0272-6386.
- Friedman, AN., Rosenberg, IH., Selhub, J., Levey, AS. & Bostom, AG. (2001). Hyperhomocysteinemia in renal transplant recipients. *Am J Transplant* . Vol.2, No.4, (April 2002), pp.308-313, ISSN: 1600-6135.
- Go, AS., Chertow, GM., Fan, D., McCulloch, CE. & Hsu, C. (1812). Chronic kidney disease and the risks of death, cardiovascular events, and hospitalization. *N Engl J Med* Vol.351; No.13, (September 2004), pp.1296-1305, ISSN: 0028-4793.
- Haffner, SM. (1978). Management of dyslipidemia in adults with diabetes. *Diabetes Care* Vol.26, Suppl 1, (January 2003), pp.S83- S86. ISSN. 0149-5992.
- Hagen, W., Födingner, M., Heinz, G., Buchmayer, H., Hörl, WH. & Sunder-Plassman, G. (1972). Effect of MTHFR genotypes and hyperhomocysteinemia on patient and graft survival in kidney transplant recipients. *Kidney Int* . Vol.78, (February 2001), pp.S253-S257, ISSN: 0085-2538.
- Harnett, JD., Foley, RN., Kent, GM., Barre, PE., Murray, D. & Parfrey, PS. (1972). Congestive heart failure in dialysis patients: prevalence, incidence, prognosis and risk factors. *Kidney Int* . Vol.47, No.3, (March 1995), pp.884-890, ISSN: 0085-2538.
- Heisel O, Heisel R, Balshaw R & Keown P. (2001). New onset diabetes mellitus in patients receiving calcineurin inhibitors: a systematic review and meta-analysis. *Am J Transplant* Vol.4, No.4, (April 2004), pp.583-595, ISSN: 1600-6135.
- Herzog, CA., Ma, JZ & Collins AJ (1812). Poor long-term survival after acute myocardial infarction among patients on long-term dialysis. *N Engl J Med*. Vol. 339, No.12, (September 1998), pp. 799-805, ISSN: 0028-4793.
- Herzog, CA., Ma, JZ. & Collins, AJ. (1981). Long term survival of renal transplant recipients in the United States after acute myocardial infarction. *Am J Kidney Dis*. Vol. 36, No. 1, (July 2000), pp.145-153, ISSN: 0272-6386.
- Hiremath, S., Ferguson, D., Doucette, S., Mulay, AV. & Knoll, GA. (2001). Renin angiotensin system blockade in kidney transplantation: a systematic review of the evidence. *Am J Transplant*. Vol.7, No.10, (October 2007), pp. 2350-2360, ISSN: 1600-6135.
- Holdaas, H., Fellström, B., Jardina, AG., Holme, I., Nyberg, G., Fauchald, P., Frønhagen-Riska, C., Madsen, S., Neumayer, H-H., Cole, E., Maes, B., Ambühl, P., Olsson,

- AG., Haertmann, A., Solbu, DO. & Pedersen, TR. on behalf of the Assessment of Lescol in renal transplantation (ALERT) Study Investigators. (1823). Effect of fluvastatin on cardiac outcomes in renal transplant recipients: a multicentre, randomised, placebo-controlled trial. *Lancet* Vol.361, No.9374, (June 2003), pp:2024-2031, ISSN: 0140-6736.
- Holdaas, H., Fellström, B., Jardine, AG. Nyberg, G., Grönhagen-Riska, C., Madsen, S., Neumayer, HH., Cole, E., Maes, B., Ambühl, P., Logan, JO., Staffler, B. & Gimpelewicz, C.(1986). Beneficial effect of early initiation of lipid-lowering therapy following renal transplantation. *Nephrol Dial Transplant* Vol.20, No.5, (May 2005), pp.974-980, ISSN: 0931-0509.
- Hornum M, Jorgensen KA, Hansen JM, Nielsen FT, Christensen KB, Mathiesen ER & Feldt-Rasmussen B. (1990). New-onset diabetes mellitus after kidney transplantation in Denmark. *Clin J Am Soc Nephrol* Vol.5, No.4, (April, 2010), pp.709-716, ISSN. 1555-9041.
- Howard, RJ., Patton, PR., Reed, AI., Hemming, AW., Van der Werf, WJ., Pfaff, WW., Srinivas, TR. & Scornik JC. (1963). The changing causes of graft loss and death after kidney transplantation. *Transplantation*. Vol. 73, No. 12, (April 2002), pp.1923-1928, ISSN: 0041-1337.
- Hricik, DE. (1981). Posttransplant hyperlipidemia: the treatment dilemma. *Am J Kidney Dis*. Vol.23, No.5, (May 1994), pp.766-771, ISSN: 0272-6386.
- Hricik, DE. (2001). Anemia after kidney transplantation-is the incidence increasing?. *Am J Transplant*. Vol.3, No.7, (July 2003), pp.771-772, ISSN.1600-6135.
- Imoagene-Oyedeji, AE., Rosas, SE., Doyle, AM., Goral, S. & Bloom, RD. (1990). Posttransplantation anemia at 12 months in kidney recipients treated with mycophenolate mofetil: risk factors and implications for mortality. *J Am Soc Nephrol*. Vol.17, No.11, (November 2006), pp.3240-7, ISSN.1046-6673.
- Jardine, AG., Holdaas, H., Fellström, B., Cole, E., Nyberg, G., Grönhagen-Riska, C., Madsen, S., Neumayer, HH., Maes, B., Ambühl, P., Olsson, AG., Holme, I., Fauchald, P., Gimpelwicz, C. & Pedersen, TR. (2001). Fluvastatin prevents cardiac death and myocardial infarction in renal transplant recipients: Pos-hoc subgroup analyses of the ALERT study. *Am J Transplant* Vol4, No.6 (June 2004), pp.988-995, ISSN: 1600-6135.
- Johnston, O., Rose, CL., Webster, AC. & Gill, JS. (1990). Sirolimus is associated with new-onset diabetes in kidney transplant recipients. *J Am Soc Nephrol* Vol.19, No.7, (July 2008), pp.1411-1418, ISSN.1046-6673.
- Kamar, N. & Rostaing, L. (1963). Negative impact of one-year anemia on long-term patient and graft survival in kidney transplant patients receiving calcineurin inhibitors and mycophenolate mofetil. *Transplantation*. Vol.85, No.8, (April 2008), pp.1120-1124, ISSN.0041-1337.
- Karthikeyan, V., Karpinski, J., Nair, RC. & Knoll G. (2001). The burden of chronic kidney disease in renal transplant recipients. *Am J Transplant*. Vol.4, No.2, (February 2004), pp.262-269. ISSN.1600-6135.
- Kasiske, BL. (1946). Risk factors for accelerated atherosclerosis in renal transplant recipients. *Am J Med*. Vol. 84, No. 6, (January 1988), pp. 985-992, ISSN: 0002-9343.
- Kasiske, BL., Guijarro, C., Massy, ZA., Wiederkehr, MR. & Ma, JZ. (1990). Cardiovascular disease after renal transplantation. *J Am Soc Nephrol*. Vol.7, No. 1, (January 1996), pp.158-165, ISSN: 1046-6673.

- Kasiske, BL., Chakkera, HA & Roel J. (1990). Explained and unexplained ischemic heart disease risk after renal transplantation. *J Am Soc Nephrol*. Vol. 11 No. 11, (September 2000a) pp. 1735-1743. ISSN. 1046-6673.
- Kasiske, BL. & Klinger, D. (1990). Cigarette smoking in renal transplant recipients. *J Am Soc Nephrol* Vol.11, No.4, (April 2000b), pp. 753-759. ISSN. 1046-6673.
- Kasiske, BL., Synder, JJ., Gilberston, D. & Matas, AJ. (2001). Diabetes mellitus after kidney transplantation in the United States. *Am J Transplant* Vol.3, No.2, (February 2003), pp.178-185, ISSN.1600-6135.
- Kasiske, BL., Anjum, S., Shah, R., Skogen, J., Kandaswamy, C., Danielson, B., O'Shaughnessy, EA., Dahl, DC., Silkensen, JR., Sahadevan, M. & Snyder, JJ. (1981). Hypertension after kidney transplantation. *Am J Kidney Dis* Vol.43, No.6, (April 2004), pp. 1071-1081, ISSN: 0272-6386.
- Kasiske, BL., Malik, MA. & Herzog, CA. (1963). Risk-stratified screening for ischemic heart disease in kidney transplant candidates. *Transplantation*. Vol. 80, No. 6, (September 2005), pp. 815-20. ISSN: 0041-1337.
- Kasiske, BL., Maclean, JR. & Snyder JJ. (1990). Acute myocardial infarction and kidney transplantation. *J Am Soc Nephrol*. Vol. 17, No.3, (March 2006), pp. 900-907, ISSN: 1046-6673.
- Kasiske, BL., de Mattos, A., Flechneer, SM., Gallon, L., Meier-Kriesche, H-U., Weir, MR. & Wilkinson, A. (2001). Mammalian target of rapamycin inhibitor dyslipidemia in kidney transplant recipients. *Am J Transplant* Vol.8, No7, (July 2008), pp. 1384-1392, ISSN: 1600-6135.
- KDIGO clinical practice guideline for the care of kidney transplant recipients. (2001). *Am J Transplant*. Vol.9, Suppl 3 (November 2009), pp.S1-155. ISSN.1600-6135.
- Kennedy, R., Case, C., Fathi, R., Johnson, D., Isbel, N. & Marwick, TH. (1946). Does renal failure cause an atherosclerotic milieu in patients with end-stage renal disease? *Am J Med*. Vol.110, No.3, (February 2001), pp.:198-204, ISSN: 0002-9343.
- Knoll, GA. (1981). Proteinuria in kidney transplant recipients: prevalence, prognosis and evidence-based management. *Am J Kidney Dis*. Vol.54, No.6 (December 2009), pp.1131-1144, ISSN: 0272-6386.
- Koomans, HE. & Ligtenberg, G. (1963). Mechanism and consequences of arterial hypertension after renal transplantation. *Transplantation* Vol.72, 6 Suppl, (September 2001), pp.S9-S12, ISSN: 0041-1337.
- Langone, AJ. & Chuang, P. (1963). Ezetimibe in renal transplant patients with hyperlipidemia resistant to HMG-CoA reductase inhibitors. *Transplantation* Vol.81, No.5, (March 2006), pp.804-807, ISSN: 0041-1337.
- Lentine, KL., Brennan, DC. & Schnitzler, MA. (1990). Incidence and predictors of myocardial infarction after kidney transplantation. *J Am Soc Nephrol*. Vol. 16, No. 2, (February 2005), pp.496-506. ISSN. 1046-6673.
- Lentine, KL., Schnitzler, MA., Abbott, KC., Li, L., Burroughs, TE., Irish, W. & Brennan, DC. (1981). De novo congestive heart failure after kidney transplantation: a common condition with poor prognostic implications. *Am J Kidney Dis* Vol.46, No.4, (October 2005), pp.720-733, ISSN. 0272-6386.
- Lentine, KL., Schnitzler, MA., Abbott, KC. Li, L., Xiao, H., Burroughs, TE., Takemoto, SK., Willoughby, LM., Gavard, JA. & Brennan, DC. (2006). Incidence, predictors, and associated outcomes of atrial fibrillation after kidney transplantation. *Clin J Am Soc Nephrol* Vol.1, No.2, (March 2006), pp.288-296, ISSN: 1555-9041.

- Lentine, KL., Rocca Rey, LA., Kolli, S., Bacchi, G., Schnitzler, MA., Abbott, KC., Xiao, H. & Brennan, DC. (2006). Variations in the risk for cerebrovascular events after kidney transplant compared with experience on the waiting list and after graft failure. *Clin J Am Soc Nephrol* Vol. 3, No.4, (July 2008a), pp. 1090-1101. ISSN. 1555-9041.
- Lentine, KL., Schnitzler, MA., Brennan, DC., Snyder, JJ., Hauptman, PJ., Abbott, KC., Axelrod, D., Salvalaggio, PR. & Kasiske, B. (2006). Cardiac evaluation before kidney transplantation: a practice patterns analysis in Medicare-insured dialysis patients. *Clin J Am Soc Nephrol*. Vol.3, No.4, (April 2008b), pp. 1115-24. ISSN. 1555-9041.
- Levin, A., Singer, J., Thompson, CR., Ross, H. & Lewis, M. (1981). Prevalent left ventricular hypertrophy in the predialysis population : identifying opportunities for intervention. *Am J Kidney Dis* . Vol.27, No.3, (March 1996), pp. 347-354, ISSN: 0272-6386.
- Ligtenberg, G., Hene, RJ., Blankestijn, PJ. & Koemans, HA (1990). Cardiovascular risk factors in renal transplant recipients: cyclosporin A versus tacrolimus. *J Am Soc Nephrol* Vol.12, No.2, (February 2001), pp.368-373, ISSN: 1046-6673..
- Lindholm, A., Albrechtsen, D., Frodin, L., Tufveson, G., Persson, NH. & Lundgran, G. (1963). Ischemic heart disease-major cause of death and graft loss after transplantation in Scandinavia. *Transplantation*. Vol. 60, No. 5, (September 1995), pp.451-457, ISSN: 0041-1337.
- Marcén, R., Pascual, J., Teruel, JL. Villafruela, JJ., Rivera, ME., Mampaso, F., Burgos, FJ. & Ortuño, J. (1963). Outcome of cadaveric renal transplant patients treated for 10 years with cyclosporine. Is chronic allograft nephropathy the major cause of graft loss?. *Transplantation*. Vol.72, No.1, (July 2001), pp.57-62, ISSN: 0041-1337.
- Marcén, R., Pascual, J., Tenorio, M., Ocaña, J., Teruel, JL., Villafruela, JJ., Fernández, M., Burgos & FJ., Ortuño, J. (1969). Chronic kidney disease in renal transplant recipients. *Transplant Proc*. Vol.37, No.9, (November 2005), pp.3718-3720, ISSN:0041- 1345.
- Marcén, R., Morales, JM., del Castillo, D., Campistol, JM., Serón, D., Valdés, F., Anaya, F., Andrés, A., Arias, M., Bustamante, J., Capdevila, L., Escuin, F., Gil-Vernet, S., González-Molina, M., Lampreave, I., Oppenheimer, F., & Pallardó, L., for the Spanish Renal Forum. (1969). Posttransplant diabetes mellitus in renal allograft recipients: a prospective multicenter study at 2 years. *Transplant Proc* Vol.38, No.10, (December 2006), pp:3530-3532. ISSN: 0041-1345.
- Marcén, R., Morales, JM., Arias, M., Fernández-Juarez, G., Fernández-Fresnedo, G., Ahdrés, A., Rodrigo, E., Pascual, J., Dominguez, B. & Ortuño, J. (1990). Ischemic heart disease after renal transplantation in patients on cyclosporine in Spain. *J Am Soc Nephrol*. Vol.17, 12 Suppl 3, (December 2006), pp.S286-S290, ISSN: 1046-6673.
- Marcén, R., Fernández, A., Pascual, J., Teruel, JL., Villafruela, JJ., Rodríguez, N., Martins, J., Burgos, FJ. & Ortuño, J. (1969). High body mass index and posttransplant weight gain are not risk factors for kidney graft and patient outcomes. *Transplant Proc* . Vol.39, No.7, (September 2007), pp.:2205-2207, ISSN. 0041-1345.
- Marcén, R., del Castillo, D., Capdevila, L., Fernandez-Fresnedo, G., Rodrigo, E., Cantarell, C., Fernandez-Rodriguez, A., López- Silva, MO., Camps, J., Aljama, P., Ortuño, J. & Arias, M. (1963). Achieving chronic kidney disease treatment targets in renal transplant recipients: results from a cross-sectional study in Spain. *Transplantation* Vol. 87, No.9, (May 2009a), pp. 1340-1346. ISSN.0041-1337.

- Marcén, R. (1971). Immunosuppressive drugs in kidney transplantation. Impact on patient survival, and incidence of cardiovascular disease, malignancy and infection. *Drugs*. Vol.69, No.16, (November 2009b), pp. 2227-2243. ISSN.0012- 6667.
- Massarweh, NN., Clayton, JL., Mangum, CA., Florman, SS. & Slakey, DP. (1963). High body mass index and short-and long- term renal allograft survival in adults. *Transplantation*. Vol.80, No.10, (November 2005), pp.1430-1434, ISSN. 0041-1337.
- Massy, ZA. (1963). Hyperlipidemia and cardiovascular disease after organ transplantation. *Transplantation*. Vol.72, suppl.6, (September 2001), pp.S13-S15, ISSN: 0041-1337.
- Massy, ZA. & Kasiske, BL. (1990). Post-transplant hyperlipidemia: mechanism and management. *J Am Soc Nephrol*. Vol.7, No.7, (July 1996), pp.971-977, ISSN: 1046-6673.
- Matas, AJ., Humar, A., Gillingham, KJ., Payne, WD., Gruessner, RWG., Kandaswamy, R., Dunn, DL., Najasrian, JS. & Sutherland, DER. (1972). Five preventable causes of kidney graft loss in the 1990s: A single-center analysis. *Kidney Int* Vol. 62, No. 2, (August 2002), pp.704-714, ISSN: 0085-2538.
- Mayer, AD., Dmitrewski, J., Squifflet, JP., Besse, T., Grabensee, B., Klein, B., Eigler, FW., Heemann, U., Pichlmayr, R., Behrend, M., Vanrenterghem, Y., Donck, J., van Hooff, J., Christiaans, M., Morales, JM., Andres, A., Johnson, RW., Short, C., Buchholz, B., Rehmert, N., Land, W., Schleibner, S., Forsythe, JL., Talbot, D., & Pohanka, E., (1963). Multicenter randomized trial comparing tacrolimus (FK506) and cyclosporine in the prevention of renal allograft rejection. A report of the European tacrolimus multicenter renal study group. *Transplantation* Vol.64, No.3 (August 1997), pp.436-443, ISSN. 0041-1337.
- Mazzaferro, S., Pascquali, M., Taggi, F., Baldinelli, M., Conte, C., Muci, ML., Pirozzi, N., Carbone, I., Francone, M. & Pugliese, F. (2006). Progression of coronary artery calcification in renal transplantation and the role of secondary hyperparathyroidism and inflammation. *Clin J Am Soc Nephrol* Vol.4, No.3 (March 2009), pp.685-690, ISSN: 1555-9041.
- McDonald, SP. & Russ, GR. (1986). Survival of recipients of cadaveric kidney transplants compared with those receiving dialysis treatment in Australia an New Zealand 1991-2001. *Nephrol Dial Transplant*. Vol.17, No.12, (December 2002) pp.2212-2219, ISSN: 0931-0509.
- McGregor, E., Jardine, AG., Murray, LS., Dargie, HJ., Rodger, RS., Junor, BJ., McMillan, MA. & Briggs JD. (1986). Pre-operative echocardiographic abnormalities and adverse outcome following renal transplantation. *Nephrol Dial Transplant*. Vol.13, No.6, (June 1998), pp.1499-505, ISSN: 0931-0509
- McGregor, E., Steward, G., Rodger, SC. & Jardine AG. (1986). Early echocardiographic changes and survival following transplantation. *Nephrol Dial Transplant*. Vol.15, No.1, (January 2000), pp.93-93, ISSN: 0931-0509.
- Meier-Kriesche, HU., Vaghela, M., Thambuganipalle, R., Friedman, G., Jacobs, M. & Kaplan, B. (1963). The effect of body mass index on long-term renal allograft. *Transplantation*. Vol.68, No.9, (November 1999), pp.1294-1297, ISSN: 0041-1337.
- Meier-Kriesche, HU., Arndorfer, JA. & Kaplan, B. (1963). The impact of body mass index on renal transplant outcomes: a significant independent risk factor for graft failure and patient death. *Transplantation*. Vol.73, No.1, (January 2002), pp.70-74, ISSN.0041-1337.

- Meier-Kriesche, HU., Baliga, R. & Kaplan, B. (1963). Decreased renal function is a strong risk factor for cardiovascular death after renal transplantation. *Transplantation* Vol.75, No.8 (April 2003), pp.1291-1205, ISSN: 0041-1337.
- Miller LW. (2001). Cardiovascular toxicities of immunosuppressive agents. *Am J Transplant.* Vol.2, No.9, (October 2002), pp.807- 818, ISSN: 1600-6135.
- Molnar, MZ., Czira, M., Ambrus, C., Kovacs, AC., Pap, J., Remport, A., Szeifer, L. & Mucsi, I. (2001). Anemia is associated with mortality in kidney-transplanted patients- a prospective cohort study. *Am J Transplant.* Vol.7, No.4, (April 2007), pp. 818-824, ISSN.1600-6135.
- Montori, VM., Basu, A., Erwin, PJ., Velosa, JA., Gabriel, SE.& Kudva, YC.(1978). Posttransplantation diabetes. A systematic review of the literature. *Diabetes Care* Vol. 25, No.3, (March, 2002), pp.538-592, ISSN: 0149-5992.
- Moore, R., Hernandez, D. & Valantine, H. (1990). Calcineurin inhibitors and post-transplant hyperlipidaemias. *Drugs Saf* Vol.24, No.10, (2001), pp.755-766.
- Morales, JM. & Dominguez-Gil, B. (1990). Impact of tacrolimus and mycophenolate mofetil combination on cardiovascular risk profile after transplantation. *J Am Soc Nephrol* Vol.17, 12 Suppl 3, (December 2006), pp.S296-S303, ISSN: 1046-6673.
- Morales, JM., Marcén, R., Andrés, A., González- Molina, M., del Castillo, D., Cabello, M., Capdevila, L., Campistol, JM., Oppenheimer, F., Serón, D., Gil-Vernet, S., Lampreave, I., Valdés, F., Anaya, F., Escuin, F., Arias, M., Pallardó, L. & Bustamente J. (1972). Renal transplantation in the modern immunosuppressive era in Spain: four-year results from a multicenter database focus on post-transplant cardiovascular disease. *Kidney Int Suppl.* Vol.111, (December 2008), pp. S94-S99, ISSN: 0085-2538.
- Murphy, SW., Foley, RN. & Parfrey, PS. (1981). Screening and treatment for cardiovascular disease in patients with chronic renal disease. *Am J Kidney Dis.* Vol.32, 5 Suppl 3, (November 1998), pp.S184-S199, ISSN: 0272-6386..
- Ojo, AO., Hanson, JA., Wolfe, RA., Leichtman, AB., Agodoa, LY. & Port FK. (1972). Long-term survival in renal transplant recipients with graft function. *Kidney Int* . Vol. 57, No.1, (January 2000), pp. 307-313, ISSN: 0085-2538.
- Ojo, AO. (1963). Cardiovascular complications after renal transplantation and their prevention. *Transplantation.* Vol.82, No.5, (September 2006), pp.603-611, ISSN: 0041-1337.
- Oliveras, A., Roquer, J., Puig, JM., Rodriguez, A., Mir, M., Orfila, A., Masramon, J. & Loveras, J. (1987). Stroke in renal transplant recipients: epidemiology, predictive risk factors and outcome. *Clin Transplant* Vol. 17, No.1, (February 2003), pp.1-8, ISSN:0902-0063.
- Opelz, G., Döhler, B. for the Collaborative Transplant Study. (2001). Improved long-term outcomes after renal transplantation associated with blood pressure control. *Am J Transplant.* . Vol.5, No.11, (November 2005), pp.2725-2731, ISSN: 1600.6135.
- Opelz, G., Zeier, M., Laux, G., Morath, C. & Döhler, B. (1990). No improvement of patients or graft survival in transplant recipients treated with angiotensin-converting enzyme inhibitors or angiotensin II type 1 receptor blockers: a collaborative transplant study report. *J Am Soc Nephrol* . Vol.17, No.11, (November 2006), pp.3257-3262, ISSN: 1046-6673.
- Paoletti, E., Amadore, M., Cassottana, P., Gherzi, M., Marsano, L. & Cannella, G. (1981). Effects of sirolimus on left ventricular hypertrophy in kidney transplant recipients:

- a 1-year nonrandomized controlled trial. *Am J Kidney Dis* . Vol.52, No.2, (August 2008), pp.324-330, ISSN: 0272-6386.
- Parfrey, PS., Harnett, JD., Foley, RN., Kent, GM., Murray, DC., Barre, PE. & Guttman. RD. (1963). Impact of renal transplantation on uremic cardiomyopathy. *Transplantation* . Vol. 60. No. 9, (November 1995), pp. 908-914, ISSN: 0041- 1337.
- Park, JM. & Luan, FL. (1990). Management of hypertension in solid-organ transplantation. *Prog Transplant*. Vol.15, No.1, (March 2005), pp. 17-22, ISSN: 1526-9248.
- Patel, RK., Mark, PB., Jonhston, N., McGregor, E., Dargie, H. & Jardine AG. (2006). Renal transplantation is not associated with regression of left ventricular hypertrophy: a magnetic resonance study. *Clin J Am Soc Nephrol* . Vol.3, No.6, (November 2008), pp.1807-1811, ISSN: 1555-9041.
- Pilmore, H., Dent, H., Chang, S., McDonald, SP. & Chadban SJ. (1963). Reduction in cardiovascular death after kidney transplantation. *Transplantation*. Vol. 89, No. 7, (April 2010), pp.851-857. ISSN: 0041-1337.
- Ridker, PM., Cushman, M., Stampfer, MJ., Tracy, RP. & Hennekens, CH. (1997). Inflammation , aspirin, and the risk of cardiovascular disease in apparently healthy men. *N Engl J Med* . Vol.336, No.14, (April 1997), pp.973-979, ISSN: 0028-4793.
- Rigatto, C., Foley, RN., Kent, GM., Guttman, R. & Parfrey, PS. (1963). Long-term changes in left ventricular hypertrophy after renal transplantation. *Transplantation* . Vol.70, No.4, (August 2000), pp. 570-575, ISSN: 0041-1337.
- Rigatto, C., Parfrey, P., Foley, R., Negrijn, C., Tribula, C. & Jeffery J. (1990). Congestive heart failure in renal transplant recipients: risk factors, outcomes, and relationship with ischemic heart disease. *J Am Soc Nephrol* . Vol.13, No.4, (April 2002), pp. 1084-1090. ISSN:1046-6673.
- Rigatto, C. (1990). Clinical epidemiology of cardiac disease in renal transplant recipients. *Semin Dial* Vol.16, No.2, (March-April 2003a), pp. 106-110, ISSN: 1525-139X.
- Rigatto, C., Foley, R., Jeffery, J., Negrijn, C., Tribula, C. & Parfrey P. (1990). Electrocardiographic left ventricular hypertrophy in renal transplant recipients: prognostic value and impact of blood pressure and anemia. *J Am Soc Nephrol* Vol. 14. No. 2, (February 2003b), pp.462-468. ISSN:1046-6673.
- Rigatto, C. (2001). Anemia, renal transplantation and the anemia paradox. *Semin Nephrol*. Vol.26, No.4, (July 2006), pp.307- 312.ISSN 0270-9295.
- Roodnat, JL., Mulder, PGH., Rischen-Vos, J., van Riemsdijk, IC., van Gelder, T., Zietse R., Izermans, JN. & Weimar, W. (1963). Proteinuria after renal transplantation affects not only graft survival but also patient survival. *Transplantation*. Vol.72, No.3, (August 2001), pp.438-444, ISSN: 0041-1337.
- Schena, FP., Pascoe, MD., Aberu, J., del Carmen, Rial, M., Oberbauer, R., Brennan, DC., Campistol, JM., Racusen, L., Polinsky, MS., Goldberg-Alberts, R., Li, H., Scarola, J. & Neylan, JF. for the sirolimus CONVERT trial study group. (1963). Conversion from calcineurin inhibitors to sirolimus maintenance therapy in renal allograft recipients: 24-months efficacy and safety results from the CONVERT trial. *Transplantation*. Vol.87, No.2 (January 2009), pp.233-242, ISSN: 0041-1337.
- Seliger, SL., Weiss, NS., Gillen, DL., Kestenbaum, B., Ball, A., Sherrard, DJ. & Stehman-Breen, CO. (1972). HMG-CoA reductase inhibitors are associated with reduced mortality in ESRD patients. *Kidney Int*. Vol.61, No.1, (January 2002), pp.297-304, ISSN:0085-2538.

- Seliger, SL., Gillen, DL., Longstreth, WT., Kestenbaum, B. & Stehman-Breen CO. (1972). Elevated risk of stroke among patients with end-stage renal disease. *Kidney Int.* Vol. 64, No.2, (August 2003), pp.603-609, ISSN:0085-2538.
- Seliger, SL., Gillen, DL., Tirschwell, D., Wasse, H., Kestenbaum, BR. & Stehman-Breen CO. (1990). Risk factors for incident stroke among patients with end-stage renal disease. *J Am Soc Nephrol* Vol. 14, No.10 (October 2003), pp.2623-2631, ISSN: 1046-6673.
- Shah, N., Al-Khoury, S., Afzali, B., Covic, A., Roche, A., Msarsh, J., Macdougall, IC. & Goldsmirhm DJA. (1963). Posttransplantation anemia in adult renal allograft recipients: prevalence and predictors. *Transplantation* Vol.81, No.8 (April 2006), pp.1112-1118, ISSN.0041-1337.
- Snyder, JJ., Kasiske, BL. & Mac Lean, R. (1990). Peripheral arterial disease and renal transplantation. *J Am Soc Nephrol* Vol.17, No.7, (July 2006), pp.2056-2068, ISSN: 1046-6673.
- Soveri, I., Holdaas, H., Jardine, A., Gimpelewicz, C., Staffler, B. & Fellström, B. (1986). Renal transplant dysfunction-importance quantified in comparison with traditional risk factors for cardiovascular disease and mortality. *Nephrol Dial Transplant* Vol.21, No.8, (August 2006), pp.2282-2289. ISSN:0931-0509.
- Stack, AG. & Bloembergen, WE. (1981). A cross-sectional study of the prevalence and clinical correlates of congestive heart failure among incident U.S. dialysis patients. *Am J Kidney Dis* Vol. 38, No. 5, (November 2001), pp. 992-1000, ISSN: 0272- 6386.
- Stuveling, EM., Hillege, HL., Bakker, SJ., Gans, RO., De Jong, PE. & de Zeeuw, D. (1972). C-reactive protein is associated with renal function abnormalities in a non-diabetic population. *Kidney Int* . Vol.63, No.2, (February 2003), pp.654-661, ISSN: 0085-2538.
- Sung, RS., Althoen, M., Howell, TA. & Merion, RM. (1963). Peripheral vascular occlusive disease in renal transplant recipients: risk factors and impact on kidney allograft survival. *Transplantation*. Vol.70, No.7, (October 2000), pp. 1049-1054, ISSN: 0041-1337.
- Teruel, JL., Rodriguez Padial, L., Quereda, C., Yuste, P., Marcen, R. & Ortuño, J. (1963). Regression of left ventricular hypertrophy after renal transplantation. A prospective study. *Transplantation*. Vol.43, No.2, (February 1987), pp.307-309, ISSN: 0041-1337
- Tucker, B., Fabbian, F., Giles, M., Thuraisingham, RC., Raine, AE. & Baker LR. (1997). Left ventricular hypertrophy and ambulatory blood pressure monitoring in chronic renal failure. *Nephrol Dial Transplant* Vol.12, No.4, (April 1997), pp. 724-728, ISSN: 0272-6376.
- Tutone, VK., Mark, PB., Steward, GA., Tan, CC., Geddes, CC. & Jardine AG. (1987). Hypertension, antihypertensive agents and outcomes following renal transplantation. *Clin Transplant* Vol.19, No. 2, (April 2005), pp.181-192, ISSN: 0902-0063.
- Van Ree, RM., de Vries, APJ., Oterdoom ,LH., The, TH., Gansevoort, RT., van der Heide, JJH., van Son, WJ., Ploeg, RJ., de Jong, PE., Gans, ROB. & Bakker, SJL. (1986). Abdominal obesity and smoking are important determinants of C-reactive protein in renal transplant recipients. *Nephrol Dial Transplant* . Vol.20, No.11 (November 2005), pp.2524-2531, ISSN: 0931-0509.
- Vanrenterghem, Y., Ponticelli, C., Morales, JM., Abramowicz, D., Baboolal, K., Eklund, B., Kliem, V., Legendre, C., Morais Sarmiento, AL. & Vincenti, F. (2001). Prevalence

- and management of anemia in renal transplant recipients: a European survey. *Am J Transplant*. Vol.3, No.7, (July 2003), pp.835-845, ISSN.1600-6135.
- Vanrenterghem, YFC., Claes, K., Montagnino, G., Fieuws, S., Maes, B., Villa, M. & Ponticelli, C. (1963). Risk factors for cardiovascular events after successful renal transplantation. *Transplantation*. Vol. 85, No.2, (January 2008), pp. 209-216, ISSN: 0041-1337.
- Vincenti, F., Jensik, SC., Filo, RS., Millar, J. & Pirsch, J. (1963). A long-term comparison of tacrolimus (FK506) and cyclosporine in kidney transplantation: evidence for improved allograft survival at five years. *Transplantation* Vol.73, No.5 (March 2005), pp. 775-782. ISSN. 0041-1337.
- Vincenti, F., Friman, S., Scheuermann, E., Rostaing, L., Jenssen, T., Campistol, JM., Uchida, K., Pescovitz, MD., Marchetti, P., Tuncer, M., Citterio, F., Wiecek, A., Chadban, S., El-Shahawy, M., Budde, K. & Goto, N. on behalf of the DIRECT investigators. (2001). Results of an international, randomised trial comparing glucose metabolism disorders and outcome with cyclosporine versus tacrolimus. *Am J Transpl* Vol.7, No.6, (June 2007), pp. 1506-1514, ISSN.1600-6135.
- Vincenti, F., Charpentier, B., Vanrenterghen, Y., Rostaing, L., Bresnahan, B., Darji, P., Massari, P., Mondragon-Ramirez, GA., Agarwal, M., Di Russo, G., Lin, CS., Garg, P. & Larsen, CP. (2001). A phase III study of belatacept-based immunosuppression regimens versus cyclosporine in renal transplant recipients (BENEFIT study). *Am J Transplant*. Vol.10, No.3, (March 2010), pp.535-546, ISSN: 1600-6135.
- Wali, RK., Wang, GS., Gottlieb, SS., Bellumkonda, L., Hansalia, R., Ramos, E., Drachenberg, C., Papadimitriou, J., Brisco, MA., Blahut, S., Fink, JC., Fisher, ML., Bartlett, ST. & Weir MR. (1983). Effect of kidney transplantation on left ventricular systolic dysfunction and congestive heart failure in patients with end-stage renal disease. *J Am Coll Cardiol*, Vol. 45, No.7 (April 2005), pp.1051-1060, ISSN: 0735-1097.
- Wang, K., Zhang, H., Li, Y., Wei, Q., Li, H., Yang, Y. & Lu, Y. (1969). Safety of mycophenolate mofetil versus azathioprine in renal transplantation: a systematic review. *Transplant Proc*. Vol.36, No7, (September 2004), pp.2068-2070, ISSN.0041-1345.
- Wang, JH. & Kasiske, BL. (1992). Screening and management of pretransplant cardiovascular disease. *Curr Opin Nephrol Hypertens*. Vol.19, No.6, (November 2010), pp.586-591, ISSN: 1473-6543.
- Webster, AC., Lee, VWS., Chapman, JR. & Craig, JC. (1963). Target of rapamycin inhibitors (sirolimus and everolimus) for primary immunosuppression of kidney transplant recipients; a systematic review meta-analysis of randomized trials. *Transplantation* Vol.81, No.9, (May 2006), pp. 1234-1248, ISSN: 0041-1337.
- Wiesbauer, F., Heinze, G., Mitterbauer, C., Harnoncourt, F., Hörl, WH. & Oberbauer, R. (1990). Statin use is associated with prolonged survival of renal transplant recipients. *J Am Soc Nephrol* Vol.19, No.11, (November 2008), pp. 2211-2218, ISSN: 1046-6673.
- Wilson, PWF. & Culleton, BF. (19981). Epidemiology of cardiovascular disease in the United States. *A J Kidney Dis*. Vol 32, No5, suppl 3. (November 1998), pp.S56-S65, ISSN: 0272-6386.
- Winkelmayer, WC., Lorenz, M., Kramar, R., Fodinger, M., Hörl, WH. & Sunder-Plassmann, G. (2001). C-reactive protein and body mass index independently predict mortality in kidney transplant recipients. *Am J Transplant*. Vol.4, No.7, (July 2004), pp.1148-1154, ISSN: 1600-6135.

- Winkelmayr, WC., Kramar, R., Curhan, GC., Chandraker, A., Endler, G., Föding, M., Hörl, WH. & Sunder Plassmann, G. (1990). Fasting plasma total homocysteine levels and mortality and allograft loss in kidney transplant recipients: a prospective study. *J Am Soc Nephrol*. Vol.16, No.1, (January 2005), pp.255-260, ISSN: 1046-6673.
- Winkelmayr, WC., Chandraker, A., Brookhart, MA., Kramar, R. & Sunder-Plassmann, G. (1986). A prospective study of anaemia and long-term outcomes in kidney transplant recipients. *Nephrol Dial Transplant*. Vol.21, No.12, (December 2006), pp.3559-3566, ISSN.0931-0509.
- Wolfe, RA., Ashby, VB., Milford, EL., Ojo, AO., Ettenger, RE., Agodoa, LY., Held, PJ. & Port, FK. (1999). Comparison of mortality in all patients on dialysis, patients on dialysis awaiting transplantation, and recipients of a first cadaveric transplant. *N Engl J Med*. Vol. 341, No.23 (December 1999), pp.1725-1730, ISSN: 0028-4793.
- Wong, BM., Huang, M., Zaltzman, JS. & Prasad, GV. (1993). Mycophenolate mofetil and C-reactive protein in renal transplant recipients. *Transplantation*. Vol.83, No.1, (January 2007), pp.48-53, ISSN: 0041-1337.
- Yorgin, PD., Scandling, JD., Belson, A., Sanchez, J., Alexander, SR. & Andreoni, KA. (2001). Late post-transplant anemia in adult renal transplant recipients. An under-recognized problem? *Am J Transplant*. Vol.2, No.5, (May 2002), pp.429-435, ISSN.1600-6135.
- Zhang, R., Leslie, B., Boudreaux, JP., Frey, D. & Reisin E. (1997). Hypertension after kidney transplantation: impact, pathogenesis and therapy. *Am J Med Sci* Vol: 325, No.4, (April 2003), pp.202-208, ISSN: 1538-2990.
- Zebe H. (1986). Atrial fibrillation in dialysis patients. *Nephrol Dial Transplant*. Vol. 16, No.5, (June 2001), pp. 765-8, ISSN: 0931- 0509.
- Zitt, N., Kollerits, B., Neyer, U., Mark, W., Heininger, D., Mayer, G., Kronenberg, F., & Lhotta, K. (1986). Cigarette smoking and chronic allograft nephropathy. *Nephrol Dial Transplant*. Vol.22, No.10, (October 2007), pp.3034-3034, ISSN. 0931-0509

IntechOpen

Understanding the Complexities of Kidney Transplantation

Edited by Prof. Jorge Ortiz

ISBN 978-953-307-819-9

Hard cover, 564 pages

Publisher InTech

Published online 06, September, 2011

Published in print edition September, 2011

Kidney transplantation is a complex field that incorporates several different specialties to manage the transplant patient. This book was created because of the importance of kidney transplantation. This volume focuses on the complexities of the transplant patient. In particular, there is a focus on the comorbidities and special considerations for a transplant patient and how they affect kidney transplant outcomes. Contributors to this book are from all over the world and are experts in their individual fields. They were all individually approached to add a chapter to this book and with their efforts this book was formed. Understanding the Complexities of Kidney Transplantation gives the reader an excellent foundation to build upon to truly understand kidney transplantation.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Roberto Marcén and Sara Jiménez (2011). Cardiovascular Diseases in Kidney Transplantation, Understanding the Complexities of Kidney Transplantation, Prof. Jorge Ortiz (Ed.), ISBN: 978-953-307-819-9, InTech, Available from: <http://www.intechopen.com/books/understanding-the-complexities-of-kidney-transplantation/cardiovascular-diseases-in-kidney-transplantation>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2011 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike-3.0 License](https://creativecommons.org/licenses/by-nc-sa/3.0/), which permits use, distribution and reproduction for non-commercial purposes, provided the original is properly cited and derivative works building on this content are distributed under the same license.

IntechOpen

IntechOpen