We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Climate Change in Spain: Phenological Trends in Southern Areas

García-Mozo, H.¹, Mestre, A.² and Galán, C.¹

¹*Dept. Botany, Ecology and Plant Physiology. University of Cordoba, Cordoba*

²*Agencia Estatal de Meteorología, Madrid
Spain*

1. Introduction

Global average surface temperatures increased by $0.6 \pm 0.2^\circ\text{C}$ over the twentieth century, and a further increase of 1.4 to 5.8°C is projected by 2100 (Figure 1). Europe recorded its warmest decade from 1990 to 1999 (IPCC, 2001; IPCC, 2007). The rate of warming during this period has been approximately double that the previous one and, greater than at any other time during the last 1,000 years. Organisms, populations and ecological communities do not, however, respond to approximated global averages. Rather, regional changes, which are highly spatially heterogeneous (Figure. 1), are more relevant in the context of ecological response to climatic change. In many regions there is an asymmetry in the warming that undoubtedly will contribute to heterogeneity in ecological dynamics across systems. Diurnal temperature ranges have decreased because minimum temperatures are increasing at about twice the rate of maximum temperatures. As a consequence, the frost-free periods in most mid- and high-latitude regions are lengthening and satellite data reveal a 10% decrease in snow cover and ice extent since the late 1960s. Changes in the precipitation regime have also been neither spatially nor temporally uniform (Figure 1). In the mid- and high latitudes of the Northern Hemisphere a decadal increase of 0.5–1% mostly occurs in autumn and winter whereas, in the sub-tropics, precipitation generally decreases by about 0.3% per decade (IPCC, 2007).

The impact of climate change, and particularly of climate warming, is being tracked in many physical and biological systems. Phenology – the timing of seasonal activities of animals and plants – is perhaps the simplest process in which to track changes in the ecology of species in response to climate change. Birds, butterflies and wild plants, in particular, include popular and easily identifiable species and thus have received considerable attention from the public. Plant phenology is seen as one of the most important bio-indicators, since trends can provide considerable temporal and spatial information regarding ongoing changes (Menzel et al., 2006). As a result many long-term phenological data sets have been collected. Studies in Europe and North America have revealed phenological trends that very probably reflect responses to recent climate change. Common changes in the timing of spring activities include earlier breeding or first singing of birds, earlier arrival of migrant birds, earlier appearance of butterflies, earlier choruses and spawning in amphibians and earlier shooting and flowering of plants. In general, spring activities have occurred progressively

earlier since the 1960s. Plant reproductive phenology is controlled by temperature, especially in tree species, and precipitation and photoperiod, especially in herbaceous ones (García-Mozo et al., 2000; 2009; Galán et al., 2005). It has been shown that plants flowering early in spring are more affected by warming than species flowering later in the year (Ahas et al., 2002; García-Mozo et al., 2002; Galán et al., 2005).

Source: IPCC 2007.

Fig. 1. Spatial variability of annual trends in temperature and precipitation since 1976 relative to 1961 to 1990 normals. a, Temperature (°C per decade); b, precipitation (% per decade).

In Spain, temperatures, especially minimum temperature, have increased over the last century by around 1.5°C the annual average (Fernández-González et al., 2005). Climate change has had a particularly marked effect on southern Spain, increasing temperatures and reducing rainfall. Rainfall has also become increasingly torrential in recent years (Pita, 2003; De Castro et al., 2005).

Most long-term phenological studies to date have focussed on northern Europe, while comparatively few have addressed the Mediterranean region (Gordo & Sanz, 2005; Peñuelas et al., 2002). The projections proposed by the Intergovernmental Panel on Climate Change (IPCC) and confirmed by the Spanish Meteorology Agency (AEMET) for the future Spanish climate, point to a continuing increase in temperatures and a decrease in annual rainfall (IPCC, 2007).

The present study, carried out by the University of Córdoba Aerobiology Research Group and the AEMET, sought to chart phenological trends in response to climate in the Andalusia region of southern Spain, and to compare *in-situ* phenological data with airborne pollen counts in the same area. For this purpose, special attention was paid to the reproductive phenophases of anemophilous species.

2. Materials and methods

2.1 Phenological data

Andalusia boasts considerable climatic and topographical diversity. Regular phenological data series dating back to the mid-1980s from four sites were selected for this study, on the basis of unbroken data series availability, presence of anemophilous species and distance from pollen traps: Bujalance and Pozoblanco, in the province of Cordoba, central Andalusia; Charcones and Huescar, in the province of Granada, a more elevated area of eastern Andalusia (Table 1).

Province	Localities	Distance	Altitude	Coordinates.	T°	Rf
Córdoba	Córdoba		123	37° 50' N, 4° 45' W	18.0	674
	Bujalance	42	222	37° 54' N, 4° 23' W	17.5	550
	Pozoblanco	86	649	38° 23' N, 4° 51' W	16.0	505
Granada	Granada		685	37° 11' N, 3° 35' W	15.5	462
	Charcones	62	1280	37° 40' N, 2° 94' W	13.9	300
	Huescar	100	1115	37° 48' N, 2.33' W	14.8	367
Province	Localities	Distance	Altitude	Coordinates.	T°	Rf
Córdoba	Córdoba		123	37° 50' N, 4° 45' W	18.0	674
	Bujalance	42	222	37° 54' N, 4° 23' W	17.5	550
	Pozoblanco	86	649	38° 23' N, 4° 51' W	16.0	505
Granada	Granada		685	37° 11' N, 3° 35' W	15.5	462
	Charcones	62	1280	37° 40' N, 2° 94' W	13.9	300
	Huescar	100	1115	37° 48' N, 2.33' W	14.8	367

Table 1. Characteristics of the study sites including distance to the province capital (km) where pollen monitoring was performed, altitude (m.a.s.l.), latitude & longitude coordinates, annual mean temperature (T°) and annual mean cumulated rainfall (Rf).

The following species were studied in the indicated dates: 2 oak species: holm oak (*Quercus ilex* subsp. *ballota* (Desf.) Samp.) from 1989 to 2008 and gall oak (*Quercus faginea* Lam.) from 1989 to 2000; olive tree (*Olea europaea* L.) from 1986 to 2008; grape (*Vitis vinifera* L.) from 1986 to 2007; and 3 Poaceae crop species: rye (*Secale cereale* L.) from 1989 to 2008, wheat (*Triticum vulgare* L.) from 1987 to 2008, and common barley (*Hordeum vulgare* L.) from 1987 to 2008. Studied periods varied depending on the site and phenophase.

The analysed species were chosen given their ecological or agricultural importance to the region. Typical vegetation of southern Spain is basically formed by Mediterranean forest and dehesas where the main arboreal genus is *Quercus* (estenopalynous genus), and specially holm-oak. This species is the first flowering oak in Andalusia, being the main species contributing to the pollen curve where there are included pollen grains from all the *Quercus* species (García-Mozo et al., 2002; Gómez-Casero et al., 2004; Gómez-Casero et al., 2007). Apart from its ecological importance, in Iberian Mediterranean ecosystems holm-oak acorn production is of vital economic importance due to acorns are the major component in the feeding systems of high-quality Iberian domestic pigs. Regards to olive tree, Spain produces 33% of the world's olive oil, and Andalusia accounts for 80% of total Spanish output being the world's leading olive-oil-producing region. The largest olive-growing areas are concentrated in the provinces of Jaén and Cordoba. Another economically important species analysed in this work is the grape. Spain is the third worldwide winemaker country counting with the largest surface devoted to vineyards, being Andalusia the sixth region in Spain. The analysed variety was "Garnacha", one of the most abundant in the region, which has a medium range flowering time. Finally, although cultivated grass species contribute in a lower percentage than wild species to the grass pollen curve (estenopalynous family), the analysed grass species selected for the present work were indicated as the crop species more represented in the Poaceae airborne pollen curve in Spain (Muñoz et al., 2000).

The following phenophases, taken directly at the field, were analysed once a week: flowering (appearance of first flowers in some individuals), fruit ripening (several ripe fruits present in several individuals), leaf unfolding (appearance of first leaves in some individuals) and leaf falling (branches have lost half their leaves).

Daily airborne pollen counts were monitored in the cities of Córdoba and Granada (Table 1). Hirst volumetric traps were used (Hirst, 1952). Daily samplings were stained and analysed by using optical microscopy following the rules laid down by the Spanish Aerobiological Network (REA) (Galán et al., 2007, http://www.uco.es/rea/infor_rea/manual_eng.pdf). Two important dates of the pollen season were analysed in the present work as indicators of the flowering period from the aerobiological data. The Pollen season Start date (PS) defined as the first day on which 1 pollen grain/m³ was recorded and the 5 following days recorded 1 or more pollen grains/m³ (García-Mozo et al., 2000), and the Pollen season Peak day (PP) defined as the day when the maximum concentration was recorded.

2.2 Climate data

Daily temperature and rainfall data (1986-2008) from the study sites were used to evaluate the impact of climate on phenology, these variables being considered the most relevant for this purpose (Peñuelas et al., 2002; Gordo & Sanz, 2005). The warming index, anomalies from average temperature, for both cities from the start of the 20th century (Figure 2) reveals a progressive rise in temperature, especially over recent years. During the study period (1986-2008), mean annual temperature increased by 4°C in Córdoba and 3.2°C in Granada.

Fig. 2. Warming Index, anomalies from average temperature, in Cordoba (a) and Granada (b) from 1917 to 2005.

2.3 Statistical analysis

Phenological events and weather data were subjected to a Spearman correlation test. Linear regression analysis was performed to calculate trend lines where R^2 values were given and also p values in order to estimate the significance of these trend lines. The Excel Microsoft

2007 and STATISTICA 6.0 software packages were used. Results were considered significant at a 95% confidence interval.

3. Results

Phenological trends for the study species are shown in Table 2. Through the studied period, the flowering dates of holm oak varied up to 60 days, with a significant average advance of 1.6 days per year (27 days during the studied period). The PP in Córdoba city better matched field data at study sites in Córdoba province than the PS date. Although the PP trend line does not show a significant statistics due to the high variability in the data, both aerobiological dates are occurring increasingly earlier, with an overall advance of 7 and 12 days respectively over the period. In this species, fruit ripening advanced by 24 days at Pozoblanco and 36 at Huescar (significant trend gradient, $p < 0.05$), leaf unfolding advanced by 25 days in *Quercus faginea* at Huescar (significant trend gradient).

In situ observations of olive flowering advanced at the three studied sites, being significant in the two last ones (Table 2): Pozoblanco (6 days), Bujalance (40 days) and Huescar (40 days). Airborne pollen data suggested a less marked advance, nevertheless the PP curve better matched field data at study sites in Córdoba province than the PS dates, except for the years 1998 to 2001 in Pozoblanco. In the case of Huescar (Granada) phenological data trend was more similar to PS dates. Also olive fruit ripening advanced significantly in the sites located in the province of Córdoba although no significant trend gradient was revealed in Huescar (Granada).

Over the period 1989-2000, *Vitis vinifera* leaf unfolding advanced an average of 10 days, and leaf life was extended up to 20 days in the case of Huescar; although an advance in the fruit ripening time was observed in Charcones (40 days) there was no significant change in the timing of fruit ripening in Huescar. Regarding herbaceous species, *Secale cereale* flowering advanced an average of 16 days between 1989 and 2006 at the Granada province sites. Field data suggested a great delay with regard to pollen dates, although the trend was similar and significant except for the Start date values. Fruit ripening in *Secale cereale* did not display any noteworthy trend between 1989 and 2008. Results for *Triticum vulgare* in Córdoba province revealed a general advance in flowering, significant in the case of field data of Pozoblanco whereas no significant trends were detected in fruit ripening. Finally phenological data for *Hordeum vulgare* indicated a flowering delay of 7 days in Pozobanco and 31 days in Bujalance. By contrast, flowering in Charcones (Granada) showed a slight advance and fruit ripening at all sites appeared largely unchanged over the study period. It is noticeable that the herbaceous species showed the less coincident trends of field phenological data and aerobiological data and the lower significance values for both types of data. In the case of *T. vulgare* and *H. vulgare* the aerobiological phase most coinciding with field phenological data was the PS curve.

Spearman analysis results showed in Table 3 indicated a negative correlation between various cumulative sums of average daily temperature and most of the flowering and fruiting events, especially in tree species. *Vitis vinifera* leaf fall displayed significant positive correlation with temperature (Tables 3b and 3c). Temperature over the preceding autumn correlated positively with most phenological phases in tree species. For some herbaceous species, there was also some correlation with temperature over the preceding months, as well as a weak but still-significant correlation with rainfall over previous months.

Site	Species	Phenophase	Gradient	R ²	p	N
Cordoba	<i>Quercus</i> spp.	Pollen season start	-0.77	0.21	0.05	17
		Pollen season peak	-0.41	0.04	0.41	17
	<i>Olea europaea</i>	Pollen season start	-0.31	0.03	0.49	20
		Pollen season peak	-0.34	0.04	0.38	20
	Poaceae	Pollen season start	-0.42	0.02	0.45	22
		Pollen season peak	-0.9	0.15	0.07	22
Pozoblanco	<i>Quercus ilex</i>	Flowering	-1.58	0.24	0.04	17
		Fruit ripening	-1.17	0.15	0.08	20
	<i>Olea europaea</i>	Flowering	-2.04	0.18	0.06	20
		Fruit ripening	-1.25	0.72	0.03	22
	<i>Triticum vulgare</i>	Flowering	-0.71	0.33	0.01	21
		Fruit ripening	-0.24	0.06	0.51	21
	<i>Hordeum vulgare</i>	Flowering	0.49	0.10	0.17	20
		Fruit ripening	0.07	0.00	0.74	21
Bujalance	<i>Olea europaea</i>	Flowering	-1.63	0.37	0.00	20
		Fruit ripening	-1.86	0.81	0.00	22
	<i>Vitis vinifera</i>	Leaf fall	0.16	0.00	0.74	15
	<i>Triticum vulgare</i>	Flowering	-0.12	0.00	0.74	21
		Fruit ripening	-0.13	0.05	0.32	21
	<i>Hordeum vulgare</i>	Flowering	1.59	0.55	0.00	21
		Fruit ripening	-0.14	0.00	0.20	21
Granada	<i>Olea europaea</i>	Pollen season start	0.43	0.19	0.07	19
		Pollen season peak	-0.08	0.00	0.57	19
	Poaceae	Pollen season start	-0.13	0.00	0.88	16
		Pollen season peak	-1.70	0.29	0.04	16
Huescar	<i>Quercus faginea</i>	Foliation	-2.30	0.46	0.01	11
		Fruit ripening	-1.79	0.84	0.00	20
	<i>Olea europaea</i>	Flowering	-1.99	0.66	0.00	19
		Fruit ripening	0.07	0.00	0.00	22
	<i>Vitis vinifera</i>	Foliation	-0.40	0.31	0.07	11
		Leaf fall	1.16	0.46	0.01	15
		Fruit ripening	-0.25	0.09	0.19	19
	<i>Secale cereale</i>	Flowering	-0.99	0.92	0.00	16
		Fruit ripening	-0.04	0.00	0.70	20
	<i>Hordeum vulgare</i>	Flowering	-0.29	0.06	0.32	20
		Fruit ripening	-0.18	0.00	0.87	20
Charcones	<i>Vitis vinifera</i>	Foliation	-1.26	0.17	0.19	11
		Fruit ripening	-0.09	0.00	0.49	19
	<i>Secale cereale</i>	Flowering	-0.95	0.42	0.01	15
		Fruit ripening	-0.01	0.00	0.99	20
	<i>Hordeum vulgare</i>	Flowering	-0.29	0.06	0.32	20
		Fruit ripening	-0.18	0.00	0.87	20

Table 2. Trends for different phenological stages. Site, species, phenological stages, gradient, R2, significance (p value) and number of years (N) are represented.

	Phase	Tmx	T	Tmn	T _{j-a}	T _{j-jn}	T _{s-n}	T _{s-n} ⁻¹	Rf	Rf _{j-a}	Rf _{j-jn}	Rf _{s-n}	Rf _{s-n} ⁻¹
<i>Quercus ilex</i>	Fl	-0.35	0.22	0.03	0.18	0.07		0.50	0.29	-0.10	0.03		-0.12
	PS	-0.12	-0.20	-0.14	-0.08	-0.16		-0.15	0.17	-0.19	-0.33		-0.46
	PP	-0.24	-0.38	-0.27	-0.23	-0.24		0.15	0.03	0.01	0.00		-0.47
	FR	-0.62	-0.41	-0.34	-0.26	-0.54	-0.15	0.34	-0.35	-0.46	-0.28	-0.42	-0.13
<i>Olea europaea</i>	Fl	-0.38	-0.22	-0.33	-0.20	-0.44		0.37	-0.29	-0.55	-0.49		-0.48
	PS	-0.53	-0.53	-0.54	-0.38	-0.51		-0.14	0.19	-0.24	-0.32		-0.62
	PP	-0.46	-0.17	-0.09	0.10	-0.35		0.09	0.22	0.32	0.33		0.08
	FR	-0.54	-0.47	-0.37	-0.31	-0.35	-0.01	0.17	-0.37	-0.56	-0.46	-0.08	-0.21
<i>Triticum vulgare</i>	Fl	-0.28	-0.21	-0.02	0.02	-0.15		0.47	0.20	-0.26	-0.10		-0.25
	PS	0.18	0.14	-0.04	-0.15	-0.05		-0.14	-0.26	-0.25	-0.33		-0.02
	PP	-0.19	-0.08	-0.04	0.19	0.06		-0.08	0.17	-0.02	0.04		0.09
	FR	-0.21	0.28	0.04	-0.05	-0.19		-0.05	-0.05	-0.04	0.01		-0.02
<i>Hordeum vulgare</i>	Fl	0.29	-0.02	-0.11	0.29	-0.03		0.03	0.30	0.23	0.07		-0.18
	PS	0.18	0.14	-0.04	-0.15	-0.05		-0.14	-0.26	-0.25	-0.33		-0.02
	PP	-0.19	-0.08	-0.04	0.19	0.06		-0.08	0.17	-0.02	0.04		0.09
	FR	-0.02	0.35	0.16	0.16	-0.09		-0.05	0.06	-0.04	0.09		0.21

Table 3. a.

	Phase	Tmx	T	Tmn	T _{j-a}	T _{j-jn}	T _{s-n}	T _{s-n} ⁻¹	Rf	Rf _{j-a}	Rf _{j-jn}	Rf _{s-n}	Rf _{s-n} ⁻¹
<i>Olea europaea</i>	Fl	-0.40	-0.45	-0.32	-0.23	-0.37		0.26	0.17	0.04	0.04		0.24
	PS	-0.17	-0.16	-0.06	-0.08	-0.18		-0.30	0.02	0.22	0.11		0.01
	PP	-0.13	0.03	0.32	0.21	0.22		0.30	0.16	0.05	0.11		-0.02
	FR	-0.57	-0.62	-0.38	-0.32	-0.38	-0.56	0.56	0.24	0.09	0.06	0.13	0.11
<i>Triticum vulgare</i>	Fl	0.04	-0.11	-0.35	-0.30	-0.30		-0.18	-0.27	-0.24	-0.20		0.23
	PS	-0.21	-0.27	-0.24	0.06	-0.23		-0.33	-0.11	-0.25	-0.30		0.22
	PP	-0.13	-0.02	0.21	0.12	0.09		0.37	0.15	-0.20	-0.12		0.03
	FR	-0.17	-0.02	-0.29	-0.03	-0.09		0.21	-0.33	-0.11	-0.19		0.17
<i>Hordeum vulgare</i>	Fl	-0.38	-0.33	-0.07	-0.12	-0.22		-0.32	-0.11	0.11	-0.02		-0.04
	PS	-0.21	-0.27	-0.24	0.06	-0.23		-0.33	-0.11	-0.25	-0.30		0.22
	PP	-0.13	-0.02	0.21	0.12	0.09		0.37	0.15	-0.20	-0.12		0.03
	FR	-0.05	0.09	-0.29	-0.25	-0.19		0.51	-0.05	0.28	-0.08		0.07
<i>Vitis vinifera</i>	LF	0.63	0.34	0.16	0.09	0.04	-0.22	0.09	0.21	0.06	0.03	0.30	0.10

Table 3. b.

	Phase	Tmx	T	Tmn	Tj-a	Tj-jn	Ts-n	Ts-n ⁻¹	Rf	Rfj-a	Rfj-jn	Rfs-n	Rfs-n ⁻¹
<i>Quercus faginea</i>	FO	-0.33	-0.40	-0.41	-0.75			0.17	-0.33	-0.13			0.03
<i>Quercus ilex</i>	PS	-0.11	-0.08	-0.04	-0.10	-0.33		0.01	0.33	0.07	-0.05		-0.61
	PP	-0.52	-0.42	-0.25	0.02	-0.22		-0.07	0.48	0.52	0.44		0.29
	FR	-0.70	-0.72	-0.64	-0.58	-0.69	-0.60	-0.31	-0.16	-0.08	-0.12	-0.03	-0.28
	FI	-0.74	-0.71	-0.59	-0.73	-0.78		-0.18	-0.25	-0.08	-0.01		-0.05
<i>Olea europaea</i>	PS	0.09	0.15	0.18	-0.01	-0.39		0.33	0.47	0.28	0.36		-0.47
	PP	-0.08	-0.27	-0.47	-0.36	-0.27		-0.36	-0.34	-0.18	-0.12		-0.26
	FR	-0.10	-0.04	0.06	0.12	-0.03	0.15	0.66	0.20	0.24	0.33	-0.31	0.12
	FI	-0.59	-0.57	-0.44	-0.63	-0.67		-0.46	0.03	0.02	0.03	-0.02	-0.17
<i>Secale cereale</i>	PS	-0.36	-0.44	-0.46	-0.26	-0.19		-0.39	-0.34	-0.41	-0.43	0.08	0.21
	PP	-0.17	-0.31	-0.44	-0.36	-0.16		-0.27	-0.23	-0.28	-0.33	0.13	-0.25
	FR	-0.52	-0.42	-0.25	0.02	-0.22		-0.07	0.48	0.52	0.44	0.20	0.29
	FI	-0.59	-0.57	-0.44	-0.63	-0.67		-0.46	0.03	0.02	0.03	-0.02	-0.17
<i>Vitis vinifera</i>	PS	-0.36	-0.44	-0.46	-0.26	-0.19		-0.39	-0.34	-0.41	-0.43	0.08	0.21
	PP	-0.17	-0.31	-0.44	-0.36	-0.16		-0.27	-0.23	-0.28	-0.33	0.13	-0.25
	FR	-0.52	-0.42	-0.25	0.02	-0.22		-0.07	0.48	0.52	0.44	0.20	0.29
<i>Vitis vinifera</i>	LF	0.27	0.21	0.09	0.18	0.20	0.25	-0.39	-0.18	-0.42	-0.45	-0.18	0.17
	FR	-0.32	-0.30	-0.24	-0.14	-0.39	-0.16	0.23	0.00	0.11	0.12	-0.24	-0.25
	FO	-0.63	-0.62	-0.48	-0.53	-0.57		0.36	-0.07	0.05	0.22		-0.04

Table 3. c.

	Phase	Tmx	T	Tmn	Tj-a	Tj-jn	Ts-n	Ts-n ⁻¹	Rf	Rfj-a	Rfj-jn	Rfs-n	Rfs-n ⁻¹
<i>Hordeum vulgare</i>	FI	-0.01	-0.20	-0.38	-0.50	-0.24		0.17	0.31	0.12	0.33		0.28
	PS	0.21	-0.01	-0.23	0.05	0.03		0.21	0.10	0.31	0.11		0.09
	PP	-0.03	-0.07	-0.08	-0.13	0.02		0.16	0.10	-0.13	0.31		0.34
	FR	-0.37	-0.43	-0.39	-0.38	-0.39		-0.30	0.42	0.38	0.22		0.28
<i>Triticum vulgare</i>	FI	-0.81	-0.80	-0.70	-0.28	-0.82		-0.03	-0.04	0.37	0.14		0.26
	PS	0.21	-0.01	-0.23	0.05	0.03		0.21	0.10	0.31	0.11		0.09
	PP	-0.03	-0.07	-0.08	-0.13	0.02		0.16	0.10	-0.13	0.31		0.34
	FR	-0.11	-0.19	-0.23	0.12	-0.35		-0.08	0.52	0.34	0.34		0.06
<i>Secale cereale</i>	FI	-0.36	-0.45	-0.43	-0.07	-0.16		0.17	0.14	-0.05	0.11		0.11
	PS	0.21	-0.01	-0.23	0.05	0.03		0.21	0.10	0.31	0.11		0.09
	PP	-0.03	-0.07	-0.08	-0.13	0.02		0.16	0.10	-0.13	0.31		0.34
	FR	-0.36	-0.41	-0.36	-0.25	-0.39		-0.24	0.43	0.40	0.19		0.19
<i>Vitis vinifera</i>	FR	-0.29	-0.24	-0.17	0.39	0.12	-0.77	-0.53	0.12	0.19	0.05	0.33	0.11
	FO	-0.32	-0.09	-0.17	-0.46	-0.32		-0.04	0.60	-0.19	0.27		-0.01

Table 3. d.

Phases: FI, flowering; PP, Pollen Start; PP, Pollen Peak; FR, Fruit Ripening; FO, Foliation; LF, Leaf Fall.
Meteorological factors: Tmx, Annual Maximum Temperature; T, Annual Mean Temperature; Tmn, Annual Minimum Temperature; Tj-a, Mean Temperature from January to April; Tj-jn, Mean Temperature from January to June; Ts-n, Mean Temperature from September to November; Ts-n-1, Mean Temperature from previous September to November; Rf, Annual Cumulated Rainfall; Rfj-a, Cumulated Rainfall from January to April; Rfj-jn, Cumulated Rainfall from January to June; Rfs-n, Cumulated Rainfall from September to November; Rfs-n-1, Cumulated Rainfall from previous September to November.

Table 3. Spearman correlation values for each species and phenophases in Pozoblanco (a), Bujalance (b), Huescar (c) and Charcones (d). p values < 0.05 are highlighted in bold.

4. Discussion

Although with some delay, probably due to the distance between field phenological stations and aerobiological stations, field phenological and aerobiological data showed similar patterns in the study species, especially in tree species. The higher differences in the herbaceous species patterns are probably influenced because cultivated grass species (as the observed in the present work) contribute in a lower percentage than wild species to the grass pollen curve (Muñoz et al., 2000; Subiza et al., 1992). In general the results indicated a general trend towards earlier onset of phenological phases, except for *Hordeum vulgare*. Shifts in leaf phenology were stronger than those previously reported for Europe by Menzel & Fabian (1999), who noted a 6-day advance in leaf unfolding and a 4.8-day delay in leaf fall over a 30-year period. The increase in growing-season length due to the earlier start recorded here also exceeds the 7 days reported for Europe between the 1960s and the 1990s, but largely coincides with the effect reported from the 1980s onwards (Keeling et al., 1996). The observed advance was also more marked than that found in North America by Bradley et al. (1999) and Abu-Asab et al. (2001). Also the observed trends are stronger than those reported in other Mediterranean areas such as California and Australia (Denisson & Roberts, 2003; Sadrasa & Monzon, 2006). The findings of the present study agree with those reported for other areas of southern Europe, in pointing to a bringing-forward of pollen-season start-dates and peak pollen counts in *Olea europaea* and *Quercus* species, and to a lesser extent in Poaceae (García-Mozo et al., 2008a; 2008b; 2009).

Although we must take into account the low significance in some of the results due to a high degree of variability in the analysed dates, the data recorded here suggest a more marked advance in southern than in north-eastern Spain, where Peñuelas et al. (2002) report during 1952-2000, an advance of 7.8 days for *Quercus ilex* flowering and 17 days for *Vitis* leaf unfolding, while flowering for *Olea europaea* and *Quercus faginea* displayed no significant trends. Data for a north region of Spain, show a lack of any apparent trend in the phenology of herbaceous crops, similarly to the results obtained in the present work.

The increase in mean annual temperatures over the study period appears to be the most marked since the mid-1970s, thus matching Mediterranean and global trends (Jones et al., 1999; IPCC, 2001). The detected increment of temperature in Spain (1.5°C) during the last century has been greater than those calculated for the world global average (0.6±0.2°C) or for Europe (0.5°C). (IPCC, 2001; IPCC, 2007). Climate change has had a particularly marked effect on southern Spain, increasing temperatures up to 4°C in last years (INM, 2002; De Castro, 2005; Ayala-Carcedo, 2004). Changes in rainfall patterns have been less pronounced, although a slight drop in annual rainfall was recorded for Andalusia over the last few years. Also during this period rainfall has also become increasingly torrential (Pita, 2003; De Castro et al., 2005).

Phenological changes displayed a strong correlation with temperature changes and especially with temperature over the months prior to the reproduction phase. The months preceding any given phenological event appeared to be the most important for the majority of the species. It was noteworthy that the previous autumn temperature influenced flowering in tree species, probably due to temperature in this period is related to vernalisation and to chilling requirements; the amount of chilling the plant receives in

autumn is related to heat accumulation in spring. The correlations between flowering and fruit ripening and rainfall were mostly slight, although the correlation was more marked for grass species – which displayed fewer phenological changes over the study period – than for tree species. The shift towards more arid conditions, owing to increased temperatures and to the slight drop in rainfall recorded in Andalusia, may strengthen this correlation in the near future (Peñuelas et al., 2002).

At the same time, the potential effects of airborne CO₂ levels on phenological phases such as flowering and fruiting cannot be ruled out (Peñuelas et al., 1995). Global climate warming seems to be the main cause of phenological changes, but other factors – including less rainfall and CO₂ enrichment – may also affect them. Field phenological and aerobiological data are important bio-indicators of the ongoing ecosystem change, and can provide a general idea of how future climate change may be the major driver of changes in biodiversity.

5. Conclusions

Flowering patterns matched aerobiological data for the study species, and particularly for trees. Preliminary results suggested a general advance in flowering, foliation, and fruit ripening for most of the species examined in this study, especially significant for flowering dates. That general advance was more marked in trees than in herbaceous species, in which flowering dates – more closely determined by water availability – displayed little change. The advance of phenological phases in trees was more evident than in the rest of the studied species. Increased temperature was the main factor affecting the advance in foliation, flowering and fruit ripening and the delay in leaf fall dates. Potential effects of the change in phenology on the growth and yield of these species can lead to an increase of crop size. Herbaceous species were more affected than tree species by changes in rainfall patterns. In conclusion, field phenological data and aerobiological data are important bio-indicators of ongoing ecosystem changes, and can provide a general idea of how future climate change may be the major driver of changes in biodiversity.

6. References

- Abu-Asab, M., Peterson, P.M., Shetler, S.G., Orly, S., 2001. Earlier plant flowering in spring as a response to global warming in Washington DC area. *Biodiversity and Conservation* 10, 597-612.
- Ahas, R., Aasa, A., Menzel, A., Fedotova, V.G., Scheifinger, H., 2002. Changes in European spring phenology. *International Journal of Climatology* 22(14), 1727-1738.
- Ayala -Carcedo, J., 2004. La realidad del cambio climático en España y sus principales impactos ecológicos y socioeconómicos. *Industria y Minería* 358, 10-29.
- Bradley, N.L., Leopold A.C., Ross J., Huffaker W., 1999. Phenological changes reflect climate change in Wisconsin. *Proceedings of the National Academy of sciences USA*, 96(17), 9701-9704.
- De Castro, M., Martín-Vide, J., Alonso S., 2005. El clima de España: pasado, presente y escenarios de clima para el siglo XXI. In: *Evaluación preliminar de los impactos en*

- España por efecto del cambio climático* (ed. Moreno Rodríguez JM), pp. 1–65. Ministerio de Medio Ambiente, Madrid.
- Dennison, P.E. & Roberts D.A. 2003. The effects of vegetation phenology on endmember selection and species mapping in southern California chaparral. *Remote Sensing of Environment* 87(2-3): 295-309.
- Fernández-González, F., Loidi, J., Moreno, J.C., del Arco, M., Fernández-Cancio, A., Galán, C., García-Mozo, H., Muñoz, J. Pérez-Badia, R., Sardinero, S. & Tellería, M., 2005. Impact on Plant Biodiversity. In: *Impacts on Climatic Change in Spain*. OCCE. Moreno, J.M. ed. Ministerio de Medio Ambiente. Madrid, Spain. 183-248 pp.
- Galán, C., García-Mozo, H., Vázquez, L., Ruiz-Valenzuela, L., Díaz de la Guardia, C. & Trigo-Pérez, M. 2005. Heat requirement for the onset of the *Olea europaea* L. Pollen season in several places of Andalusia region and the effect of the expected future climate change. *International Journal of Biometeorology*, 49(3), 184–188.
- Galán, C., Cariñanos P., Alcázar P., Dominguez E., 2007. Quality Manual. Spanish Aerobiology Network (REA). Servicio Publicaciones Universidad de Córdoba, Córdoba, Spain. http://www.uco.es/rea/infor_rea/manual_eng.pdf
- García-Mozo, H., Galán, C., Gómez-Casero, M.T. & Dominguez, E., 2000. A comparative study of different temperature accumulation methods for predicting the *Quercus* pollen season start in Córdoba (Southwest Spain). *Grana* 39: 194-199.
- García-Mozo, H., Galán, C., Aira, M.J., Belmonte, J., Díaz de la Guardia, C. Fernández, D. Gutiérrez, A.M., Rodríguez, FJ., Trigo, M.M., Dominguez-Vilches, E. 2002. Modelling start of oak pollen season in different climatic zones in Spain. *Agricultural and Forest Meteorology*, 110, 247-257.
- García-Mozo H., Chuine I., Aira M.J., Belmonte J., Bermejo D., Díaz de la Guardia C., Elvira B., Gutiérrez M., Rodríguez-Rajo J., Ruiz L., Trigo M.M., Tormo R., Valencia R., Galán C., 2008a. Regional phenological models for forecasting the start and peak of the *Quercus* pollen season in Spain. *Agricultural and Forest Meteorology* 148,372-380.
- García-Mozo, H., Orlandi, F., Galan C., Fornaciari, M., Romano, B., Ruiz, L., Díaz de la Guardia, C., Trigo, M.M., Chuine, I., 2008b. Olive flowering phenology variation between different cultivars in Spain and Italy: modelling analysis. *Theoretical and Applied Climatology* 95:385:395.
- García-Mozo H., Galán, C., Belmonte, J., Bermejo. D., Candau, P., Díaz de la Guardia., C., Elvira, B., Gutiérrez, M., Jato, V., Silva, I., Trigo, M.M., Valencia, R., Chuine, I., 2009. Predicting the start and peak dates of the Poaceae pollen season in Spain using process-based models. *Agriculture and Forest Meteorology* 149, 256-262.
- Gómez-Casero, M.T., Hidalgo, P., García-Mozo, H., Domínguez, E., & Galán, C. 2004. Pollen biology in four mediterranean *Quercus* species. *Grana*, 43:1-9.
- Gómez-Casero, M.T., Galán, C. & Domínguez-Vilches, E., 2007. Flowering phenology of Mediterranean *Quercus* species in different locations (Córdoba, SW Iberian Peninsula). *Acta Botánica Malacitana* 32. 127-146

- Gordo, O., Sanz, J.J., 2005. Phenology and climate change: a long-term study in a Mediterranean locality. *Oecologia* 146,484-495.
- Hirst, J., 1952. An automatic volumetric spore-trap. *Ann. Appl. Biol.* 36, 257-265.
- Instituto Nacional de Meteorología (INM), 2002. Valores normales y estadísticos de observatorios meteorológicos principales (1971-2000). Vols. 1-5.
- IPCC, 2001. Climate Change 2001: The Scientific Basis. Contribution of Working Group I to the Third Assessment Report of the Intergovernmental Panel on Climate Change [Houghton, J.T., Y. Ding, D.J. Griggs, M. Noguer, P.J. van der Linden, X. Dai, K. Maskell, and C.A. Johnson (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 881pp.
- IPCC, 2007: Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment. Report of the Intergovernmental Panel on Climate Change [Solomon, S., D. Qin, M. Manning, Z. Chen, M. Marquis, K.B. Averyt, M. Tignor and H.L. Miller (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 996 pp.
- Jones P.D., New M., Parker D.E., Martin S., Rigor I.G. 1999. Surface air temperature and its changes over the past 150 years. *Reviews of Geophysics* 37:173-199.
- Keeling C.D., Chin J.F.S., Whorf, T.P. 1996. Increased activity of northern vegetation inferred from atmospheric CO₂ measurements. *Nature*, 382:146-149.
- Menzel A., Fabian P. 1999. Growing season extended in Europe. *Nature*, 397: 659.
- Menzel, A., Sparks, T. H., Estrella, N., Koch, E., Aasa, A., Ahas R., Alm-Kübler, K., Bissolli, P., Braslavská, O., Briede, A., Chmielewski, F. M., Crepinsek, Z., Curnel, Y., Dahl Å., C., Defila, Donnelly, A., Filella Y., Jatzak K., Måge, F., Mestre, A., Nordli, Ø., Peñuelas J., Pirinen, P., Remišová, V., Scheffinger, H., M., Striz, A., Susnik, A., Van Vliet, J. H., Wielgolaski F. E. & Züst A.S.Z. 2006. European phenological response to climate change matches the warming pattern. *Global Change Biology* 12 (10): 1969-1976.
- Moreno, J.M. 2005. Impacts on Climatic Change in Spain. OCCE. ed. Ministerio de Medio Ambiente. Madrid, Spain. 183-248 pp.
- Muñoz, A.F., Silva, I. & Tormo, R. 2000. The relationships between Poaceae pollination levels and cereal yields. *Aerobiologia* 16: 281-286.
- Peñuelas J., Biel C., Estiarte M. 1995. Growth, biomass allocation, and phenology of pepper plants submitted to elevated CO₂ and different nitrogen and water availabilities. *Photosynthetica*, 31:91-99.
- Peñuelas, J., Filella, I., Comas, P. 2002. Changed plant and animal life cycles from 1952 to 2000 in the Mediterranean region. *Global Change Biology* 8:531-544.
- Pita, M.F. 2003. El Clima de Andalucía. In: López-Ontiveros, A. (ed.), *Geografía de Andalucía*. Ariel, Barcelona, Spain, pp. 137-174.
- Romberger J.A. 1963. Meristems growth and development in woody plants. USDA Technical Bulletin, N° 1292. U.S. Government Printing Office, Washington DC, 214.
- Sadrasa V.O. & Monzon J.P. 2006. Modelled wheat phenology captures rising temperature trends: Shortened time to flowering and maturity in Australia and Argentina *Field Crops Research* 99(2-3):136-146.

Subiza, J., Maseillo, J.M., Subiza, J.L., Jerez, M., Hinojosa, M. and Subiza, E., 1992. Prediction of annual variations in atmospheric concentrations of grass pollen. A method based on meteorological factors and grain crop estimates. Clin. Exp. Allergy 22, 540-546.

IntechOpen

IntechOpen

Climate Change - Socioeconomic Effects

Edited by Dr Houshan Kheradmand

ISBN 978-953-307-411-5

Hard cover, 454 pages

Publisher InTech

Published online 09, September, 2011

Published in print edition September, 2011

This book shows some of the socio-economic impacts of climate change according to different estimates of the current or estimated global warming. A series of scientific and experimental research projects explore the impacts of climate change and browse the techniques to evaluate the related impacts. These 23 chapters provide a good overview of the different changes impacts that already have been detected in several regions of the world. They are part of an introduction to the researches being done around the globe in connection with this topic. However, climate change is not just an academic issue important only to scientists and environmentalists; it also has direct implications on various ecosystems and technologies.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

García-Mozo, H., Mestre, A. and Galán, C. (2011). Climate Change in Spain: Phenological Trends in Southern Areas, Climate Change - Socioeconomic Effects, Dr Houshan Kheradmand (Ed.), ISBN: 978-953-307-411-5, InTech, Available from: <http://www.intechopen.com/books/climate-change-socioeconomic-effects/climate-change-in-spain-phenological-trends-in-southern-areas>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2011 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike-3.0 License](https://creativecommons.org/licenses/by-nc-sa/3.0/), which permits use, distribution and reproduction for non-commercial purposes, provided the original is properly cited and derivative works building on this content are distributed under the same license.

IntechOpen

IntechOpen