We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Global Heating Threatens the `Iiwi (*Vestiaria coccinea*), Currently a Common Bird of Upper Elevation Forests in Hawaii

Anthony Povilitis

*Life Net Nature, Willcox, Arizona
USA*

1. Introduction

The `Iiwi is one of 17 surviving Hawaiian honeycreepers (Fringillidae: Drepanidinae) of 37 species known historically and 55 extant prior to human arrival on Hawaii (Pratt 2009). Its closest relative is the extinct Hawaii Mamo (*Drepanis pacifica*) (Pratt 2005). Disease and habitat loss are primary reasons for the decline of Hawaiian honeycreepers and other native forest birds. Extinctions continue to this day, with the most recent being the Poo-uli (*Melamprosops phaeosoma*) in 2004.

The `Iiwi, a scarlet bird with black wings and tail, and a long curved, salmon-colored bill, is generally placed in the monotypic genus *Vestiaria*. It is a largely nectarivorous species that occurs commonly in closed canopy, high-stature native forests above 1500 m elevation (Fancy and Ralph 1998). `Iiwi breed and winter primarily in mesic and wet forests dominated by native 'ōhi'a (*Metrosideros polymorpha*) and koa (*Acacia koa*) trees (Scott et al. 1986). They often travel widely in search of 'ōhi'a flowers and are important 'ōhi'a pollinators (Mitchel et al. 2005). The birds respond to seasonal flowering patterns, often moving to lower elevations where they are exposed to deadly disease (Pratt 2005). The `Iiwi uses its long bill to extract nectar from decurved corollas of Hawaiian lobelioids, which have become far less common on Hawaii over the past century (Smith et al. 1995).

Female `Iiwi typically lay two eggs, and they alone are thought to incubate eggs and brood young (Mitchel et al. 2005). But males provision females with food off the nest. Breeding takes place predominantly from February to June, and is usually associated with peak flowering of 'ōhi'a (Fancy and Ralph 1998).

For native Hawaiians, the `Iiwi and other forest birds have a spiritual nexus. Feathered objects represented gods, ancestors, and divine lineage (Amante-Helweg and Conant 2009). Red feathers of clothing, such as cloaks, capes, and helmets, were predominantly from `Iiwi. Once a familiar sight on all main Hawaiian Islands, the `Iiwi remains an icon of Hawaii's native forests.

Today `Iiwi occur in higher elevation habitats largely free of avian disease, to which the species is highly susceptible. With climate change, these refugia may be lost entirely as pathogens and vectors advance upslope in response to higher ambient temperatures. This prognosis points to the needs for swift remedial action by responsible U.S. federal and State of Hawaii authorities to prevent the `Iiwi from joining the tragically long list of extinct or feared extinct Hawaiian birds (Banko and Banko 2009).

2. Population status

The ʻIiwi occurs on the Hawaiian islands of Kauai, Oahu, Maui, Molokai, and Hawaii (Gorresen et al. 2009). Once widely distributed in native forests on all major Hawaiian Islands, it is now mostly restricted to elevations above 1250 m because of avian diseases and habitat loss elsewhere (Warner 1968, Scott et al. 1986, Fancy and Ralph 1998, Pratt 2005).

<i>Kauai</i> -- ʻIiwi numbers decreased by 62%, from 26,000 ± 3,000 to 9,985 ± 960, between the 1970s and 2000 (Foster et al. 2004, Gorresen et al. 2009); ʻIiwi range contracted from 140 to 110 sq km, consistent with a shift in its low elevation boundary from ~900 m to >1,100 m.
<i>Oahu</i> – Few, if any, birds remain; 8 individuals dispersed in 3 isolated locations were reported in 1994-1996 (VanderWerf and Rohrer 1996).
<i>Molokai</i> -- Few birds (1-3) were detected from 1988-2004 (Reynolds and Snetsinger 2001, Gorresen et al. 2009), contrasting with 12 in 1979 (Scott et al. 1986).
<i>Maui</i> -- About 19,000 ± 2,000 individuals occurred in restricted upper elevation habitats of east Maui (Scott et al. 1986); ~ 180 ± 150 birds were reported in isolated west Maui prior to 1980 (Scott et al. 1986); the west Maui population persists today at a very low number (Gorresen et al. 2009).
<i>Hawaii Island</i> – 340,000 ± 12,000 birds were estimated in higher elevation range; ~1,000 birds in lower elevation Kohala and Puna areas (Scott et al. 1986); overall downward trends are evident in recent decades (Camp et al. 2009a, Gorresen et al. 2009); of 10 study locations, ʻIiwi appear now absent at one, declining at 5, stable at 3, with no estimate for 1 (Gorresen et al. 2009).
Regional breakout of data for Hawaii Island:
<u>Northeast area:</u> For the Hakalau Forest National Wildlife Refuge (Hakalau Unit; 1,500-2,000 m elevation) population trend data vary from stable (over a 21-year period) to declining (during a recent 9-year period), except for increasing numbers in limited newly restored upper elevation habitat (Camp et al. 2009a). Recent ʻIiwi numbers were estimated at ~61,000 birds.
<u>Central windward area:</u> ʻIiwi frequency decreased 54% between late 1970s and 1986-2000 periods in National Park and Hamakua areas, with specific study area declines and evidence of upward range contraction (Gorresen et al. 2005, Camp et al. 2009b); ʻIiwi showed pronounced decline at lower elevations (East Rift, <1,000 m elev., and ʻŌlaʻa, ~1,200-1,400 m, 1977-1994 data); modest declines (Kūlani-Keauhou, 1,500-2,000 m, 1977-2003 data) or stability (Mauna Loa Strip, ~1,500-2,000 m, 1977-1994 data) at higher elevations.
<u>Southeast area:</u> Lower ʻIiwi density in the Kaʻū area (2002 and 2005 data) than previously (1976 and 1993 data) (Gorresen et al. 2009); recent estimate of ~ 78,000 birds, with 60% occurring above 1,500 m (Gorresen et al. 2007).
<u>Leeward (western) area:</u> ʻIiwi densities have dramatically declined in the Hualālai and Kona regions (1997-2000); they are decreasing at lower elevations (<1,500 m; Kona Forest Unit, Hakalau Forest National Wildlife Refuge); stable only at upper elevations (Gorresen et al. 2009); ʻIiwi range is contracting upslope, with few occurrences below 1,100 m during the breeding season (Camp et al. 2002).

Table 1. ʻIiwi population estimates for Hawaiian islands.

`Iiwi are declining everywhere in Hawaii, except at high elevation on east Maui Island and northeast Hawaii Island (Gorresen et al. 2009) (Table 1), and population extinctions are impending throughout the Islands (Banko and Banko 2009).

On Kauai, in the western portion of the species' range, `Iiwi numbers have declined sharply (Table 1). Risk of extirpation from the island is of immediate concern because of severely diminished disease-free habitat. Oahu, Molokai, and the isolated western area of Maui have small remnant `Iiwi populations at high risk of extinction (Gorresen et al. 2009). `Iiwi are gone from nearby Lanai. These four areas comprise the central portion of the species' geographic range.

On east Maui and the Island of Hawaii, forming the eastern part of the species' range, `Iiwi populations are restricted to high elevations (Table 1). While some populations are still large, they are at risk of fragmentation and decimation resulting from the spread of avian disease driven by climate warming (Pratt et al. 2009).

The `Iiwi population of Hawaii has been estimated at 360,000 birds (Pratt et al. 2009), with the vast majority of birds (~90%) occurring on Hawaii Island (Scott et al. 1986). Declines in `Iiwi abundance corresponding with reduced lower elevation range since the early 1970s are consistent with anticipated impacts of mosquito borne disease (Foster et al. 2004). The population trend is downward on all islands, with some stability in high elevation areas (Pratt et al. 2009). Climate change is now setting the stage for widespread disease transmission at the highest elevations on Maui and Hawaii Island (Benning et al. 2002; LaPointe et al. 2005).

`Iiwi pairs are reported to produce on average only 1.33 chicks per year, reflecting low productivity characteristic of Hawaiian forest birds in general (Woodworth and Pratt 2009). However, the `Iiwi has the lowest annual survivorship reported ($55\% \pm 12$ SE for adults and $9\% \pm 5$ for juveniles) for any extant species of honeycreeper, reflecting the impact of malaria and avian pox and/or low re-sighting probabilities (Fancy and Ralph 1998; Pratt 2005).

`Iiwi populations have suffered from fragmentation as well as reduced size and range. Small population units are at risk of extinction from random demographic fluctuations, localized catastrophes (severe storms, wild fire, disease outbreaks, volcanism, etc.), inbreeding depression, and genetic drift (Primack 2006).

3. Forest habitat

Most of the `Iiwi's original forest habitat has been cleared for food crops, livestock grazing, tree plantations, and land development, with habitat losses since human settlement ranging from 52% on Hawaii Island to 85% on O'ahu (Fancy and Ralph 1998). The amount of habitat available to the `Iiwi and other forest birds has declined over the past few decades as many areas become dominated by invasive non-native species (Price et al. 2009). On the island of Hawaii, additional forest habitat loss results from land development, logging, and conversion to livestock pasture.

`Iiwi habitat across Hawaii is primarily threatened by destruction and adverse modification by feral pigs and other exotic ungulates (goats, sheep, mouflon, deer, cattle) (USFWS 2006, Pratt et al. 2009). Alien animals destroy forest understory vegetation, eliminate food plants for birds, create mosquito breeding sites through ground disturbances, provide openings on the forest floor for weeds, transport weed seeds to native forests, cause soil erosion, disrupt seedling regeneration of native plants, and girdle young trees (Fancy and Ralph 1998; Pratt 2005; USFWS 2006). Spread of exotic ungulates that are especially difficult to contain (i.e.,

axis deer on Maui and Molokai, black-tailed deer on Kauai, and mouflon sheep on Hawaii Island) represent a growing threat to `Iiwi habitat as these high-jumping species invade areas even with fencing designed to exclude feral pigs and goats (Price et al. 2009). Browsing and soil compaction by feral pigs, goats, and deer in Molokai has reduced 'ōhi'a forest to grassy scrubland (Hess 2008).

Hawaiian forests are severely modified by invasive alien plant species that displace native plants used by foraging and nesting birds (Scott et al. 1986; Foster et al. 2004) and increase the frequency of forest fires (Pratt et al. 2009). Herbivory by the introduced black rat on the flowers and fruits of native plants may also reduce food resources for native birds and impact regeneration of native plants (Banko and Banko 1976). Introduced predatory insects also may reduce or eliminate specialized native insects that are needed for pollination of plants important to `Iiwi.

Introduced species of insects and birds can compete with native birds for food and other resources. `Iiwi may face competition from Japanese White-eye (*Zosterops japonicas*) (Mountainspring and Scott 1985), a malaria resistant species, whose numbers have increased at least on Kauai over the past 30 years (Foster et al. 2004). Negative correlations between `Iiwi and Japanese White-eye densities may stem from competition for limited nectar resources (Fancy and Ralph 1998). There are no current efforts to control competing species within the recovery areas of endangered forest birds (USFWS 2006). Habitat degradation by non-native mammals, plants, and invertebrates will likely continue to result in loss, modification, and curtailment of `Iiwi habitat and range.

4. Climate change

The `Iiwi survives in habitat largely free of avian malaria (*Plasmodium relunctum*) and bird pox (*Aviapoxvirus*). Such habitat is currently limited to 8.9 ha (22 acres) on Kauai, 2,632 ha (6,500 acres) on Maui, and 6,478 ha (16,000 acres) on Hawaii Island, with virtually none on Oahu and Molokai (Pratt et al. 2009). The elevational advance of these pathogens driven by climate change immediately endangers the smaller `Iiwi populations on Kauai, Oahu, Molokai, and west Maui, and threatens the larger ones on east Maui and Hawaii Island.

4.1 `Iiwi is highly vulnerable to disease

Avian disease is a primary reason for the decline of `Iiwi and other Hawaiian honeycreepers (Pratt 2005, Atkinson and LaPointe 2009). Warner (1968) demonstrated high susceptibility of honeycreepers that died from avian malaria and bird pox after experimental exposure to mosquito infested lower elevations where the birds were absent. Van Ripper et al. (1986) also provided experimental evidence of high susceptibility of `Iiwi to avian malaria. More recently, Atkinson et al. (1995) experimentally exposed several species of honeycreepers to a single bite of a malaria infected mosquito and found that effects were most severe in `Iiwi with significantly higher mortality and clear manifestations of malaria disease at death. `Iiwi were infected by either single (low-dose) or multiple (high-dose) mosquito bites. Mortality in both groups was significantly higher than in uninfected controls, reaching 100% of high-dose birds and 90% (9 of 10) in low-dose birds.

While some individual `Iiwi are known to have at least temporarily survived malaria, there is no evidence of population level tolerance or resistance to the disease. Atkinson et al. (1995) found that the one `Iiwi that survived malaria after a single experimental bite from

an infected mosquito did not develop new parasitemia after multiple bites from infected mosquitoes. This indicated that `Iiwi are capable of an immunological response at least to the administered strain of malaria. Freed et al. (2005) discovered tolerance to malaria in two wild `Iiwi that successfully bred 2-years post infection. However, broken head feathers in these birds suggested physiological costs of malarial tolerance that could reduce survivorship of wild birds. Studies of experimentally infected birds indicate that tolerant birds likely retain chronic infection for life (Atkinson et al. 2001, Valkiunas 2005). Challenges to the immune system by stress or excessive energy expenditure can result in recrudescence of a chronic infection to higher parasitemia levels (Freed et al. 2005). Infected birds lose weight and suffer malaria related pathologies (Atkinson et al. 2001), and would be expected to be more susceptible than healthy birds to predation, competition, avian pox, unfavorable weather, and other stressors. A comparison of infection incidence in `Iiwi and other Hawaii forest birds suggests that few `Iiwi survive exposure in the wild (Atkinson et al. 2005).

It is uncertain if exiting larger populations of `Iiwi on Maui and Hawaii Island could evolve tolerance rapidly enough to avoid extinction from increased malaria parasitism. This would depend on exposure rapidity, the extent of current disease tolerance, if any, the virulence of *Plasmodium* strains, patterns of selection and genetic drift, rates of evolution in hosts, vector, and pathogen, and other factors. The avian disease system on Hawaii would be further complicated if new reservoir hosts or vectors enter the picture (Atkinson and LaPointe 2009).

Lethal effects of avian poxvirus have also been experimentally demonstrated in Hawaiian honeycreepers (Jarvi et al. 2008). Freed et al. (2005) found a dead `Iiwi in the field with massive poxvirus sores on its ankles. The bird also tested positive for malaria. A significantly high proportion of Hawaiian forest birds with avian pox also had chronic malaria, suggesting interaction between the two diseases (Atkinson et al. 2005).

The downward trajectory of `Iiwi populations (Table 1) indicates a pattern of decline similar to Hawaiian forest birds already acknowledged to be endangered and very vulnerable to disease, and dissimilar to populations of the unlisted Amakihi (*Hemignathus* spp.) (Shehata et al. 2001, Woodworth et al. 2005) and Apapane (*Himatione sanguine*) (Atkinson et al. 2005) which have shown some disease resistance and population persistence at lower elevations.

Among the most endangered Hawaiian bird species, the `Ō`ū, (*Psittirostra psittacea*), like the `Iiwi, was widespread on all main islands across a wide range of habitats a century ago (USFWS 2006). However, `Ō`ū primarily inhabited the lower to mid-elevation forests where the impact of introduced mosquito-borne diseases was first manifested. Today, the `Ō`ū is probably extinct. Similar widespread exposure of `Iiwi to avian diseases can be expected in coming decades as a consequence of climate change.

4.2 Disease will spread over `Iiwi range as ambient temperatures rise

Avian malaria in Hawaii has been mostly confined to elevations below 1500 m (van Riper et al. 1986) where cool temperatures limit mosquito presence and development of the malaria parasite (LaPointe 2000). Recent climate modeling, however, has projected avian malaria to reach elevations up to or beyond 1900 m within this century, affecting most if not all remaining forest bird habitat (Benning et al. 2002).

Benning et al. (2002) modeled changes in malaria prevalence for Hawaiian honeycreepers at high quality habitat sites, assuming a 2° Celsius (C) increase in regional temperatures (based

on International Panel on Climate Change 2007 projections; see Meehl et al. 2007). Current low-risk habitat diminished by 57% (665 to 285 ha) at the Hanawi Natural Area Reserve, Maui. Low-risk habitat at the Hakalau National Wildlife Refuge on Hawaii Island declined by 96% (3,120 to 130 ha). On Kauai (the Alakai Swamp), currently with little or no malaria free habitat, a 2° C warming placed most habitat (84%) at highest risk for malaria infection in native birds. Current mean ambient temperatures are believed to already allow limited disease transmission throughout Kauai as all `Iiwi habitat occurs below 1600 m elevation (LaPointe et al. 2005).

The effects of a 2° C warming would almost certainly eliminate the small `Iiwi populations from the lower-elevation islands of Kauai, Molokai, and Oahu, and from West Maui. Larger populations on East Maui and Hawaii Island would be expected to decline severely in a manner corresponding to decreases (~60-96%) in high elevation, disease-free refuges (Atkinson and LaPointe 2009).

The prognosis for `Iiwi and many other native forest birds appears worse than indicated by the Benning et al. (2002) model. The model assumed an increase of 2° C above current temperature, corresponding to ~2.7° C increase above pre-industrial levels. However, recent analysis of global heating indicates that temperature increases in Hawaii and elsewhere are unlikely to be limited to 2° C in this century. Increases in global temperature are currently on a trajectory to reach 2° C (above pre-industrial levels) by mid-century and about 5° C by 2100 (Meinshausen et al. 2009, Sokolov et al. 2009). Global greenhouse gas emissions would need to be halved by 2050 (from 1990 levels) to keep near the 2° C level with a high probability (55-88%) (Meinshausen et al. 2009). Unfortunately, under current multi-national policies regarding greenhouse gas emissions, there is virtually no chance of limiting heating to 2° C even with full policy implementation (Rogelj et al. 2009). For Hawaii, only a low global emissions scenario would likely keep temperature increases to 2° C (Karl et al. 2009).

An added concern is the risk of abrupt increases in global temperature unaccounted for in most modeled climate projections (Lovelock 2009). For example, a global climate model used by Sokolov et al. (2009) did not fully incorporate positive feedbacks that may occur, for example, if increased temperatures cause a large-scale melting of permafrost in arctic regions and subsequently release large quantities of methane, a very potent greenhouse gas (Rice 2009). If these positive feedback loops should occur, and evidence is mounting that they will (McCarthy 2010), temperatures are likely to increase to an even greater degree in Hawaii.

For Hawaii, Giambelluca et al. (2008) document a long-term increase in temperature and an accelerated rate of increase over the past few decades consistent with global trends (0.04° C/decade over an 88-year period, and about 0.2° C/decade since 1975). Moreover, since 1975 higher elevation temperatures exceeded average warming (a 0.27° C/decade increase) with steepest increases in minimum (night time) temperature (near 0.5° C/decade), which is likely the most limiting for malaria transmission. The recent surface temperature trend in Hawaii is only slightly lower than the overall global trend. Similar surface warming has been detected elsewhere in the Pacific, and is associated with an increase in sea surface temperatures, upper ocean heat content, and sea level height (Richards and Timmermann 2008).

In Hawaii, the upper limit of mosquito presence appears to have increased substantially, from about 600 m in the late 1960s to 1100-1500 m in recent decades (Pratt 2005). Freed et al. (2005) reported that prevalence of malaria in Hawaiian forest birds at 1900 m on the island of Hawai'i more than doubled over a decade. A highly significant increase of malaria in `Iiwi was associated with much warmer summertime air temperatures. The 13° C threshold for malaria development projected for 1900 m sites by the conservative Benning et

al. (2002) model was surpassed in 2001 by a wide margin (4.4° C; Freed et al. 2005). Measured temperatures were believed to exceed model expectations because the site was strongly affected by the island's trade wind inversion layer related to tropical air circulation. The altitude of the inversion has averaged 1900 m, above which cooler, drier conditions prevail (*Atlas of Hawaii*, 3rd edition). The response of the inversion layer to climate heating is uncertain (Pounds et al. 1999, Loope and Giambelluca 1998). If the inversion layer rises, disease epizootics could become commonplace at higher elevations with devastating short-term consequences for `Iiwi. If the inversion falls, and higher temperatures become associated with high-elevation drought, the effects would be very damaging to upper elevation Hawaiian forests and ultimately to surviving honeycreepers including the `Iiwi (Benning et al. 2002). Given that scenario, or if the inversion layer remains stable, high-elevation forest bird populations may be squeezed between expanding disease transmission from lower elevations and the upper limits of suitable habitat (Atkinson and LaPointe 2009). Hawaii may see an increased frequency of heavy rain events and increased rainfall during summer months (Karl et al. 2009), conditions that, along with increased temperature, are likely to facilitate breeding of malaria-carrying mosquitoes (Ahumada et al. 2004). At the same time, overall annual precipitation for the Hawaiian Islands may decline (Chu and Chen 2005) thereby affecting habitat quality (e.g., 'ōhi'a forest) for the `Iiwi.

4.3 Confounding population stressors and threats

Ectoparasites, particularly chewing lice (Phthiraptera), may impact `Iiwi by increasing morbidity and reducing the ability of birds to survive environmental challenges. Freed et al. (2008) documented an explosive increase in the prevalence of chewing lice in all bird host species at a study site on Hawaii Island. The number of major fault bars in wing and tail feathers, a sign of nutritive stress, was correlated with intensity of infection, suggesting an indirect cost to parasitized birds. Poorer body condition preceded the outbreak indicating the synergistic effect of multiple stressors on forest birds. At a minimum, chewing lice will increase food requirements of hosts. This indirect cost may be especially relevant because it can affect the ability of birds to mount a sufficient immune defense against diseases like avian malaria and pox. Chewing lice may also directly contribute to bird mortality (Freed et al. 2008).

Additional risks to `Iiwi from disease include potential introductions of West Nile virus, new avian malaria vectors (such as temperate varieties of *Culex quinquefasciatus*), or biting midges (Culicoides) that transmit avian diseases.

Introduced rats are serious predators on adults and nests of Hawaiian forest birds, and are abundant in high elevation habitats (Atkinson 1977, Scott et al. 1986, Fancy and Ralph 1998, VanderWerf and Smith 2002). Feral cats, introduced small Indian mongoose, and the native Short-eared Owl and introduced Barn Owl may also impact native Hawaiian birds (Scott et al. 1986; Kowalsky et al. 2002). Predator control efforts generally have not been conducted over areas large enough to result in significant improvement in the status of imperiled forest birds (USFWS 2006). Logistical and other obstacles to predator control can be great, especially in rugged bird habitat.

Epizootics involving avian malaria or other pathogens could quickly eliminate remaining `Iiwi from the lower elevation islands of Kauai, Oahu, and Molokai, and from west Maui in the near term, and could diminish and fragment `Iiwi populations on higher elevation east Maui and Hawaii Island. There is currently no habitat on Kauai, Oahu, and Molokai where mean ambient temperature entirely restricts malaria development (Benning et al.

2002). These islands are vulnerable to avian malaria at all elevations on a more or less ongoing basis. A recent avian malaria outbreak on Hawaii Island was associated with increases in summertime temperatures related to tropical inversion layer conditions (Freed et al. 2005). Outbreaks of malaria can be triggered by warm periods linked to inversion layer dynamics or El Niño events, and will likely intensify and persist longer with ongoing climate change.

Hurricanes are known for their devastating effects on island birds (Foster et al. 2004). They reduce habitat by blowing down trees and by creating forest openings that facilitate the spread of invasive alien plants. The ʻIiwi decline on Kauai after a 1992 hurricane may have partially resulted from the birds seeking substitute nectar resources at lower elevations where risk of malaria transmission is highest (Foster et al. 2004).

Hurricanes are likely to intensify in a warmer climate (Meehl et al. 2007) in terms of wind speeds and precipitation, though the number of storms may be fewer (Bengtsson et al. 2007). Infectious mosquitoes can be carried upslope in strong winds, a probable factor in malaria outbreaks on Hawaii above 1900 m elevation (Freed et al. 2005).

On Hawaii Island, volcanism presents a potential threat to substantial acreage of forest bird habitat. For example, a large portion of the Upper Waiākea Forest Reserve, location of some of the last observations of ʻŌʻū and considered prime habitat for the species, was inundated by the 1984 Mauna Loa lava flow which destroyed thousands of acres of forest and created a treeless corridor over 1 km wide (USFWS 2006).

5. Conservation

Current regulations by the U.S. government and the State of Hawaii are inadequate to conserve high elevation forests needed to buffer the ʻIiwi and other susceptible forest birds against the upslope advance of avian diseases driven by global heating. While some progress has been made to re-forest former upper elevation habitat areas with native trees and reduce or eliminate harmful alien species from existing ones, huge tracts of land needed for forest bird conservation in Hawaii remain degraded or without native tree cover (USFWS 2009). A preponderance of lands intended for forest bird recovery are not managed conservation lands (Pratt et al. 2009). Management actions identified in existing forest bird conservation plans have not been implemented at ecologically relevant scales, and successful efforts to restore higher elevation forests must occur across tens of thousands of areas, not hundreds (Scott 2009). On the Island of Maui, for example, more than half of the lands identified for forest bird recovery remain without native forests, have only remnant forest patches, or are dominated by introduced tree species and other alien vegetation (personal observation). Yet restoration of high elevation koa/ʻōhiʻa forest to protect native birds is clearly a stated conservation priority (Scott et al. 1986, USFWS 2006)

At current rates, reforestation and forest enhancement efforts for Hawaiian forest birds will not achieve habitat conservation goals in time to build and expand populations robust enough to withstand avian malaria and other consequences of climate change. Of over 140 actions for forest bird recovery relating to reforestation and securing recovery areas (USFWS 2006), 61% have not begun, 37% are ongoing, and only 2% are complete or partially so (USFWS 2009). Likewise, of more than 160 actions designed to reduce or eliminate exotic ungulates and mammalian bird-predators, 71% are not yet underway, 27% are ongoing, and less than 2 % are complete or partially complete.

Poor political and policy decisions are responsible for the current inadequacy of management to prevent forest bird extinctions. The problem includes conflicting

management goals and policies, most notably involving state forest lands (USFWS 2006, 2009), and failure to provide necessary funding (Leonard 2008).

Leonard (2009) discusses political obstacles to saving Hawaiian forest birds, including a state mandate to provide public hunting opportunities of exotic ungulates even where incompatible with conservation of native birds. Actions such as fencing and ungulate control for bird conservation may result in the loss of hunting areas, which is very controversial within his state agency (Leonard 2008). Even proposals for protecting limited forest in areas of little or no public access receive fierce opposition from local hunters (San Nicolas 2010). Native forest restoration is also hampered by agency decisions favoring exotic tree species or leasing for livestock (USFWS 2006).

In terms of addressing climate change, existing international and U.S. regulatory goals to reduce global greenhouse gas emissions are clearly inadequate to safeguard the 'Īiwi against climate related extinction. As discussed, severe shrinkage of habitat absent of or at low risk of avian disease is expected with a 2° C rise in ambient temperature. While the 2009 U.N. Climate Change Conference in Copenhagen called on countries to hold the increase in global temperature below 2° C, the *non-binding* "Copenhagen Accord" that emerged from the conference fell way short of that goal. A summary by the Pew Center (2010) of four analytical reviews of the Accord found that collective national pledges to cut greenhouse gas emissions are inadequate to achieve the 2° C goal, and instead suggest emission scenarios leading to a 3 to 3.9° C warming.

Economic growth is the most significant factor driving projected increases in carbon dioxide emissions, as the world continues to rely on fossil fuels for most of its energy use (USEIA 2009). Yet the prevailing economic and political framework for the U.S. and most other countries is to maximize growth as a priority. In a high growth scenario, world carbon dioxide emissions increase at an average rate of 1.8 percent annually from 2006 to 2030, as compared with 1.4 percent under standard assumptions (USEIA 2009).

The United States is responsible for over 20% of worldwide carbon dioxide emissions (USEIA 2004). While the U.S. Environmental Protection Agency (EPA) currently has some authority to regulate greenhouse gas emissions, the agency bends under political pressure (Bravender and Samuelsohn 2010) and has not set targets or standards to protect the 'Īiwi or other wildlife. Prospects for regulations within the foreseeable future adequate to stem the climate-change threat to the 'Īiwi are very poor. For example, the U.S. Congress has failed to pass climate change legislation and, as of early 2011, is considering bills to block EPA's limited authority to regulate greenhouse gas emissions (New York Times 2011).

The nation's top wildlife agency, the US Fish and Wildlife Service (USFWS), is focused on a climate adaptation strategy for wildlife in general but with little, if any, emphasis on regulation of greenhouse gas emissions (USFWS 2010). The agency has been urged to promote reductions in emissions while expediting upper elevation habitat restoration in conservation plans for endangered Hawaiian forest birds (Povilitis and Suckling 2010).

The 'Īiwi is not included on the USFWS list of endangered species and therefore does not merit the conservation provisions of the U.S. Endangered Species Act (ESA), such as the protection of its "critical habitat." Also, like other Hawaiian honeycreepers, it is not protected under the U.S. Migratory Bird Treaty Act. In 2010, a formal request was made to the USFWS to list the 'Īiwi under the ESA and designate and protect critical habitat for the species (Center for Biological Diversity 2010). While the listing of species endangered by climate change is controversial because of an overall backlog of listing requests, new

administrative strategies or procedures will be needed if the U.S. is to fulfill its commitment to safeguard endangered wildlife (Woody 2011).

The ESA defines the term “critical habitat” to mean specific areas essential to the conservation of the species which may require special management considerations or protection (ESA 1973). Critical habitat designation would legally ensure that U.S. government actions avoid jeopardizing the species and promote its conservation. Designation would alert federal and state agencies and private landowners to the need for habitat management and restoration actions in areas essential for the ‘Iiwi, and require remedies to institutional conflicts that undermine habitat conservation. Specific measures to reduce the climate change/disease threat include reforestation, elimination or control of alien species inimical to the survival of ‘Iiwi, and special measures to monitor and reduce (or eliminate) occurrence of avian malaria vectors. Programs to re-establish native forests, reduce rat depredation, control weeds, and fence out and remove ungulates are essential for forest bird recovery in high elevation habitats that serve as native bird refugia (Gorresen et al. 2005). Reducing mortality in key habitat areas, such as that caused by rodent predation, may lessen the threat from disease by improving survival and reproduction of any birds with disease tolerance or natural immunity (VanderWerf and Smith 2002). The evolutionary acceleration of disease resistance through rodent control is possible (USFWS 2006).

Critical habitat should include all areas needed to provide sufficient forested habitat to support viable or potentially viable ‘Iiwi populations on Kauai, Oahu, Molokai, Maui, and Hawaii Island, as each island represents a significant portion of the species’ natural range. This should include areas on Maui and Hawaii Island above the current limit of tree growth to accommodate any forest expansion resulting from climate change. Critical habitat designation for ‘Iiwi would extend habitat protection to other listed endangered Hawaiian birds, where ranges overlap since most currently listed forest birds do not have critical habitat designations.

6. Conclusion

The best available science indicates that global warming will allow avian diseases to spread throughout most or all of the ‘Iiwi’s geographic range. The ‘Iiwi is highly vulnerable to avian diseases and cannot sustain itself where disease prevails. ‘Iiwi in the central portion of the species’ range (Oahu, Molokai, and west Maui) are critically endangered because of small population sizes and exposure to malaria. Those to the west (on Kauai) are severely threatened as disease free habitat is fast disappearing. ‘Iiwi in the eastern portion of the range (east Maui and Hawaii Island) face further population declines and eventual extinction with ongoing climate change. The ‘Iiwi merits protection under the US Endangered Species Act that includes designation and conservation of critical habitat.

7. References

- Ahumada, J. A., D.A. LaPointe and M. D. Samuel. 2004. Modeling the population dynamics of *Culex quinquefasciatus* (Diptera: Culicidae) along an elevational gradient in Hawaii. *Journal of Medical Entomology* 41(6):1157-1170.

- Amante-Helweg, V.L.U. and S. Conant. 2009. Hawaiian culture and forest birds in Conservation Biology of Hawaiian Forest Birds: Implications for Island Avifauna. T. K. Pratt, C.T. Atkinson, P. C. Banko, J. D. Jacobi, B. L. Woodworth (eds.). Yale University Press. New Haven.
- Atkinson, I. A. E. 1977. A reassessment of factors, particularly *Rattus rattus* L., that influenced the decline of endemic forest birds in the Hawaiian Islands. *Pacific Science* 31:109-133.
- Atkinson, C. T., R. J. Dusek, and J. K. Lease. 2001. Serological responses and immunity to superinfection with avian malaria in experimentally-infected Hawai'i 'Amakihi. *Journal of Wildlife Diseases* 37:20-27.
- Atkinson, C. T. and D. A. LaPointe. 2009. Ecology and pathogenicity of avian malaria and pox. Chapter 9 in Conservation Biology of Hawaiian Forest Birds: Implications for Island Avifauna. Carter T. Atkinson, Paul Christian Banko, James D. Jacobi, Bethany Lee Woodworth. Yale University Press. New Haven.
- Atkinson, C.T., J. K. Lease, R. J. Dusek, and M. D. Samuel. 2005. Prevalence of pox-like lesions and malaria in forest bird communities on leeward Mauna Loa volcano, Hawaii. *Condor* 107:537-546.
- Atkinson, C. T., K. L. Woods, R. J. Dusek, L. Sileo, and W. M. Iko. 1995. Wildlife disease and conservation in Hawaii: Pathogenicity of avian malaria (*Plasmodium relictum*) in experimentally infected 'Iiwi (*Vestiaria coccinea*). *Parasitology* 111:S59-S69.
- Banko, W. E. and P. C. Banko. 1976. Role of food depletion by foreign organisms in historical decline of Hawaiian forest birds. Pages 29-43 in Proceedings of 1st conference in Natural sciences, Hawai'i Volcanoes National Park, Hawai'i.
- Banko, W. E. and P. C. Banko. 2009. Historic decline and extinction in Conservation Biology of Hawaiian Forest Birds: Implications for Island Avifauna. T. K. Pratt, C.T. Atkinson, P. C. Banko, J. D. Jacobi, B. L. Woodworth (eds.). Yale University Press. New Haven.
- Bengtsson, L., K. I. Hodges, M. Esch, N. Keenlyside, L. Kornblueh, J.-J. Luo, and T. Yamagata. 2007. How may tropical cyclones change in a warmer climate? *Tellus* 59A:539-561.
- Benning, T. L., D.A. LaPointe, C.T. Atkinson, and P. M. Vitousek. 2002. Interactions of climate change with biological invasions and land use in the Hawaiian Islands: Modeling the fate of endemic birds using a geographic information system. *Proceedings of the National Academy of Sciences* 99(22):14246-14249.
- Bravender, R. and D. Samuelsohn. 2010. EPA's gradual phase in of GHG Regs garners qualified praise from Senators. *New York Times*. Feb. 23.
- Camp, R. J., P. M. Gorresen, B. L. Woodworth, and T. K. Pratt 2002. Preliminary analysis of forest bird survey data for national parks in Hawaii: Hawaii Volcanoes National Park 1977-2000. Interim Report, May 2002. Hawaii Forest Bird Interagency Database Project, Pacific Island Ecosystems Research Center, Biological Resources Division, U.S.G.S.
- Camp, R. J., T. K. Pratt, P.M. Gorresen, J. J. Jeffrey, and B. L. Woodworth. 2009a. Passerine bird trends at Hakalau Forest National Wildlife Refuge, Hawai'i. Hawai'i

- Cooperative Studies Unit Technical Report HCSU-011. University of Hawai'i at Hilo.
- Camp, R. J., B. L. Woodworth, T. K. Pratt, C. Collins, and H. Howitt. 2009b. The Hawaii bird interagency database project.
http://biology.usgs.gov/pierc/HFBIDPSite/HCC_2000_Poster.pdf Accessed 9-22-09
- Center for Biological Diversity. 2010. Petition to list the 'I'iwi (*Vestiaria coccinea*) as threatened or endangered under the U.S. Endangered Species Act. Tucson, Arizona, U.S.A.. August 24.
- Chu, P.-S., and H. Chen, 2005: Interannual and interdecadal rainfall variations in the Hawaiian Islands. *Journal of Climate* 18:4796-4813.
- Fancy, S. G. and C. J. Ralph. 1998. 'I'iwi (*Vestiaria coccinea*) in A. Poole and F. Gill, eds. *The Birds of North America*, No. 327. The Birds of North America, Inc., Philadelphia, PA, U.S.A.
- Foster, J. T., E. J. Tweed, R. Camp, B. L. Woodworth, C. Adler, and T. Telfer. 2004. Long-term population changes of native and introduced birds in the Alaka'i Swamp, Kaua'i. *Conservation Biology* 18(3):716-725.
- Freed, L. A., R. L. Cann, M. L. Goff, W. A. Kuntz, and G. R. Bodner. 2005. Increase in avian malaria at upper elevation in Hawaii. *Condor* 107:753-764.
- Freed, L. A., M. C. Medeiros, and G.R. Bodner. 2008. Explosive increase in ectoparasites in Hawaiian forest birds. *Journal of Parasitology* 94(5):1009-1021.
- Giambelluca, T. W., H.F. Diaz, and M. S. A. Luke. 2008. Secular temperature changes in Hawaii. *Geophysical Research Letters* 35:L12702.
- Gorresen, P.M., R. J. Camp, and T. K. Pratt. 2007. Forest Bird Distribution, Density and Trends in the Ka'u Region of Hawai'i Island. Open-File Report 2007-1076.
- Gorresen, P.M., R.J. Camp, T. K Pratt, and B. L. Woodworth. 2005, Status of forest birds in the central windward region of Hawai'i Island: population trends and power analyses: U.S. Geological Survey, Biological Resources Discipline, Open-File Report 2005-1441.
- Gorresen, P. M., R. J. Camp, M. H. Reynolds, B. L. Woodworth, and T. K. Pratt. 2009. Status and trends of native Hawaiian songbirds. Chapter 5 in *Conservation Biology of Hawaiian Forest Birds: Implications for Island Avifauna*. T. K. Pratt, C.T. Atkinson, P. C. Banko, J. D. Jacobi, B. L. Woodworth (eds.). Yale University Press. New Haven.
- Harvell, C. D., C. E. Mitchell, J. R. Ward, S. Altizer, A. P. Dobson, R. S. Ostfeld, and M. D. Samuel. 2002. Climate warming and disease risks for terrestrial and marine biota. *Science* 296:2158-2162.
- Hess, S.C. 2008. Wild sheep and deer in Hawaii – a threat to fragile ecosystems. USGS, Hawaii National Park, Hawaii.
- Jarvi, S. I., D. Triglia, A. Giannoulis, M. Farias, K. Bianchi, and C. T. Atkinson. 2008. Diversity, origins and virulence of Avipoxviruses in Hawaiian Forest Birds. *Conservation Genetics* 9:339-348.
- Karl, T. R., J. M. Melillo, and T. C. Peterson (editors). 2009. *Global climate change impacts in the United States*. Cambridge University Press, New York.

- Kowalsky, J. R., T. K. Pratt, & J. C. Simon. 2002. Prey taken by feral cats (*Felis catus*) and barn owls (*Tyto alba*) in the Hanawi Natural Area Reserve, Maui, Hawai'i. 'Elepaio 62:129-131.
- LaPointe, D. A. 2000. Avian malaria in Hawai'i: the distribution, ecology, and vector potential of forest-dwelling mosquitoes. Ph.D. dissertation, University of Hawaii at Manoa, Honolulu, HI.
- LaPointe, D., T. L. Benning, and C. Atkinson. 2005. Avian malaria, climate change, and native birds of Hawaii. In *Climate change and biodiversity* (T.E. Lovejoy and L. Hannah eds). Yale University Press, New Haven.
- Leonard, D. L. 2008. Recovery expenditures for birds listed under the US Endangered Species Act: the disparity between mainland and Hawaiian taxa. *Biological Conservation* 141:2054-2061.
- Leonard, D. L. 2009. Social and political obstacles to saving Hawaiian birds: Realities and remedies. Chapter 24 in *Conservation Biology of Hawaiian Forest Birds: Implications for Island Avifauna*. T. K. Pratt, C.T. Atkinson, P. C. Banko, J. D. Jacobi, B. L. Woodworth (eds.). Yale University Press. New Haven.
- Loope, L.L. and T.W. Giambelluca. 1998. Vulnerability of island tropical montane cloud forests to climate change, with special reference to East Maui, Hawaii. *Climatic Change* 39:503-517.
- Lovelock, J. 2009. *The vanishing face of Gaia*. Basic Books. New York.
- McCarthy, J. 2010. Reflections on: our planet and its life, origins, and futures. *Science* 326:1646-1655.
- Meehl, G. A., T. F. Stocker, W. D. Collins, P. Friedlingstein, A. T. Gaye, J. M. Gregory, A. Kitoh, R. Knutti, J. M. Murphy, A. Noda, S. C. B. Raper, I. G. Watterson, A. J. Weaver, and Z.-C. Zhao, 2007: Global climate projections. In: *Climate Change 2007: The Physical Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change* [Solomon, S., D. Qin, M. Manning, Z. Chen, M. Marquis, K.B. Averyt, M. Tignor, and H.L. Miller (eds.)]. Cambridge University Press, Cambridge, UK, and New York, pp. 747-845.
- Meinshausen, M., N. Meinshausen, W. Hare, S. C. B. Raper, K. Frieler, R. Knutti, D. J. Frame, and M.R. Allen. 2009. Greenhouse-gas emission targets for limiting global warming to 2 degrees C. *Nature* 458:1158-1162.
- Mitchell, C, C. Ogura, D. W. Meadows, A. Kane, L. Strommer, S. Fretz, D. Leonard, and A. McClung. 2005. *Hawaii's Comprehensive Wildlife Conservation Strategy*. Department of Land and Natural Resources. Honolulu, Hawai'i.
- Mountainspring, S., and J. M. Scott. 1985. Interspecific competition among Hawaiian forest birds. *Ecological Monographs* 55:219-239.
- New York Times. 2011. Global Warming. Science Topics. Updated: Jan. 13, 2011 <http://topics.nytimes.com/top/news/science/topics/globalwarming/index.html>
- Pew Center on Global Climate Change. 2010. <http://www.pewclimate.org/copenhagenaccord> Accessed May 5.
- Pounds, J. A., M. P. L. Fogden and J. H. Campbell. 1999. Biological response to climate change on a tropical mountain. *Nature* 398:611-615.

- Povilitis, A. and K. Suckling. 2010. Addressing climate change threats to endangered species in U.S. recovery plans. *Conservation Biology* 24(2):372-376.
- Pratt, D. H. 2005. *The Hawaiian honeycreepers*. Oxford University Press. Oxford.
- Pratt, T. K. 2009. Origins and evolution. Chapter 1 in *Conservation Biology of Hawaiian Forest Birds: Implications for Island Avifauna*. T. K. Pratt, C.T. Atkinson, P. C. Banko, J. D. Jacobi, B. L. Woodworth (eds.). Yale University Press. New Haven.
- Pratt, T. K., C. T. Atkinson, P. C. Banko, J. D. Jacobi, B. L. Woodorth, and L. A. Mehrhoff. 2009. Can Hawaiian forest birds be saved? Chapter 25 in *Conservation Biology of Hawaiian Forest Birds: Implications for Island Avifauna*. T. K. Pratt, C.T. Atkinson, P. C. Banko, J. D. Jacobi, B. L. Woodworth (eds.). Yale University Press. New Haven.
- Price, J.P., J. D. Jacobi, L.W. Pratt, F.R. Warshauer, and C.W. Smith. 2009. Protecting forerist bird populations across landscapes. Chapter 16 in *Conservation Biology of Hawaiian Forest Birds: Implications for Island Avifauna*. T. K. Pratt, C.T. Atkinson, P. C. Banko, J. D. Jacobi, B. L. Woodworth (eds.). Yale University Press. New Haven.
- Primack, R. B. 2006. *Essentials of Conservation Biology*. 4th Edition. Sinauer Associates Sunderland, MA.
- Reynolds, M. H. and T. J. Snetsinger. 2001. The Hawai'i rare bird search 1994-1996. Pp. 133-143 in J. M. Scott, S. Conant, and C. van Riper, III, eds. *Evolution, ecology, conservation, and management of Hawaiian birds: a vanishing avifauna*. Studies in Avian Biology No. 22. Cooper Ornithological Society. Allen Press, Lawrence, KS, U.S.A.
- Rice, D. 2009. Global warming may be twice as bad as previously expected. *USA Today*. May 21.
- Richards, K. and A. Timmermann. 2008. *Climate change and biodiversity in Melanesia. Regional Climate Change Projections for the Southwestern Pacific with a focus on Melanesia* International Pacific Research Center, School of Ocean and Earth Science and Technology, University of Hawaii CCBM Paper 1. Bishop Museum Technical Report 42(1).
- Rogelj, J., B. Hare, J. Nabel, K. Macey, M. Scheffer, K. Markmann, and M. Meinshausen. 2009. Halfway to Copenhagen, no way to 2 C. *Nature Reports* 3:81-83.
- San Nicolas, C. 2010. Hunters take aim at plan to preserve forest. *The Maui News*. April 30. <http://www.mauinews.com/page/content.detail/id/531038.html?nav=10>
- Scott, J. M. 2009. Forward. in *Conservation Biology of Hawaiian Forest Birds: Implications for Island Avifauna*. T. K. Pratt, C.T. Atkinson, P. C. Banko, J. D. Jacobi, B. L. Woodworth (eds.). Yale University Press. New Haven.
- Scott, J. M., S. Mountainspring, F. L. Ramsey, and C.B. Kepler. 1986. *Forest bird communities of the Hawaiian Islands: their dynamics, ecology, and conservation*. Studies in Avian Biology No. 9. Cooper Ornithological Society. Allen Press. Lawrence.

- Shehata, C. L., L. A. Freed, and R. L. Cann. 2001. Changes in native and introduced bird populations on O'ahu: infectious diseases and species replacement. *Studies in Avian Biology* 22:264-273.
- Smith, T. B., L. A. Freed; J. Kaimanu Lepson and J. H. Carothers. 1995. Evolutionary Consequences of Extinctions in Populations of a Hawaiian Honeycreeper. *Conservation Biology* 9(1):107-113.
- Sokolov, A. P., P. H. Stone, C. E. Forest, R. Prinn, M. C. Sarofim, M. Webster, S. Paltsev, C. A. Schlosser, D. Kicklighter, S. Dutkiewicz, J. Reilly, C. Wang, B. Felzer, and H. D. Jacoby. 2009. Probabilistic forecast for 21st century climate based on uncertainties in emissions (without policy) and climate parameters. *Journal of Climate* DOI: 10.1175/2009JCLI2863.1.
- U.S. Endangered Species Act (ESA) 1973. U.S. Code 16, chap. 35.
- U.S. Energy Information Administration (USEIA). 2004. International energy outlook. <http://www.eia.doe.gov/oiaf/ieo/emissions.html>. Accessed May 5.
- U.S. Energy Information Administration (USEIA). 2009. International energy outlook. <http://www.eia.doe.gov/oiaf/ieo/emissions.html>. Accessed May 5.
- U.S. Fish and Wildlife Service (USFWS). 2006. Revised Recovery Plan for Hawaiian Forest Birds. Region 1, Portland.
- U.S. Fish and Wildlife Service (USFWS) 2009. USFWS website. Accessed 16 September 2009. <https://ecos.fws.gov/roar/pub/planImplementationStatus.action?documentId=1001509>
- U.S. Fish and Wildlife Service (USFWS). 2010. National Fish and Wildlife Climate Adaptation Strategy. <http://www.fws.gov/nfwcaspvpgp.html> Accessed 5 May 2010.
- Valkiunas, G. 2005. Avian malaria parasites and other haemosporidia. CRC Press, Boca Raton.
- VanderWerf, E. A., and J. L. Rohrer. 1996. Discovery of an 'I'iwi population in the Ko'olau Mountains of O'ahu. *'Elepaio* 56(4):25-28.
- VanderWerf, E. A., and D. G. Smith. 2002. Effects of alien rodent control on demography of the O'ahu 'Elepaio, an endangered Hawaiian forest bird. *Pacific Conservation Biology* 8:73-81.
- van Riper, C. III, S. G. van Riper, M. L. Goff, and M. Laird. 1986. The epizootiology and ecological significance of malaria in Hawaiian land birds. *Ecological Monographs* 56:327-344.
- Warner, R. E. 1968. The role of infectious diseases in the extinction of the endemic Hawaiian avifauna. *Condor* 70:101-120.
- Woodworth, B. L., C. T. Atkinson, D. A. LaPointe, P. J. Hart, C. S. Spiegel, E. J. Tweed, C. Henneman, J. LeBrun, T. Denette, R. DeMots, K. L. Kozar, D. Triglia, D. Lease, A. Gregor, T. Smith, and D. Duffy. 2005. Host population persistence in the face of introduced vector-borne diseases: Hawaii amakihi and avian malaria. *Proceedings of the National Academy of Sciences (USA)* 102:1531-1536
- Woodworth, B.L. and T. K. Pratt. 2009. Life history and demography. Chapter 8 in *Conservation Biology of Hawaiian Forest Birds: Implications for Island Avifauna*.

T. K. Pratt, C.T. Atkinson, P. C. Banko, J. D. Jacobi, B. L. Woodworth (eds.). Yale University Press.

Woody, G. 2011. Wildlife at Risk Face Long Line at U.S. Agency. New York Times. 20 April.

IntechOpen

IntechOpen

Climate Change - Geophysical Foundations and Ecological Effects

Edited by Dr Juan Blanco

ISBN 978-953-307-419-1

Hard cover, 520 pages

Publisher InTech

Published online 12, September, 2011

Published in print edition September, 2011

This book offers an interdisciplinary view of the biophysical issues related to climate change. Climate change is a phenomenon by which the long-term averages of weather events (i.e. temperature, precipitation, wind speed, etc.) that define the climate of a region are not constant but change over time. There have been a series of past periods of climatic change, registered in historical or paleoecological records. In the first section of this book, a series of state-of-the-art research projects explore the biophysical causes for climate change and the techniques currently being used and developed for its detection in several regions of the world. The second section of the book explores the effects that have been reported already on the flora and fauna in different ecosystems around the globe. Among them, the ecosystems and landscapes in arctic and alpine regions are expected to be among the most affected by the change in climate, as they will suffer the more intense changes. The final section of this book explores in detail those issues.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Anthony Povillitis (2011). Global Heating Threatens the `I`iwi (*Vestiaria coccinea*), Currently a Common Bird of Upper Elevation Forests in Hawaii, *Climate Change - Geophysical Foundations and Ecological Effects*, Dr Juan Blanco (Ed.), ISBN: 978-953-307-419-1, InTech, Available from: <http://www.intechopen.com/books/climate-change-geophysical-foundations-and-ecological-effects/global-heating-threatens-the-i-iwi-vestiaria-coccinea-currently-a-common-bird-of-upper-elevation-for>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2011 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike-3.0 License](https://creativecommons.org/licenses/by-nc-sa/3.0/), which permits use, distribution and reproduction for non-commercial purposes, provided the original is properly cited and derivative works building on this content are distributed under the same license.

IntechOpen

IntechOpen