We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Effects of High Night Temperature on Crop Physiology and Productivity: Plant Growth Regulators Provide a Management Option

Abdul Razack Mohammed and Lee Tarpley
Texas AgriLife Research and Extension Center, Beaumont, Texas
USA

1. Introduction

Crop production is experiencing both increases in the frequency and intensity of high night temperatures (HNT) (IPCC, 2007). The HNT threatens the sustainability of crop production both currently and in the future. Recent meteorological data indicated faster increases in night temperatures (NT) than day temperatures (Alward et al., 1999). Experimental evidence also indicated the importance of assessing the effects of NT separately from the day temperature, as a smaller increase in NT can show a drastic decrease in crop production (Peng et al., 2004). Studies have indicated decreased crop yields as a result of HNT (Hall, 1992; Mohammed & Tarpley, 2009a). Apart from HNT, climate change will also increase a variety of environmental stresses affecting crop production, namely heat, drought, salinity, and relative humidity, hence farming in the future climate will have to be better adapted to a range of abiotic and biotic stresses. A long-term approach to negate the effects of abiotic stress is to develop stress-tolerant cultivars. The short-term approach includes the use of agrochemicals, especially those with plant growth regulator (PGR) capabilities, for the prevention and/or amelioration of various environmental stresses, including heat stress. Studies focusing on day temperature stress and, more recently, on NT stress have increased, but, studies looking at the effects of PGR under heat stress are rare (Ashraf & Foolad, 2007). This chapter primarily focuses on the effects of HNT on crop production and the beneficial effects of PGR (glycine betaine [GB] and salicylic acid [SA]) application against heat stress on crop production with special emphasis on rice (*Oryza sativa* L.) production.

2. Night temperature and its impact on crop production

Global climate change is a dynamic process affecting global air temperature, oceanic temperature, rainfall, wind and quality of incoming solar radiation. Global circulation models predict 1.4 to 5.8 °C rise in global temperature because of projected increase in the concentrations of all greenhouse gases by the end of the 21st century (Intergovernmental Panel on Climate Change [IPCC], 2007). Much of this increase in average daily temperature is projected to be due to an increase in NT. Night temperatures are expected to increase at a faster rate than day temperatures due to less radiant heat loss because of increased

cloudiness (Alward et al., 1999). In some cases, increases in NT have been documented. For example, the scientists at the International Rice Research Institute (IRRI) reported an increase of 1.13 °C in NT over a period of 25 years (1979-2003) in the Philippines (Peng et al., 2004). The HNT increased at a rate of 0.18 °C per decade over a period of 45 years (1950-1995) in Libya (Jones et al., 1999).

Long- and short-term episodes of heat stress are predicted to occur more frequently as a result of global warming, affecting many aspects of crop growth and development, reducing crop yield and decreasing crop quality. Many studies have reported the impacts of long- and short-term temperature stresses on crop production. However, most of the studies assume no difference in the influence of day versus night temperature (Peng et al., 2004). High night temperature is known to decrease yield in several crops like cereals [rice, wheat (*Triticum aestivum* L.) and corn (*Zea Mays* L.)], legumes [cowpea (*Vigna unguiculata* L. Walp.)], oil seeds [soybean (*Glycine max* L. Merr.) and sunflower (*Helianthus annuus* L.)], vegetables [tomato (*Solanum lycopersicum* L.)], fruits [apples (*Malus domestica* L.) and pineapple (*Ananas comosus* L.)] and fibers [cotton (*Gossypium hirsutum* L.)] (Chang, 1981; Gibson & Mullen, 1996; Izquierdo et al., 2002; Kondo & Takahashi, 1989; Loka & Oosterhuis, 2010; Mohammed & Tarpley, 2009a; Neales et al., 1980; Peters et al., 1971; Seddigh & Jolliff, 1984; Warrag & Hall, 1984; Willits & Peet, 1998).

High night temperature decreases crop production by decreasing photosynthetic function, sugar and starch content (Loka & Oosterhuis, 2010; Turnbull et al., 2002), increasing respiration rate (Mohammed & Tarpley, 2009b), suppressing floral bud development (Ahmed & Hall, 1993), causing male sterility and low pollen viability and hastening crop maturity (Mohammed & Tarpley, 2009a; Seddigh and Jolliff, 1984). Another effect of HNT that can contribute to reduced crop yield is decreased antioxidant capacity of the plants. Under normal physiological conditions, the toxic effects of reactive oxygen species (ROS) are minimized by enzymatic and non-enzymatic antioxidants. Under stress conditions, oxidant levels can overwhelm the antioxidant levels leading to cell damage. The increased production of ROS [oxide radical (O_2^-), H_2O_2 , and the hydroxyl radical ($-OH$)], or the plant's decreased ability to neutralize ROS, as a result of heat stress negatively affects many physiological processes in plants, thus decreasing yield.

2.1 Methodology for determining rice plant response to high night temperature stress

Plant height and the numbers of tillers and leaves were recorded at harvest. Daily observations were made for the appearance of panicles. At harvest, leaves were separated from the stems and leaf area was measured using a CI-251 area meter (CID Inc., Camas, Washington, USA). During the rice plant vegetative stage, leaf photosynthetic rate, internal CO_2 concentration and respiration rate were measured using LI-6400 portable photosynthesis system (LI-COR Inc., Lincoln, Nebraska, USA). Leaf nitrogen content was determined using a FP-528 Nitrogen/Protein analyzer (LECO Corporation, St. Joseph, Michigan, USA) and was expressed as percentage (%; w/w). Total chlorophyll content, pollen germination and spikelet fertility were determined using the procedures from Mohammed & Tarpley (2011a), and relative injury to the membrane and total antioxidant capacity were determined using procedures from Mohammed & Tarpley (2009b).

2.2 Effects of high night temperature on plant morphology

Plant morphology deals with plant development, form, and structure. In a narrow sense, plant morphology refers to the external appearance of the plant. For example plant height

and the numbers of tillers, leaves, panicles and grains are referred as rice morphological parameters. The effects of HNT on plant morphology vary from negative to positive, including no effect. Seddigh & Jolliff (1984) reported that HNT promoted early vegetative growth and hastened physiological maturity, but did not affect morphological characteristics such as plant height, number of auxiliary branches and number of nodes in soybean. The HNT has no effect on rice (cultivar Cocodrie which is a japonica cultivar) plant height and number of tillers but can increase number of leaves and leaf area (Fig. 1). However, Cheng et al. (2009) and Yoshida et al. (1981) reported reduction in plant height, number of tillers and total biomass for rice cultivar 'IR 72' (indica cultivar) as a result of HNT. Similar results were seen in redstem stork's bill (*Erodium cicutarium* L.) for which HNT decreased number of branches, leaf area and plant dry matter (Blackshaw & Entz, 1995). The decreased production of plant dry matter was directly correlated with decreased production of photosynthates.

Fig. 1. Effects of high night temperature on rice morphology. Each bar represents average $\pm$ S.E. Different letters indicate means are significantly different at the $P<0.05$ level.

2.3 Effects of high night temperature on plant physiology

Plant physiology deals with the functioning of plants and is closely related with plant morphology. Fundamental processes such as photosynthesis, respiration, plant water and nutrition status, plant hormone functions and translocation of photosynthates are encompassed in plant physiology. Photosynthesis is a complex process and leaf photosynthetic rates depend upon leaf chlorophyll and nitrogen content, photosystems, stomatal characteristics and enzyme activities. The environmental stresses have a direct effect on leaf photosynthetic rates and an indirect effect through their effects on leaf

chlorophyll and nitrogen content, stomatal characteristics and enzyme activities. The HNT had positive and negative effects, including no effect, on the following day's photosynthetic rates (Frantz et al., 2004; Mohammed & Tarpley 2009a). Rice leaf photosynthetic rates showed negative correlation with HNT (Fig. 2). The decrease in photosynthetic rates as a result of HNT was accompanied with decrease in leaf chlorophyll and leaf nitrogen content in rice (Fig. 2). Premature loss of chlorophyll as a result of heat stress has been noticed in many crop species (Guo et al., 2006; Reynolds et al., 1994). Damage to the photosynthetic electron transport particularly at the site of photosystem II (Havaux and Tardy, 1996) and inhibition of photosynthetic CO₂ fixation (Sayed et al., 1989; Yamane et al., 1997) as a result of heat stress can also reduce photosynthetic rates, thus leading to decrease in yield. Apart from decreases in photosynthetic rates, increased respiration rates can also decrease the crop yield.

Plant respiration is one of the important processes in terms of understanding plant growth and development when subjected to stress. Increase in respiration from climate warming is of serious concern as respiratory processes can consume a larger portion of total photosynthates (Paembonan et al., 1992). On average, the carbon lost from respiratory metabolism within an individual plant ranges between 30 and 70% of the carbon gained through photosynthesis (Peterson & Zelitch, 1982). Respiration is typically partitioned into the functional components of construction (growth), maintenance and ion uptake to facilitate our understanding of the impact of the environment on respiratory processes (Amthor, 1986; Farrar, 1985; Lambers, 1985). Maintenance respiration is mainly associated with turnover of proteins and lipids and maintenance of ion concentrations across membranes (Penning de Vries, 1975). It is the most responsive to environmental changes among the functional components of respiration (Ryan, 1991). At high temperatures, the cost of maintenance increases to support protein turnover and to maintain active ion fluxes across the membranes (Penning de Vries, 1975), thereby increasing maintenance respiration. An increase in NT from 27 °C to 32 °C increased respiration rates by 40% in rice leaves (Fig. 2). Previous studies on rice, cotton, lettuce (*Lactuca sativa* L.), tomato and soybean indicated an increase in maintenance respiration with warmer nights (Frantz et al., 2004; Loka & Oosterhuis, 2010; Mohammed & Tarpley, 2009b). Hence, HNT can stimulate respiration rates, thereby negatively affecting the yield (Zheng et al., 2002). The other consequence of increased respiration is increased production of ROS.

The production of ROS is an unavoidable consequence of aerobic respiration, with the majority of the ROS produced in photosynthetic tissue (mostly leaves) in the dark by mitochondrial electron transport chain activity (miETC) (McDonald & Vanlerberghe, 2005). The production of ROS by mitochondria has been shown to increase in many plants as a result of biotic and abiotic stresses (McDonald & Vanlerberghe, 2005; Moller, 2001). An increase in NT increases maintenance respiration and thus increases production of ROS. Physiological injury due to heat stress has been associated with increases in oxidative damage to the membrane in plant species (Larkindale & Knight, 2002). Plants increase maintenance respiration to support repair mechanisms of the membranes due to oxidative damage (Amthor & McCree, 1990). Thus, an increase in respiration occurs 1) with an increase in temperature (Huang et al., 1998) and 2) to support repair mechanisms of the membranes due to oxidative damage (Amthor & McCree, 1990). In addition, ROS interferes with photosynthesis and respiration by disrupting water, ion, and organic solute movement across plant membranes by affecting membrane stability (Christiansen, 1978).

Destabilized membranes are leaky membranes, thus interpreting the amount of electrolytic leakage from the membrane can be used as an indicator of cell membrane stability (Sullivan & Ross, 1979). Many studies have examined stability loss of the membranes subjected to environmental stresses by measuring electrolytic leakage from the membranes (Blum & Ebrercon, 1981; Dexter, 1956; Ibrahim and Quick, 2001; Ismail and Hall, 1999; Sullivan, 1972). An increase in NT from 27 °C to 32 °C increased injury to the membrane by 60% in rice leaves (Fig. 2). Similar results of increased injury to the membrane as a result of heat stress were seen in many crop species (Ibrahim & Quick, 2001; Ismail & Hall, 1999,, Martineau et al., 1979; Mohammed & Tarpley, 2009b). The properties of the photosynthetic system, including key enzymes and thylakoid membrane activities depend on the stability of membranes (Björkman et al., 1980). Moreover, it is well known that a functional cell-membrane system is central to crop productivity and acclimation of plants to high temperature (Raison et al., 1980). Mohammed & Tarpley (2009b) and Reynolds et al. (1994) positively associated membrane stability with yield performance under heat-stressed conditions in rice and wheat.

Fig. 2. Effects of high night temperature on rice physiology. Each bar represents average $\pm$ S.E. Different letters indicate means are significantly different at the $P<0.05$ level.

2.4 Effects of high night temperature on yield and yield-related parameters

The reproductive phase is relatively more sensitive than the vegetative phase to heat stress in many crop species (Hall, 1992). Moreover, differential temperature sensitivity for reproductive and vegetative growth has been reported in rice (Baker et al., 1992), soybean (Baker et al., 1989), wheat (Mitchell et al., 1993) and many other crops. Warrag & Hall (1984) reported that HNT induced male sterility and excessive floral abscissions in cowpea, whereas high day temperature had no adverse effect. The HNT can decrease crop yields by decreasing crop growth duration, suppressing floral bud development and decreasing pollen production and viability (Ahmed & Hall, 1993; Mohammed & Tarpley, 2009a; Prasad et al., 1999; Prasad et al., 2006). Increase in NT from 27 °C to 32 °C decreased crop growth duration by 2 days, as indicated by emergence of 1st panicle, in rice (Fig. 3). Decreased crop duration due to HNT decreases the time for carbohydrate accumulation, thus decreasing the yield (Cantarero et al., 1999). Badu-Apraku et al. (1983) attributed dramatic yield loss under high temperatures during the period of grain filling to sucrose availability (Afuakwa et al., 1984) and activity levels of enzymes involved in starch and sugar metabolism (Singletary et al., 1994). Moreover, the suppression of floral buds and flowering under HNT was also attributed to a shortage of photosynthetic assimilates supplied to the floral buds (Guinn, 1974), and/or an inability of floral buds to mobilize carbohydrates under heat stress (Dinar & Rudich, 1985). Decreased pollen germination due to heat stress has been noticed in many crop species (Hall, 1992; Matsui et al., 2001; Mohammed & Tarpley, 2009a). This decrease in pollen germination at high temperatures is due to poor anther dehiscence and pollen reception (Prasad et al., 2006), reduced pollen swelling and decreased anther pore size (Matsui & Kagata, 2003).

Fig. 3. Effects of high night temperature on rice yield-related parameters. Each bar represents average $\pm$ S.E. Different letters indicate means are significantly different at the $P<0.05$ level.

In most cereal crops, spikelet fertility, described as the ratio between filled grains and total grains, decreased with decrease in pollen germination. Increase in NT from 27 °C to 32 °C decreased spikelet fertility by 70% in rice (Fig. 3). Apart from pollen germination, hormonal balance in the sink (Micheal & Beringer, 1980) and/or availability and transportability of photosynthates to the sink from the source (Afuakwa et al., 1984) and/or inability of floral buds to mobilize carbohydrates (Dinar & Rudich, 1985) and/or altered activities of starch and sugar biosynthesis enzymes (Keeling et al., 1994; Singletary et al., 1994) can govern the spikelet fertility at high temperatures. The capacity to survive heat stress varies with plant species, genotype and developmental stage. The PGR provide an option to protect the plants from heat stress, thus providing sustainability for crop productivity.

3. Plant growth regulators

Genetic improvement and the modification of agronomic practices can be beneficial for crop adaptation to future climate conditions. However, these approaches are more applicable for long-term episodes of heat stress. The use of agrochemicals, especially those with PGR capabilities, for the prevention and/or amelioration of various environmental stresses, such as those resulting from global climatic change, is a viable approach to make crop production more resilient to short-term episodes of heat stress. Plant growth regulators are natural or synthetic chemical compounds that are used to promote or inhibit plant growth and development or alter specific physiology or metabolic factors. Like plant hormones, small concentrations of PGRs are enough to influence plant growth and development. In recent years, numerous PGRs have been developed to promote crop production under stress conditions. Glycine betaine, SA, vitamin E, proline and choline are some of the agrochemicals which can induce stress-tolerance (thermotolerance, drought tolerance, cold tolerance and/or salinity tolerance) in various crop plants. In this chapter, we focus on the beneficial effects of GB and SA application against heat stress on crop production.

Oxygen is potentially toxic to all organisms because metabolism and environmental stresses generate ROS from the oxygen. Under normal physiological conditions, the toxic effects of ROS are minimized by enzymatic and non-enzymatic antioxidants (Kreiner et al., 2002). Under stress conditions, oxidant levels overwhelm the antioxidant levels leading to cell damage (Kreiner et al., 2002). Plants exposed to environmental stress, such as heat, cold, drought and salinity produce ROS, which damage macromolecules and cell membranes (Zhang & Kirkham, 1996). Angiosperms possess several enzymatic and non-enzymatic scavenging systems to minimize deleterious effects of ROS. These include lipid-soluble antioxidants (e.g. α -tocopherol and β -carotene), water-soluble reactants (e.g. ascorbic acid and glutathione), and enzymatic antioxidants (e.g. superoxide dismutase, catalase and enzymes of the ascorbate and glutathione cycle) (Zhang & Kirkham, 1996). Glycine betaine and SA are synthesized in the plants and play important but different, roles in preventing oxidative damage to the membranes (Bowler et al., 1992; Demiral & Turkan, 2004) and inducing thermotolerance in plants subjected to stresses (Ashraf & Foolad, 2007; Caldas et al., 1999; Farooq et al., 2008a, 2008b; Larkindale & Huang, 2004; Mohammed & Tarpley, 2009b; Pan et al., 2006; Raskin, 1992). The commercial availability of GB and SA provides potential crop-management options to reduce crop heat stress events.

Glycine betaine, an important osmoprotectant is an amino-acid derivative that is naturally synthesized and accumulates under stress conditions in some plants (Bohnert & Jensen, 1996; Demiral & Turkan, 2004). It enhances stress tolerance by protecting enzymes (Paleg et

al., 1981), photosystem II (Allakhverdiev et al., 1996), membrane integrity and antioxidant status of the plant (Mohammed & Tarpley, 2009b). Glycine betaine may also protect plants from environmental stresses indirectly via its role in signal transduction and ion homeostasis (John, 2002; Yilmaz, 2004). It regulates signal transduction and ion homeostasis by affecting MAP kinases, and expression of many genes related to signal transduction (John, 2002). Exogenous application of GB improved growth, survival and induced stress tolerance in many crop species (Diaz-Zorita et al., 2001; Harinasut et al., 1996; Rajasekaran et al., 1997) and also increased total antioxidant capacity and antioxidant enzyme activities (Farooq et al., 2008a; Mohammed & Tarpley, 2009b).

Salicylic acid is an endogenous plant growth regulator of phenolic nature and is considered to be a potent plant hormone because of its diverse regulatory roles in plant metabolism (Raskin, 1992). It plays an important role in the regulation of plant growth, development and responses to abiotic and biotic stresses (Raskin, 1992; Senaratna et al., 2000; Yalpani et al., 1994) by affecting photosynthetic rate, stomatal conductance, transpiration, ion uptake and transport, respiration rates, membrane stability, antioxidant capacity and antioxidant enzyme activities (Farooq et al., 2008a; Harper & Balke, 1981; Khan et al., 2003; Mohammed & Tarpley, 2009b). In addition, SA is considered to be an important signaling molecule involved in stress tolerance. It also promotes pathogenesis-related (PR) gene expression and induces systemic acquired resistance (SAR) in plants (Ryals et al., 1994). Besides providing disease resistance to the plants, SA can modulate plant responses to a wide range of oxidative stresses (Shirasu et al., 1997). It plays an essential role in preventing oxidative damage in plants by detoxifying superoxide radicals (Bowler et al., 1992) and by altering antioxidant capacity and antioxidant enzyme activities in plants (Chen et al., 1997; Dat et al., 1998; Fodor et al., 1997; Larkindale & Huang, 2004; Mohammed & Tarpley, 2009b; Rao et al., 1997; Wang & Li, 2006). Plants pre-treated with SA have shown increased thermotolerance (Larkindale & Knight, 2002).

3.1 Effects of plant growth regulators on plant morphology and physiology under high night temperature

Exogenous application of PGRs on plants under high night temperatures have shown positive, negative or no effect on plant morphology. Exogenous foliar application of GB or SA did not affect plant height in rice (Fig. 4). However, Farooq et al., (2008a, 2008b) state that seed priming of maize (*Zea mays* L.) with GB or SA increased shoot length, root length and biomass production. Increase in plant height as a result of SA application is due to increased cell enlargement, endoreduplication and/or cell division (Kang et al., 2007; Vanacker et al., 2007).

Glycine betaine and SA have profound effects on crop production through their effects on crop physiology. Foliar application of GB or SA increases leaf photosynthetic rates and decreases internal CO₂ concentration, respiration rates and membrane injury in rice (Fig. 5). Similar results were seen in other crop species and growth conditions where application of GB or SA increased photosynthetic rates and decreased respiration rates and injury to the membranes (Farooq et al., 2008a; Khan et al., 2010; Zhou et al., 1999).

In rice, application of GB or SA did not affect leaf chlorophyll or nitrogen content (Fig. 5). However, studies have shown that exogenous application of GB stabilizes pigments and prevents water oxidation and photooxidation (Cha-um et al., 2006, Sakamoto et al. 1998). Stabilization of pigments (chlorophyll and carotenoids) aids in light energy capture as required for photosynthesis. Several studies have reported that application of GB aids in

protecting the photosynthetic machinery of the plant organelles by stabilizing the ultrastructure of the chloroplast, photosystem-II reaction centers and maintaining the oxygen-evolving machineries (Allakhverdiev et al., 2003; Busheva & Apostolova, 1997; Lee et al., 1997; Makela et al., 2000; Pospisil & Dau 2000; Sakamoto & Murata, 2000). On the other hand SA increases photosynthetic efficiency by increasing pigment concentrations and protecting photosystem function via activation of antioxidant defense systems as a signaling molecule (Dat et al., 1998; Ervin et al., 2005; Senaratna et al., 2001). The other physiological parameters such as respiration rates, transpiration rate and injury to leaf membrane decreased with application of GB or SA (Mohammed & Tarpley, 2011b). Studies have reported antitranspiration effects of GB and SA applications in crop species suggesting that GB and SA play an important role in controlling stomatal opening and closing under stress conditions (Cha-um et al., 2006; Khan et al., 2010; Larque-Saavedra, 1978). The most important effect of PGR's on crop production is through their ability to maintain the integrity of the membranes. The functional cell-membrane system is central to crop productivity. Stable membranes facilitate free movement of ions and water across the membrane and provide a favorable environment for enzymes and thylakoid membrane activities (Björkman et al., 1980). Several studies have reported that GB or SA application can maintain membrane integrity under stress conditions (Bohnert & Jensen, 1996; Farooq et al., 2008a, 2008b; Hussain et al., 2008; Mohammed & Tarpley, 2011b; Pan et al., 2006) by up regulating antioxidant enzyme activities and total antioxidant capacity (Demiral & Türkan, 2004; Farooq et al., 2008a, 2008b; He et al., 2005; Mohammed & Tarpley, 2009b).

Fig. 4. Effects of glycine betaine and salicylic acid on rice morphology under high night temperature. Each bar represents average $\pm$ S.E. Different letters indicate means are significantly different at the $P<0.05$ level.

Fig. 5. Effects of glycine betaine and salicylic acid on rice physiology under high night temperature. Each bar represents average $\pm$ S.E. Different letters indicate means are significantly different at the $P < 0.05$ level.

3.2 Effects of plant growth regulators on plant antioxidant enzyme activities and total antioxidant capacity under high night temperature

The antioxidant status of the plant is closely associated with its stress tolerance (Smirnoff, 1995). The severity of ROS-induced damage depends upon the antioxidant status of the plant and activities of antioxidant enzymes such as superoxide dismutase (SOD), ascorbate peroxidase (AP), catalase (CAT) and glutathione reductase (GR). Many studies have shown an increase in endogenous antioxidant levels as a result of exogenous application of GB or SA (Chen et al., 1997; Diaz-Zorita et al., 2001; Fang et al., 2009; Mohammed & Tarpley, 2009b). In rice, exogenous application of GB or SA increased total antioxidant capacity under high night temperature (Fig. 6). These increases in the levels of endogenous antioxidant are due to increases in antioxidant enzyme activities (Demiral & Türkan, 2004; Farooq et al., 2008a, 2008b; He et al., 2005; Khan et al., 2010).

Fig. 6. Effects of glycine betaine and salicylic acid on total antioxidant capacity under high night temperature. Each bar represents average $\pm$ S.E. Different letters indicate means are significantly different at the $P < 0.05$ level.

3.3 Effects of plant growth regulators on yield and yield-related parameters under high night temperature

Rice phenology under high night temperature was not altered by exogenous application of GB or SA as indicated by 1st panicle emergence date (Fig. 7). However, previous studies by Oota (1975) and Datta & Nanda (1985) showed that SA application induces early flowering in duckweeds (*Lemna gibba* L.) and chenna millet (*Panicum miliaceum* L.). Exogenous application of GB or SA increased spikelet fertility in rice (Fig. 7). The increase in spikelet fertility was due to increased photosynthesis, and decreased respiration and injury to the membrane stability as a result of GB or SA application. Exogenous application of GB or SA increases stability of the membranes as a result of increased antioxidant capacity of the plants. The enhanced membrane stability is responsible for increased photosynthesis and decreased respiration, which in turn increases crop production. In a nutshell, exogenous applications of GB or SA have been shown to increase stress tolerance, and often enhance plant growth and final crop yield under stress conditions, including HNT.

4. Conclusion

Yield, the final manifestation of all the physiological processes, decreased due to high night temperatures. The decrease in crop yields as a result of HNT was due to increased respiration and decreased photosynthesis and membrane stability. The decrease in photosynthesis was associated with decreases in photosynthesis-governing parameters such as stomatal conductance, internal CO_2 concentration, leaf chlorophyll content and leaf nitrogen, as a result of HNT. Moreover, decreased crop growth duration, pollen production and viability also contributed to decreased yields under HNT. Exogenous application of PGRs such as GB and SA increased production of photosynthates and decreased consumption of photosynthates and injury to the membrane, thereby increasing crop productivity under HNT. The exogenous application of GB and SA increases antioxidant levels, thereby protecting the membranes and enzymes against heat-induced ROS-mediated degradation, thus increasing crop productivity. However, there is a significant variation in

Fig. 7. Effects of glycine betaine and salicylic acid on rice yield-related parameters under high night temperature. Each bar represents average $\pm$ S.E. Different letters indicate means are significantly different at the $P < 0.05$ level.

response to exogenous application of PGRs such as GB and SA among different plant species. This varied response among the plant species is due to 1) plant quantitative response to exogenous application of PGRs such as GB and SA at different stages of plant development, 2) plant exhibition of different responses to PGRs such as GB and SA at different stages of plant development and 3) variation in the effective concentration of a particular PGR among species. The PGRs such as GB and SA provide an option to protect the plants from different stresses like high day temperatures, high night temperature, drought and salinity, however, over-application of the PGRs such as GB and SA may be toxic to some plants resulting in growth inhibition or yield reduction.

5. Acknowledgment

We thank Texas Rice Belt Warehouse and Texas Rice Research Foundation for their support throughout this work.

6. References

Afuakwa, J.J.; Crookston, R.K. & Jones, R.J. (1984). Effect of Temperature and Sucrose Availability on Kernel Black Layer Development in Maize. *Crop Science*, Vol. 24, pp. 285-288, ISSN 0011-183X.

- Ahmed, F.E. & Hall, A.E. (1993). Heat Injury during Early Floral Bud Development in Cowpea. *Crop Science*, Vol. 33, pp. 764-767, ISSN 0011-183X.
- Allakhverdiev, S.I.; Feyziev, Y.M.; Ahmed, A.; Hayashi, H.; Aliev, J.A.; Kimlov, V.V.; Murata, N. & Carpentier, R. (1996). Stabilization of Oxygen Evolution and Primary Electron Transport Reactions in Photosystem II Against Heat Stress with Glycinebetaine and Sucrose. *Journal of Photochemistry and Photobiology*, Vol. 34, pp. 149-157, ISSN 0031-8655.
- Alward, R.D.; Detling, J.K. & Milchunas, D.G. (1999). Grassland Vegetation Changes and Nocturnal Global Warming. *Science*, Vol. 283, pp. 229-231, ISSN 0036-8075.
- Amthor, J.S. & McCree, K.J. (1990). Carbon Balance of Stress Plants: A Conceptual Model for Integrating Research Results, In: *Stress Response in Plants: Adaptation and Acclimation Mechanism*, R.G. Alscher & J.R. Cumming (Eds.), 1-15, Alan R. Liss, New York, USA.
- Amthor, J.S. (1986). Evolution and Applicability of a Whole Plant Respiration Model. *Journal of Theoretical Biology*, Vol. 122, pp. 473-490, ISSN 0022-5193.
- Ashraf, M. & Foolad, M.R. (2007). Roles of Glycine Betaine and Proline in Improving Plant Abiotic Stress Resistance. *Environmental and Experimental Botany*, Vol. 59, pp. 206-216, ISSN 0098-8472.
- Badu-Apraku, B.; Hunter, R.B., & Tollenaar, M. (1983). Effect of Temperature During Grain Filling on Whole Plant and Grain Yield in Maize (*Zea mays* L.). *Canadian Journal of Plant Science*, Vol. 63, pp. 357-363, ISSN 0008-4220.
- Baker, J.T.; Allen, L.H. Jr. & Boote K.J. (1992). Response of Rice to Carbon Dioxide and Temperature. *Agricultural and Forest Meteorology*, Vol. 60, pp. 153-166, 0168-1923.
- Baker, J.T.; Allen, L.H., Jr.; Boote, K.J.; Jones, P. & Jones, J.W. (1989). Response of Soybean to Air Temperature and Carbon Dioxide Concentration. *Crop Science*, Vol. 29, pp. 98-105, ISSN 0011-183X.
- Blackshaw, R.E. & Entz, T. (1995). Day and Night Temperature Effects on Vegetative Growth of *Erodium cicutarium*. *Weed Research*, Vol. 35, pp. 471-476, ISSN 0043-1737.
- Blum, A. & Ebercon, A. (1981). Cell Membrane Stability as a Measure of Drought and Heat Tolerance in Wheat. *Crop Science*, Vol. 21 pp. 43-47, ISSN 0011-183X.
- Bohnert, H.J. & Jensen, R.G. (1996). Strategies for Engineering Water-stress Tolerance in Plants. *Trends in Biotechnology*, Vol. 14, pp. 89-97, ISSN 0167-7799.
- Bowler, C.; Montagu, M.V. & Inze, D. (1992). Superoxide Dismutase and Stress Tolerance. *Annual Review of Plant Physiology and Plant Molecular Biology*, Vol. 43, pp. 83-116, ISSN 1040-2519.
- Busheva, M. & Apostolova, E. (1997). Influence of Saccharides and Glycinebetaine on Freezing of Photosystem II-enriched Particles: a Chlorophyll Fluorescence Study. *Photosynthetica*, Vol. 34, pp. 591-594, ISSN 0300-3604.
- Caldas, T.; Demont-Caulet, N.; Ghazi, A. & Richarme, G. (1999). Thermoprotection by Glycine Betaine and Choline. *Microbiology*, Vol. 145, pp. 2543-2548, ISSN 1350-0872.
- Cantarero, M.G.; Cirilo, A.G. & Andrade, F.H. (1999). Night Temperature at Silking Affects Kernel Set in Maize. *Crop Science*, Vol. 39, pp. 703-710, ISSN 0011-183X.
- Cha-um, S.; Supaibulwatana, K. & Kirdmanee, C. (2006). Water Relation, Photosynthetic Ability and Growth of Thai Jasmine Rice (*Oryza sativa* L. ssp. indica cv. KDML105)

- to Salt Stress by Application of Exogenous Glycinebetaine and Choline. *Journal of Agronomy and Crop Science*, Vol. 192, pp. 25-36, ISSN 0931-2250.
- Chen, Z.; Iyer, S.; Caplan, A.; Klessig, D.F. & Fan, B. (1997). Differential Accumulation of Salicylic Acid and Salicylic Acid-sensitive Catalase in Different Rice Tissues. *Plant Physiology*, Vol. 114, pp. 193-201, ISSN 0032-0889.
- Cheng, W.; Sakai, H.; Yagi, K. & Hasegawa, T. (2009). Interactions of Elevated [CO₂] and Night Temperature on Rice Growth and Yield. *Agricultural and Forest Meteorology*, Vol. 149, pp. 51-58, ISSN 0168-1923.
- Christiansen, M.N. (1978). The Physiology of Plant Tolerance to Temperature Extremes. In: *Crop Tolerance to Suboptimal Land Conditions*, G.A. Jung (Ed.), 173-191, American Society of Agronomy, Madison, Wisconsin, USA.
- Dat, J.F.; Lopez-Delgado, H.; Foyer, C.H. & Scott, I.M. (1998). Parallel Changes in H₂O₂ and Catalase During Thermotolerance Induced by Salicylic Acid or Heat Acclimation in Mustard Seedlings. *Plant Physiology*, Vol. 116, pp. 1351-1357, ISSN 0032-0889.
- Datta K.S. & Nanda K.K. (1985). Effect of Some Phenolic Compounds and Gibberellic Acid on Growth and Development of Cheena Millet (*Panicum miliaceum* L.). *Indian Journal of Plant Physiology*, Vol. 28, pp. 298-302, ISSN 0019-5502.
- Demiral, T. & Türkan, I. (2004). Does Exogenous Glycinebetaine Affect Antioxidative System of Rice Seedlings Under NaCl Treatment? *Journal of Plant Physiology*, Vol. 161, pp. 1089-1100, ISSN 0176-1617.
- Dexter, S.T. (1956). Evaluation of Crop Plants for Winter Hardiness. *Advances in Agronomy*, Vol. 8, pp. 203-209, ISSN 0065-2113.
- Diaz-Zorita, M.; Fernandez-Canigia, M.V. & Grosso, G.A. (2001). Application of Foliar Fertilizers Containing Glycinebetaine Improved Wheat Yields. *Journal of Agronomy and Crop Science*, Vol. 186, pp. 209-215, ISSN 0931-2250.
- Dinar, M. & Rudich, J. (1985). Effect of Heat Stress on Assimilate Partitioning in Tomato. *Annals of Botany*, Vol. 56, pp. 239-249, ISSN 0305-7364.
- Ervin, E.H.; Zhang, X. & Schmidt, R.E. (2005). Exogenous Salicylic Acid Enhances Post-Transplant Success of Heated Kentucky Bluegrass and Tall Fescue Sod. *Crop Science*, Vol. 45, pp. 240-244, ISSN 0011-183X.
- Fang, C-X.; Xiong, J.; Qiu, L.; Wang, H-B.; Song, B-Q.; He, H-B.; Lin, R-Y. & Lin, W-X. (2009). Analysis of Gene Expressions Associated with Increased Allelopathy in Rice (*Oryza sativa* L.) Induced by Exogenous Salicylic Acid. *Plant Growth Regulation*, Vol. 57, pp. 163-172, ISSN 0167-6903.
- Farooq, M.; Aziz, T.; Hussain, M.; Rehman, H.; Jabran, K. & Khan, M.B. (2008a). Glycinebetaine Improves Chilling Tolerance in Hybrid Maize. *Journal of Agronomy and Crop Science*, Vol. 194, pp. 152-160, ISSN 0931-2250.
- Farooq, M.; Aziz, T.; Basra, S.M.A.; Cheema, M.A. & Rehman, H. (2008b). Chilling Tolerance in Hybrid Maize Induced by Seed Priming with Salicylic Acid. *Journal of Agronomy and Crop Science*, Vol. 194, pp. 161-168, ISSN 0931-2250.
- Farrar, J.F. (1985). The Respiratory Source of CO₂. *Plant Cell and Environment*, Vol. 8, pp. 427-438, ISSN 0140-7791.

- Fodor, J.; Gullner, G.; Adam, A.L.; Barna, B.; Komives, T. & Kiraly, Z. (1997). Local and Systemic Responses of Antioxidants to Tobacco Mosaic Virus Infection and to Salicylic Acid in Tobacco. *Plant Physiology*, Vol. 114, pp. 1443-1451, ISSN 0032-0889.
- Frantz, J.M.; Cometti, N.N. & Bugbee, B. (2004). Night Temperature has a Minimal Effect on Respiration and Growth in Rapidly Growing Plants. *Annals of Botany*, Vol. 94, pp. 155-166, ISSN 0305-7364.
- Gibson, L.R., & Mullen, R.E. (1996). Influence of Day and Night Temperature on Soybean Seed Yield. *Crop Science*, Vol. 36, pp. 98-104, ISSN 0011-183X.
- Guinn, G. (1974). Abscission of Cotton Floral Buds as Influenced by Factors Affecting Photosynthesis and Respiration. *Crop Science*, Vol. 14, pp. 291-293, ISSN 0011-183X.
- Guo, Y.P.; Zhou, H.F. & Zhang, L.C. (2006). Photosynthetic Characteristics and Productive Mechanism against Photooxidation during High Temperature Stress in Two Citrus Species. *Scientia Horticulturae*, Vol. 108, pp. 260-267, ISSN 0304-4238.
- Hall, A.E. (1992). Breeding for Heat Tolerance. *Plant Breeding Reviews*, Vol. 10, pp.129-168, ISSN 0730-2207.
- Harinasut, P.; Tsutsui, K.; Takabe, T.; Nomura, M.; Takabe, T. & Kishitani, S. (1996). Exogenous Glycinebetaine Accumulation and Increased Salt-tolerance in Rice Seedlings. *Bioscience Biotechnology and Biochemistry*, Vol. 60, pp. 366-368, ISSN 0916-8451.
- Harper, J .R. & Balke, N.E. (1981). Characterization of the Inhibition of K⁺ Absorption in Oats Roots by Salicylic Acid. *Plant Physiology*, Vol. 68, pp. 1349-1353, ISSN 0032-0889.
- Havaux, M. & Tardy, F. (1996). Temperature-dependent Adjustment of the Thermal Stability of Photosystem 2 in vivo: Possible Involvement of Xanthophylls-Cycle Pigments. *Planta*, Vol. 198, pp. 324-333, ISSN 0032-0935.
- He, Y.; Liu, Y.; Cao, W.; Huai, M.; Xu, B. & Huang, B. (2005). Effects of Salicylic Acid on Heat Tolerance Associated with Antioxidant Metabolism in Kentucky Bluegrass. *Crop Science*, Vol. 45, pp. 988-995, ISSN 0011-183X.
- Huang, B., Liu, X. & Fry J.D. 1998. Shoot Physiological Responses of Two Bentgrass Cultivars to High Temperature and Poor Soil Aeration. *Crop Science*, Vol. 38, pp. 1219-1224, ISSN 0011-183X.
- Ibrahim, A.M.H. & Quick, J.S. (2001). Heritability of Heat Tolerance in Winter and Spring Wheat. *Crop Science*, Vol. 41, pp. 1401-1405, ISSN 0011-183X.
- IPCC (Intergovernmental Panel on Climate Change) (2007). Climate Change 2007: Impacts, Adaptation and Vulnerability. In: *Contribution of Working Group II to Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, M.L. Parry; O.F. Canziani; J.P. Palutikof; P.J. van der Linden; & C.E. Hanson (Eds.), 1000, Cambridge University Press, Cambridge, UK.
- Ismail, A.M. & Hall, A.E. (1999). Reproductive-stage Heat Tolerance, Leaf Membrane Thermostability and Plant Morphology in Cowpea. *Crop Science*, Vol. 39, pp. 1762-1768, ISSN 0011-183X.
- Izquierdo, N.; Aguirrezábal, L.; Andrade, F. & Pereyra, V. (2002). Night Temperature Affects Fatty Acid Composition in Sunflower Oil Depending on the Hybrid and the Phenological Stage. *Field Crops Research*, Vol. 77, pp. 115-126, ISSN 0378-4290.

- John, E. (2002). Candidate Effector and Regulator Genes Activated by Glycine Betaine in Arabidopsis. Available from <http://abstracts.aspb.org/pb2002/public/P63/0153.html>.
- Jones, P.D.; New, M.; Parker, D.E.; Martin, S. & Rigor, I.G. (1999). Surface Air Temperature and its Changes over the Past 150 Years. *Reviews of Geophysics*, Vol. 37, pp. 173–199.
- Kang, G.Z., Z.X. Wang, K.F. Xia, and G.C. Sun, 2007: Protection of Ultrastructure in Chilling-stressed Banana Leaves by Salicylic Acid. *Journal of Zhejiang University Science*, Vol. 8, pp. 277-282, ISSN 1009-3095.
- Keeling, P.L.; Banisadr, R.; Barone, L.; Wasserman, B.P. & Singletary, G.W. (1994). Effect of Temperature on Enzymes in the Pathway of Starch Biosynthesis in Developing Wheat and Maize Grain. *Australian Journal of Plant Physiology*, Vol. 21, pp. 807-827, ISSN 0310-7841.
- Khan, N.A.; Syeed, S.; Masood, A.; Nazar, R. & Iqbal, N. (2010). Application of Salicylic Acid Increases Contents of Nutrients and Antioxidative Metabolism in Mungbean and Alleviates Adverse Effects of Salinity Stress. *International Journal of Plant Biology*, Vol. 1, pp. e1, ISSN 2037-0156.
- Khan, W.; Prithiviraj, B. & Smith, A. (2003). Photosynthetic Responses of Corn and Soybean to Foliar Application of Salicylates. *Journal of Plant Physiology*, Vol. 160, pp. 485-492, ISSN 0176-1617.
- Kondo, S. & Takahashi, Y. (1989). Relation between Early Drop of Apple Fruit and Ethylene Evolution under High Night-temperature Conditions. *Journal of Japanese Society for Horticultural Science*, Vol. 58, pp. 1-8, ISSN 1882-3351.
- Kreiner, M.; Harvey, L.M. & McNeil, B. (2002). Oxidative Stress Response of a Recombinant *Aspergillus niger* to Exogenous Menadione and H₂O₂ Addition. *Enzyme and Microbial Technology*, Vol. 30, pp. 346-353, ISSN 0141-0229.
- Lambers, H. (1985). Respiration in Tack Plants and Tissues: Its Regulation and Dependence on Environmental Factors, Metabolism and Invaded Organism. In: *Higher Plant Cell Respiration. Encyclopedia of Plant Physiology*, R. Douce & D.A. Day (Eds.), New series, Vol. 18, Springer-Verlag, Berlin, Germany.
- Larkindale, J. & Huang, B. (2004). Thermotolerance and Antioxidant Systems in *Agrostis stolonifera*: Involvement of Salicylic Acid, Abscissic Acid, Calcium, Hydrogen Peroxide and Ethylene. *Journal of Plant Physiology*, Vol. 161 pp. 405-413, ISSN 0176-1617.
- Larkindale, J. & Knight, M.R. (2002). Protection Against Heat Stress-induced Oxidative Damage in Arabidopsis Involves Calcium, Abscissic Acid, Ethylene, and Salicylic Acid. *Plant Physiology*, Vol. 128, pp. 682-695, ISSN 0032-0889.
- Larque-Saavedra, A. (1978). The Antitranspirant Effect of Acetylsalicylic Acid on *Phaseolus vulgaris*. *Plant Physiology*, Vol. 43, pp. 126-128, ISSN 0032-0889.
- Lee, C.B.; Hayashi, H. & Moon, B.Y. (1997). Stabilization by Glycinebetaine of Photosynthetic Oxygen Evolution by Thylakoid Membranes from Synechococcus PCC7002. *Molecular Cell*, Vol. 30, pp. 296-299, ISSN 1097-2765.
- Loka, D.A., Oosterhuis, D.M., 2010. Effect of High Night Temperatures on Cotton Respiration. ATP Levels, and Carbohydrate Content. *Environmental and Experimental Botany*, Vol. 68, pp. 258-263, ISSN 0098-8472.

- Mäkelä, P.; Karkkainen, J. & Somersalo, S. (2000). Effect of Glycine Betaine on Chloroplast Ultrastructure, Chlorophyll and Protein Content, and RUBPCO Activities in Tomato Grown under Drought or Salinity. *Biologia Plantarum*, Vol. 3, pp. 471-475, ISSN 0006-3134.
- Martineau, J.R.; Specht, J.E.; Williams, J.H. & Sullivan, C.Y. (1979). Temperature Tolerance in Soybeans. I. Evaluation of a Technique for Assessing Cellular Membrane Thermostability. *Crop Science*, Vol. 19, pp. 75-78, ISSN 0011-183X.
- Matsui, T. & Kagata, H. (2003). Characteristics of Floral Organs Related to Reliable Self Pollination in Rice (*Oryza sativa* L.). *Annals of Botany*, Vol. 91, pp. 473-477, ISSN 0305-7364.
- McDonald, A.E. & Vanlerberghe, G.C. (2005). Alternative Oxidase and Plastoquinol Terminal Oxidase in Marine Prokaryotes of the Sargasso Sea. *Gene*, Vol. 349, pp. 15-24, ISSN 0378-1119.
- Micheal, G. & Beringer, H. (1980). The Role of Hormones in Yield Formation. In: *Physiological Aspects of Crop Productivity*, G. Micheal & H. Beringer (Eds.), 85-116, Proceedings of the 15th Colloquium of the International Potash Institute, Bern, Switzerland.
- Mitchell, R.A.C.; Mitchell, V.J.; Driscoll, S.P.; Franklin, J. & Lawlor, D.W. (1993). Effects of Increased CO₂ Concentration and Temperature on Growth and Yield of Winter Wheat at Two Levels of Nitrogen Application. *Plant, Cell and Environment*, Vol. 16, pp. 521-529, ISSN 0140-7791.
- Mohammed, A.R. & Tarpley, L. (2011a). Morphological and physiological responses of nine southern U.S. rice cultivars differing in their tolerance to enhanced ultraviolet-B radiation. *Environmental and Experimental Botany*, Vol. 70, pp. 174-184, ISSN 0098-8472.
- Mohammed, A.R. & Tarpley, L. (2011b). Characterization of Rice (*Oryza sativa* L.) Physiological Responses to α -Tocopherol, Glycine Betaine or Salicylic Acid Application. *Journal of Agriculture Science*, Vol. 3, pp. 3-13, ISSN 0021-8596.
- Mohammed, A.R. & Tarpley, L. (2009a). High Nighttime Temperatures Affect Rice Productivity through Altered Pollen Germination and Spikelet Fertility. *Agricultural and Forest Meteorology*, Vol. 149, pp. 999-1008, ISSN 0168-1923.
- Mohammed, A.R. & Tarpley, L. (2009b). Impact of High Nighttime Temperature on Respiration, Membrane Stability, Antioxidant Capacity, and Yield of Rice Plants. *Crop Science*, Vol. 49, pp. 313-322, ISSN 0011-183X.
- Moller, I.M. (2001). Plant Mitochondria and Oxidative Stress: Electron Transport, NADPH turnover, and Metabolism of Reactive Oxygen Species. *Annual Review of Plant Physiology and Plant Molecular Biology*, Vol. 52, pp. 561-591, ISSN 1040-2519.
- Neales, T.F.; Sale, P.J.M. & Meyer, C.P. (1980). Carbon Dioxide Assimilation by Pineapple Plants, *Ananas comosus* (L.) Merr. II. Effects of Variation of the Day/Night Temperature Regime. *Australian Journal of Plant Physiology*, Vol. 7, pp. 375-385, ISSN 0310-7841.
- Oota, Y. (1975). Short-day Flowering of *Lemna gibba* G3 Induced by Salicylic Acid. *Plant Cell and Physiology*, Vol. 16, pp. 1131-1135, ISSN 0032-0781.

- Paembonan, S.A.; Hagihara, A. & Hozumi, K. (1992). Long-term Respiration in Relation to Growth and Maintenance Processes of the Aboveground Parts of a Hinoki Forest Tress. *Tree Physiology*, Vol. 10, pp. 101-110, ISSN 0829-318X.
- Paleg, L.G.; Douglas, T.J.; van Daal, A. & Keech, D.B. (1981). Proline, Betaine and Other Organic Solutes Protect Enzymes Against Heat Inactivation. *Australian Journal of Plant Physiology*, Vol. 8, pp. 107-114, ISSN 0310-7841.
- Pan, Q.; Zhan, J.; Liu, H.; Zhang, J.; Chen, J.; Wen, P. & Huang, W. (2006) Salicylic Acid Synthesized by Benzoic Acid 2 Hydroxylase Participates in the Development of Thermotolerance in Pea Plants. *Plant Science*, Vol. 171, pp. 226-233, ISSN 0168-9452.
- Peng, S.; Huang, J.; Sheehy, J.E.; Laza, R.C.; Visperas, R.M.; Zhong, X.; Centeno, G.S.; Khush, G.S. & Cassman, K.G. (2004). Rice Yields Decline with Higher Night Temperature from Global Warming. *Proceedings of the National Academy of Sciences of the United States of America* Vol. 101, pp. 9971-9975, ISSN 0027-8424.
- Penning de Vries, F.W.T. (1975). The Cost of Maintenance Process in the Cell. *Annals of Botany*, Vol. 39, pp. 77-92, ISSN 0305-7364.
- Peterson, R.B. & Zelitch, I. (1982). Relationship between Net CO₂ Assimilation and Dry Weight Accumulation in Field-grown Tobacco. *Plant Physiology*, Vol. 70, pp. 677-685, ISSN 0032-0889.
- Pospisil, P. & Dau, H. (2000). Chlorophyll Fluorescence Transients of Photosystem II Membrane Particles as a Tool for Studying Photosynthetic Oxygen Evolution. *Photosynthesis Research*, Vol. 65, pp. 41-52, ISSN 0166-8595.
- Prasad, P.V.V.; Boote, K.J.; Allen Jr, L.H.; Sheehy, J.E. & Thomas, J.M.G. (2006). Species, Ecotype and Cultivar Differences in Spikelet Fertility and Harvest Index of Rice in Response to High Temperature Stress. *Field Crops Research*, Vol. 95, pp. 398-411, ISSN 0378-4290.
- Prasad, P.V.V.; Craufurd, P.Q. & Summerfield, R.J. (1999). Fruit Number in Relation to Pollen Production and Viability in Groundnut Exposed to Short Episodes of Heat Stress. *Annals of Botany*, Vol. 84, pp. 381-386, ISSN 0305-7364.
- Rajasekaran, L.R.; Kriedemann, P.E.; Aspinall, D. & Paleg, L.G. (1997). Physiological Significance of Proline and Glycinebetaine: Maintaining Photosynthesis during NaCl Stress in Wheat. *Photosynthetica*, Vol. 34, pp. 357-366, ISSN 0300-3604.
- Rao, M.V.; Paliyath, G.; Ormrod, P.; Murr, D.P. & Watkins, C.B. (1997). Influence of Salicylic Acid on H₂O₂ Production, Oxidative Stress, and H₂O₂-metabolizing Enzymes. *Plant Physiology*, Vol. 115, pp. 137-149, ISSN 0032-0889.
- Raskin, I. (1992). Role of Salicylic Acid in Plants. *Annual Review of Plant Physiology and Plant Molecular Biology*, Vol. 43, pp. 439-463, ISSN 1040-2519.
- Reynolds, M.P.; Balota, M.; Delgado, M.I.B.; Amani, I. & Fisher, R.A. (1994). Physiological and morphological traits associated with spring wheat yield under hot irrigated conditions. *Australian Journal of Plant Physiology*, Vol. 21, pp. 717-730, ISSN 0310-7841.
- Ryals, J.; Uknes, S. & Ward, E. (1994). Systematic Acquired Resistance. *Plant Physiology*, Vol. 104, pp. 1109-1112, ISSN 0032-0889.
- Ryan, M.G. (1991). Effects of Climate Change on Plant Respiration. *Ecological Applications*, Vol. 1, pp. 157-167, ISSN 1051-0761.

- Sakamoto, H. & Murata, N. (1998). Metabolic Engineering of Rice Leading to Biosynthesis of Glycinebetaine and Tolerance to Salt and Cold. *Plant Molecular Biology*, Vol. 38, pp. 1011-1019, ISSN 0167-4412.
- Sayed, O.H.; Earnshaw, M.J. & Emes, M.J. (1989). Photosynthetic Response of Different Varieties of Wheat to High Temperature. *Journal of Experimental Botany*, Vol. 40, pp. 633-638, ISSN 0022-0957.
- Seddigh, M. & Jolliff, G.D. (1984). Night Temperature Effects on Morphology, Phenology, Yield and Yield Components of Indeterminate Field-grown Soybean. *Agronomy Journal*, Vol. 76, pp. 824-828, ISSN 0002-1962.
- Senaratna, T.; Touchell, D.; Bunn, T. & Dixon, K. (2000). Acetyl salicylic acid (aspirin) and salicylic acid induce multiple stress tolerance in bean and tomato plants. *Plant Growth Regulation*, Vol. 30, pp. 157-161, ISSN 0167-6903.
- Shirasu K.; Nakajima H. & Rajashekar K. (1997). Salicylic Acid Potentiates an Agonist-dependent Gain Control that Amplifies Pathogen Signal in the Activation of Defense Mechanisms. *Plant Cell*, Vol. 9, pp. 261-270, ISSN 1040-4651.
- Singletary, G.W.; Banisadr, R. & Keeling, P.L. (1994). Heat Stress during Grain Filling in Maize: Effects on Carbohydrate Storage and Metabolism. *Australian Journal of Plant Physiology*, Vol. 21, pp. 829-841, ISSN 0310-7841.
- Smirnoff, N. (1995). Antioxidant Systems and Plant Response to the Environment. In: *Environment and Plant Metabolism: Flexibility and Acclimation*, N. Smirnoff (Ed.), 217-243, Bios Scientific Publishers, Oxford, UK.
- Sullivan, C.Y. (1972). Mechanisms of Heat and Drought Resistance in Grain Sorghum and Methods of Measurement. In: *Sorghum in the Seventies*, N.G.P. Rao & L.R. House (Eds.), 247-264, Oxford and IPH Publishing Company, New Delhi, India.
- Sullivan, C.Y. & Ross, W.M. (1979). Selecting for Drought and Heat Resistance in Grain Sorghum. In: *Stress Physiology in Crop Plants*, H. Mussell, H. & R.C. Staples (Eds.), 263-281, John Wiley & Sons, New York, USA.
- Turnbull, M.H.; Murthy, R. & Griffin, K.L. (2002). The Relative Impacts of Daytime and Night-time Warming on Photosynthetic Capacity in *Populus deltoides*. *Plant Cell and Environment*, Vol. 25, pp. 1729-1737, ISSN 0140-7791.
- Vanacker, H.; Lu, H.; Rate, D.N. & Greenberg, J.T. (2001). A Role for Salicylic Acid and NPR1 in Regulating Cell Growth in Arabidopsis. *Plant Journal*, Vol. 28, pp. 209-216, ISSN 0960-7412.
- Warrag, M.O.A. & Hall, A.E. (1984). Reproductive Responses of Cowpea (*Vigna unguiculata* (L.) Walp.) to Heat Stress. II. Responses to Night Air Temperature. *Field Crops Research*, Vol. 8, pp. 17-33, ISSN 0378-4290.
- Willits, D.H. & Peet, M. (1998). The Effect of Night Temperature on Greenhouse Grown Tomato Yields in Warm Climates. *Agricultural and Forest Meteorology*, Vol. 92, pp. 191-202, ISSN 0168-1923.
- Yalpani, N.; Enyedi, A.J.; Leon, J. & Raskin, I. (1994). Ultraviolet Light and Ozone Stimulate Accumulation of Salicylic Acid, Pathogen-related Proteins and Virus Resistance in Tobacco. *Planta*, Vol. 193, pp. 372-376, ISSN 0032-0935.

- Yamane, Y.; Kashino, Y. & Koike, H. (1997). Increase in Fluorescence F_0 Level and Reversible Inhibition of Photosystem 2 Reaction by High-temperature Treatments in Higher Plants. *Photosynthetic Research*, Vol. 52, pp. 57-64.
- Yilmaz, J.L. (2004). Abiotic Stress Tolerance: Metabolic and Physiological Effects of Compatible Solutes and Calmodulin on *E. coli* and Tobacco. In: *Pure and Applied Biochemistry*, J.L. Yilmaz (Ed), 1-40, Lund University, Lund, Sweden.
- Yoshida, S. (1981). Physiological Analysis of Rice Yield. In: *Fundamentals of Rice Crop Science*, S. Yoshida (Ed.), 231-251. International Rice Research Institute, Los Banos, Philippines.
- Zhang, J. & Kirkham, M.B. (1996). Antioxidant Responses to Drought in Sunflower and Sorghum Seedlings. *New Phytologist*, Vol. 132, pp. 361-373, ISSN 0028-646X.
- Zheng, S.H.; Nakamoto, H.; Yoshikawa, K.; Furuya, T. & Fukuyama, M. (2002). Influence of High Night Temperature on Flowering and Pod Setting in Soybean. *Plant Production Science*, Vol. 5, pp. 215-218, ISSN 1343-943X.
- Zhou, X.M., Mackenzie, A.F., Madramootoo, C.A., & Smith, D.L.J. (1999). Effects of Some Injected Plant Growth Regulators, With or Without Sucrose, on Grain Production, Biomass and Photosynthetic Activity of Field-grown Corn Plants. *Journal of Agronomy and Crop Science*, Vol. 183, pp. 103-110, ISSN 0931-2250.

IntechOpen

Global Warming Impacts - Case Studies on the Economy, Human Health, and on Urban and Natural Environments

Edited by Dr. Stefano Casalegno

ISBN 978-953-307-785-7

Hard cover, 290 pages

Publisher InTech

Published online 03, October, 2011

Published in print edition October, 2011

This book addresses the theme of the impacts of global warming on different specific fields, ranging from the regional and global economy, to agriculture, human health, urban areas, land vegetation, marine areas and mangroves. Despite the volume of scientific work that has been undertaken in relation to each of these issues, the study of the impacts of global warming upon them is a relatively recent and unexplored topic. The chapters of this book offer a broad overview of potential applications of global warming science. As this science continues to evolve, confirm and reject study hypotheses, it is hoped that this book will stimulate further developments in relation to the impacts of changes in the global climate.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Abdul Razack Mohammed and Lee Tarpley (2011). Effects of High Night Temperature on Crop Physiology and Productivity: Plant Growth Regulators Provide a Management Option, *Global Warming Impacts - Case Studies on the Economy, Human Health, and on Urban and Natural Environments*, Dr. Stefano Casalegno (Ed.), ISBN: 978-953-307-785-7, InTech, Available from: <http://www.intechopen.com/books/global-warming-impacts-case-studies-on-the-economy-human-health-and-on-urban-and-natural-environments/effects-of-high-night-temperature-on-crop-physiology-and-productivity-plant-growth-regulators-provid>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2011 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](https://creativecommons.org/licenses/by/3.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen