We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Comparative Evaluation of Different Techniques for Aflatoxin Detoxification in Poultry Feed and Its Effect on Broiler Performance

T. Mahmood¹, T.N. Pasha¹ and F.M. Khattak²

¹Department of Food and Nutrition,
University of Veterinary and Animal Sciences, Lahore,

²Avian Science Research Centre, SAC, Edinburgh,

¹Pakistan

²UK

1. Introduction

Aflatoxins (AF), the toxic secondary metabolites produced by *Aspergillus flavus* and *Aspergillus parasiticus*, are a major concern in the poultry production. AF metabolites are stable and fairly resistant compounds to degradation (Dalvi, 1986; Park, 2002; Desphande, 2002; Lesson *et al.*, 1995; Feuill, 1996). These metabolites are usually produced during the growth of the *Aspergillus flavus*, *Aspergillus parasiticus* and *Aspergillus niger* on certain foods and feedstuffs under favourable conditions of moisture, temperature and aeration (Goto *et al.*, 1997; Dutta and Das, 2001). Their toxicity depends on several factors including its concentration, the duration of exposure, the species, sex, age, and health status of animals (Jewers, 1990). Contamination of AF in feed causes aflatoxicosis in poultry that is characterised by reduced feed intake, decreased weight gain, poor feed utilization (Tedesco *et al.*, 2004; Bailey *et al.*, 2006; Shi *et al.*, 2006, 2009), increased susceptibility to environmental and microbial stresses, and increased mortality (Leeson *et al.*, 1995). AF can also cause productive deterioration which is associated with changes in biochemical and hematological parameters (Denli *et al.*, 2004; Basmacioglu *et al.*, 2005; Bintvihok and Kositcharoenkul, 2006), liver and kidney abnormalities, and impaired immunity, which is able to enhance susceptibility to some environmental and infectious agents (Ibrahim *et al.*, 2000; Oguz *et al.*, 2003). AF has been reported to have effect on metabolism in poultry by decreasing the activities of several enzymes that are important in the digestion of starch, proteins, lipids and nucleic acids. Consequently, the activities of serum glutamate pyruvate transaminase, serum glutamate oxaloacetate transferase and γ -glutamyl transferase are increased, primarily indicating hepatic damage (Devegowda and Murthy, 2005). AF is also known to interfere with metabolism of vitamin D, iron and copper and can cause leg weakness (Khajarern and Khajarern, 1999). Severe economic losses have been reported in the poultry industry due to aflatoxicosis (Kubena *et al.*, 1991, 1995). Ultimately, the transmission of AF and its metabolites from feed to animal edible tissues and products, such as liver and eggs, becomes a potential hazard for human health.

The occurrence of mycotoxin in nature is considered a global problem. However, in certain regions of the world, some of the mycotoxins are produced more commonly than others. Several *invitro* and *invivo* studies conducted in India, Pakistan, Egypt & South Africa suggested that AF are often present in substantial levels in mixed feed & ingredients (Devegowda and Murthy, 2005). Although, AF in feed and food is considered to be a major concern in warm and humid climatic regions of the world, however, caution must be exercised even in colder regions, when using feedstuffs imported from warm and humid countries.

With increasing knowledge and awareness of AF as a potent source of health hazards to both man and farm animals, producers, researchers and government organizations are making great effort to develop effective preventive management and decontamination technologies to minimise the toxic effects of AF content in foods and feedstuffs. In order to reduce the toxic and economic impact of mycotoxins, established regulations and legislative limits have been set for AF in poultry feed. Many countries follow a maximum permissible level of 20ppb for AF in poultry feed (CAST, 2003; FAO, 1995).

Appropriate pre and post-harvest contamination can be reduced by using appropriate agricultural practices. However, the contamination is often unavoidable and still remains a serious problem associated with many important agricultural commodities, which emphasizes the need for a suitable process to inactivate the toxin. Besides the preventive management, several approaches have been employed including physical (feed mill techniques, blending, extraction, irradiation, and heating), chemical (acids, bases, alkali treatments and oxidizing agents) biological treatments (certain species of fungi and bacteria) and solvent extraction to detoxify AF in contaminated feeds and feedstuffs (Coker *et al.*, 1986; Piva *et al.*, 1995; Parlat *et al.*, 1999). Since the beginning of 1990s, the adsorbent-based studies have also been reported to be effective in removing AF from contaminated feed and minimise the toxicity of AF in poultry (Ibrahim *et al.*, 2000). Among several adsorbents commercially available in the market, Zeolites (Miazzo *et al.*, 2000), bentonites (Rosa *et al.*, 2001, Pasha *et al.*, 2007, 2008) and clinoptilolite (CLI), (Oguz and Kurtoglu, 2000; Oguz *et al.*, 2000 a, b), were preferred because of their high binding capacities for AF and their reducing effect on AF-absorption from the gastrointestinal tract.

All these methods cannot be used in practical feed manufacturing, because of the limitation of the nutrients decomposition, non availability of commercial methods and their residual effects. The increasing number of reports on detoxification of AF in poultry feed using different techniques has given rise to a demand for practical and economical detoxification procedures. Some of the physical treatments are reported to be relatively costly and may also remove or destroy essential nutrients in feed. Whereas, chemical methods are considered to be time consuming, expensive as they mostly require suitable reaction facilities, and are reported to have deteriorating residual effects on animal health (Coker, 1979; Coker *et al.*, 1985). Certain legal implications are also associated with the use of different detoxifying methods. For example, European community (EC) is in favour of use of physical decontamination processes and sorting procedures. However, neither the use of chemical decontamination processes, nor the mixing of batches with the aim of decreasing the level of contamination below the maximum tolerable level is legal within the European Union (Avantaggiato *et al.*, 2005). Although several mycotoxin detoxifying or adsorbing techniques have been assessed independently however, limited information is available on the comparison of different techniques. To further understand the mechanisms of aflatoxin and detoxification of poultry feed by heat treatment (extrusion) and added adsorbents

(Sodium bentonite or Mycofix®Plus), a study was conducted to compare different detoxification techniques and to further investigate its effect on broiler performance.

2. Materials and methods

2.1 Birds and diet

Two hundred day-old commercial broiler chicks were randomly distributed to 5 dietary treatments with 4 replicates of 10 chicks each. During the first 21 days, all birds were fed on diet 1 which was the basal starter ration without any aflatoxin contamination (AF) and detoxifying treatment (DT). The ingredient composition of the basal diet is presented in Table 1. Experimental diets were prepared by replacing maize with contaminated maize having 70 ppb AF (Treatment 2) and were subjected to different DT (Treatment 3 to 5).

Ingredient	Starter (%)	Grower (%)
Maize	50.70	60.00
Rice tips	10.00	10.00
Corn gluten meal 60 %	5.00	3.00
Soybean meal	10.00	6.00
Guar meal	5.00	5.00
Cotton seed meal	7.00	7.00
Rape seed meal	7.00	7.00
Fish meal	3.00	
Molasses	1.00	
Di-calcium Phosphate	1.00	1.00
Lysine	0.30	0.35
Calculated nutrient composition		
M E (kcal/ kg)	3000	3100
Crude protein	22.00	19.00
Crude fiber	4.53	4.48
Lysine	1.08	0.93
Methionine	0.49	0.45
Cystine	0.27	0.27
Met+Cys	0.88	0.80
Linoleic acid	1.21	1.36
Calcium	1.00	0.92
Phosphorous total	0.75	0.69
Phosphorus available	0.44	0.41

Table 1. Ingredient and nutrient composition of basal starter and grower diet of broilers.

On day 21, birds were fed on one of five experimental diets. Experimental diets were fed from day 21 to day 42 of the trial. Feed and water was available on *ad libitum* basis. All the birds were vaccinated against Newcastle Disease (N.D.), with Lasota strain eye droppings at day 7 and with oil based vaccine (intra muscular) at day 21 of the experiment.

2.2 Experimental design

The experimental design consists of five dietary treatments; 1 (0 ppb AF & no DT); 2 (70 ppb AF & no DT); 3 (70 ppb AF & DT by Extrusion); 4 (70 ppb & DT by Mycofix®Plus); 5 (70 ppb & DT by Sodium bentonite).

2.3 Aflatoxin production and analysis

Aflatoxins were produced by the inoculation of fungus on corn as described by (Lillehoj, *et al.*, 1974) with some amendments. Fermentation was carried out in 1-liter Erlenmeyer flasks containing 50g of whole corn kernels. 25 ml distilled water was added to the corn (50 g) in the Erlenmeyer flasks, and the mixture was allowed to stand for 2 hrs with frequent shaking. The flasks were tightly plugged with cotton and autoclaved at 121°C at 15 psi for 15 min and cooled at room temperature. They were then inoculated with 3ml spore suspension in a sterile environment, placed on an orbital shaker at 200rpm and incubated at 28 °C. At 24 and 48hr, sterile water (3-5ml) was added in the flask, quantity of water adjusted in a manner that individual kernels do not adhere with each other.

If the corn did pack in clumps, the material was loosened by vigorous shaking, and if required, clumps were smashed with the help of a sterile rod within sterile environment to make sure that individual kernel should be kept free from others. On 7-8d the flasks were again autoclaved at 121°C at 15 psi for 15 min, and placed in a hot oven at 60 °C for 24hr till all the moisture was removed.

The AF containing corn Kernels were grinded to powdered form and was quantitatively evaluated by direct competitive enzyme link immuno-sorbent assay (ELISA) described by Barabolak (1977) using (RIDASCREEN® FAST Aflatoxin Total) kit and mixed in feed according to the calculation to get the desirable level of aflatoxin (70 ppb) in the feed. The prepared experimental diets were analysed again using ELISA technique to confirm the AF levels.

2.4 Methods used for detoxification of AF

2.4.1 By extrusion

The AF containing corn was passed through the extruder, following the procedure described by Grehainge *et al.*, 1983. The corn in the experimental starter and grower diets was replaced by extruded corn (Treatment 3).

2.4.2 By Mycofix

A commercially available Mycofix® plus was added in the experimental diet at the recommended dose rate of 0.5 Kg per ton of feed (Treatment 4).

2.4.3 By Sodium Bentonite (SB)

The source and composition of SB used was same as described previously (Pasha *et al.*, 2007). The supplemental SB was added in the experimental diet at 1% of the feed (Treatment 5).

2.5 Sampling

Body weights and feed intake per pen was recorded weekly and mortality was recorded as it occurred. At day 28, 35 and 42 of bird's age, five birds per replicate were randomly selected for estimation of antibody titers against Newcastle disease (ND). The blood samples (3 ml) were collected from wing vein. The blood serum was separated and analysed by Haemagglutination inhibition (HI) method described by Sever (1962). After blood collection, birds were humanely killed and bursa of Fabricius was removed and weighed.

3. Statistical analysis

The results (pen means) were subjected to one-way ANOVA as a complete randomized design (CRD) using Genstat 11 for window. Treatment means were compared by the Tukey's test and statistical significance was accepted at $P < 0.05$

4. Results and discussion

4.1 Growth performance

The body weight gain, feed intake and FCR values showed no differences between different treatments (Tables 2). This lack of difference in performance between treatments could either be due to lower level of AF (70ppb) or shorter administrative period (day 21 to 42 of age). Similarly no change in production parameters have been reported previously in several studies (Ouz *et al.*, 2000b, 2003; Ortatatli *et al.*, 2005; Magnoli *et al.*, 2008a,b, 2011) when birds were fed diets low in toxin (50 to 100 ppb AF) for a period of 46 days. The results from the present study are not in agreement with other studies where significantly reduced body weights were observed when birds were exposed to higher dietary AF (400 and 600 ppb AF). The depression in growth upon feeding AF was attributed to reduced protein and energy utilization (Smith and Hamilton, 1970; Lanza *et al.* 1980; Doerr *et al.*, 1983; Dalvi and Ademoyero, 1984; Verma *et al.*, 2002) which impaired nutrient absorption and reduced pancreatic digestive enzyme production (Osborne and Hamilton, 1981) and consequently reduced appetite (Sharline *et al.*, 1980). Similarly, significant depressions in body weight gain were also recorded in broilers given diets containing 1 and 2 mg/kg of AF (1000 to 2000 ppb) at 4 and 7 weeks of age. Several studies have also shown that dietary AF adversely affected the growth of broilers in a dose-dependent manner (Johri and Sadagopan, 1989; Espada *et al.*, 1992; Beura *et al.*, 1993). A similar reductions in weight gain were also observed in broilers at dietary AF contents of 0.75 mg/kg (750 ppb) and above and these depression in body weight in toxin fed groups were reported to be dose dependent (Reddy *et al.*, 1984).

Treatment	DESCRIPTION	Weight Gain (g)	Feed Intake (g)	FCR
1	(0 ppb AF & no DT)	1566	3221	2.05
2.	(70 ppb AF & no DT)	1527	3215	2.10
3	70 ppb AF + DT by extrusion	1533	3249	2.11
4	70 ppb AF + DT with Mycofix®Plus	1544	3233	2.09
5	70 ppb AF + DT with Sodium bentonite	1564	3244	2.07
	SED	24.94	24.51	0.048
	P-value	0.444	0.608	1.00

Means within a column were not different (P < 0.05). Tukeys T test was used for means separation; SED – standard error of the difference.

Table 2. Weight gain, feed intake, and FCR of birds fed different experimental diets.

In contrast to our results, reduced feed intake and poor feed efficiency in broilers has also been reported in birds fed diets containing AF at 2, 4 and 6 weeks of age when level of dietary AF was higher than 100 ppb (Sharline *et al.* 1980; Huff and Doerr, 1981; Nandkumar *et al.* 1984; Rajasekhar Reddy *et al.* 1982; Johri and Majmudar, 1990; Verma *et al.*, 2004). These authors have suggested that the reduced appetite during aflatoxicosis could be due to impaired liver metabolism caused by the liver damage. The reason for these differences in performance compared to present study could be due to different type, age and dose of AF. It is likely that juvenile birds may respond differently to the same dose of AF in diet as their physiological needs and capacity to absorb is higher compared to older birds.

It is suggested that extrusion-cooking is an efficient process used for eliminating some of the naturally occurring food toxins. The process involves high temperature (up to 250°C), short time (usually 1-2 min), high pressures (up to 25 mPa) and low water contents (below 30) and has been used to eliminate some of the toxins in food and feed ingredients (Harper, 1989; Fast, 1991; Kohlwey et al. 1995;). It has also been used as a kind of bioreactor to decontaminate AF. In spite the fact that feed moisture, barrel temperature and the die diameter are identified as different variables that can influence AF reduction during extrusion cooking, a reduction in total AF content (up to 84%) has been reported when artificially contaminated peanut meal was detoxified using extrusion cooking (Grehaigne *et al.*, 1983; Cheftel, 1986, 1989). Similarly, commercial adsorbents are also reported to bind the aflatoxins in feed and prevent its absorption in animal gastrointestinal tract (Ramos and Hernandez, 1997). Numerous studies have shown the effectiveness of these agents to bind aflatoxins in vitro (Huff *et al.*, 1992; Diaz *et al.*, 2002). Mycofix, is one of the adsorbent that can be added in poultry feed and is claimed to neutralize moderate levels of aflatoxin (up to 2500-3500 ppb) in poultry feed. Mycofix deactivates aflatoxin with its polar functional group, due to AF fixation to adsorbing components in Mycofix, with stable binding capacity. Adsorption starts in the oral cavity during salivation and continues in stomach and gut. The fixed mycotoxin being unable to enter the blood and subsequently excreted in faeces after 98% adsorption of AF by Mycofix (Biomim®, 2000). Similarly, incorporation of SB in poultry diet is another proved adsorbent, to have high AF binding capacities both in vitro (Magnoli *et al.*, 2008a) and in vivo (Rosa *et al.*, 2001; Magnoli *et al.*, 2008b). Bentonites are basically clays with strong colloidal properties that absorb water rapidly, which results in swelling and a manifold increase in volume, giving rise to a thixotropic, gelatinous substance (Bailey *et al.*, 1998). Bentonites are composed of hydrated aluminosilicates of sodium (Na), potassium (K), calcium (Ca), and occasionally iron, magnesium, zinc, nickel, etc. They have a high negative charge and are balanced by cations such as Mg, K, and Na, therefore, they do not react with food/feed ingredients and act as inert material due to their neutral pH or slightly alkaline nature. However, the adsorption ability of these clays varies from one geological deposit to another.

In the current study, neither extrusion nor any of the adsorbent (Mycofix®Plus and SB) resulted in any significant improvement in birds performance. The reason for lack of significant effects for DT methods used could probably be due to the performance of the birds on the AF containing diets (treatment 2). If the diet containing 70ppb AF (Treatment 2) had negatively influenced performance, it would be expected that DT methods used would restore or improve production. Therefore, this result does not imply that the DT methods used are not effective but rather indicate that birds exposed to higher levels of AF in diets are more likely to be benefited from the detoxified feed.

4.2 Antibody titre

The means of antibody titre (HA) against Newcastle disease (ND) showed no difference ($P>0.05$) between treatments when analysed at 21, 28, 35 and 42 day of the trial (Table 3).

The presence of AF in the feed is reported to decrease vaccinal immunity and may therefore lead to the occurrence of disease even in properly vaccinated flocks (Lesson *et al.*, 1995). Aflatoxins have been associated to have immunosuppressive effect due to direct inhibition of protein synthesis, including those with specific functions such as immunoglobulins IgG, IgA, inhibition of migration of macrophages, interference with the haemolytic activity of

complement, reduction in the number of lymphocytes through its toxic effect on the Bursa of Fabricius and impairment of cytokines formation by lymphocytes (Tung *et al.*, 1975; Creppy *et al.*, 1979, Devegowda and Murthy, 2005). In present study, no difference ($P>0.05$) in HA titres was observed when treatment with AF (treatment 2) was compared with all other treatment groups suggests that birds exposed to 70ppb AF in diet do not show any signs of immunosuppression. Similarly, Gabal and Azzam, (1998) suggested that prolonged administration of AF at the low levels do not markedly change the hematological and serological parameters of broiler chickens, but may cause relevant lesions in liver and renal tissues. Moreover, the metabolism of broilers seems to be more adapted to high concentrations of aflatoxin in the feed when administered from 21 to 42 d of age, when compared with data reported from similar experiments conducted with broilers aging 1 to 21 d and with other species such as turkey poult (Pier *et al.*, 1979; Campbell *et al.*, 1988; Gabal and Azzam, 1998).

Treatment	Description	HA		
		Day 28	Day 35	Day 42
1	0 ppb AF & no DT	14.9	257.6	184.0
2	70 ppb AF & no DT	13.7	222.9	138.9
3	70 ppb AF + DT by extrusion	17.1	268.9	168.9
4	70 ppb AF + DT with Mycofix®Plus	13.9	237.2	168.9
5	70 ppb AF + DT with Sodium bentonite	16.5	245.3	184.4
	SED	4.65	23.6	17.65
	P-Value	0.924	0.396	0.143

Means within a column were not different ($P < 0.05$). Tukey’s test was used for means separation; SED – standard error of the difference.

Table 3. Effects of experimental diets on haemagglutination titres (HA) against ND at day 28, 35 and 42 of the trial.

In contrast to our study dietary AF has been reported to cause vaccine failure as indicated by significantly reduced ($P<0.05$) antibody titres against Newcastle disease vaccine when birds were fed diets containing 2000 to 3000 ppb AF (Rathore *et al.*, 1987; Mangat *et al.*, 1988; Viridi *et al.*, 1989; Ghosh *et al.*, 1990; Bakshi, 1991; Mohiuddin and Reddy, 1993; Sharma, 1993; Mohiudin, 1993). Similarly, Azzam and Gabal (1998) reported that even low levels of dietary AF (200 ppb) can cause reduction in antibody titers to vaccines for Newcastle disease, infectious bronchitis, and infectious bursal disease in layers, when fed for a longer period (40 weeks). This difference in response to HA titres results could be attributed to the higher inclusion levels of dietary AF (200 to 3000 ppb) used in these studies.

The use of feed adsorbents is considered the most promising and economical approach for reducing mycotoxicosis in animals. The beneficial effect of Mycofix® has been reported to ameliorate the negative effect of AF on IBDV antibody titres and the effects are attributed to the presence of phytogetic substances, a hepatoprotective flavolignins (silymarin) in Mycofix, which prevents toxins from entering the liver cell membranes, and as it contains the terpenoid complexes, which reduce inflammations and protect the mucous membranes (Biomin®, 2000). Similarly, Ibrahim *et al.*, (2000) reported that SB is also effective in ameliorating the suppressive effect of AF on the HI-titer in chicks vaccinated against Newcastle disease and the best result was obtained when SB was added at a rate of 0.4% of feed to the AF-containing diets. This effect was attributed to the role of SB as a sequestering

agent against AF present in the diet through reducing its bioavailability in the gastrointestinal tract (Araba and Wyatt, 1991). However, in the present study, no differences ($P>0.05$) in ELISA titres were observed when birds fed AF diet (Treatment 2) were compared with all other treatment groups. This result further support our growth performance results and indicates that diets low in AF (70 ppb) do not depress broiler growth and vaccinal immunity.

4.3 Bursal body weight ratio (BBR)

The sensitivity of the immune system to mycotoxin-induced immunosuppression arises from the vulnerability of the continually proliferating and differentiating cells that participate in immune mediated activities and regulate the complex communication network between cellular and humoral components. AF are reported to inhibits the histological development and functional maturation of lymphoid organs (Celik *et al.*, 2000). Morphological evidence to explain the immunosuppressive effects of AF (2500 ppb) was documented by Celik *et al.* (2000) in broiler chickens after 21 days of feeding and the major signs were reduction in the weights of lymphoid organs including bursa of Fabricius, spleen and thymus. Similarly, Verma *et al.*, (2004) reported a significant decrease in the relative weight of the bursa of Fabricius when birds were exposed to diets having 2000 ppb AF. Similar reduction in BBR and moderate histopathological changes have been reported in broilers (Giambrone *et al.*, 1985; Marquez and Hernandez, 1995), laying hens (Dafalla *et al.*, 1987), ducks (Sell *et al.*, 1998; Khajareru and Khajareru, 1999) and wild turkeys (Quist *et al.*, 2000) when birds were fed diets having various levels of AF (100 to 500 ppb). In addition, vacuolation of liver cells and cellular depletion in the follicle medulla of the bursa Fabricii has been reported to be produced as an indication of aflatoxicosis by feeding lower levels of AF (100 ppb) over a long-term period of 42 days (Espada *et al.*, 1992). In contrast, the present study indicated no significant difference ($P>0.05$) in BBR between different treatment groups (Table 4). However, it cannot be concluded from the present investigation whether 70 ppb AF level in broiler diet can cause aflatoxicosis in broilers, as no significant difference ($P>0.05$) was observed when different response parameters tested were compared to the those of AF contaminated diet. This difference in results probably could be due to differences in age or genetic strain of birds, nutritional status, and source of mycotoxins, exposure time, vaccination schedule, serologic technique and management practices used in these studies.

Treatment	Description	BBR
1	0 ppb AF & no DT	1.69
2	70 ppb AF & no DT	1.54
3	70 ppb AF + DT by extrusion	1.77
4	70 ppb AF + DT with Mycofix®Plus	1.60
5	70 ppb AF + DT with Sodium bentonite	1.70
	SED	0.77
	P-value	0.998

Means within a column were not different ($P < 0.05$). Tukeys T test was used for means separation; SED – standard error of the difference..

Table 4. Effect of experimental diets on the average mean bursal body weight ratio (BBR) of birds at 42 days of age.

5. Conclusion

The manifestation and magnitude of a AF related response depends upon dose of AF, time period that the determined dose is exposed to the animal and interactions (such as age of animal, nutritional status at the time of AF exposure, presence of multiple mycotoxins in the diets etc). Prevention and control of AF in the poultry production chain requires the knowledge and consideration of all factors influencing mycotoxin formation in the field and during the storage of feedstuffs. The results from the current study demonstrated that growth performance and immune response was not depressed when broilers aged 21 to 42 days were exposed to diets containing 70ppb AF. However, methods of DT compared did not result in any significant improvement ($P>0.05$) in any of the response parameter. Further studies are recommended to evaluate the efficacy of the detoxifying agents by using a factorial designs that include a non-contaminated diet and a contaminated diet, both with and without DT.

6. References

- Araba, M & Wyatt, RD. (1991). Effects of sodium bentonite hydrated aluminosilicate (Novasil) and charcoal on aflatoxicosis in broiler chickens. *Poultry Science*, 1(70): 6-11.
- Avantaggiato, G., Solfrizzo, M., Visconti, A. (2005). Recent advances on the use of adsorbent materials for detoxification of *Fusarium* mycotoxins. *Food additives and Contaminants*, 22: 379-388.
- Azzam A. H. & Gabal M. A. (1998). Aflatoxin and immunity in layer hens. *Avian Pathology*; 27:570-577.
- Bailey, C. A., Latimer, G. W, A., Barr, C., Wigle W. L., Haq, A. U., Balthrop, J. E. & Kubena, L. F. (2006). Efficacy of montmorillonite clay (NovaSil PLUS) for protecting full-term broilers from aflatoxicosis. *Journal of Applied Poultry Science*, 15:198-206
- Bailey, R.H., Kubena, L.F., Harvey, R.B., Buckley, S.A. & Rottinghaus, G.E. (1998). Efficacy of various inorganic sorbents to reduce the toxicity of aflatoxin and T-2 toxin in broiler chickens. *Poultry Science*, 77, 1630-1632.
- Bakshi, C.S. (1991). Studies on the effect of graded dietary levels of aflatoxin on immunity in commercial broilers. *M.V.Sc. Thesis*, Indian Veterinary Research Institute, Izatnagar, U.P., India.
- Barabolak, R.J. (1977). Improved procedure for quantitative determination of aflatoxin in corn and wet milled corn products. *AOAC*, 60 (2): 308.
- Basmacioglu, H., Oguz, H., Ergul, M., Col, R. & Birdane, Y. O. (2005). Effect of dietary esterified glucomannan on performance, serum biochemistry and haematology in broilers exposed to aflatoxin. *Czech Journal of Animal Science*, 50: 31-39.
- Beura, C.K., Johri, T.S., Sadagopan, V.R. & Panda, B.K. (1993). Interaction of dietary protein level on dose response relationship during aflatoxicosis in commercial broilers. I. Physical responses livability and nutrient retention. *Indian Journal of Poultry Science*, 28: 170-177.
- Bintvihok, A., and. Kositcharoenkul, S. (2006). Effect of dietary calcium propionate on performance, hepatic enzyme activities and aflatoxin residues in broilers fed a diet containing low levels of aflatoxin B₁. *Toxicon*, 47:41-46

- Biomin®. (2000) Mycofix® Plus 3.0. A modular system to deactivate mycotoxins. *Biomin*, GTI GmbH. Herzogenburg, Austria.
- Campbell, M.L., May, D., Huff, W.E. & Doer, J.A. (1988). Evaluation of immunity of young broiler chickens during simultaneous aflatoxicosis and ochratoxicosis. *Poultry Science*, 62: 2138-2144
- Celik, I., Oguz, H., Demet, O., Donmez, H.H., Boydak, M. and Sur, E. (2000). Efficacy of polyvinylpyrrolidone in reducing the immunotoxicity of aflatoxin in growing broilers. *British Poultry Science*, 41: 430-439
- Cheftel, J. C. (1986). Nutritional effects of extrusion-cooking. *Food Chemistry*. 20. 262-285.
- Cheftel, J. C. (1989). Extrusion cooking and food safety. In *Extrusion Cooking* Eds. Mercier, C., Linko, P. and Harper, J.M. pp. 435-462. St Paul, Minnesota, USA: American Association of Cereal Chemists.
- Coker, R.D. (1979). Aflatoxin : past, present and future.- *Tropical Science*, 21: 143-162.
- Coker, R.D., Jewers, K. & Jones, B.D. (1986). The treatment of aflatoxin contaminated commodities. *International Biodeterioration*, 22S, pp. 103-108.
- Coker, R.D., Jewers, K., Jones, B.D. (1985). The destruction of aflatoxin by ammonia : practical possibilities. *Tropical Science*, 25: 139-154.
- Council for Agricultural Science and Technology (CAST). (2003). *Mycotoxins: Risks in plant, animal and human systems*. Task Force Report No 139. Ames, Iowa, USA.
- Creppy, E.E., Lugnier, A.A.J., Fasiolo, F., Heller, K., Roschenthaler, R. & Dirheimer, G. (1979). In vitro inhibition of yeast phenylalanyl-t-RNA synthetase by ochratoxin A. *Chemical and Biological Interactions*, 24: 257-261.
- Dafalla, R., Yagi, A.I. & Adam, S.E.I. (1987). Experimental aflatoxicosis in hybro-type chicks; sequential changes in growth and serum constituents and histopathological changes. *Veterinary and Human Toxicology* 29: 222-225.
- Dalvi, R. R. & A. A. Ademoyero. (1984). Toxic effects of aflatoxin B₁ in chickens given feed contaminated with *Aspergillus flavus* and reduction of the toxicity by activated charcoal and some chemical agents. *Avian Disease*, 28: 61-69
- Dalvi, R.R. (1986). An overview of aflatoxicosis of poultry. Its characteristics, prevention and reduction. *Veterinary Research and Communication*, 10: 429-443.
- Denli, M., Okan, F. & Doran, F. (2004). Effect of conjugated linoleic acid (CLA) on the performance and serum variables of broiler chickens intoxicated with aflatoxin B₁. *South African Journal Of Animal Sciences*, 34: 97-103
- Desphande, S. S. (2002). Fungal toxins. In S. S. Desphande (Ed.), *Handbook of food toxicology*, New York: Marcel Decker. 387-456.
- Devegowda, G & Murthy, T.N.K. (2005). Mycotoxins: Their effects in poultry and some practical solutions. In. *The Mycotoxin Blue book*, Ed: Diaz, D.E. pp: 25-57. Nottingham University press, United Kingdom:
- Diaz, D. E., Hagler, W. M, Hopkins, B. A. & Whitlow, L. M. (2002). Aflatoxin binders I: In vitro binding assay for aflatoxin B₁ by several potential sequestering agents. *Mycopathologia*, 156: 223-226
- Doerr, J.A., Huff, W.E., Webeck, C.J., Chhaloupka, G.W., May, J.D. & Merkley, J.W. (1983) Effects of low level chronic aflatoxicosis in broiler chickens. *Poultry Science*, 62: 1971 – 1977.
- Dutta, T.K. & Das, P.I. (2001). Isolation of aflatoxigenic strains of aspergillus and detection of aflatoxin B₁ from feeds in India, *Mycopathologia*, 151: 29-33.

- Espada, Y., Domingo, M., Gomez, J. & Calvo, M.A. (1992). Pathological lesions following an experimental intoxication with aflatoxin B1 in broiler chickens. *Research in Veterinary Science*, 53: 275–279.
- Fast, R.B. (1991) Manufacturing technology of ready-to-eat cereals. In *Breakfast Cereal and How They Are Made*, Eds. Fast, R.B. and Caldwell, E.F. pp. 15±42. St Paul, Minnesota, USA: American Association of Cereal Chemists.
- Feuell, A. (1996). Aflatoxins in groundnuts. Part 9: Problems of detoxification. *Tropical Science*, 8: 61-70.
- Food and Agricultural Organization.(1995). Worldwide Regulations for Mycotoxins. *FAO Food and Nutrition paper 64*, FAO, Vialledella Terme di Caracalla, 00100, Rome, Italy.pp.43.
- Gabal, M. A. and Azzam, A. H. (1998). Interaction of aflatoxin in the feed and immunization against selected infectious diseases in poultry. II. Effect on one-day-old layer chicks simultaneously vaccinated against Newcastle disease, infectious bronchitis and infectious bursal disease', *Avian Pathology*, 27(3): 290 – 295
- GenStat. 11th edn for windows. Hemel Hempstead, UK: VSN International Ltd.
- Ghosh, R.C., Chauhan, H.V.S. & Roy, S. (1990). Immunosuppression in broilers under experimental aflatoxicosis. *British Veterinary Journal*, 146: 457 – 462.
- Giambrone, J.J., Diener, U.L., Davis, N.D., Panangala, V.S. & Hoerr, F.J. (1985). Effects of aflatoxin on young turkeys and broilers chickens. *Poultry Science*, 64, 1678–1684.
- Goto, T., Peterson, S. W., Ito, Y. & Wilkins, D. T. (1997). Mycotoxin producing ability of *Aspergillus tamari*. *Mycotoxins*, 44: 17-20.
- Grehaigne, B., Chouvel, H., Pina, M., Graille, J. & Cheftel, J.C. (1983). Extrusion cooking of aflatoxin containing peanut meal with and without addition of ammonium hydroxide. *Lebensmittel-Wissenschaft und-Technologie*, 16: 317–322.
- Harper, J.M. (1989). Food extruders and their applications. In *Extrusion Cooking*, Ed. Mercier, C., Linko, P. and Harper, J.M. pp. 1±16. St Paul, Minnesota, USA: American Association of Cereal Chemists.
- Huff, W.E. & Doerr, J.A. (1981). Synergism between aflatoxin and ochratoxin A in broiler chickens. *Poultry Science*, 60: 550 – 555.
- Huff, W.E., Kubena, L.F., Harvey, R.B. & Phillips, T.D. (1992). Efficacy of hydrated sodium aluminosilicate to reduce the individual and combined toxicity of aflatoxin and ochratoxin A. *Poultry Science*, 71: 64 – 69.
- Ibrahim, I.K., Shareef, A.M., Al-Joubory, K.M.T. (2000). Ameliorative effects of sodium bentonite on phagocytosis and Newcastle disease antibody formation in broiler chickens during aflatoxicosis. *Research in Veterinary Science*, 69: 119–122.
- Jewers, K. (1990). Mycotoxins and their effect on poultry production. *Options Méditerranéennes, Sér. A/No. 7*:195–202.
- Johri, T.S. & Majumdar, S. (1990). Effect of methionine, choline, BHT, supplemented aflatoxic diets, *Proc. XIII National Symposium of Indian Poultry Science.*, Assoc, held at Bombay on 20-22 Dec., 1990.
- Johri, T.S. & Sadagopan, V.R. (1989). Aflatoxin occurrence in feedstuffs and its effect on poultry production. *Journal of Toxicology and Toxin Reviews*, 8: 281 – 287.
- Khajareen, J. & Khajareen, S. (1999). Positive effects of Mycosorb against aflatoxicosis in ducklings and broilers. In: Poster presentation at *Alltech's 15th Annual Symposium on Biotechnology in the Feed Industry*, Lexington, KY.

- Kohlwey, D.E., Kendall, J.H. & Mohindra, R.B. (1995). Using the physical properties of rice as a guide formulation. *Cereal Food World*, 40: 728-732.
- Kubena, L.F., Edrington, T.S., Kamps-Holtzapple, Harvey, C. R., Elissalde, M.H. & Rottinghaus, G.E. (1995). Effect of feeding fumonisin B1 present in *Fusarium moniliforme* culture material and aflatoxin singly and in combination to turkey poults. *Poultry Science*, 74: 1295-1303.
- Kubena, L.F., Huff, W.E., Harvey, R.B., Yersin, A.G., Elissad, M.H., Witzel, D.A., Giroir, L.E., Philips, T.D. & Petersen, H.D. (1991). Effect of a hydrated sodium calcium aluminosilicate on growing turkey poults during aflatoxicosis. *Poultry Science*, 70: 1823-1930.
- Lanza, G.M., Washburn K.W. & Wyatt, R.D. (1980). Variation with age in response of broilers to aflatoxin. *Poultry Science*, 59(2): 282-288.
- Leeson, S., Diaz, G., & Summers, J.D. (1995). Aflatoxins. In: Leeson, S., Diaz, G., Summers, J.D. (Eds.), *Poultry Metabolic Disorders and Mycotoxins*. pp. 248-279. University Books, Canada, Ont.
- Lillehoj, E. B., Garcia, W. J. & Lambrow M. (1974). *Aspergillus flavus* infection and aflatoxin production in corn: influence of trace elements. *Applied Microbiology*, 28 (5): 763-767.
- Magnoli, A.P., Cavaglieri, L., Magnoli, C., Monge, J.C., Miazzo, R., Peralta, M. F., Salvano, M., Rosa, R.C.A., Dalcero, A. & Chiacchiera, S.M. (2008a). Bentonite performance on broiler chickens fed with diets containing natural levels of aflatoxin B₁. *Revue de Medecine Veterinaire*, 30: 55-60.
- Magnoli, A.P., Monge, M.P., Miazzo, R.D. , Cavaglieri, L.R., Magnoli, C.E., Merkis C.I., Cristofolini, A.L., Dalcero, A.M., & Chiacchiera, S.M. (2011). Monensin affects the aflatoxin-binding ability of a sodium bentonite. *Poultry Science*, 90: 48-52.
- Magnoli, A.P., Tallone, L., Rosa, R.C.A., Dalcero A.M., Chiacchiera S.M., & Torres Sanchez R.M. (2008b). Structural characteristic of commercial bentonites used as detoxifier of broiler feed contamination with aflatoxin. *Applied Clay Science*, 40: 63-71.
- Mangat, A.P.S., Gill, B.S. & Maiti, N.K. (1988). Immunosuppressive effect of low level of aflatoxin against Ranikhet disease in chicken. *Indian Journal of Comparative Microbiology, Immunology and Infectious Diseases*, 10: 25 – 30.
- Marquez, R.N.M. & Hernandez, T.R. (1995). Aflatoxin adsorbent capacity of two Mexican aluminosilicates in experimentally contaminated chick diet. *Food Additives and Contaminants*, 12: 431-433.
- Miazzo, R., Rosa, C.A., De Queiroz Carvalho, E.C., Magnoli, C., Chiacchiera, S.M., Palacio, G., Saenz, M., Kikot, A., Basaldella, E. & Dalcero, A. (2000). Efficacy of synthetic zeolite to reduce the toxicity of aflatoxin in broiler chicks. *Poultry Science*, 79: 1-6.
- Mohiuddin, S.M. & Reddy, V.M. (1993). Immunosuppressive effect of aflatoxin in bursectomised chicks against Ranikhet disease vaccine. *Indian Journal of Animal Sciences*, 63: 279 – 280.
- Mohiudin, S M. (1993). Effects of aflatoxin on immune response in viral diseases. *Poultry adviser*, 26: 63-66
- Nandakumar Reddy, D. Rao, P.V., Reddy, V.R. & Yadgiri, B. (1984). Effect of selected levels of dietary aflatoxin on the performance of broiler chicken. *Indian Journal of Animal Science*. 54(1): 68-73.
- Oguz, H. & Kurtoglu, V. (2000). Effect of clinoptilolite on fattening performance of broiler chickens during experimental aflatoxicosis. *British Poultry Science*, 41: 512-517.

- Oguz, H., Hadimli, H.H., Kurtoglu, V., & Erganis, O. (2003). Evaluation of humoral immunity of broilers during chronic aflatoxin (50 and 100 ppb) and clinoptilolite exposure. *Revue de Medicine Veterinaire* 154: 483–486.
- Oguz, H., Kecec, T., Birdane, Y.O., Onder, F. & Kurtoglu, V. (2000a). Effect of clinoptilolite on serum biochemical and haematological characters of broiler chickens during experimental aflatoxicosis. *Research in Veterinary Science*, 69: 89–93.
- Oguz, H., Kurtoglu, V. & Coskun, B. (2000b). Preventive efficacy of clinoptilolite in broilers during chronic aflatoxin (50 and 100 ppb) exposure. *Research in Veterinary Science*, 69: 197–201.
- Ortatatli, M., Ouz, H., Hatipoglu, F. & Karaman, M. (2005). Evaluation of pathological changes in broilers during chronic aflatoxin (50 and 100 ppb) and clinoptilolite exposure. *Research Veterinary Science*, 78: 61–68.
- Osborne, D.J. & Hamilton, P.B. (1981). Decreased pancreatic digestive enzymes during aflatoxicosis. *Poultry Science*, 60: 1818–1821.
- Park, DL. (2002). Mycotoxins and food safety. In: *Advances in experimental medicine and biology* Eds: DeVries JW; Trucksess MW; Jackson LS 504: 173–179
- Parlat, S.S., Yildiz, A.O. & Oguz, H. (1999). Effect of clinoptilolite on fattening performance of Japanese quail (*Coturnix coturnix japonica*) during experimental aflatoxicosis. *British Poultry Science*, 40: 495–500.
- Pasha, T.N., Farooq M.U., Khattak, F.M., Jabbar, M.A., & Khan, A.D. (2007). Effectiveness of sodium bentonite and two commercial products as aflatoxin absorbents in diets for broiler chickens. *Animal Feed Science & Technology*, 132:103–110.
- Pasha, T.N., Mahmood, A., Khattak, F.M., Jabbar, M.A., & Khan, A.D. (2008). The effect of feed supplementation with different sodium bentonite treatments on broiler performance. *Turkish Journal of Veterinary and Animal Sciences*, 32: 245–248.
- Pier, A.C., Richard, J.L. & Thurston, J.R. (1979). The influence of mycotoxins on resistance and immunity. In *Interaction of Mycotoxins in Animal Production*, (pp. 96–117). Washington, DC: National Academy of Sciences.
- Piva, G., Galvano, F.P.F., Pietri, A. & Piva, A. (1995). Detoxification methods of aflatoxins. *A Review of Nutrition Research*.15: 767–776.
- Quist, C.F., Bounous, D.I., Kilburn, J.V., Nettles, V.F. & Wyatt, R.D. (2000.) The effect of dietary aflatoxin on wild turkey poults. *Journal of Wildlife Disease*, 36: 236–444.
- Rajashekhara Reddy, A., Reddy, V.R., Rao, P.V. & Yadgiri, B. (1982). Effect of experimentally induced aflatoxicosis on the performance of commercial chicks. *Indian Journal of Animal Science*, 52: 405–410.
- Ramos, A. J., and E. Hernandez. (1997). Prevention of aflatoxicosis in farm animals by means hydrated sodium calcium aluminosilication addition to feedstuffs. A review. *Animal. Feed Science & Technology*, 65: 197–206.
- Rathore, B.S., Verma, K.C., Singh, S.D. & Khera, S.S. (1987) Epidemiological studies on Ranikhet disease vaccine failure in chickens. *Indian Journal of Comparative Microbiology, Immunology and Infectious Diseases*, 8: 175–177.
- Reddy, P.S., Reddy, C.V., Reddy, V.R. & Rao, P.V. (1984) Occurrence of aflatoxin in some feed ingredients in three geographical regions of Andhra Pradesh. *Indian Journal of Animal Science*, 54: 235–238.
- Rosa, C.A., Miazzi, R., Magnoli, C., Salvano, M., Chiac, S.M., Ferrero, S., Saenz, M., Carvalho, E.C. & Dalcero, A. (2001). Evaluation of the efficacy of bentonite from the

- south of Argentina to ameliorate the toxic effects of aflatoxin in broilers. *Poultry Science*, 80: 139–144.
- Sell, S., Xu, K.L., Huff, W.E., Kubena, L.F., Harvey, R.B. & Dunsford, H.A. (1998). Aflatoxin exposure produces serum a fetoprotein elevations and marked oval cell proliferation in young male Pekin ducklings. *Pathology* 30: 34–39.
- Sever, J. L. (1962). Application of a microtechnique to viral serological investigations. *Journal of Immunology*, 88; 320–329.
- Sharline, K.S.B., B.J. Howarth & R.D. Wyatt. (1980). Effect of dietary aflatoxin on reproductive chicks. *Poultry Science*, 72: 651–657.
- Sharma R. P. (1993). Immuntotoxicity of mycotoxins. *Journal of Dairy Science*, 76: 892– 897.
- Shi, Y. H., Xu, Z. R., Feng, J. L. & Wang, C. Z. (2006). Efficacy of modified montmorillonite nanocomposite to reduce the toxicity of aflatoxin in broiler chicks. *Animal Feed Science Technology*. 129: 138–148
- Shi, Y., Xu, Z., Sun, Y., Wang, C.& Feng, J. (2009). Effects of two different types of motmorillonite on growth performance and serum profiles of broiler chicks during aflatoxicosis. *Turkish Journal of Veterinary and Animal Science*, 33: 15–20.
- Smith, J.W. & Hamilton, P.B. (1970). Aflatoxicosis in broiler chickens. *Poultry Science*, 49: 207 – 215.
- Tedesco, D., Steidler S., Galletti, S., Taneni, M., Sonzogni O.& Ravarotto, L.(2004). Efficacy of silymarin-phospholipid complex in reducing the toxicity of aflatoxin B₁ in broiler chicks. *Poultry Science*, 83:1839–1843
- Tung, H.T., Wyatt, R.D., Thaxton, P. & Hamilton, P.B. (1975) Concentration of serum proteins during aflatoxicosis. *Toxicology and Applied Pharmacology*, 34: 320 – 326.
- Verma, J., Johri, T. S. , Swain, B. K. and Ameena, S. (2004) 'Effect of graded levels of aflatoxin, ochratoxin and their combinations on the performance and immune response of broilers. *British Poultry Science*, 45 (4):512-518.
- Verma, J., B. K. Swain, and T. S. Johri. (2002). Effect of various levels of aflatoxin and ochratoxin A and combinations thereof on protein and energy utilisation in broilers. *Journal of Science Food and Agriculture*, 82:1412–1417.[
- Virdi, J.S., Tiwari, R.P., Saxena, M., Khanna, V., Singh, G., Saini, S.S. & Vadehra, D.V. (1989) Effect of aflatoxin on immune system of the chick. *Journal of Applied Toxicology*, 9: 271-275.

Aflatoxins - Detection, Measurement and Control

Edited by Dr Irineo Torres-Pacheco

ISBN 978-953-307-711-6

Hard cover, 364 pages

Publisher InTech

Published online 21, October, 2011

Published in print edition October, 2011

This book is divided into three sections. The section called Aflatoxin Contamination discusses the importance that this subject has for a country like the case of China and mentions examples that illustrate the ubiquity of aflatoxins in various commodities. The section Measurement and Analysis, describes the concept of measurement and analysis of aflatoxins from a historical perspective, the legal, and the state of the art in methodologies and techniques. Finally the section entitled Approaches for Prevention and Control of Aflatoxins on Crops and on Different Foods, describes actions to prevent and mitigate the genotoxic effect of one of the most conspicuous aflatoxins, AFB1. In turn, it points out interventions to reduce identified aflatoxin-induced illness at agricultural, dietary and strategies that can control aflatoxin. Besides the preventive management, several approaches have been employed, including physical, chemical biological treatments and solvent extraction to detoxify AF in contaminated feeds and feedstuffs.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

T. Mahmood, T.N. Pasha and F.M. Khattak (2011). Comparative Evaluation of Different Techniques for Aflatoxin Detoxification in Poultry Feed and Its Effect on Broiler Performance, Aflatoxins - Detection, Measurement and Control, Dr Irineo Torres-Pacheco (Ed.), ISBN: 978-953-307-711-6, InTech, Available from: <http://www.intechopen.com/books/aflatoxins-detection-measurement-and-control/comparative-evaluation-of-different-techniques-for-aflatoxin-detoxification-in-poultry-feed-and-its->

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2011 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](https://creativecommons.org/licenses/by/3.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen