We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Multi-Trial Ecotoxicological Diagnostic Tool in Cetacean Skin Biopsies

Maria Cristina Fossi and Letizia Marsili
*Department of Environmental Sciences,
 University of Siena, Siena,
 Italy*

1. Introduction

In the last 20 years there has been growing concern about potential hazard to cetaceans occasioned by multiple stress factors due to bioaccumulation of anthropogenic contaminants combined with infectious diseases, food depletion and climate change. In this context, the main aim of this chapter is to present the development and the application of a non-lethal multi trial biomarker tool applied to cetacean skin biopsies to diagnose “toxicological stress syndromes” related to multiple human pressures. Here we present a multidisciplinary approach to detect presence and effects of anthropogenic contaminants combined with general stressors using a wide range of diagnostic markers (molecular and gene expression biomarkers, immunological and nutritional status) with contaminant analysis, for a complete assessment of health status of cetaceans. The chapter is subdivided in three sections: in section one we describe the main threats faced by cetaceans focusing on the case study of Mediterranean Sea; in the second part we will describe the sensitivity of the multi trial biomarker diagnostic tool in skin biopsy cetacean samples; and in the third part we will describe the application of the proposed multidisciplinary approach to explore the effects of multiple human pressures on cetacean health status (field applications), comparing fin whale (*Balaenoptera physalus*) (mysticete) population of Mediterranean compared to the one of Sea of Cortez (Mexico). Final development of this interdisciplinary diagnostic methodology, embedded in a statistical Expert System, will provide exhaustive information about “toxicological status” of cetaceans as a powerful frontier approach for the conservation of relevant species of sea biodiversity.

2. Section one - Threats faced by cetaceans: The case study of Mediterranean sea

Threats facing cetaceans have changed through time. While overkill from hunting was the most obvious and immediate threat to some species and populations during much of the 20th century, the relative importance of other threats, particularly bycatch in fisheries and contamination, has increased dramatically during the last few decades. It is difficult to distinguish the effects of one threat from those of another when multiple threats are acting simultaneously; this is particularly evident in the Mediterranean area. Exposure to contaminants combined with other human impacts can affect survivorship, recruitment,

reproductive success, mutation rates and migration and may play a significant role in the partitioning of genetic variation among high stress exposed and less stress exposed populations (Whitehead et al., 2003). Taking into account this complex pattern, international institutions like IWC and ACCOBAMS have recently encouraged research for development of suites of sensitive non-lethal tools in skin biopsies of free-ranging animals, to define the “health status” of threatened cetacean species (Pollution 2000+, IWC Project). Threats faced by cetaceans can be schematized below.

2.1 Pollution: Bioaccumulation and effects of anthropogenic contaminants

Mediterranean top predators, and particularly cetacean odontocetes, accumulate high concentrations of anthropogenic contaminants such as organochlorine compounds (OCs), polycyclic aromatic hydrocarbons (PAHs) and trace elements. The pollutant burden is lower at the lowest levels of the food chain and becomes higher as the position in the food web increases. Important factors for bioaccumulation are also metabolic rates, gender, age and adipose tissue concentration. The ecotoxicological risk of some species is also related to their “biochemical vulnerability” to lipophilic xenobiotics due to the low capacity of their detoxifying enzymatic systems (Tanabe & Tatsukawa, 1992; Fossi et al., 1997). Since the incidence of pathology in these species is closely related to the level of pollution in their environments, bacterial and viral infections and contaminants must be considered from a holistic point of view. Mass mortalities of dolphins and seals have occurred in particularly polluted areas (such as the Mediterranean Sea), when levels of OCs, PAHs and heavy metals reached very high levels. It is reported that lipophilic contaminants cause immune and reproductive dysfunctions (Hammond et al., 2005). Some contaminants, such as OCs are also known as endocrine disrupting chemicals (EDCs) (Fossi et al., 2003). In fact several examples suggest that exposure to levels of OC insecticides and PCBs commonly detected in Mediterranean odontocetes has affected endocrine function and reproduction in other marine mammal species (Fossi & Marsili, 2003). Moreover there is growing concern about accumulation and effects of emerging contaminants and polybrominated diphenyl ethers (PBDEs, a major family of flame retardants) in the food chain. PBDEs are lipophilic, persistent and toxic to fauna and humans (Alaee et al., 2003). The highest levels of PBDEs have been found in top marine predators, including Mediterranean odontocetes. Also phthalates (found in numerous commercial products as additives and used primarily to soften polyvinyl chloride - PVC) are a relevant and emerging cause of concern for these species, considered as potential EDCs. Some phthalates have been linked to reproductive effects and altered hormone levels, however the effects and concentrations of these compounds have not been assessed yet. Another important class of contaminants for cetaceans is PAHs, derived from both natural (e.g., oil spill, forest fires, natural petroleum seeps) and anthropogenic sources (e.g., combustion of fossil fuels, use of oil for cooking and heating, coal burning). Studies have shown that PAHs with four or more rings can induce dioxin-like activity and weak estrogenic responses (Villeneuve et al., 2002). As pointed out by the 2010 scientific expedition undertaken by the program Mediterranean Endangered, new priorities in marine environment contamination are the so-called microplastics (usually defined as plastic particles smaller than 5 mm) (Barnes et al., 2009). Micro debris floating on the Mediterranean Sea have reached 115,000 particles per km² with a maximum of 892,000 particles. Impacts of microplastics to organisms and the environment are largely unknown. More than 180 species have been documented to absorb plastic debris including planktophagous species. For example, laboratory study has shown that krill species *Euphasia*

pacifica ingested 20µm polyethylene fragments. Clearly, considerably reducing microplastic inputs must be a priority of this decade for the biodiversity conservation in the EU strategy.

2.2 Diseases: Immunosuppression and infectious diseases

Cetaceans are normally susceptible to a large plethora of pathogenic organisms. For example, the family *Cyamidae* comprises 28 species of parasitic crustaceans, all of which live exclusively on the cetaceans skin (Haney et al., 2004). The majority of these infective organisms are repelled by innate host defences that include nonimmunological anatomical and physiological barriers (e.g. mucociliary blanket), antimicrobial factors (e.g. lactoferrin, defensins, immunolysozyme) and immunological effector cells (e.g. neutrophils, eosinophils, macrophages and natural killer cells). Several studies have suggested that contaminants could have immunosuppressive effects on marine mammals (Beckmen et al., 2003). Since 1987, at least 8 epidemics of *Morbillivirus* infection (MI) have caused mass mortality of various free-living pinniped and cetacean populations worldwide. Along the coast of Spain and Italy new cases of MI were reported in 2007 and 2008. Another intriguing issue regards the synergic effects, associated with chronic exposure to a number of environmental pollutants, such as OCs and trace elements, as previously mentioned. In fact, it is also unknown whether and how these chemicals contribute towards modulating the pathogenic and pathogenetic activity primarily displayed by sea mammal MI (Di Guardo et al., 2005).

2.3 Bycatch

Several species of marine mammals are at risk of extinction from being captured as bycatch in commercial fisheries. In the Mediterranean Sea sperm whales (*Physter macrocephalus*), striped dolphins (*Stenella coeruleoalba*), and short-beaked common dolphins (*Delphinus delphis*) are affected by pelagic driftnets and gillnets. The only mitigation measure applied today is the ban of swordfish driftnet fishery (Young & Iudicello, 2007). Other approaches that have been developed and implemented to address this problem, include devices and gear changes, time and area closures and moratoria. The scientific committee of ACCOBAMS and the IWC indicate the bycatch as one of the major threats to marine mammals. Between 1999 and 2004, SOS Grand Bleu counted 50,000 deaths due to sword fish “spadare”.

2.4 Shipping, collision and noise pollution

All cetacean species, from fin whales to bottlenose dolphins (*Tursiops truncatus*) and from pelagic to coastal species are potentially subjected to collisions with boats, even if most collisions go unreported. Several cases of collisions between fin whales and large ships are reported from stranding cetaceans and from photo-identification data (Panigada et al., 2006). Cetaceans need acoustic signals to navigate, find and capture prey, and to locate mates, social partners and predators. Heavy vessel traffic, seismic testing, dredging, and drilling for oil and gas cause noise travelling thousands of km underwater. There is no doubt that cetaceans react to noise, but it is extremely difficult to quantify the effects and establish thresholds of disturbance. It is reported that use of mid-frequency long-range active sonar by navy caused the stranding of 11 Cuvier's beaked whales (*Ziphius cavirostris*) on the coastline of Greece (Frantzis, 1998).

2.5 Food depletion

Fisheries can have a major impact upon the status and distribution of various cetaceans. In the Mediterranean Sea, the decline in high-order marine predators feeding on epipelagic

prey (cetaceans, large pelagic fish) is consistent with the hypothesis of prey depletion, presumably due to intensive exploitation of fish stocks (over-fishing). In fact the combination of some 50,000–100,000 small gillnet fishing boats, plus large bottom trawlers, has depleted numerous fish, crustacean, and mollusc populations. In eastern Ionian Sea, the short-beaked common dolphin shows evidence of decline correlated with food depletion (Bearzi et al., 2006).

2.6 Climate change

It is generally agreed that the Mediterranean region is an area sensitive to global climate change, due to its position on the edge of the climatologically determined Hadley cell, which makes it a transition area between two very different climate regimes to the North and the South (Li et al., 2006). Many global and regional models simulate several degrees of warming (from 3 to 7°C) in the Mediterranean by the end of the 21st century. The potential impacts of climate change include drought, decline of water quality, floods, and change in seawater temperature/salinity, the rising of sea levels and reduction of biodiversity. This would pose several threats to cetaceans, mainly linked to changes in the distribution of sufficient quality and quantity prey of marine mammals (Simmonds & Nunny, 2002). Climate-related effects may combine with other factors such as pollution and disease to have significant impacts on cetacean populations.

2.7 Genetic erosion

Studies to date have indicated a complex pattern of population genetic structure for most cetacean species investigated. Seasonal patterns of movement and the possibility of extremely large scale dispersal, or local isolation (sympatric or parapatric) between populations, generate a mosaic of genetic diversity that cannot easily be determined by an intuitive assessment of geography. Although the Mediterranean basin is closed and has the heavily industrialized coasts, it hosts several cetacean species biologically differentiated between pelagic (*Stenella coeruleoalba*) and coastal (*Tursiops truncatus*) odontocetes and the resident mysticete (*Balaenoptera physalus*). Previous studies showed evidence of clear genetic differentiation between Atlantic and Mediterranean populations of striped dolphin and found much lower genetic variation of Mediterranean specimens (Garcia-Martinez et al., 1999). Gene flow through the Gibraltar Strait appears to be limited and easily prevents the genetic exchange between the Atlantic and Mediterranean cetacean populations. For the determination of population structure of wild species, genetic information about population subdivision has practical applications for short- and long-term strategies for animal management, conservation and protection (Hoelzel, 1998) and genetic diversity appears to be a major factor determining the success of a species in harsh environments. High genetic diversity within a population increases the chance for survival of at least some individuals, and possible changes in the environment, due to natural causes or anthropogenic pressures, can be better tolerated. Exposure to contaminants combined with other human impacts can affect survivorship, recruitment, reproductive success, mutation rates and migration, and may play a significant role in partitioning genetic variation among high stress exposed and less stress exposed populations (Whitehead et al., 2003). This is particularly true for populations inhabiting a closed basin characterized by considerable concentrations of environmental contaminants (OCs), orders of magnitude higher than those detected in oceanic environments. Bioaccumulation of pollutants, combined with other factors, such as

immunosuppression, infectious diseases, climate change, and food depletion could soon lead to a further additional decrease in total genetic variability of Mediterranean cetacean populations, posing alarming threat to these species.

3. Section two - Skin biopsy for diagnosis of anthropogenic threats

Cetacean skin biopsies are biological material suitable for the hazard assessment of free-ranging cetaceans (Fossi et al., 2010). Our research team firmly believe that the use of cetacean skin biopsies, as non-lethal biological material for the hazard assessment, is a powerful procedure to screen a large number of samples, with a minimal disturb to animals. In this chapter integument biopsy will be described as diagnostic tool for the comprehensive diagnosis of multiple stress factors, health status and genetic population variability. The diagnostic set developed in our lab and described in this chapter will enable to detect: i) presence of contaminants, ii) exposure to contaminants, iii) reproductive alteration, iv) genotoxicity, v) immunosuppression, vi) general stress, vii) feeding ecology (Fig.1).

Fig. 1. Skin biopsy for diagnosis of anthropogenic threats

3.1 Sampling of integument biopsies

Samples of skin biopsy or integument biopsy (epidermis, dermis and blubber) can be obtained from free-ranging dolphin (es. *Tursiops truncatus*, *Stenella coeruleoalba*) from different study areas using an aluminium pole armed with biopsy tips (0.7 cm ø, 3.0 cm length) or with a crossbow and darts. Integument biopsies from large odontocete (es. *Physeter macrocephalus*) or mysticete (*Balaenoptera physalus*) can be obtained using a crossbow

and darts armed with tips (0.9 cm ø, 4.0 cm length). A Barnett Wildcat II crossbow with a 150-pound test bow, using a biopsy dart with modified stainless steel collecting tip and floater, can be used to collect biopsies. To avoid the possibility of infection, the bolt tip need to be sterilised before shooting. Biopsy samples can be taken between the dorsal fin and the upper part of the caudal peduncle. The procedure consists in approaching the specimen at low-to-moderate speed and shooting the dart at a range of 20-50 m. The skin biopsy will be immediately placed in liquid nitrogen or stored in cell medium (CITES Nat. IT025IS, Int. CITES IT 007).

3.2 Detection of anthropogenic contaminants

Analysis of contaminant levels provide important data about exposure of cetaceans, that can be correlated with the biomarker data for a complete evaluation of their toxicological stress. The blubber layer of cetacean integument biopsy is the target tissue for the analysis of lipophilic and non-lipophilic contaminants in free-ranging populations (Fossi et al., 2000). A set of major POPs, trace elements and emerging contaminants can be analysed in skin biopsies of different cetacean species. The methods are reported below.

Trace elements: For trace elements analysis, lyophilised and homogenised samples can be acid-digested and analysed for trace elements using Atomic Absorption Spectrometry and Emission Spectrometry with FIMS-AAS (Flow Injection Mercury System) for determining Hg, THGF-AAS (Transversely Heated Graphite Furnace) and ICP-AES (Inductively Coupled Plasma).

Organochlorines: The analytical method used for quantitative and qualitative analysis of HCB, DDTs, PCDDs, PCDFs and PCBs will be High Resolution Capillary Gas chromatograph equipped with an electron capture detector (63Ni ECD)(AGILENT 6890/N), according to the U.S. Environmental Protection Agency (EPA) 8081/8082 modified (Marsili & Focardi, 1996). The GC has a SPB-5 bonded phase in a fused silica capillary column, 30 m long.

PAHs: Levels of PAHs and PAH fingerprint will be evaluated by High Performance Liquid Chromatography (Waters 600 HPLC) with Fluorescence Detector (Waters 474 Scanning Fluorescence Detector) and UV Detector (Waters 2487 Dual ÷ Absorbance Detector); PAH separation will be performed using a reversed phase column with an acetonitrile/water gradient (Marsili et al., 1997).

PBDEs: To detect PBDE, sample extracts will be analyzed on a GC/MS system (HP 6890 gas chromatograph coupled to an HP 5973 low-resolution mass spectrometer) using both EI and negative-chemical ionization (NCI) on an HP-5MS (5% phenyl methyl siloxane) capillary column, according to Pettersson et al. (2004).

Emerging contaminants (Pthalates): Several analytical methods have been developed for determining phthalates by GC (Feng et al., 2005), and HPLC (De Orsi et al., 2006) using different preconcentration techniques (Li et al., 2008).

3.3 Markers of exposure

A set of biomarkers of exposure to anthropogenic contaminants can be investigated in cetaceans skin biopsies (Fig.1). Particular attention will be paid to:

Cytochrome P450 (CYP) - Cytochrome P450 is the most important metabolic/detoxifying enzyme system in mammals. It is substrate-inducible and substrate-specific. CYP1A and CYP2B have been detected in cetacean skin and the induction of these isoforms was found after exposure to lipophilic contaminants both *in vitro* and in field studies (Fossi et al., 2006;

Montie et al., 2008; Hooker et al., 2008). Induction of CYP isoforms can thus be considered as a powerful biomarker of exposure. The induction of different CYP enzymes can be investigated by western blot (WB), immunohistochemistry and quantitative real-time PCR (qRT-PCR) (Spinsanti et al., 2006, 2008).

Nuclear receptors (AhR) - The cytosolic aryl hydrocarbon receptor (AhR) is a ligand inducible nuclear transcription factor which can induce the transcription of xenobiotic metabolizing enzymes such as CYP1A. AhR gene expression was found to be significantly induced in skin of chloracne patients subject to long-term exposure to dioxins and dibenzofurans (Tang et al., 2008). Presence and modulation of AhR in cetacean skin samples will be here investigated by qRT-PCR.

3.4 Markers of reproductive alterations

Endocrine Disrupting Chemicals (EDCs) are a structurally diverse group of compounds that may adversely affect the health of humans and wildlife, or their progenies, by interaction with the endocrine system (Colborn, 1998). Many of the known EDCs are estrogenic, affecting particularly reproductive functions. A main mechanism of endocrine disruption occurs through receptor-mediated mechanisms or the cross-talk of these receptors with the AHR. In skin biopsy we can investigate the gene expression of steroid hormone receptors and any possible alterations due to contaminants.

Steroid hormone receptors. Estrogen receptor(s) (ER), androgen receptor(s) (AR) and progesterone receptor(s) (PR) presence can be investigated in biopsy samples of cetaceans by using WB and immunohistochemistry techniques. Gene expression will also be investigated by qRT-PCR. Steroid hormone receptors are known to be induced in various species and tissues by specific contaminants.

Steroid hormones and steroidogenic enzymes. A set of key genes (Hilscherova et al., 2004) involved in steroid hormone synthesis in humans can be investigated in cell cultures and skin biopsy samples by qRT-PCR and WB, to explore the expression and its potential alterations due to toxicants. We will explore the up/down-regulation of some genes including: CYP11A (cholesterol side-chain cleavage), CYP11B1 (steroid 11 β -hydroxylase), CYP11B2 (aldosterone synthetase), CYP17 (steroid 17 α HSD1-hydroxylase), CYP19 (aromatase) and CYP21B2 (steroid 21-hydroxylase).

3.5 Markers of genotoxicity

DNA integrity is essential for the correct transmission of genetic information to the next generation. Chemicals and/or physical agents that have the potential to cause loss of the DNA's structural and functional integrity are defined genotoxics (Shugart, 2000). Following failing of the repair mechanism, structural changes in DNA become irreversible (apoptosis, chromosome aberrations) and the cells may exhibit altered functions, from uncontrolled proliferation cells to carcinogenesis (Lee & Steinert, 2003). Finally the damage caused by genotoxic compounds leads to DNA mutations and possible alterations in subsequent generations (Gil & Pla, 2001). In this chapter we will present how to evaluate the presence of DNA damage and the efficiency of cells repair mechanism by comet assay. Diffusion assay will be used to evaluate apoptosis in cells.

The Comet Assay or single cell gel electrophoresis (SCGE) is a sensitive, rapid and inexpensive method for measuring DNA damage in individual cells. As reported by Frenzilli et al. (1999) the single cells are embedded in agarose on microscope slides, lysed to

remove the majority of the proteins, electrophoresed, then stained with Sybr Safe in order to visualize the DNA. As result, the DNA of undamaged cells appears as a spherical mass occupying the cavity formed by the lysed cell. On the contrary, cells with DNA damage form a comet image, due to the amount of DNA fragments: smaller is the fragment size, greater is their number and the percentage of DNA that migrates in an electric field. The assay can be performed to examine DNA single strand breaks or to measure double strand breaks in individual cells.

Markers associated with apoptosis: to detect the apoptotic cells, a comet assay modified protocol will be used (Singh et al., 1988). This test, as the comet assay, is relatively easy to apply to most eukaryotic cell types, it is sensitive for detecting low levels of DNA damage or multiple classes of DNA damage and require a small cells number (Dhawan et al., 2009).

Lipid peroxidation: it involves a set of chain reactions by ROS because of their double bonds. Lipid peroxidation is the result of interactions of lipidic radicals and/or formation of non radical species by lipid proxy radicals. The resulting LOOH (lipid hydroperoxide) can easily decomposes into several reactive species, including lipid alkoxyl radicals (LO), aldehydes, lipid epoxides and alcohols. Most of these products are toxic and active mutagens. Peroxidised membranes become rigid and lose permeability and integrity (Valavanidis et al., 2006). Lipid peroxidation will be investigated in skin biopsies by measuring levels of MDA using a spectrophotometric test.

3.6 Diagnostic markers of immunosuppression

Mammal skin is an important immune organ, displaying various defence mechanisms which can be divided into three major functional compartments: epithelial defence, innate-inflammatory immunity and adaptive immunity (Meyer et al., 2007). In this biological material we will develop and validate a set of biomarkers in cultured fibroblasts and skin biopsies to monitor the immunological status of cetaceans. The Major Histocompatibility Complex (MHC) is a set of molecules displayed on cell surfaces that are responsible for lymphocyte recognition and "antigen presentation". These molecules control the immune response through recognition of "self" and "non-self" and are of two types: Class I and Class II molecules. In the Class I molecules there is a family of polymorphic genes, named MIC genes, where MICA and MICB are functional. MIC genes are mainly expressed in endothelial cells and fibroblasts. MIC molecules are considered to be stress-induced antigens that are recognized by cytotoxic T cells and natural killer (NK) cells. The evaluation of MICA protein expression in cetaceans can be used to evaluate the status of the immune system of the different species and will be investigated by WB, immunohistochemistry, qRT-PCR and indirect immunofluorescence (Marsili et al., 2008).

3.7 Diagnostic markers of general stress

Cetaceans are exposed to a diverse array of multiple stressors and here we propose a suite of skin biomarkers to assess general stress.

Heat-shock proteins (HSPs): A number of environmentally and chemical agents are known to induce a set of cellular stress proteins, the heat-shock proteins (HSPs) (Nover, 1991; Nover & Scharf, 1991). Their amount increases rapidly when cells are subjected or exposed to a wide variety of stresses such as heat, xenobiotics or drugs; pathological stimuli such as viral, bacterial and parasitic infections, inflammation and autoimmunity (Wu & Tanguay, 2006). In skin biopsy we can explore the expression of HSPs by WB and qRT-PCR (Cao et al., 1999; Rossner et al., 2003).

Cortisol alteration: Cortisol is mainly involved in carbohydrate, lipid, and protein metabolism and serves as an indicator of an animal's state of well-being, as its levels increase during times of distress ("stress hormone"). We can measure levels of cortisol in the blubber biopsies of cetaceans by enzyme immuno-assays. In fact adipocytes are equipped with a functional G6PT-hexose-6-phosphate dehydrogenase-11 β -hydroxysteroid dehydrogenase type 1 system that is a potential target for modulating local glucocorticoid activation (Marcolongo et al., 2007).

3.8 Diagnostic markers of feeding ecology

Assessing the diet and trophic position of cetaceans is important for understanding the ecology of marine food webs. Decreased quantity and quality of prey, due mainly to over-fishing, have been cited as factors promoting population decline of Mediterranean cetaceans (Bearzi et al., 2006). Feeding ecology is a fundamental aspect in the understanding, management and conservation of free-ranging marine mammals. Nutritional status of marine mammals is a factor that can limit reproductive output and thus population growth. However, foraging of large cetaceans can be difficult to assess when direct observation is not possible, and the use of chemical feeding ecology (i.e., stable isotope and fatty acid analysis) has become increasingly important (Dehn et al., 2006). Stable isotopes of carbon and nitrogen, evaluated using Isotope Ratio Mass Spectrometer (Herman et al., 2005), have been established as powerful tools in animal ecology. They occur naturally, and nitrogen isotope ratios of prey are reflected in tissues of the consumer, with slight enrichment occurring at each trophic step (Kelly, 2000). Stable carbon isotopes are generally used to provide information on spatial habitat use and carbon sources rather than trophic relationships as they enrich in consumer tissues only to a minor degree (Burton & Koch, 1999). An increase of $\delta^{15}\text{N}$ values has been shown in specimens in particular conditions such as nutritional stress (Fuller et al., 2005) or disease (Katzenberg & Lovell 1999). Thus, measurements of the fatty acid composition of the blubber of marine mammals, evaluated using quadruple gas chromatography/mass spectrometry (Krahn, 2004), allow the identification of the diet of these animals. The differences either between individuals or between populations (Møller et al., 2003) or between species (Borobia et al., 1995), are a direct function of their prey fatty acid composition.

3.9 Diagnostic markers for genetic erosion and population studies

Accurate screening of current overall genetic variability of Mediterranean cetaceans and its constant monitoring over the years would identify potential environmental threats and interspecies differences in susceptibility to contaminants.

DNA microsatellites: DNA microsatellites are short (10-50 copies) tandem repeats of mono- to hexa-nucleotide units, assumed to be randomly distributed throughout nDNA, cpDNA and mtDNA. SSRs can appear as simple, interrupted and compound repeats, and are usually distributed across non-coding (and, less often, coding) regions of the genome. Irrespective of hypotheses about their evolutionary mechanisms (slipped-strand mispairing (slippage) error during DNA replication of a single DNA double helix or recombination), SSRs have been applied as a highly variable and powerful genetic marker in areas ranging from ancient and forensic DNA studies to population and conservation genetics (Jarne and Lagoda, 1996). Nucleotide sequence analysis of the genes encoded in the mitochondrial genome (mtDNA) constitutes an additional powerful tool of modern phylogeography and molecular population genetics (Bickham et al., 2000; Gaspari et al., 2007). mtDNA is located

outside the nucleus and it encodes for a subset of the proteins necessary for the mitochondrion itself. The number of copies of mitochondrial genes greatly outnumbers that of nuclear encoded genes. The mitochondrial genome is inherited solely through the maternal line in a clonal fashion and usually does not undergo recombination. Moreover, its rapid rate of base substitution makes the mtDNA genome an ideal marker for establishing levels of variability between and among populations or species (Natoli et al., 2004, 2005).

4. Section three - The case study of the multi-trial diagnostic tool in fin whale (*Balaenoptera physalus*) skin biopsies of the Pelagos Sanctuary (Mediterranean Sea) and the Gulf of California (Mexico)

In this section we describe the application of the proposed multidisciplinary approach to explore the effects of multiple human pressures on cetacean health status (field applications), comparing *Balaenoptera physalus* cetacean population of Mediterranean with the Sea of Cortez population. As previously mentioned (section one) over the past decades, there has been a growing concern regarding the potential threat to Mediterranean cetaceans from persistent organic pollutants such as organochlorine compounds (OCs) (Fossi et al., 2006) and polybrominated diphenyl ethers (PBDEs). Cetaceans of the Gulf of California (Sea of Cortez – Mexico) are reputed to be less exposed to anthropogenic pressure. To date, OC concentrations have been investigated in only three marine mammal species from the Gulf of California (Niño-Torres et al., 2009). The main objective of this case study (Fossi et al. 2010) was to develop and apply a set of sensitive non-lethal diagnostic biomarkers to skin biopsies of fin whales to evaluate the toxicological status of this mysticete in the Pelagos Sanctuary (Ligurian, Corsica and North Tyrrhenian Seas) and in the Gulf of California. We propose a “multi-trial diagnostic tool”, combining molecular biomarkers and gene expression with the analysis of OCs, PAHs and PBDEs.

Mediterranean fin whale – The fin whale is the only mysticete that is regularly found in the Mediterranean Sea (Bérubé et al., 1998), facing a number of anthropogenic threats, such as chemical and acoustic pollution, entanglement in fishing gear and disturbance and collisions from commercial and pleasure boats.

Gulf of California fin whale – Fin whales are permanent residents of the Gulf of California. This population of approximately 610 animals (Urbán-Ramírez et al., 2005) is considered one of the most isolated in the world (Bérubé et al., 1998), and it constitutes a unique and separate conservation unit vulnerable to anthropogenic effects. Although the Gulf of California is considered one of the most pristine and bio-diverse areas of the world (hosting 36 species of marine mammals), increasing human activity is beginning to affect it.

4.1 Experimental design

4.1.1 Sampling

Integument biopsies were obtained from free-ranging fin whales in the Pelagos Sanctuary (n = 12, 6 males and 6 females) and the Gulf of California (n = 5, 3 males and 2 females) during the summer of 2008, using biopsy darts launched with a crossbow (CITES Nat. IT 025IS, Int. CITES IT 007). Sex was determined according to Bérubé and Palsbøll (1996).

4.1.2 Skin biopsy as diagnostic tool

To validate this “multi-trial diagnostic tool”, a two-phase experimental protocol was followed (Fig. 2). In the first phase of the project (field studies), we applied a multi-

disciplinary methodology to explore the effects of the exposure of Mediterranean and Mexican fin whales to anthropogenic contaminants, using skin biopsies as a diagnostic tool and combining the analysis of molecular biomarkers (Western Blot (WB) of CYP1A1 and CYP2B) and gene expression (qRT-PCR of HSP70, ERa, AHR, E2F-1) with the analysis of OC, PAH and PBDE residues in subcutaneous blubber. In the second phase (in vitro experiments), whale biopsy slices were treated with mixtures of OCs, as an innovative tool for the study of intra-species sensitivity to various classes of environmental contaminants.

Fig. 2. Experimental design and sampling areas – validation of a “multi-trial diagnostic tool” to evaluate the toxicological status of fin whales in the Pelagos Sanctuary (Mediterranean Sea) and the Gulf of California (Sea of Cortez – Mexico). A two-phase experimental protocol was followed: first phase: field studies; second phase: in vitro experiments (Fossi et al, 2010).

4.1.3 Results

The two populations of fin whales (first phase results) showed differences in contaminant levels and biomarker responses. Higher levels of PCBs, DDTs, OCs-EDCs and PAHs were found in both male and female (PAHs $p < 0.1$) Mediterranean fin whales in comparison to the Cortez specimens (Fig. 3a-d), confirming the high toxicological stress to which the fin whale population in the Pelagos Sanctuary is exposed.

Levels of low-brominated PBDEs were higher in samples from the Sea of Cortez (Fig. 3e), ranging from 282 to 30,506 ng/g dw, while samples from the Mediterranean sea showed lower average levels. The most abundant congener was PBDE 47. In general, samples from the Sea of Cortez had a major number of detected congeners, such as 47, 100, 99, 154 and 153.

Fig. 3. First phase results – contaminant levels (PCBs (A), DDTs (B), OCs-EDCs (C), PAHs (D), PBDEs (E)) and biomarker responses WB of CYP1A1 (F), CYP2B (G), gene expression (qRT-PCR) of ER α (H), E2F-1 (I), HSP70 (J), AHR (K) in skin biopsies of specimens from the two populations of fin whales (*= $p < 0.1$; **= $p < 0.05$). Pelagos Sanctuary (n = 12, males = 6; females = 6); Sea of Cortez (n = 5, males = 3; females = 2). (Fossi et al, 2010).

Exploring molecular biomarker responses, the induction of CYP1A1 in Mediterranean male whales, if compared to males from the Gulf of California (Fig. 3f), can be related to the presence of high levels of planar compounds, such as coplanar PCBs, and PAHs (Fig. 3a and 3d). A statistically significant positive correlation (ρ Spearman = 0.73, p = 0.003) was found between total PAHs and CYP1A1 induction in male (n = 11) specimens.

A lack of CYP2B induction, despite high levels of lipophilic contaminants, was evident in both male and female Mediterranean whales, even though the differences between the Mediterranean and Mexican specimens are not statistically significant (Fig. 3g). On the other hand, a preliminary warning signal is represented by the high induction of CYP2B in the Mexican fin whales (Fig. 2g), but further investigations are needed.

Exploring gene expression biomarker responses (Fig. 2h–2k), the E_R and E2F-1 genes were up-regulated in the specimens from the Pelagos Sanctuary with respect to those from Mexico for both males (E_R: 3.6_-fold; E2F-1: 1.7_-fold) and females (E_R: 1.3_-fold; E2F-1: 2.4_-fold) (p < 0.05). These data suggest, in the first case (E_R), high exposure to EDC compounds such as OCs-EDCs and, in the second case (E2F-1), the presence of apoptosis processes as a sign of toxicological stress in the Mediterranean population. In contrast, the expression of HSP70 is higher in the Mexican male specimens than that of the Pelagos Sanctuary (1.5_-fold), whereas the Pelagos females exhibit an overexpression of the HSP70 gene with respect to the Mexican specimens (1.9_-fold). The AHR gene is slightly up-regulated in the Mexican specimens (p < 0.05 in males).

The results of the second phase (in vitro tests) show marked differences in CYP1A1 and CYP2B induction by OCs in the whale biopsy slices of the two populations (male specimens), with higher sensitivity responses in the Mexican mysticetes. A dose-dependent induction of CYP1A1 was detected only in biopsy slices from Cortez specimens (0.01 μ g/ml = 2.6-fold, 0.1 μ g/ml = 3.6-fold and 1 μ g/ml = 4.4-fold with respect to the control). The in vitro tests showed no induction of CYP1A1 and CYP2B for the male Mediterranean whales (slices).

In conclusion, this “multi-trial diagnostic tool”, applied to skin biopsies, underlined differences in OC, OC-EDC, PBDE and PAH levels and molecular and gene expression biomarker responses between the two populations. The presence of a higher “toxicological stress” in the Pelagos population is highlighted by warning signals such as CYP1A1 induction and the up-regulation of E_R and E2F-1 genes, combined with a lack of CYP2B induction in both field and in vitro experiments. Moreover, particular concern arises from the high levels of low-brominated PBDEs found in the Mexican whale specimens. Future development of this methodology could provide a statistical system for obtaining more complete information about the “toxicological stress syndrome” in cetaceans, providing a predictive model for hazards in susceptible areas targeted by increasing tourism, such as the Gulf of California (Fossi et al, 2010).

5. Conclusion

The data reported in this chapter confirm that cetacean skin biopsies are a biological material suitable for the hazard assessment of free-ranging cetaceans. This chapter shows that the use of cetacean skin biopsies, as non-lethal biological material for the hazard assessment, is a powerful procedure to screen a large number of samples, with a minimal disturb to animals. Here integument biopsy is proposed as diagnostic tool for the

comprehensive diagnosis of multiple stress factors, health status and genetic population variability. The methods proposed could be used to:

- support in ranking the threatened levels for cetaceans populations/species;
- as an operative tool, suitable for both scientific and decision making purposes supporting in identification of proper mitigation measures;
- find application on two main EU areas of interest: Convention on Biological Diversity (CBD) (2010-2020) - EU Marine Strategy.

The final outcome of these studies will be the development of a prototype DSS - Decision Supporting System for ranking toxicological risk of cetacean populations/species based on integration and analysis of data from the above diagnostic approach. Such a scheme is currently not available to the scientific community nor to policy makers worldwide, neither in the terrestrial nor marine environment. In case of success these achievements are susceptible to be applied on a wider-scale to other marine threatened species.

6. Acknowledgements

The authors particularly thanks the precious collaborators Silvia Maltese, Cristina Panti, Giacomo Spinsanti, Silvia Casini, and Daniele Coppola. The projects described in this chapter were partially supported by the Italian and Mexican Ministry of Foreign Affair and the Italian Ministry of Environment.

7. References

- Alaee, M., Arias, P., Sjodin, A. & Bergman, Å. (2003). An overview of commercially used brominated flame retardants, their applications, their use patterns in different countries/regions and possible modes of release. *Environment International*, Vol. 29, No. 6, pp. 683-689, ISSN 0160-4120.
- Barnes, D.K.A., Galgani, F., Thompson, R.C. & Barlaz, M. (2009). Accumulation and fragmentation of plastic debris in global environments. *Philosophical Transactions of the Royal Society B (The Royal Society)*, Vol. 364, No. 1526, pp. 1985-1998, ISSN 0962-8436.
- Bearzi, G., Politi, E., Agazzi, S. & Azzellino, A. (2006). Prey depletion caused by overfishing and the decline of marine megafauna in eastern Ionian Sea coastal waters (central Mediterranean). *Biological Conservation*, Vol. 127, No. 4, pp. 373-382, ISSN 0006-3207.
- Beckmen, K.B., Blake, J.E., Ylitalo, G.M., Stott, J.L. & O'Hara, T.M. (2003). Organochlorine contaminant exposure and associations with hematological and humoral immune functional assays with dam age as a factor in free-ranging northern fur seal pups (*Callorhinus ursinus*). *Marine Pollution Bulletin*, Vol. 46, No. 5, pp. 594-606, ISSN 0025-326X.
- Berubè, M. & Palsboll, P. (1996). Identification of sex in cetaceans by multiplexing with three ZFX and ZFY specific primers. *Molecular Ecology*, Vol. 5, pp. 283-287, ISSN: 0962-1083.
- Bérubé, M., Aguilar, A., Dendanto, D., Larsen, F., Notarbartolo di Sciara, G., Sears, R., Sigurjonsson, J., Urbán, R. & Palsbøll, P. (1998). Population genetic structure of North Atlantic, Mediterranean Sea and Sea of Cortez fin whales, *Balaenoptera*

- physalus* (Linnaeus, 1758); analysis of mitochondrial and nuclear loci. *Molecular Ecology*, Vol. 7, pp. 585-599, ISSN: 0962-1083.
- Bickham, J.W., Sandhu, S., Hebert, P.D., Chikhi, L. & Athwal, R. (2000). Effects of chemical contaminants on genetic diversity in natural populations: implications for biomonitoring and ecotoxicology. *Mutation Research*, Vol. 463, No. 1, pp. 33-51, ISSN 0027-5107.
- Borobia, M., Gearing, P.J., Simard, Y., Gearing, J.N. & Beland, P. (1995). Blubber fatty acids of finback and humpback whales from the Gulf of St. Lawrence. *Marine Biology*, Vol.122, pp. 341-353, ISSN: 0025-3162.
- Burton, R.K. & Koch, P.L. (1999). Isotopic tracking of foraging and long-distance migration in northeastern Pacific pinnipeds. *Oecologia*, Vol.119, pp. 578-585, ISSN 0029-8549.
- Cao, Y., Ohwatari, N., Matsumoto, T., Kosaka, M., Ohtsuru, A. & Yamashita, S. (1999). TGF-beta1 mediates 70-kDa heat shock protein induction due to ultraviolet irradiation in human skin fibroblasts. *Pflügers Archiv: European Journal of Physiology*, Vol. 438, No. 3, pp. 239-244, ISSN 0031-6768.
- Colborn, T., Smolen, M.J. & Rolland, R. (1998). Environmental neurotoxic effects: the search for new protocols in functional teratology. *Toxicology and Industrial Health*, Vol.14, pp. 9-23, ISSN 1477-0393.
- De Orsi, D., Gagliardi, L., Porrà, R., Berri, S., Chimenti, P., Granese, A. Carpani, I. & Tonelli, D. (2006). An environmentally friendly reversed-phase liquid chromatography method for phthalates determination in nail cosmetics. *Analytica Chimica Acta*, Vol. 555, pp. 238-241, ISSN 0003-2670
- Dehn, L.A., Follmann, E. H., Rosa, C., Duffy, L. K., Thomas, D. L., Bratton, G. R., Taylor, R. J. & O'Hara, T. M. (2006). Stable isotope and trace element status of subsistence-hunted bowhead and beluga whales in Alaska and gray whales in Chukotka. *Marine Pollution Bulletin*, Vol.52, pp. 301-319, ISSN 0025-326X.
- Dhawan, A., Bajpayee, M. & Parmar, D. (2009). Comet assay: a reliable tool for the assessment of DNA damage in different models. *Cell Biology And Toxicology*, Vol. 25, No. 1, pp. 5-32, ISSN 0742-2091.
- Di Guardo, G., Marruchella, G., Agrimi, U. & Kennedy, S. (2005). Morbillivirus infections in aquatic mammals: a brief overview. *Journal of Veterinary Medicine Series A-Physiology Pathology Clinical Medic*, Vol. 52, No. 2, pp. 88-93, ISSN 0931-184X.
- Feng, Y.L., Zhua, J.P. & Robert, S. (2005). Development of a headspace solid-phase microextraction method combined with gas chromatography mass spectrometry for the determination of phthalate esters in cow milk. *Analytica Chimica Acta*, Vol.538, pp. 41-48, ISSN 0003-2670.
- Fossi, M.C., Marsili, L., Junin, M., Castello, H., Lorenzani, J.A., Casini, S., Savelli, C. & Leonzio, C. (1997). Use of nondestructive biomarkers and residue analysis to assess the health status of endangered species of pinnipeds in the southwest Atlantic. *Marine Pollution Bulletin*, Vol. 34, No. 3, pp. 157-162, ISSN 0025-326X.
- Fossi, M.C., Marsili, L., Neri, G., Casini, S., Bearzi, G., Politi, E., Zanardelli, M. & Panigada, S. (2000). Skin biopsy of Mediterranean cetaceans for the investigation of

- interspecies susceptibility to xenobiotic contaminants. *Marine Environmental Research*, Vol. 50, No. 1-5, pp. 517-521, ISSN 0141-1136.
- Fossi, M.C. & Marsili, L. (2003). Endocrine disrupters in aquatic mammals. *Pure and Applied Chemistry*, Vol. 75, No. 11-12, pp. 2235-2247, ISSN 0033-4545.
- Fossi, M.C., Marsili, L., Neri, G., Natoli, A., Politi, E. & Panigada, S. (2003). The use of non-lethal tool for evaluating toxicological hazard of organochlorine contaminants in Mediterranean cetaceans: new data ten years after the first paper published in MPB. *Marine Pollution Bulletin*, Vol. 46, pp. 964-971, ISSN 0025-326X.
- Fossi, M.C., Marsili, L., Casini, S. & Bucalossi, D. (2006). Development of new-tools to investigate toxicological hazard due to endocrine disruptor organochlorines and emerging contaminants in Mediterranean cetaceans. *Marine Environmental Research*, Vol. 62, pp. 200-204, ISSN 0141-1136.
- Fossi, M. C., Urban, J., Casini, S., Maltese, S., Spinsanti, G., Panti, C., Porcelloni, S., Panigada, S., Lauriano, G., Nino-Torres, C., Rojas-Bracho, L., Jimenez, B., Munoz-Arnanz, J. & Marsili, L. (2010). A Multi-Trial diagnostic tool in fin whale (*Balaenoptera physalus*) skin biopsies of the Pelagos Sanctuary (Mediterranean Sea) and the Gulf of California (Mexico). *Marine Environmental Research*, Vol. 69, No. 1, pp. 17-20, ISSN 0141-1136.
- Frantzis, A. (1998). Does acoustic testing strand whales? *Nature*, Vol. 392, pp. 29, ISSN 0028-0836.
- Frenzilli, G., Scarcelli, V., Taddei, F. & Nigro, M. (1999). Adaption of SCGE as a candidate for monitoring marine ecosystems. *Neoplasma*, Vol. 46, pp. 6-7, ISSN 0028-2685.
- Fuller, B.T., Fuller, J.L., Sage, N.E., Harris, D.A., O'Connell, T.C. & Hedges, R.E.M. (2005). Nitrogen balance and $\delta^{15}\text{N}$: why you're not what you eat during nutritional stress. *Rapid Communications in Mass Spectrometry*, Vol. 19, pp. 2481-2728, ISSN 0951-4198.
- Garcia Martinez, J., Moya, A., Raga, J.A. & Latorre, A. (1999). Genetic differentiation in the striped dolphin *Stenella coeruleoalba* from European waters according to mitochondrial DNA (mtDNA) restriction analysis. *Molecular Ecology*, Vol. 8, pp. 1069-1073, ISSN 0962-1083.
- Gaspari, S., Azzellino, A., Airoidi, S. & Hoelzel, A.R. (2007). Social kin associations and genetic structuring of striped dolphin populations (*Stenella coeruleoalba*) in the Mediterranean Sea. *Molecular Ecology*, Vol. 16, No. 14, pp. 2922-2933, ISSN 0962-1083.
- Gil, F. & Pla, A. (2001). Biomarkers as biological indicators of xenobiotic exposure. *Journal of Applied Toxicology*, Vol. 21, pp. 245-255, ISSN 0260-437X.
- Hammond, J.A., Hall, A.J. & Dyrynda, L. (2005). Comparison of polychlorinated biphenyl (PCB) induced effects on innate immune functions in harbour and grey seals. *Aquatic Toxicology*, Vol. 74, No. 2, pp. 126-138, ISSN 0166-445X.
- Haney, T.A., De Almeida, A.O. & Reis, M.S.S. (2004). A new species of Cyamid (Crustacea: Amphipoda) from a stranded cetacean in Southern Bahia, Brazil. *Bulletin of Marine Science*, Vol. 75, No. 3, pp. 409-421, ISSN 0007-4977.

- Herman, D.P., Burrows, D.G., Wade, P.R., Durban, J.W., LeDuc, R.G., Matkin, C.O. & Krahn, M.M. (2005). Feeding ecology of eastern North Pacific killer whales from fatty acid, stable isotope, and organochlorine analyses of blubber biopsies. *Marine Ecology. Progress Series*, Vol. 302, pp. 275-291, ISSN 0171-8630.
- Hilscherová, K., Jones, P.D., Gracia, T., Newsted, J.L., Zhang, X., Sanderson, Yu, R.M.K., Wu, R.S.S. & Giesy, J.P. (2004). Assessment of the effects of chemicals on the expression of ten steroidogenic genes in the H295R cell line using real-time PCR. *Toxicological Sciences*, Vol.81, pp. 78-89, ISSN 1096-6080.
- Hoelzel, A.R. (1998). Genetic structure of cetacean populations in sympatry, parapatry and mixed assemblages; implications for conservation policy. *Journal of Heredity*, Vol. 89, pp. 451-458, ISSN 0022-1503.
- Hooker, S.K., Metcalfe, T.L., Metcalfe, C.D., Angell, C.M., Wilson, J.Y., Moore, M.J. & Whitehead, H. (2008). Changes in persistent contaminant concentration and CYP1A1 protein expression in biopsy samples from northern bottlenose whales, *Hyperoodon ampullatus*, following the onset of nearby oil and gas development. *Environmental Pollution*, Vol. 152, No. 1, pp. 205-216, ISSN 0269-7491.
- Jarne, P. & Lagoda, P.J.L. (1996). Microsatellites, from molecules to populations and back. *Trends in Ecology and Evolution*, Vol. 11, pp. 424-429, ISSN 0169-5347.
- Katzenberg, M.A. & Lovell, N.C. (1999). Stable isotope variation in pathological bone. *International Journal of Osteoarchaeology*, Vol.9, pp.316-324, ISSN 1047-482X.
- Kelly, J.F. (2000). Stable isotopes of carbon and nitrogen in the study of avian and mammalian trophic ecology. *Canadian Journal of Zoology*, Vol.78, pp. 1-27, ISSN 0008-4301.
- Krahn, M.M., Herman, D.P., Ylitalo, G.M., Sloan, C.A., Burrows, D.G., Hobbs, R.C., Mahoney, B.A., Yanagida, G.K., Calambokidis, J. & Moore, S.E. (2004). Stratification of lipids, fatty acids and organochlorine contaminants in blubber of white whales and killer whales. *Journal of Cetacean Research and Management*, Vol.6, pp. 175-189, ISSN 1561-0713.
- Lee, R.F. & Steinert, S. (2003). Use of the single cell gel eletrophoresis/comet assay for detecting DNA damage in aquatic (marine and freshwater) animals. *Mutation Research*, Vol. 544, pp. 43-64, ISSN 0921-8262.
- Li, L., Bozec, A., Somot, S., Beranger, K., Bouruet-Aubertot, P., Sevault, F. & Crepon, M. (2006). Chapter 7 Regional atmospheric, marine processes and climate modelling, In: P. Lionello, P. Malanotte-Rizzoli & R. Boscolo, (Eds), *Developments in Earth and Environmental Sciences*, Elsevier, Vol. 4, Mediterranean, pp. 373-397, ISSN 1571-9197, ISBN 9780444521705.
- Li, J., Can, Y., Shia, Y., Moua, S. & Jiang, G. (2008). Analysis of phthalates via HPLC-UV in environmental water samples after concentration by solid-phase extraction using ionic liquid mixed hemimicelles. *Talanta*, Vol.74, No.4, pp. 498-504, ISSN 0039-9140.
- Marcolongo, P., Piccirella, S., Senesi, S., Wunderlich, L., Gerin, I., Mandl, J., Fulceri, R., Bánhegyi, G. & Benedetti, A. (2007). The glucose-6-phosphate transporter-hexose-6-phosphate dehydrogenase-11 β -hydroxysteroid dehydrogenase type 1 system of the adipose tissue. *Endocrinology*, Vol. 148, No. 5, pp. 2487-2495.

- Marsili, L. & Focardi, S. (1996). Organochlorine levels in subcutaneous blubber biopsies of fin whales (*Balaenoptera physalus*) and striped dolphins (*Stenella coeruleoalba*) from the Mediterranean Sea. *Environmental Pollution*, Vol. 91, No. 1, pp. 1-9.
- Marsili, L., Fossi, M.C., Casini, S., Savelli, C., Jimenez, B., Junin, M. & Castello, H. (1997). Fingerprint of polycyclic aromatic hydrocarbons in two populations of southern sea lions (*Otaria flavescens*). *Chemosphere*, Vol. 34, No. 4, pp. 759-770, ISSN 0045-6535.
- Marsili, L., Fossi, M.C., Bucalossi, D., Porcelloni, S., Maltese, S. & Casini, S. (2008). Use of immunofluorescence technique in cultured fibroblasts from Mediterranean cetaceans as new “in vitro” tool to investigate effects of environmental contaminants. *Marine Environmental Research*, Vol. 66, No. 1, pp. 151-153, ISSN 0141-1136.
- Meyer, T., Stockfleth, E. & Christophers, E. (2007). Immune response profiles in human skin. *British Journal of Dermatology*, Vol. 157, No. 2, pp. 1-7, ISSN 1365-2133.
- Møller, P., Born, E.W., Dietz, R., Ruzzante, D., Haug, T. & Øien, N. (2003). Differences in fatty acid composition of blubber in minke whales (*Balaenoptera acutorostrata*) from Greenland and the NE Atlantic. *Journal of Cetacean Research and Management*, Vol.5, n.2, pp.115-124, ISSN 1561-0713.
- Montie, E.W., Fair, P.A., Bossart, G.D., Mitchum, G.B., Houde, M., Muir, D.C.G., Letcher, R.J., McFee, W.E., Starczak, V.R., Stegeman, J.J. & Hahn, M.E. (2008). Cytochrome P4501A1 expression, polychlorinated biphenyls and hydroxylated metabolites, and adipocyte size of bottlenose dolphins from Southeast United States. *Aquatic Toxicology*, Vol. 86, pp. 397-412, ISSN 0166-445X.
- Natoli, A., Peddemors, V.M. & Hoelzel, A.R. (2004). Population structure and speciation in the genus *Tursiops* based on microsatellite and mitochondrial DNA analyses. *Journal of Evolutionary Biology*, Vol. 17, No. 2, pp. 363-375, ISSN 1420-9101.
- Natoli, A., Birkun, A., Aguilar, A., Lopez, A. & Hoelzel, A.R. (2005). Habitat structure and the dispersal of male and female bottlenose dolphins (*Tursiops truncatus*). *Proceedings. Biological Sciences / The Royal Society*, Vol. 272, No. 1569, pp. 1217-1226, ISSN 0962-8452.
- Nover, L. (1991). Inducers of hsp synthesis: heat shock and chemical stressors. In: *Heat Shock Response*, Nover L. (Ed.), CRC Press, Boca Raton, FL, pp. 5-40.
- Nover, L. & Scharf, K.D. (1991). Heat shock proteins. In: *Heat Shock Response*, Nover L. (Ed.), CRC Press, Boca Raton, FL, pp. 41-128.
- Panigada, S., Pesante, G., Zanardelli, M., Capoulade, F., Gannier, A. & Weinrich, M.T. (2006). Mediterranean fin whales at risk from fatal ship strikes. *Marine Pollution Bulletin*, Vol. 52, No. 10, pp. 1287-1298, ISSN 0025-326X.
- Pettersson, A., van Bavel, B., Engwall, M. & Jimenez, B. (2004). Polybrominated diphenylethers and methoxylated tetrabromodiphenylethers in cetaceans from the Mediterranean Sea. *Archives of Environmental Contamination and Toxicology*, Vol. 47, No. 4, pp. 542-550, ISSN 0090-4341.

- Rossner, P.Jr, Binkova, B. & Sram, R.J. (2003). Heat shock proteins hsp32 and hsp70 as biomarkers of an early response? *Mutation Research*, Vol. 542, pp. 105 --116, ISSN 0921-8262.
- Shugart, L.R. (2000). DNA damage as a biomarker of exposure. *Ecotoxicology*, Vol. 9, pp. 329-340, ISSN 0963-9292.
- Simmonds, M. & Nunny, L. (2002) Cetacean Habitat Loss and Degradation in the Mediterranean Sea. Accobams secretariat, Monaco, February 2002, section 7, 23 p.
- Singh, N.P., McCoy, M.T., Tice, R.R. & Sch, E.L. (1988). A simple technique for quantitation of low levels of DNA damage in individual cells. *Experimental Cell Research*, Vol. 175, No.1, pp. 184-191, ISSN 0014-4827.
- Spinsanti, G., Panti, C., Lazzeri, E., Marsili, L., Casini, S., Frati, F. & Fossi, M.C. (2006). Selection of reference genes for quantitative RT-PCR in striped dolphin (*Stenella coeruleoalba*) skin biopsies. *BMC Molecular Biology*, Vol. 7, No.32, ISSN 1471-2199.
- Spinsanti, G., Panti, C., Bucalossi, D., Marsili, L., Casini, S., Frati, F. & Fossi, M.C. (2008). Selection of reliable reference genes for qRT-PCR studies on cetacean fibroblast cultures exposed to OCs, PBDEs, and 17 β -estradiol. *Aquatic Toxicology*, Vol. 87, pp. 178-186, ISSN 0166-445X.
- Tanabe, S. & Tatsukawa, R. (1992). Chemical modernization and vulnerability of cetaceans: increasing toxic threat of organochlorine contaminants. In: *Persistent Pollutants in Marine Ecosystems*, C.H. Walker & D.R. Livingstone, (Eds), 161-177, Pergamon Press Ltd, Oxford, ISBN 0-08-041874-0.
- Tang, N.J., Liu, J., Coenraads, P.J., Dong, L., Zhao, L.J., Ma, S.W., Chen, X., Zhang, C.M., Ma, X.M., Wei, W.G., Zhang, P. & Bai, Z.P. (2008). Expression of AhR, CYP1A1, GSTA1, c-fos and TGF- α in skin lesions from dioxin-exposed humans with chloracne. *Toxicology Letters*, Vol. 177, No. 3, pp. 182-187, ISSN 0378-4274.
- Urbán-Ramírez, J., Rojas-Bracho, L., Guerrero-Ruíz, M., Jaramillo-Legorreta, A. & Findley, L.T. (2005). Identification of sex in Cetaceans by multiplexing with ZFX and ZFY specific primer. In: Cartron, J.E., Ceballos, G., Felger, R.S. (Eds.), *Biodiversity Ecosystems and Conservation in Northern Mexico*. Oxford University Press, New York, pp. 276-297, ISBN 9780195348125.
- Valavanidis, A., Vlahogianni, T., Dassenakis, M. & Scoullos, M. (2006). Molecular biomarkers of oxidative stress in aquatic organisms in relation to toxic environmental pollutants. *Ecotoxicology and Environmental Safety*, Vol. 64, pp. 178-189, ISSN 0147-6513.
- Villeneuve, D.L., Khim, J.S., Kannan, K. & Giesy, J.P. (2002). Relative potencies of individual polycyclic aromatic hydrocarbons to induce dioxinlike and estrogenic responses in three cell lines. *Environmental Toxicology*, Vol. 17, pp. 128-137, ISSN 1520-4081.
- Whitehead, A., Anderson, S.L., Kuivila, K.M., Roach, J.L. & May, B. (2003). Genetic variation among interconnected populations of *Catostomus occidentalis*: implications for distinguishing impacts of contaminants from biogeographical structuring. *Molecular Ecology*, Vol. 12, pp. 2817-2833, ISSN 0962-1083.

- Wu, T. & Tanguay, R.M. (2006). Antibodies against heat shock proteins in environmental stresses and diseases: friend or foe? *Cell Stress Chaperones*, Vol. 11, No. 1, pp. 1-12, ISSN 1355-8145.
- Young, N.M. & Iudicello, S. (2007). Worldwide Bycatch of Cetaceans. An evaluation of the most significant threats to cetaceans, the affected species and the geographic areas of high risk, and the recommended actions from various independent institutions. U.S. Dep. Commerce, *NOAA Technical Memorandum*, NMFS-OPR-36 July 2007, 236pp.

Skin Biopsy - Perspectives

Edited by Dr. Uday Khopkar

ISBN 978-953-307-290-6

Hard cover, 336 pages

Publisher InTech

Published online 02, November, 2011

Published in print edition November, 2011

Skin Biopsy - Perspectives is a comprehensive compilation of articles that relate to the technique and applications of skin biopsy in diagnosing skin diseases. While there have been numerous treatises to date on the interpretation or description of skin biopsy findings in various skin diseases, books dedicated entirely to perfecting the technique of skin biopsy have been few and far between. This book is an attempt to bridge this gap. Though the emphasis of this book is on use of this technique in skin diseases in humans, a few articles on skin biopsy in animals have been included to acquaint the reader to the interrelationship of various scientific disciplines. All aspects of the procedure of skin biopsy have been adequately dealt with so as to improve biopsy outcomes for patients, which is the ultimate goal of this work.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Maria Cristina Fossi and Letizia Marsili (2011). Multi-Trial Ecotoxicological Diagnostic Tool in Cetacean Skin Biopsies, Skin Biopsy - Perspectives, Dr. Uday Khopkar (Ed.), ISBN: 978-953-307-290-6, InTech, Available from: <http://www.intechopen.com/books/skin-biopsy-perspectives/multi-trial-ecotoxicological-diagnostic-tool-in-cetacean-skin-biopsies>

INTech
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2011 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](https://creativecommons.org/licenses/by/3.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen