We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Mycoflora and Biodiversity of Black Aspergilli in Vineyard Eco-Systems

Cinzia Oliveri and Vittoria Catara

Università degli Studi di Catania

*Dipartimento di Scienze delle Produzioni Agrarie ed Alimentari, Catania
Italy*

1. Introduction

Environmental conditions in vineyard eco-systems are of particular interest because they can influence the fungal populations associated with grapes, fungal-plant interactions, and production of secondary metabolites, including mycotoxins. Some fungal species are pathogenic to grapevines, infecting the roots, canes, leaves and fruit (Hewitt, 1988; Tournas & Katsoudas, 2005). Grape contamination by different moulds occurs during vineyard pre-harvesting, harvesting and grape processing (Magnoli et al., 2003). Moulds commonly isolated from grapes are *Alternaria*, *Cladosporium* and *B. cinerea*, the latter causing bunch rot. Pathogenic and opportunistic species of *Fusarium*, *Penicillium* and *Aspergillus* can also colonize inducing grape disease.

In heavily infected fruit, moulds alter chemical composition and mould enzymes adversely affect wine flavor and colour as well as yeast growth during alcoholic fermentation (Fleet, 1999; Fleet, 2001). Some vineyard fungal species are capable of producing toxic secondary metabolites (mycotoxins) in infected tissue, which may contaminate grapes and grape products such as wine, grape juice and dried vine fruit (reviewed in Nielsen et al., 2009). The mycotoxins of greatest significance include aflatoxins, citrinin, patulin and ochratoxin A (OTA) and recently fumonisin B₂ (FB₂) (Frisvad et al., 2007; Logrieco et al., 2010; Morgensen et al., 2010a, 2010b; Susca et al., 2010). The most important mycotoxin in grapes and the grape-wine chain is OTA first reported by Zimmerli and Dick (1996). It has nephrotoxic, carcinogenic (2B group) (IARC, 1993), teratogenic and immunotoxic effects (Abarca et al., 2001; Castegnaro & Pfohl-Leszkowicz, 2002, Da Rocha et al., 2002; Pfohl-Leszkowicz et al., 2002; Petzinger & Weidenbach, 2002; Vrabcheva et al., 2000). Thereafter, several authors reported OTA contamination in wine and the presence of OTA-producing fungi in grapes in different wine-growing areas around the world (Battilani et al., 2006; Leong et al., 2007; Medina et al., 2005; Sage et al., 2004; Tjamos et al., 2004). The contamination of grapes with OTA can occur in the field, even without visible symptoms, while the grapes are still on the vine (Serra et al. 2006).

2. Occurrence and biodiversity of black aspergilli from grapes

2.1 Role in Ochratoxin A contamination

Several surveys conducted in the Mediterranean, South America, and Australia reported fungal species belonging to *Aspergillus* section *Nigri* (also known as Black Aspergilli-BA) as

the major responsible for OTA contamination in grape (reviewed in Perrone et al., 2007). Several BA have been isolated from grape or from vineyard soil/air such as *A. niger* aggregate (namely *A. niger* sensu stricto, *A. tubingensis*, *A. foetidus*, and *A. brasiliensis*), *A. carbonarius* and the uniseriate species *A. aculeatus*, *A. japonicus*, and *A. uvarum* (Medina et al., 2005; Perrone et al., 2008). Only a few species produce OTA among the *A. niger* aggregate, *A. carbonarius*, and *A. japonicus* (Battilani et al., 2003a). *A. ochraceus* (belonging to section *Circumdati*), although able to produce OTA, have only occasionally been isolated from grape. The most frequently occurring species are *A. niger* aggregate and *A. carbonarius*, respectively, although the highest percentage of OTA-producing strains has been detected in the latter species (Serra et al., 2005).

OTA contamination of dried vine fruit was also found to be due to black aspergilli in Europe, including Spain, Hungary and other parts of the world such as Argentina and Australia (Varga & Kozakiewicz, 2006). In spite of the higher incidence of species belonging to the *A. niger* aggregate found in vineyards, only 5-10 % of *A. niger* OTA-producing strains were detected, whereas more than 50% and, in some studies up to 100%, in *A. carbonarius* (Battilani et al. 2006; Heenan et al., 1998; Perrone et al., 2006a; Serra et al., 2005). Other *Aspergillus* species, such as *A. helicotrix*, *A. ellipticus* and *A. heteromorphus*, *A. ochraceus* are rare (Bau et al., 2005).

2.2 Isolation and Identification tools

Black aspergilli are isolated and identified at genus and species level by morphological criteria: colour, density and colony appearance (layer colour, wrinkled, umbilical, thick or flat) and microscope observation (conidial head, conidiophore and conidia characters) in accordance with appropriate keys (Klick, 2002; Klick and Pitt, 1988; Pitt & Hocking, 1999, 2009; Samson et al., 2004, 2007). The taxonomy of *Aspergillus* section *Nigri* is widely studied but although identification at section level is quite easy, at species level it is much more complex since morphologically taxa differences are very subtle requiring taxonomic expertise. For macromorphological observations, Czapek yeast autolysate (CYA), malt extract autolysate (MEA), Czapek yeast autolysate with 5% NaCl (CYAS) agar, yeast extract-sucrose (YES) agar, oatmeal agar (OA) and Czapek agar (CZA) are used (Samson et al., 2004). Some differential growth media e.g. DYSG Agar, Coconut Cream Agar (Heenan et al., 1998) and MEA-B (Pollastro et al., 2006) may facilitate the recognition of ochratoxigenic black aspergilli (Samson et al., 2007). Useful physiological features are very good growth and sporulation at 37 °C as well as growth and acid production on CREA agar (Samson et al., 2004). Species can be identified by micromorphological analysis of the fungal structures by light microscopy. Scanning Electron Microscopy (SEM) is helpful for vesicle observation which is necessary for distinguishing between uniseriate (i.e. *A. aculeatus*, *A. japonicus*) and biseriate species (i.e. *A. carbonarius*, *A. ibericus*, *A. niger*) and conidia ornamentation which can distinguish between *A. niger* aggregate, *A. carbonarius* and *A. ibericus* (Serra et al., 2006; Varga et al., 2000) (Table 1).

As for other fungal species studies based on molecular sequence analysis of ribosomal and ubiquitous genes (ITS, IGS, calmodulin, β -tubulin, elongation factor) and polymorphisms by obtained Amplified Fragment Length Polymorphism (AFLP), Random Amplified Polymorphic DNA (RAPD) and microsatellites, have been performed for *Aspergilli* isolated from grapes. These studies have provided useful information on the taxonomy of BA and methods for their detection, identification and monitoring (reviewed in Abarca et al., 2004; Geiser et al., 2007; Niessen et al., 2005; Perrone et al., 2007, 2009; Samson et al., 2007).

Species	Conidiophore	Conidial size (µm)	OTA production
<i>A. japonicus</i> / <i>A. aculeatus</i>	Uniseriate	4–5	Negative
<i>A. niger</i> aggregate	Biseriate	3–5	Positive (low %)
<i>A. sclerotioniger</i>	Biseriate	5–6	Positive
<i>A. carbonarius</i>	Biseriate	7–9	Positive (high %)
<i>A. ibericus</i>	Biseriate	5–7	Negative

Table 1. Some characteristics of the main black *Aspergillus* species (from Serra et al., 2006)

PCR primers for detecting BA target the generic sequences for identifying or characterizing the fungus taxonomy at the intraspecific level or the sequence of genes involved in mycotoxin biosynthesis which are not necessarily able to distinguish fungal species (Bau et al., 2005; Dao et al., 2005; Perrone et al., 2007; Sartori et al., 2006; Schmidt et al., 2003, 2004; Serra et al., 2005).

Species-specific primers based on ITS sequence differences were developed for *A. ellipticus*, *A. heteromorphus*, *A. japonicus*, *A. niger* (Gonzales-Salgado et al., 2005), *A. carbonarius* and *A. ochraceus* (Patiño et al., 2005). Some others were developed for *A. carbonarius* identification based on SCAR primers (Pollastro et al., 2003; Pelegrinelli-Fungaro et al., 2004), whereas PCR-RFLP analysis is necessary to distinguish the strains of *Aspergillus niger* aggregate into two groups: *A. niger* and *A. tubingensis* (N and T type) (Accensi et al., 2001, Gonzalez-Salgado et al., 2005).

More primers have been developed for the genes encoding the polyketide synthases (PKSs) involved in OTA biosynthesis in both *Aspergillus* and *Penicillium* (Ayoub et al., 2010; Atoui et al., 2006; Dao et al., 2005; O’Callaghan et al., 2003).

Recent studies have reported the use of degenerate primers targeting the ketosynthase domain (KS) which identified a new pks gene from *A. carbonarius* (ACpks) (Atoui et al., 2006; Gallo et al., 2009). By screening *Aspergillus* isolates with ACpks specific primers, ACpks homologues appeared to be present in *A. sclerotioniger* and *A. ibericus* which are closely related to *A. carbonarius*.

A duplex real-time PCR assay for simultaneously detecting members of the *Aspergillus niger* aggregate and *A. carbonarius* was developed by López-Mendoza et al. (2009) and Selma et al. (2009). They targeted the beta -ketosynthase and acyl transferase domains of the poliketide synthase of *A. carbonarius* and the *A. niger* aggregate, the assay allowing preferential amplification at greater concentrations providing a fast and accurate tool to monitor, OTA-producing species in grapes in a single reaction. These approaches gave rise to molecular diagnostic assays based on expression profiling and which determine the molecular triggers controlling OTA biosynthesis in *Aspergillus* spp.

More recently real-time/quantitative PCR (qPCR) protocols have detected and quantified ochratoxigenic fungi, developed using constitutive genes (González-Salgado et al., 2009; Morello et al., 2007; Mulè et al., 2006) or genes involved in toxin biosynthesis (Atoui et al., 2007; Schmidt et al., 2004; Selma et al., 2009).

2.3 European biodiversity monitoring

DNA-based fingerprinting techniques such as AFLP, RFLP, RAPD, ap-PCR, and the sequencing of subgenomic DNA fragments have drastically improved the understanding of the occurrence and biodiversity of *Aspergillus* spp. in grapes and vineyards worldwide.

Sequencing techniques were primarily useful at the species identification level, whereas fingerprinting techniques were exploited at the intraspecific level (Ferracin et al., 2009; Dachoupakan et al., 2009; Perrone et al., 2007). Although all these studies contribute to analysing the species composition and genetic diversity of grape mycobiota, no genotypical differences could be established between OTA producers and non producers (Ilic et al., 2001, 2004; Martinez-Culebraz et al., 2009; Chiotta et al., 2011). Moreover, there was no correlation between genotype, the ability to produce OTA and geographical origin (Niessen et al., 2005). Surveys conducted in Europe during the four-year EU project 'Wine-Ochra Risk' (QLK1-CT 2001-01761) indicated a significant correlation between the incidence of grape infected by black aspergilli, as potentially OTA producer, at harvest and in climatic conditions and geography (latitude and longitude); there was increasing incidence from West to East and North to South (Battilani et al., 2006). Aspergilli in vineyards varied depending on years and geographic areas: France, Greece and Israel were the areas with the highest incidence, followed by South Italy, Spain and Portugal (Abarca et al., 2001; Battilani et al., 2006; Guzev et al., 2006; Logrieco et al., 2007; Otteneder & Majerus., 2000; Sage et al., 2002). In countries with colder temperate climates such as Germany, Northern Hungary, the Czech Republic as well as the northern parts of Portugal, France and Italy, BA has not often been isolated from grapes, although sometimes OTA has been detected in wines. The identification of OTA producing *Penicillium* species from grapes in Northern Italy and France suggests they could be responsible for contamination in these regions (Battilani et al., 2001; Rousseau, 2004).

Surveys in 107 vineyards in the Mediterranean basin have identified four main *Aspergillus* populations: *A. carbonarius*, *A. tubingensis*, *A. niger*, and a group of *Aspergillus* 'uniseriate' isolates morphologically indistinguishable from *A. japonicus* and *A. aculeatus*. The latter could be clearly distinguished by molecular tools such as AFLP, RFLP and sequence analyses (Bau et al. 2006; Perrone et al. 2006a, 2006b). Highest genetic variability was observed in the *A. niger* group due to its complexity and the difficulty of identifying it at species level by both AFLP and the sequencing of calmodulin and β -tubulin subgenomic fragments (Perrone et al., 2007). In Australian vineyards, Leong et al. (2007) documented the dominance of *A. niger* over *A. carbonarius* and *A. aculeatus*. Polyphasic studies using macro- and micromorphology, secondary metabolite profiles, partial sequences of β -tubulin, calmodulin and ITS genes, and AFLP analysis led to the description of a new *Aspergillus* species: *A. ibericus*, which is closely related to *A. carbonarius* but unable to produce OTA (Serra et al. 2006) and *A. brasiliensis* belonging to the *A. niger* aggregate (Varga et al., 2007), both isolated from grapes in the Iberian Peninsula; *A. uvarum*, morphologically very similar to *A. japonicus* and *A. aculeatus*, but clearly distinct by the molecular analysis of grapes samples in Portugal, Italy, France, Israel, Greece and Spain.

In Italy, field surveys studied the fungi associated with grapes and their ability to produce OTA in different grape-growing areas (as regards grape variety and farming methods) in the north and south of the country (Battilani et al., 2002, 2006). Analysis of these grape samples revealed that *A. niger* aggregate was the prevalent species and *A. carbonarius* was mostly found in Southern Italy and Sicily (Lucchetta et al., 2010; Oliveri, 2007). *A. carbonarius* was never dominant at different growth stages, or in different geographical areas and years, but it was confirmed as the key fungus because of the high percentage of strong OTA producing isolates in the population.

In sixteen vineyards located in 13 provinces (including Modena, Imola, Ravenna, Brindisi and the warmest places such as Trapani and Ragusa), the effect of geographic area on fungal flora was confirmed, even though a major role was played by meteorological conditions,

both on fungal colonisation and the OTA content in bunches. BA were present in bunches from setting, colonising most berries at early veraison (Battilani et al., 2006). The detection of isolates belonging to the *A. niger* aggregate, *A. carbonarius* and uniseriate varied with growth stage. At setting and berry pea-sized stages, more than 50% of isolates belonged to uniseriate; starting from early veraison, the *A. niger* aggregate became dominant (about 50%) whereas *A. carbonarius* was around 20% from pea-size to harvesting. The region with the highest percentage of grapes berries colonised by the *A. niger* aggregate was Veneto, while the lowest was in central Emilia Romagna. The highest incidence of *A. carbonarius* was detected in Puglia and the lowest in Emilia Romagna and Veneto. The number of OTA-producing strains among BA, isolated in each vineyard at different growth stages, was generally very limited (an encouraging result) (Battilani et al., 2006).

Molecular techniques to investigate strain variation in toxigenic and non-toxigenic black *Aspergillus* spp. showed that isolates of *A. carbonarius* and *A. niger* clustered into species groups, however, within species, strains displaying similar degrees of toxigenicity did not cluster together when characterized by RAPD techniques (Ilic et al., 2001, 2004). The profile of the *Aspergillus terreus* species isolated from dried grapes, analysed by RAPD, indicated great genomic diversity (Narasimhan & Asokan, 2010). Martínez-Culebras et al. (2009) recently carried out a study on ochratoxigenic mycobiota in grapes by ap-PCR sequence analysis of the ITS and IGS regions and their ability to produce OTA. Based on ap-PCR profiles, derived from two microsatellite primers, three main groups were obtained by UPGMA cluster analysis corresponding to *A. carbonarius*, *A. niger* and *A. tubingensis*. The cophenetic correlation values corresponding to ap-PCR UPGMA analysis showed higher genetic variability in *A. niger* and *A. tubingensis* than in *A. carbonarius*. In addition, no genotypical differences could be established between OTA producers and non-producers in all the species analysed. Regarding uniseriate black aspergilli, low divergence was found between *A. aculeatus* and *A. uvarum*. OTA-production seems to be strain related since it was found in different clusters, with either ap-PCR or IGS-ITS phylogenetic analysis.

2.4 Epidemiology in vineyard

Black aspergilli are affected by several factors in the vine environment, i.e., grape status, the number of damaged grape berries, meteorological conditions, vineyard location, the cropping system as well as chemical treatments (Battilani et al., 2003b, 2006; Belli et al., 2005; 2007a; Blesa et al., 2006; Clouvel et al., 2008; Hocking et al., 2007; Leong et al., 2006). Generally fungi have been detected in vineyards and on grapes from setting. However, grape aspergilli increase gradually, reaching their maximum values at the beginning of veraison and ripening (Battilani et al., 2002). As *Aspergillus* species are not considered primary pathogens, various grape damage, such as attack by other fungi or mechanical injury, dramatically increases the risk of fungal infection by these species and OTA contamination (Serra et al., 2006; Belli et al., 2007b). However, grape damage due to insects, birds or other fungal infections, is the primary factor affecting the development of the disease and OTA accumulation in grapes (Cozzi et al., 2006).

Some Australian studies have demonstrated that vineyard soil at a depth of 0–5 cm beneath the vines is the primary reservoir of black aspergilli (Clarke et al., 2003; Kazi et al. 2004; Leong et al. 2006). Concentrations were also higher in the soil directly beneath the vines compared to the inter-row area. It is postulated that air movement deposits spores from the soil onto the grapes berry surfaces, because BA spores in air samples were higher closer to

the soil (Kazi et al., 2003a). Soil temperature could also affect the incidence of *A. carbonarius* in the soil; the optimal temperature for spore survival was around 25 °C, with counts decreasing at 15 °C and 35°C. Survival at 40 °C was poor (Kazi et al., 2003b, 2004).

Agronomic practices and biological and chemical treatments have been found to reduce BA colonization and OTA levels in grapes and grape-derived products (reviewed in Varga & Kozakiewicz, 2006).

3. Biodiversity in a restricted geographical area: A case study from the Mount Etna Slopes

3.1 Vineyard mycobiota

Most studies address the biodiversity of grape mycoflora and above all of BA in countries and in wine producing areas. Recently we performed a two-year survey in a restricted geographical area to assess the population density and biodiversity of the mycoflora associated with the grapes, air and soil of the vineyards on the slopes of Mount Etna (eastern Sicily, Italy) where there is a long tradition of grape cultivation. This area is characterized by a temperate Mediterranean climate, with an average annual rainfall of 800 mm and high day/night temperature fluctuations. Moreover, the area is characterized by old and autochthonous wine-grape cultivars, i.e. Etna rosso DOC (Nerello mascalese, Nerello Cappuccio) and Etna Bianco (Carricante, Catarratto).

Special emphasis was made on toxigenic fungal species, e.g. *Aspergillus* Sect. *Nigri* spp. and *Penicillium* spp. (Oliveri, 2006; Oliveri et al., 2008). It is postulated that air movement deposits soil fungal spores onto grapes, because their incidence in air samples increases closer to the soil. So, healthy grape soil beneath the vines was ecologically monitored and its air was sampled at two different heights at the pea stage, early veraison and ripening using plating methods. Spore-producing filamentous fungi were detected, identified at the genus level, and then the *Aspergillus* and *Penicillium* strains were isolated and identified at the species level (Fig. 1).

Fig. 1. Scanning electron microscopy pictures of a) *A. carbonarius*, *A. niger* b) and c) *A. japonicus* spores, where spore ornamentation differences are clearly seen (bar = 1 µm) (Zeiss, DSM 940).

The most frequent genera isolated from air, soil and grapes by increasing order were *Aspergillus*, *Penicillium*, *Cladosporium* and *Rhizopus*.

Fungi belonging to *Aspergillus* spp. were present in all the sampled matrices from the pea stage and they were predominant to *Penicillium* spp. from early veraison to ripening. Fungi from 6 genera were isolated on grapes at the ripening stage in eight different vineyards (Fig. 2).

Population densities of *Aspergillus* spp. in grape wash water ranged between 4.5×10^3 and 1.2×10^4 cfu ml⁻¹.

The most frequent *Penicillium* species isolated from the vineyard eco-system were *P. chrysogenum*, *P. expansum* and *P.olsonii* (Tab. 2). *P. verrucosum* was isolated from only one soil sample. Among the *Aspergillus* species, the most frequent were from section *Nigri*; the level of contamination by *A. ochraceus* and *A. flavus* was low.

Fig. 2. Population composition (%) detected in 8 vineyards (V1-V8) from 20 grape samples, during the ripening stage and over a 2-year survey.

Genus	Species	n. samples ^a		
		air (224)	Soil (112)	Grapes (160)
<i>Aspergillus</i>	<i>niger</i> aggregate	125	94	128
	<i>carbonarius</i>	43	49	38
	<i>flavus</i>	18	21	1
	<i>ochraceus</i>	8	20	1
<i>Penicillium</i>	<i>aurantiogriseum</i>	4	3	n.d.
	<i>chrysogenum</i>	50	55	n.d.
	<i>expansum</i>	63	22	15
	<i>italicum</i>	29	19	14
	<i>olsonii</i>	n.d.	7	n.d.
	<i>verrucosum</i>	n.d.	1	n.d.

n.d.= not detected
^a in parenthesis the total number of analyzed samples for each source

Table 2. *Aspergillus* and *Penicillium* spp. isolated from air, soil and grape samples over a two year survey.

3.2 Black Aspergilli and OTA producers

Black aspergilli appeared in all the tested samples, their incidence being higher at early veraison and ripening (Oliveri et al., 2006; Oliveri, 2006) (Fig.3 a-c). According to macro- and micromorphological characteristics, they were identified and classified into two main groups, *A. niger* aggregate and *A. carbonarius* (Fig.3 g-n). *A. ochraceus* has occasionally been detected in grape samples (Fig.3 d-f). A subset of 66 strains was selected for further analysis. PCR assays supported the morphological identification. *A. niger*, *A. carbonarius* and *A. japonicus* were identified by target sequences for each species according to assays described by González-Salgado et al. (2005) and Patiño et al. (2005). In order to characterise the species in *A. niger* aggregate, i.e. *A. niger* and *A. tubingensis*, very difficult to differentiate by classical morphological criteria, the RFLP analysis with *RsaI* was performed. This differentiation is very important to as to avoid overestimating toxicological contamination and related risks. A primer annealing site or restriction nuclease cleavage site was further confirmed by ITS sequencing which also confirmed the identity of the isolates (Oliveri et al., 2008). The OTA production of isolates belonging to *A. carbonarius*, *A. niger*, *A. tubingensis* and *A. japonicus* was assessed by enzyme-linked immunosorbent assay (Oliveri et al., 2006b, 2008). 56% of strains were shown to produce OTA. *A. carbonarius* isolates were the strongest OTA producers with some of them producing high concentrations of OTA (>40 ppb).

3.3 Intraspecific variability

A fAFLP protocol was used to assess specific and intraspecific variability (Oliveri et al., 2008). In agreement with other studies (Perrone et al., 2006a, 2006b), the AFLP technique generated enough polymorphism to differentiate between and within the species of black aspergilli. *A. niger*, *A. tubingensis*, *A. carbonarius* and *A. japonicus* strains were clearly differentiated, although *A. niger* strains clustered into two different groups. Intraspecific variability didn't correlate with the isolate origin. In fact isolates from different vineyards either of grape or the environment could also cluster in the same or in different clusters. Perrone et al. (2006a, 2006b) analyzed representative strains from the main wine producing

Fig. 3. Fungal colonies isolated from grape (a) , soil (b) and air (c) samples from vineyard. Colony morphologies and light microscopy pictures of conidiphores and conidia of representative isolates belonging to *Aspergillus ochraceus* (d-f), *A. carbonarius* (g - i), *A. niger* (l - n).

MPVCT: Micothèque of Institute of Plant Pathology, University of Catania, Italy; USA; MUCL: Micothèque de L'Université Catholique de Lovain, Belgium
* Reference strains

Fig. 4. UPGMA dendrogram obtained from fAFLP analysis with the selective primer pair E-AT (Cy5-labelled) and M-CT on 17 isolates (10 *A. niger*, 3 *A. tubingensis*, 4 *A. carbonarius*) and 2 reference strains isolated from grape samples in the same vineyard. Fragments between 50 and 600 bp were analysed with PHYLIP® v. 3.66 software

European countries (Italy, France, Spain, Portugal, Greece and Israel) and the four main groups were obtained by AFLP clustering analysis of the strains, three of them showing a well-defined homogeneous population/species with intraspecific homology higher than 48%: *A. carbonarius*, *A. tubingensis* and *Aspergillus* 'uniseriate'. The fourth cluster, called *A. niger* 'like', showed low homology with *A. niger* 'type strain' and high internal heterogeneity. The intra-population variability of *Aspergillus* Section *Nigri* strains isolated

from grape samples within the same vineyard proved that mixed populations of *A. niger* and *A. carbonarius* were present and most of them were OTA producers (Fig. 4). As for larger population studies, no correlation was found between genotypes and mycotoxin production (Martinez-Culebraz et al., 2009).

4. Conclusion

This chapter summarizes data on grape mycoflora, toxigenic fungi and mycotoxin contamination at the pre-harvesting, harvesting and processing stages. Grape rotting and spoilage can be caused by a variety of fungal species, including *Botrytis cinerea*, *Penicillium*, *Aspergillus*, *Alternaria* and *Cladosporium*. In recent years, black *Aspergillus* species (Section *Nigri*) and in particular *A. carbonarius* and *A. niger* aggregate have been described as the main source of grape contamination with the mycotoxin ochratoxin A. In this chapter, we highlighted how *Aspergillus* species distribution on European grapes may occur and vary in relation to meteorological conditions and geographical areas and several studies have shown an increase in the amount of OTA in warmer climates. The literature on various molecular methods used for species identification is reviewed and a critical evaluation of the usefulness of various techniques and genomic loci for the species identification of black aspergilli is presented. Reports of the occurrence of black aspergilli in vineyards and their potential toxigenicity must be reconsidered on the basis of the wide molecular biodiversity found within morphologically undistinguishable strains of this section. Mycotoxin production is a characteristic of the species, so by studying the species intraspecific biodiversity can predict potential mycotoxin hazards. Different isolates belonging to the black aspergilli species showed varying abilities to produce OTA so it becomes utmost importance to guarantee a quality control of the grapes and grape derived products, through accurate contaminant mycoflora identification.

5. References

- Abarca, M.L., Accensi, F., Bragulat, M.R. & Cabañes, F.J. (2001). Current importance of ochratoxin A-producing *Aspergillus* spp. *Journal of Food Protection*, Vol. 64, pp. 903–906, ISSN 0362-028X
- Abarca, M.L., Accensi, F., Cano, J., Cabañes, F.J. (2004). Taxonomy and significance of black aspergilli. *Antonie Van Leeuwenhoek*, Vol. 86, pp. 33–49, ISSN 0003-6072
- Accensi, F., Cano, J., Figuera, L., Abarca, M.L. & Cabañes, F.J. (1999). New PCR method to differentiate species in the *Aspergillus niger* aggregate. *FEMS Microbiology Letters*, Vol. 180, No.2, pp. 191–196, ISSN 0378-1097
- Atoui, A., Dao, P., Mathieu, F. & Lebrihi, A. (2006). Amplification and diversity analysis of ketosynthase domains of putative polyketide synthase genes in *Aspergillus ochraceus* and *Aspergillus carbonarius*. *producers of ochratoxin A. Molecular Nutrition & Food Research*, Vol. 50, pp. 448–493, ISSN 1613-4133
- Atoui, A., Mathieu, F. & Lebrihi, A. (2007). Targeting a polyketide synthase gene for *Aspergillus carbonarius* quantification and ochratoxin A assessment in grapes using real-time PCR. *International Journal of Food Microbiology*, Vol. 115, pp. 313–318, ISSN 0168-1605

- Ayoub, F., Reverberi, M., Ricelli, A., D'Onghia, A. M. & Yaseen, T. (2010). Early detection of *Aspergillus carbonarius* and *A. niger* on table grapes: a tool for quality improvement. *Food Additives and Contaminants*, Vol. 27, No. 9, pp. 1285-1293, ISSN 0265-203X
- Battilani, P., Giorni, P. & Pietri, A. (2001). Role of cultural factors on the content of ochratoxin A in grape. *Journal of Plant Pathology*, Vol. 83, p. 231, ISSN 0929-1873
- Battilani, P., & Pietri, A. (2002). Ochratoxin A in grapes and wine. *European Journal of Plant Pathology*, Vol. 108, pp. 639-643, ISSN 0929-1873
- Battilani, P., Pietri, A., Bertuzzi, T., Languasco, L., Giorni, P. & Kozakiewicz, Z. (2003). Occurrence of ochratoxin A-producing fungi in grapes grown in Italy. *Journal of Food Protection*, Vol. 66, pp. 633-636, ISSN 0362-028X
- Battilani, P., Giorni, P. & Pietri, A. (2003b). Epidemiology and toxin-producing fungi and ochratoxin A. *European Journal of Plant Pathology*, Vol. 109, pp. 723-730, ISSN 0929-1873
- Battilani, P., Barbano, C., Marin, S., Sanchis, V., Kozakiewicz, Z. & Magan, N. (2006). Mapping of *Aspergillus* Section *Nigri* in Southern Europe and Israel based on geostatistical analysis. *International Journal of Food Microbiology*, Vol. 111, pp. S72-S82, ISSN 0168-1605
- Bau, M., Castellà, G., Bragulat, M.R. & Cabañes F.J. (2005). DNA-based characterization of ochratoxin-A-producing and non-producing *Aspergillus carbonarius* strains from grapes. *Research in Microbiology*, Vol. 156, pp. 375-381, ISSN 0923-2508
- Bellí, N., Mitchell, D., Marín, S., Alegre, I., Ramos, A. J., Magan, N. & Sanchis, V., (2005). Ochratoxin A-producing fungi in Spanish wine grapes and their relationship with meteorological conditions. *European Journal of Plant Pathology*, Vol. 113, pp. 233-239, ISSN 0929-1873
- Bellí, N., Marín, S., Argilés, E., Ramos, A.J. & Sanchis, V. (2007a). Effect of chemical treatments on ochratoxinogenic fungi and common mycobiota of grapes (*Vitis vinifera*). *Journal of Food Protection*, Vol. 70, No.1, pp. 157-163, ISSN 0362-028X
- Bellí, N., Marín, S., Coronas, I., Sanchis, V. & Ramos, A.J. (2007b). Skin damage, high temperature and relative humidity as detrimental factors for *Aspergillus carbonarius* infection and ochratoxin A production in grapes. *Food Control*, Vol. 18, pp. 1343-1349, ISSN 0956-7135
- Blesa, J., Soriano, J. M., Moltó, J. C. & Mañes, J. (2006). Factors affecting the presence of ochratoxin A in wines. *Critical Reviews in Food Science and Nutrition*, Vol. 46, pp. 473-478, ISSN 1040-8398.
- Castegnaro, M. & Pfohl-Leszkowicz, A. (2002). Les mycotoxines: Contaminations omniprésentes dans l'alimentation animales et humaines. La sécurité alimentaire du consommateur, Lavoisier, Tec & doc., Moll & Moll (Eds.), Paris.
- Chiotta, M.L., Reynoso, M.M., Torres, A.M., Combina, M. & Chulze, S.N. (2011). Molecular characterization and toxigenic profile of *Aspergillus* section *Nigri* populations isolated from the main grape-growing regions in Argentina. *Journal of Applied Microbiology*, Vol. 110, No.2, pp. 445-454, ISSN 1364-5072
- Clarke, K., Emmett, R.W., Kazi, B.A., Nancarrow, N. & Leong, S., (2003). Susceptibility of dried grape varieties to berry splitting and infection by *Aspergillus carbonarius* *Proceedings of 8th International Congress of Plant Pathology*, vol. 2, p. 140, Christchurch, New Zealand

- Clouvel, P., Bonvarlet, L., Martinez, A., Lagouarde, P., Dieng, I. & Martin, P. (2008). Wine contamination by ochratoxin A in relation to vine environment. *International Journal of Food Microbiology*, Vol. 123, pp. 74-80, ISSN 0168-1605
- Cozzi, G., Pascale, M., Perrone, G., Visconti, A. & Logrieco, A. (2006) Effect of *Lobesia botrana* damages on black aspergilli rot and ochratoxin A content in grapes. *International Journal of Food Microbiology* Vol. 111, No. S1, pp. S88-S92
- Da Rocha R., Palacios, V., Combina, M., Fraga, M.E., De Oliveira Rekson, A., Magnoli, C.E. & Dalcero, A.M. (2002) Potential ochratoxin A producers from wine grapes in Argentina and Brazil. *Food Additives and Contaminants* 19, pp. 408-414.
- Dachoupakan, C., Ratomahenina, R., Martinez, V., Guiraud, J.P., Baccou, J.C. & Schorr-Galindo, S. (2009). Study of the phenotypic and genotypic biodiversity of potentially ochratoxigenic black aspergilli isolated from grapes. *International Journal of Food Microbiology*, Vol. 132, No. 1, pp. 14-23, ISSN 0168-1605
- Dao, H.P., Mathieu, F. & Lebrihi, A. (2005). Two primer pairs to detect OTA producers by PCR method. *International Journal of Food Microbiology*, Vol. 104, pp. 61-67, ISSN 0168-1605
- Ferracin, L.M., Frisvad, J.C., Taniwaki, M.H., Iamanaka, B.T, Sartori, D., Schapovaloff, M.E. & Pelegrinelli Fungaro, M.H. (2009). Genetic relationships among strains of the *Aspergillus niger* Aggregate. *Brazilian Archives of Biology and Technology*, Vol. 52, pp. 241-248, ISSN 1516-8913
- Fleet, G.H. (1999) Microorganisms in food ecosystems. *International Journal of Food Microbiology*, Vol. 50, pp. 101-117, ISSN 0168-1605
- Fleet, G.H. (2001) Wine. In: *Food Microbiology Fundamentals and Frontiers*, 2nd edn, Doyle, M.P., Beuchat, L.R. and Montville, T.J. (Ed.), 747-772. Washington, DC: ASM Press.
- Frisvad, J.C., Smedsgaard, J., Samson, R.A., Larsen, T.O. & Thrane, U. (2007). Fumonisin B2 production by *Aspergillus niger*. *Journal of Agricultural and Food Chemistry*, Vol. 14;55, No.23, pp. 9727-32, ISSN 0021-8561
- Gallo, A., Perrone, G., Solfrizzo, M., Epifani, F., Abbas, A., Dobson, A.D.W. & Mulè, G. (2009). Characterisation of a pks gene which is expressed during ochratoxin A production by *Aspergillus carbonarius*. *International Journal of Food Microbiology*, Vol. 129, (October 2009) pp. 8-15, ISSN 0168-1605
- Geiser, D.M., Klich, M.A., Frisvad, J.C., Peterson, S.W., Varga, J. & Samson R.A. (2007). The current status of species recognition and identification in *Aspergillus*. *Studies in Mycology*, Vol. 59, pp. 1-10, ISSN 0166-0616
- González-Salgado, A., Patiño, B., Vázquez, C. & González-Jaén, M.T. (2005). Discrimination of *Aspergillus niger* and other *Aspergillus* species belonging to section *Nigri* by PCR assays. *FEMS Microbiology Letters*, Vol. 245, pp. 353-361, ISSN 0378-1097
- González-Salgado, A., Patino, B., Gil-Serna, J., Vazquez, C. & González -Jaen, M.T. (2009). Specific detection of *Aspergillus carbonarius* by SYBRs Green and TaqMans quantitative PCR assays based on the multicopy ITS2 region of the rRNA gene. *FEMS Microbiology Letters*, Vol. 295, pp. 57-66, ISSN 0378-1097
- Guzev, L., Danshin, A., Ziv, S. & Lichter, A. (2006). Occurrence of ochratoxin A producing fungi in wine and table grapes in Israel. *International Journal of Food Microbiology*, Vol. 111, pp. S67-S71, ISSN 0168-1605

- Heenan, C.N., Shaw, K.J. & Pitt, J.I. (1998) Ochratoxin A production by *Aspergillus carbonarius* and *A. niger* isolates and detection using coconut cream agar. *Journal of Food Mycology*, Vol. 1, pp. 67–72.
- Hewitt, W.B. (1988). Berry rots and raisin moulds. In: Pearson, R.G., Goheen, A.C. (Eds.), *Compendium of Grape Diseases*. The American Phytopathological Society, pp. 26–28, APS Press, ISBN 92-9043-153-9. St. Paul, Minnesota.
- Hocking, A.D., Su-lin, L. Leong, S.L., Kazi, B.A., Emmett, R.W., Eileen, S. & Scott E.S. (2007). Fungi and mycotoxins in vineyards and grape products. *International Journal of Food Microbiology*, Vol. 119, No.1-2, (October 2007), pp. 84-88, ISSN 0168-1605
- IARC (1993) IARC Monographs on the Evaluation of Carcinogenic Risks to Humans (1993). Ochratoxin A. In: Some Naturally Occurring Substances: Food Items and Constituents, Heterocyclic Aromatic Amines and Mycotoxins. IARC Press, Lyon, France Vol. 56, pp. 489–521.
- Ilic, Z., Pitt, J.I. & Carter, D.A. (2001). Identifying genes associated with ochratoxinA production in *Aspergillus carbonarius* and *Aspergillus niger*. *Proceedings of 21st Fungal Genetics Conference*, Pacific Grove, USA, March 13–18, 2001
- Ilic, Z., Pitt, J.I., & Carter, D.A., (2004). Genetic diversity and ochratoxin production in the black *Aspergilli*. Australian Mycological Society, joint meeting with the Australian Society for Microbiology Annual Scientific Meeting, Sydney, Australia, September 27–28, 2004
- Kazi, B.A., Emmett, R.W., Clarke, K. & Nancarrow, N., (2003a). Black *Aspergillus* mould in Australia. *Proceedings of 8th International Congress of Plant Pathology*, Vol. 2, p. 119, Christchurch, New Zealand
- Kazi, B.A., Emmett, R.W., Nancarrow, N. & Clarke, K. (2003b). Effects of temperature, moisture and/or irrigation on the survival of *Aspergillus carbonarius* in soil. *Proceedings of 8th International Congress of Plant Pathology*, Vol. 2, p. 140, Christchurch, New Zealand
- Kazi, B.A., Emmett, R.W., Nancarrow, N. & Clarke, K. (2004). Incidence of *Aspergillus carbonarius* in Australian vineyards. In: Ophel-Keller, K., Hall, B. (Eds.), 75–76, 3rd Australasian Soil-borne Disease Symposium, South Australian Research and Development Institute, Adelaide, Australia
- Klich, M.A. & Pitt, J.I. (1988). A Laboratory Guide to Common *Aspergillus* Species and Their Teleomorphs. CSIRO Division of Food Processing, North Ryde, New South Wales, Australia.
- Klich, M.A (2002). Identification of common *Aspergillus* Species. Centraalbureau Voor Schimmelcultures, Utrecht, The Netherlands.
- Leong, S.L., Hocking, A.D., Pitt, J.I., Kazi, B.A., Emmett, R.W. & Scott, E.S. (2006). Australian research on ochratoxigenic fungi and ochratoxin A. *International Journal of Food Microbiology*, Vol. 111, pp. S10-S17, ISSN 0168-1605
- Leong, S.L., Hocking, A.D. & Scott, E.S. (2007). *Aspergillus* producing ochratoxin A: isolation from vineyard soils and infection of Semillon bunches in Australia. *Journal of Applied Microbiology*, Vol. 102, pp. 124–133, ISSN 1364-5072.
- Logrieco, A., Moretti, A., Perrone, G. & Mulè, G. (2007). Biodiversity of complexes of mycotoxigenic fungal species associated with *Fusarium* ear rot of maize and *Aspergillus* rot of grape. *International Journal of Food Microbiology*, Vol. 119, pp. 11–16, ISSN 0168-1605

- Logrieco, A., Ferracane, R., Visconti, A. & Ritieni, A. (2010). Natural occurrence of fumonisin B₂ in red wine from Italy. *Food Additives and Contaminants*, Vol. 27, No. 8, pp. 1136-1141, ISSN 0265-203X
- López-Mendoza, M.C., Crespo-Sempere, A. & Martínez-Culebras P.V. (2009). Identification of *Aspergillus tubingensis* strains responsible for OTA contamination in grapes and wine based on the acyl transferase domain of a polyketide synthase gene. *International Journal of Food Science & Technology*, Vol. 44, No. 11, (November 2009), pp. 2147-2152, ISSN 0950-5423
- Lucchetta, G., Bazzo, I. Cortivo, G. dal Stringher, L., Bellotto, D., Borgo, M. & Angelini, E. (2010). Occurrence of black aspergilli and ochratoxin A on grapes in Italy. *Toxins*, Vol. 2, No. 4, pp. 840-855, ISSN 2072-6651
- Martínez-Culebras, P.V., Crespo-Sempere, A., Sánchez-Hervás, M., Elizaquivel, P., Aznar, R., Ramón, D. (2009). Molecular characterization of the black *Aspergillus* isolates responsible for ochratoxin A contamination in grapes and wine in relation to taxonomy of *Aspergillus* section *Nigri*. *International Journal of Food Microbiology*, Vol. 132, No. 1, (June 2009), pp. 33-41, ISSN 0168-1605
- Magnoli, C.E., Violante, M., Combina, M., Palacio, G. & Dalcero, A.M. (2003). Mycoflora and ochratoxin-producing strains of *Aspergillus* section *Nigri* in wine grapes in Argentina. *Letters in Applied Microbiology*, Vol. 37, pp. 179-184, ISSN 0266-8254
- Medina, A., Mateo, R., López-Ocaña, L., Valle-Algarra, F.M. & Jiménez, M. (2005). Study of Spanish grape mycobiota and ochratoxin A production by isolates of *Aspergillus tubingensis* and other members of *Aspergillus* Section *Nigri*. *Applied and Environmental Microbiology* 71, pp. 4696-4702, ISSN 0099-2240
- Morello, L.G., Sartori, D., De Oliveira Martinez, A.L., Vieira, M.L.C., Taniwaki, M.H. & Pelegrinelli Fungaro, M.H. (2007). Detection and quantification of *Aspergillus westerdijkiae* in coffee beans based on selective amplification of β -tubulin gene by using real-time PCR. *International Journal of Food Microbiology*, Vol. 119, pp. 270-276, ISSN 0168-1605
- Mogensen, J. M., Frisvad, J. C., Thrane, U. & Nielsen, K.F. (2010a). Widespread occurrence of the mycotoxin fumonisin B₂ in wine. *Journal of Agricultural and Food Chemistry*, Vol. 58, No. 8, pp. 4853-4857, ISSN 0021-8561
- Mogensen, J. M. Larsen, T. O. & Nielsen, K. F. (2010b). Production of fumonisin B₂ and B₄ by *Aspergillus niger* on grapes and raisins. *Journal of Agricultural and Food Chemistry*, Vol. 58, No. 2, pp. 954-958, ISSN 0021-8561
- Mulè, G., Susca, A., Logrieco, A., Stea, G. & Visconti, A. (2006). Development of a real-time PCR assay for the detection of *Aspergillus carbonarius* in grapes. *International Journal of Food Microbiology*, Vol. 111, Supplement 1, (September 2006), pp. S28-S34, ISSN 0168-1605
- Narasimhan B. & Asokan M. (2010). Genetic variability of *Aspergillus terreus* from dried grapes using RAPD-PCR. *Advances in Bioscience and Biotechnology*, Vol. 1, No. 4, (October 2010) pp. 345-353, ISSN 2156-8456
- Niessen, L., Schmidt, H., Mühlencoert, E., Farber, P. & Karolewicz, A. (2005). Advances in the molecular diagnosis of ochratoxin A-producing fungi. *Food Additives and Contaminants*, Vol. 22, No. 4, pp. 324-334, ISSN 0265-203X

- Nielsen, K.F., Mogensen, J.M., Johansen, M., Larsen, T.O. & Frisvad, J.C. (2009). Review of secondary metabolites and mycotoxins from the *Aspergillus niger* group. *Analytical and Bioanalytical Chemistry*, Vol. 395, No.5, pp. 1225-1242, ISSN 1618-2642
- O'Callaghan, J., Caddick, M.X. & Dobson, A.D. (2003). A polyketide synthase gene required for ochratoxin A biosynthesis in *Aspergillus ochraceus*. *Microbiology*, Vol. 149, pp. 3485-3491, ISSN 1350-0872
- Oliveri, C. (2006). Caratterizzazione morfologica e molecolare di specie ocratossigene di *Aspergillus* sez. *Nigri* isolate in vigneti dell'Etna ed esperienze di lotta biologica. PhD thesis, University of Catania, 179 pp.
- Oliveri, C., Fardella, M., Grimaldi, V. & Catara, V. (2006a). Characterization of OTA-producing *Aspergillus* spp., strains isolated from sicilian vineyards. *Journal of Plant Pathology*, Vol. 88, No.3, pp. S51-S52, ISSN 1125-4653
- Oliveri, C., Fardella, M., Grimaldi, V. & Catara, V. (2006b). Characterization of OTA-producing *Aspergillus* spp. strains isolated from sicilian vineyards *Journal of Plant Pathology*, Vol. 88, No.3, p. S52.
- Oliveri, C., Torta, L. & Catara, V. (2008). A polyphasic approach to the identification of ochratoxin A-producing black *Aspergillus* isolates from vineyards in Sicily. *International Journal of Food Microbiology*, Vol. 27, (June 2008), pp. 147-154, ISSN 0168-1605
- Ottener, H. & Majerus, P. (2000). Occurrence of ochratoxin A (OTA) in wines: influence of the type of wine and its geographical origin. *Food Additives and Contaminants*, Vol. 17, pp. 793-798, ISSN 0265-203X
- Patiño, B., Gonzáles-Salgado, A., Gonzáles-Jaén, M.T. & Vázquez, C. (2005). PCR detection assays for the ochratoxigen-producing *Aspergillus carbonarius* and *A. ochraceus* species. *International Journal of Food Microbiology*, Vol. 104, pp. 207-214, ISSN 0168-1605
- Pelegrielli-Fungaro, M.H., Vissotto, P.C., Sartori, D., Vilas-Boas, L.A., Furlaneto, M C. & Taniwaki, M.H. (2004). A molecular method for detection of *Aspergillus carbonarius* in coffee beans. *Current Microbiology*, Vol. 49, pp. 123-127, ISSN 0343-8651
- Perrone, G., Mulè, G., Susca, A., Battilani, P., Pietri, A. & Logrieco, A. (2006a). Ochratoxin A production and AFLP analysis of *Aspergillus carbonarius*, *Aspergillus tubingensis*, and *Aspergillus niger* strains isolated from grapes in Italy. *Applied and Environmental Microbiology*, Vol. 72, pp. 680-685, ISSN 0099-2240
- Perrone, G., Susca, A., Epifani, F. & Mulè, G. (2006b). AFLP characterization of Southern Europe population of *Aspergillus* Section *Nigri* from grapes. *International Journal of Food Microbiology*, Vol. 111, pp. S22-S27, ISSN 0168-1605
- Perrone, G., Susca, A., Cozzi, G., Ehrlich, K., Varga, J., Frisvad, J.C., Meijer, M., Noonim, P., Mahakarnchanakul, W. & Samson, R.A. (2007). Biodiversity of *Aspergillus* species in some important agricultural products. *Studies in Mycology* 59, 53-66, ISSN 0166-0616
- Perrone, G., Varga, J., Susca, A., Frisvad, J.C., Stea, G., Kocsubé, S., Tóth, B., Kozakiewicz, Z. & Samson, R.A. (2008). *Aspergillus uvarum* sp. nov., an uniseriate black *Aspergillus* species isolated from grapes in Europe. *International Journal of Systematic and Evolutionary Microbiology*, Vol.58, (2008), pp. 1032-1039, ISSN 14665026
- Petzinger, E. & Weidenbach, A. (2002). Mycotoxins in the food chain: the role of ochratoxins. *Livestock Production Science*, Vol. 76, (2002), pp. 245-250, ISSN 0301-6226.

- Pfohl-Leszkowicz, A., Petkova-Bocharova, T., Chernozemsky, I.N. & Castegnaro, M. (2002). Balkan endemic nephropathy and associated urinary tract tumours: a review on aetiological causes and the potential role of mycotoxins. *Food additives and Contaminants*, Vol. 19, No. 3, (2002), pp.282-302, ISSN 0265-203X
- Pitt, J.I. & Hocking, A.D. (1999). *Fungi and Food Spoilage*, 2nd edn. Aspen Publishers, Inc., Gaithersburg, Md.
- Pitt, J.I. & Hocking, A.D. (2009). *Fungi and Food Spoilage*, 3rd edn. Springer-Verlag New York Inc., 04.09.2009, pp.540, ISBN 0387922067
- Pollastro, S., De Miccolis Angelini, R., Abbatecola, A., De Guido, M.A. & Faretra, F. (2005). Real-Time PCR for quantitative detection of *Aspergillus carbonarius* in grapes and musts. *Proceedings of International Workshop: Ochratoxin A in grapes and wine: prevention and control*, p. 43, Marsala (TP), Italy, October 20-21, 2005
- Pollastro, S., De Miccolis, R.M. & Faretra, F. (2006). A new semi-selective medium for the ochratoxigenic fungus *Aspergillus carbonarius*. *Journal of Plant Pathology*, Vol. 88, pp. 107-112, ISSN 1125-4653
- Rousseau, J., 2004. Ochratoxin A in wines: Current knowledge. Retrieved from Vinidea.net, Wine Internet Technical Journal.
- Sartori, D., Furlaneto, M.C., Martins, M.K., Ferreira de Paula, M.R., Pizzirani-Kleiner, A.A., Taniwaki, M.H. & Pelegrinelli-Fungaro, M.H. (2007). PCR method for the detection of potential ochratoxin-producing *Aspergillus* species in coffee beans. *Research in Microbiology*, Vol.157, 350-354, ISSN 0923-2508.
- Sartori, D., Taniwaki, M.T., Iamanaka, B., & Pelegrinelli Fungaro M.H. (2010). Molecular Diagnosis of Ochratoxigenic Fungi. (2010) Part 1, pp. 195-212
- Sage, L., Krivobok, S., Delbos, E., Seigle-Murandi, F. & Creppy, E.E. (2004). Fungal microflora and ochratoxin A production in grapes and musts from France. *Agricultural and Food Chemistry*, Vol. 50, pp. 1306-1311, ISSN 0021-8561
- Sage, L., Garon, D., Seigle-Murandi, F. (2004). Fungal microflora and ochratoxin A risk in French vineyards. *Journal of Agricultural and Food Chemistry*, Vol. 52, pp. 5764-5768, ISSN 0021-8561
- Samson, R.A., Hoekstra, E.S. & Frisvad, J.C., 2004. In: Samson, R.A., Hoekstra, E.S. (Eds.), *Introduction to Food- and Airborne Fungi*, 7th ed. Centraalbureau Voor Schimmelcultures, Utrecht, The Netherlands. 389 pp.
- Samson, R.A., Noonim, P., Meijer, M., Houbaken, J., Frisvad, J.C. & Varga, J. (2007). Diagnostic tools to identify black aspergilli. *Studies in Mycology*, Vol. 59, pp. 129-145, ISSN 0166-0616
- Schmidt, H., Ehrmann, M., Vogel, R. F., Taniwaki, M. H. & Niessen, L. (2003). Molecular Typing of *Aspergillus ochraceus* and construction of species specific SCAR-primers based on AFLP. *Systematic & Applied Microbiology*, Vol. 26, pp.138-146, ISSN 0723-2020
- Schmidt, H., Bannier, M., Vogel, R.F. & Niessen, L. (2004). Detection and quantification of *Aspergillus ochraceus* in green coffee by PCR. *Letters in Applied Microbiology*, Vol. 38, pp. 464-469, ISSN 0266-8254
- Selma, M.V., Martinez-Culebras, P.V., Elizaquivel, P. & Aznar, R. (2009). Simultaneous detection of the main black aspergilli responsible for ochratoxin A (OTA) contamination in grapes by multiplex real-time polymerase chain reaction. *Food Additives and Contaminants*, Vol. 26, No.2, pp. 180-188, ISSN 0265-203X

- Serra, R., Braga, A. & Venâncio, A. (2005). Mycotoxin-producing and other fungi isolated from grapes for wine production, with particular emphasis on ochratoxin A. *Research in Microbiology*, Vol. 156, pp. 515–521, ISSN 0923-2508
- Serra, R., Cabañes, J., Perrone, G., Kozakiewicz, Z., Castellá, G., Venâncio, A. & Mulè, G. (2006). *Aspergillus ibericus*: a new species of the Section *Nigri* isolated from grapes. *Mycologia*, Vol. 98, No.2, pp. 295–306, ISSN 0027-5514
- Serra, R., Mendonca, C. & Venâncio, A. (2006). Fungi and ochratoxin A detected in healthy grapes for wine production. *Letters in Applied Microbiology*, Vol. 42, pp. 42–47, ISSN 0266-8254
- Susca, A., Proctor, R.H., Mulè, G., Stea, G., Ritieni, A., Logrieco, A. & Moretti, A. (2010). Correlation of Mycotoxin Fumonisin B₂ Production and Presence of the Fumonisin Biosynthetic Gene *fum8* in *Aspergillus niger* from Grape. *Journal of Agricultural and Food Chemistry*, Vol. 58, No.6, pp. 9266–9272, ISSN 0021-8561
- Tournas, V.H. & Katsoudas, E. (2005). Mould and yeast flora in fresh berries, grapes and citrus fruits. *International Journal of Food Microbiology* Vol. 105, pp.11–17, ISSN 0168-1605
- Tjamos, S.E., Antoniou, P.P., Kazantzidou, A., Antonopoulos, D.F., Papageorgiou, I. & Tjamos, E.C. (2004). *Aspergillus niger* and *Aspergillus carbonarius* in Corinth raisin and wine-producing vineyards in Greece: population composition, ochratoxin A production and chemical control. *Journal of Phytopathology*, Vol. 152:, pp. 250–255, ISSN 0931-1785
- Tjamos, S.E., Antoniou, P.P. & Tjamos, E.C. (2006). *Aspergillus* spp., distribution, population composition and ochratoxin A production in wine producing vineyards in Greece. *International Journal of Food Microbiology*, Vol. 111, pp. S61–S66, ISSN 0168-1605
- Varga, J., Kevei F., Hamari, Z., Toth, B., Teren, J., Croft, J.H. & Kozakiewicz, Z. (2000). Genotypic and phenotypic variability among black aspergilli. In: Samson RA, Pitt JI, eds. *Integration of modern taxonomic methods for Penicillium and Aspergillus classification*. Amsterdam, the Netherlands: Harwood Academic Publishers. pp. 397–411.
- Varga J & Kozakiewicz Z. (2006). Ochratoxin A in grapes and grape-derived products. *Trends in Food Science & Technology*, Vol. 17, pp. 72–81. ISSN 0924-2244
- Varga, J., Kocsubé, S., Tóth, B., Frisvad, J.C., Perrone, G., Susca, A., Meijer, M. & Samson, R.A. (2007). *Aspergillus brasiliensis* sp. nov., a biseriolate black *Aspergillus* species with world-wide distribution. *International Journal of Systematic and Evolutionary Microbiology*, Vol. 7, pp. 1925–1932, ISSN 1466-5026
- Vrabcheva, T., Usleber, E., Dietrich, R. & Märtilbauer, E. (2000). Co-occurrence of ochratoxin A and citrinin in cereals from Bulgarian villages with a history of Balkan endemic nephropathy. *Journal of Agricultural and Food Chemistry*, Vol. 48, (2000), pp. 2483–2488, ISSN 0021-8561
- Zimmerli, B. & Dick, R. (1996). Ochratoxin A in table wine and grape-juice: occurrence and risk assessment. *Food Additives and Contaminants*, Vol. 13, pp. 655–668, ISSN 0265-203X

The Dynamical Processes of Biodiversity - Case Studies of Evolution and Spatial Distribution

Edited by PhD. Oscar Grillo

ISBN 978-953-307-772-7

Hard cover, 366 pages

Publisher InTech

Published online 02, December, 2011

Published in print edition December, 2011

Driven by the increasing necessity to define the biological diversity frame of widespread, endemic and threatened species, as well as by the stimulating chance to describe new species, the study of the evolutive and spatial dynamics is in constant execution. Systematic overviews, biogeographic and phylogenetic backgrounds, species composition and distribution in restricted areas are focal topics of the 15 interesting independent chapters collected in this book, chosen to offer to the reader an overall view of the present condition in which our planet is.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Cinzia Oliveri and Vittoria Catara (2011). Mycoflora and Biodiversity of Black Aspergilli in Vineyard Eco-Systems, The Dynamical Processes of Biodiversity - Case Studies of Evolution and Spatial Distribution, PhD. Oscar Grillo (Ed.), ISBN: 978-953-307-772-7, InTech, Available from: <http://www.intechopen.com/books/the-dynamical-processes-of-biodiversity-case-studies-of-evolution-and-spatial-distribution/mycoflora-and-biodiversity-of-black-aspergilli-in-vineyard-eco-systems>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2011 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](https://creativecommons.org/licenses/by/3.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen