We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Anticancer and Antimicrobial Potential of Plant-Derived Natural Products

Wamidh Hadi Talib

*Department of Clinical Pharmacy and Therapeutics, University of Applied Science,
Jordan*

1. Introduction

Plant use in treating diseases is as old as civilization (Fabricant & Farnsworth, 2001) and traditional medicines are still a major part of habitual treatments of different maladies (Alviano & Alviano, 2009). In recent times and due to historical, cultural, and other reasons, folk medicine has taken an important place especially in developing countries where limited health services are available. However, the absence of scientific evaluation of medicinal plants to validate their use may cause serious adverse effects (Souza et al., 2004).

Plants are considered as one of the main sources of biologically active materials. Recent records reported that medicinal herbs are used by 80% of the people living in rural areas as primary healthcare system (Sakarkar & Deshmukh, 2011). It has been estimated that about 50% of the prescription products in Europe and USA are originating from natural products including plants or their derivatives (Cordell, 2002; Newman et al., 2003). Out of the 250,000 – 500,000 plant species on earth, only 1-10 % have been studied chemically and pharmacologically for their potential medicinal value (Verpoote, 2000). In the Middle East region 700 species of the identified plants are known for their medicinal values (Azaizeh et al., 2006).

In spite of the recent domination of the synthetic chemistry as a method to discover and produce drugs, the potential of bioactive plants or their extracts to provide new and novel products for disease treatment and prevention is still enormous (Raskin et al., 2002). Compared with chemical synthesis, plant derived natural products represent an attractive source of biologically active agents since they are natural and available at affordable prices (Ghosh et al., 2008). Also plants derived agents may have different mechanisms than conventional drugs, and could be of clinical importance in health care improvement (Eloff et al., 1998). Plant materials might be bioactive secondary metabolites that have the potential to treat different afflictions. Examples of these compounds include phenols, phenolic glycosides, unsaturated lactones, sulphur compounds, saponins, cyanogenic glycosides and glucosinolates (Mukherjee et al., 2001; Quiroga et al., 2001). Such plant derived natural products are the main focus of many scientists to develop new medication for different diseases like cancer and microbial infection.

Cancer is a group of diseases characterized by uncontrolled growth and spread of abnormal cells (Karp, 1999). The high mortality rate among cancer patients is an indication of the

limited efficiency of the current therapies including radiation, chemotherapy and surgery (Xu et al., 2009). Cancer development is a multi-step process including induction of genetic instability, abnormal expression of genes, abnormal signal transduction, angiogenesis, metastasis, and immune evasion (Boik, 2001). For many years, scientists were searching for miracle cures for cancer using chemically synthesized or natural pure compounds. In the last few decades, research has been focused on the use of natural products such as crude plant extracts or a combination of different phytochemicals for cancer therapy; this trend is based upon: first, the synergistic effect of the different plant metabolites in the crude extract, second, is the multiple points of intervention of such extracts (Neergheen, 2009). This is one of the many faces of using plants in the quest of controlling different diseases. Another face is the use of such plant products in controlling microbial resistance spread. As a result of the uncontrolled use of many antibiotics, their efficiency is being threatened by the emergence of microbial resistance to existing chemotherapeutic agents (Cowan, 1999; Pareke & Chanda, 2007). Bacterial strains such as methicilin-resistant *Staphylococcus aureus* (MRSA), pencillin-resistant *Streptococcus pneumonia* (PRSP), and Vancomycin-resistant enterococci (VRE) in addition to the development of multidrug-resistant (MDR) bacterial strains (Alanis, 2005) are just few examples that made the search for new and novel bioactive substances among the first priorities in the search for antitumor, antibacterial, and antifungal substances (Ficker et al., 2005).

Realizing all the aforementioned, it is clear that there is a pressing need to participate in the search for new and novel bioactive agents that would help in providing new avenues in fighting diseases and reducing suffering. This chapter will provide information about selected plants that have a potential to provide new anticancer and/or antimicrobial agents.

2. The history of traditional medicine

For thousands of years and in different parts of the world, medicinal plants have been used to treat different diseases (Palombo, 2009). Fossil records documented the use of medicinal plants by humans before 60,000 years (Fabricant & Farnsworth, 2001). Nowadays plants continue to be the major source of medicine in rural regions of developing countries (Chitme et al., 2003) and it has been estimated that about 80% of peoples in developing countries are still using medicinal plants for their health care (Kim, 2005).

The Eastern region of the Mediterranean has been characterized by high inventory of medicinal herbs used by local traditional healers to treat different ailments (Azaizeh et al., 2006). Research on medicinal plant treasures is based on present day and historical systems of traditional and local medicine (Al-Qura'n, 2009). During the Ottoman Empire and following the Byzantine traditions, hospitals were run by physician who used pharmacists to gather medicinal plants and prepare remedies originating from classical Greek and folk medicinal practice (Littlewood et al., 2002).

A comprehensive study on practitioners and herbalist in Palestine revealed that approximately 129 plant species are still prescribed to treat different diseases including liver, digestive tract, respiratory system, skin, cancer and other diseases (Azaizeh et al., 2003). On the other hand, the high diversity of plant species in Jordan encouraged many to study the distribution and use of medicinal plants in this country. More than 100 herbalists were interviewed in an extensive survey for Jordanian medicinal plants indicated the

presence of 150 herbal plants used in folkloric medicine (Abu-Irmaileh & Afifi, 2003). Seventy nine plant species are still in use in traditional medicine in the Showbak region (south of Jordan) while forty six are part of the popular medicine in the Ajloun Heights region (north of Jordan) and some of the plants are used in both regions (Aburjai et al., 2007; Al-Qura'n, 2009). Most of the practitioners were not licensed and have no scientific information about medicinal plants (Abu-Irmaileh & Afifi, 2003; Azaizeh et al., 2003).

The diversity of plants in the Mediterranean region is declining, were recent estimates reported less than 200-250 plant species are used to treat different ailments in the Arab traditional medicine (Said et al., 2002; Abu-Irmaileh & Afifi, 2003; Saad et al., 2005), compared to more than 700 species which were identified for their medicinal uses in previous decades (Azaizeh et al., 2006). The high rate of plant extinction on the earth necessitates an increase in the efforts to study plant natural products for their potential to provide treatment for different afflictions.

3. Cancer biology

DNA damage causes conversion of normal cell into a cancer cell. Cancer cells lack the ability to communicate with their neighboring cells. The first cancer cell starts to divide producing daughter cells, which in turn divide to produce more and more cancer cells. As cancer cells divide, they develop malignant characteristics including metastasis, immune system evasion, and induction of blood vessels formation (angiogenesis). Continuous cell division of cancer cells lead to the formation of tumors. In solid tumors, blood vessels become structurally and functionally abnormal; this abnormality leads to heterogeneous blood flow which creates chronically hypoxic and acidic regions in the core of the solid tumor (Brown & Wilson, 2004). These hypoxic regions lead to the activation of angiogenesis and cell survival genes in addition to other genes that induce drug resistant (Chen et al., 2003). Furthermore, the low pH microenvironment of cancer cells in the tumor core may prevent the active uptake of some anticancer drugs (Mahoney et al., 2003).

The two traditional therapies (chemotherapy and radiation) are not greatly efficient in treating hypoxic cancer cells (Tannock et al., 1998). The killing effect of ionizing radiation depends on the presence of oxygen which is absent or very low in the tumor core and the poor vascularization minimizes the delivery of chemotherapeutic agents (Brown & Wilson, 2004). This makes the poorly vascularized regions of tumors a major obstacle to effective treatment and opens the door to other therapies that may use different mechanisms to targets highly resistant cancer cells.

4. Oncogenes and tumor suppressor genes

Two sets of genes are controlling cancer development. Oncogenes are the first set of genes and are involved in different cell activities including cell division. However, overexpression of these genes transforms a normal cell into a cancer cell. On the other hand, the second set of genes (tumor suppressor genes) inhibits cancer cell formation by different mechanisms. Tumor suppressor genes are underexpressed in cancer cells while, oncogenes are overexpressed. Table 1 summarizes the main oncogenes and tumor suppressor genes and their role in cancer development. Oncogenes and their products represent good targets for cancer therapy. Other targets include enzymes involved in cell division like topoisomerases

that unwind the DNA during replication. The diversity of plant derived natural products can provide therapeutic products attacking different targets in cancer cells.

Oncogenes	Functions of their proteins
<i>Bcl-2</i>	Inhibits apoptosis and protect cancer cell from free radicals.
<i>c-myc</i>	Initiate cell division and inhibits differentiation.
<i>HER-2/neu(c-erb-2)</i>	Facilitates signal transduction, expressed in 33% of breast cancers.
<i>MDM2</i>	Protect cancer cells from apoptosis by binding and inhibiting <i>p⁵³</i> (tumor suppressor gene that induces apoptosis in damaged cells).
<i>ras</i>	Facilitate tumor invasion by stimulation of collagenase production and inhibits apoptosis by increasing <i>MDM2</i> expression.
<i>fos</i> and <i>jun</i>	Promote uncontrolled proliferation by their participation in cell cycle initiation.
Tumor suppressor genes	Functions of their proteins
<i>Bax</i>	An inducer of apoptosis
<i>Cx32</i> , <i>Cx 43</i> and other connexin producing genes	Inhibits carcinogenesis by restoring communication between cells through gap junctions.
<i>p⁵³</i>	Initiates DNA repair and induce apoptosis in cells that cannot be repaired.
Reference: Boik, 2001	

Table 1. The main oncogenes and tumor suppressor genes.

5. The use of plants in cancer therapy

Cancer is one of the major causes of death and the number of cancer patients is in continuous rise. Every year 2-3 % of deaths recorded world wide arise from different types of cancer (Madhuri & Pandey, 2009). The available treatment methods include surgery, chemotherapy, and radiation (Tannock et al., 1998). The increasing costs of conventional treatments (chemotherapy and radiation) and the lack of effective drugs to cure solid tumors encouraged people in different countries to depend more on folk medicine which is rooted in medicinal plants use (Wood-Sheldon et al., 1997). Such plants have an almost unlimited capacity to produce substances that attract researchers in the quest for new and novel chemotherapeutics (Reed & Pellecchia, 2005). Although some plant products are used in cancer therapy, plant derived anticancer agents represent only one-fourth of the total treatments options. Since 1961, nine compounds originating from plants have been approved for use in cancer therapy in the United States. These agents are vinblastine, vincristine, navelbine, etoposide, teniposide, taxol, taxotere, topotecan, and irinotecan (Lee, 1999). In an extensive study, the anticancer properties of 187 plant species were evaluated. Among them, only 15 species have been used to treat cancer clinically (Kintzios, 2006). It was observed that different plants contain different bioactive compounds and these vary with area, climate and mode of agricultural practice if they are not present in wild environment.

Herbivory, pathogens and competition are the driving forces that induce plant species to develop chemical defense compounds. These plant origin compounds are good models for elucidation of their functional roles in medication and treatment of different afflictions (Wood-Sheldon et al., 1997). For example, the lignin in the roots of *Anthriscus sylvestris* showed an insecticidal activity (Kozawa et al., 1982). Poisonous plants exposed to frequent grazing by animals are commonly rich in alkaloids which have many biological activities including anticancer potential (Kintzios, 2006). However, the growth regulatory properties of some plant metabolites allow them to act as chemotherapeutical agents. Flavonoids from *Scutellaria baicalensis* act on cyclin-dependent kinases to inhibit cancer cell proliferation (Dai & Grant, 2003; Chang et al., 2004).

Thirteen distinct groups of plant-derived natural products with antitumor properties were documented (Kintzios, 2006). Among them, alkaloids (Facchini, 2001), phenylpropanoids (Dixon & Paiva, 1995) and terpenoids (Trapp & Croteau, 2001) are well known for their antitumor potentials.

An integrated part of cancer cell development is the resistance to programmed cell death (apoptosis). Re-establishment of apoptosis in cancer cells is a target mechanism for anticancer agents (Joshi et al., 1999). Some plant-derived products are known to selectively induce apoptosis in cancer cells, which represent the ideal property for successful anticancer agents (Hirano et al., 1995). Identifying the mode of action of anticancer agents of plant origin provide helpful information for their future use. Thus it is important to screen the apoptotic potential of plants either in their crude extract form or as pure compounds (Tarapadar et al., 2001). Due to their multiple intervention strategies, crude plant extracts have been proposed to prevent, arrest, or reverse the cellular and molecular processes of carcinogenesis (Neergheen et al., 2009).

Since the distribution of bioactive compounds differs according to the plant used, different solvents were used to extract these compounds from different plants. The methanol extract of *Scutellaria orientalis* showed potent anti-leukemic activity against HL-60 cell line (Ozmen et al., 2010). The water extract of *Rheum officinale* exhibited significant antiproliferative activity by inducing apoptosis in MCF-7 and A549 cell lines (Li et al., 2009). A potent antiproliferative activity was also reported for the hexane extract of *Casearia sylvestris* stem bark against different cancer cell lines (Mesquita et al., 2009) and the butanol extract of *Pfuffia paniculata* demonstrated high cytotoxic activity against MCF-7 cell line (Nagamine et al., 2009). Additionally, the *Physalis minima* chloroform extract induced apoptosis in human lung adenocarcinoma cell line (Leong et al., 2009). Out of 76 Jordanian plant species, the ethanolic extracts of *Inula graveolens*, *Salvia dominica*, *Conyza canadiensis* and *Achillea santolina* showed potent antiproliferative activity against MCF-7 cell line (Abu-Dahab & Afifi, 2007). The aqueous methanol of *Ononis hirta* and *Inula visocsa* showed high ability to selectively target MCF-7 cancer cells and induced apoptosis (Talib & Mahasneh, 2010a).

A substantial progress has been made in the treatment of cancer since the early years of anticancer drug research. An example of successful anticancer plant products are the vinca alkaloids which were isolated from *Catharanthus roseus*. The first identified vinca alkaloids were Vincristine and Vinblastine (Duflos et al., 2002). The antitumor activities of these compounds involve binding to tubulin and disruption of mitotic spindle assembly (Nobili et al., 2009). Myelosuppression and neurotoxicity were reported as side effects of vinca

alkaloids (Yun-San et al., 2008). Another example of plant products in cancer therapy is taxol which was extracted from the bark of *Taxus brevifolia* (Wani et al., 1971). Its cytotoxic activity is mediated by stabilizing microtubules rather than destabilizing them (Horwitz et al., 1993). Since taxol drugs have low solubility, they are administered together with solvents which cause some adverse effects like hypersensitivity and neuropathies (Onetto et al., 1993). Other anticancer plant products mediate their activities by inhibiting the enzyme DNA topoisomerase. Plants under this category include camptothecins extracted from the Chinese tree *Camptotheca acuminata* (Wall et al., 1966) and Podophyllotoxins extracted from the roots of the Indian plant *Podophyllum peltatum* (Nobili et al., 2009).

In spite of the success of previously mentioned anticancer plant products, the development of multi-drug resistance in cancer chemotherapy still one of the major problems (Xu et al., 2009). To avoid this problem, researchers focused on other targets including starving cancer cells by targeting the process of angiogenesis (formation of new blood vessels) which represent an attractive target since tumors depend on angiogenesis for survival and metastasis (Griggs et al., 2001). Some plant derived agents showed promising anti-angiogenic activity such as stilbene combretsatin- A4 which was isolated from *Combretum caffrum* (Young & Chaplin, 2004). Another strategy to target multi-drug resistance cancers is to use a combination of chemotherapy with other strategies including anaerobic bacteria which experimentally showed promising results in targeting solid tumors (Dang et al., 2001). Many plants exhibited promising anticancer activities (Table 2). Such plants may provide compounds that can change the list of drugs available for cancer treatment in the future.

Family	Species	Compounds
Anacardiaceae	<i>Rhus succedanea</i>	Flavones, aldehydes
Annonaceae	<i>Annona chrimola</i> , <i>A. muricata</i> , <i>A. senegalensis</i> ,	Annonaceous acetogenins
	<i>A. reticulate</i> , <i>A. squamosa</i> , <i>A. bullata</i>	(lactones)
	<i>Goniothalamus gardneri</i> , <i>G. amuyon</i> and <i>G. giganteus</i>	lactones
	<i>Polyathia barnesii</i>	Clerodane diterpenes
	<i>Xylopia aromatica</i>	Annonaceous acetogenins
Apiaceae	<i>Anthriscus sylvestris</i>	Lignans
	<i>Seseli mairei</i>	Polyacetylenes
Apocynaceae	<i>Alstonia scholaris</i> R. BR.	Extract, indole alkaloids
	<i>Ervatamia divaricata</i> , <i>E. microphylla</i> , <i>E. heyneana</i>	Alkaloids
	<i>Plumeria rubra</i>	Alkaloids (iridoids), lignans
Araliaceae	<i>Dendropanax arboreus</i>	Oxylipins (linoleic acid derivatives)
	<i>Panax ginseng</i> , <i>P. quinquefolius</i> , <i>P. vietnamensis</i>	Saponins, polysaccharides, polyacetylenic alcohols

Aristolochiaceae	<i>Aristolochia elegans</i> , <i>A. versicolor</i>	Sesquiterpene lactones
Asclepiadaceae	<i>Calotropis procera</i> , <i>C. gigantea</i>	Glycosides
Asteraceae	<i>Neurolaena lobata</i>	Sesquiterpene lactones
Bigoniaceae	<i>Kigelia pinnata</i>	Dichloromethane extracts
Burseraceae	<i>Bursera simaruba</i> , <i>B. permollis</i> , <i>B. morelensis</i> , <i>B. microphylla</i> , <i>B. klugii</i> , <i>B. schlechtendalii</i>	Lignans
Caesalpinaceae	<i>Caesalpinia sappan</i>	Ethyl acetate extracts
Campanulaceae	<i>Platycodon grandiflorum</i>	Polysaccharides
Capparaceae	<i>Polanisia dodecandra</i>	Flavonols
Celastraceae	<i>Glyptopetalum sclerocarpum</i>	Terpenoids
	<i>Maytenus boaria</i> , <i>M. guangsiensis</i> , <i>M. ovatus</i> , <i>M. senegalensis</i> , <i>M. wallichiana</i> and <i>M. emarginata</i>	Triterpenes
Combretaceae	<i>Bucida buceras</i>	Flavanones, diterpenes
	<i>Terminalia arjuna</i>	Flavones, tannins
Compositae	<i>Eupatorium cannabinum</i>	Lactones
	<i>E. cuneifolium</i> , <i>E. rotundifolium</i> , <i>E. semiserratum</i> , <i>E. altissimum</i>	Flavones
	<i>Hellenium microcephalum</i> , <i>H. hoopesii</i>	Sesquiterpene lactones
	<i>Xanthium strumarium</i>	Alkaloids
	<i>Inula viscosa</i>	Flvonoids and alkaloids
	<i>Inula graveolens</i> , <i>Achillea santolina</i> , <i>Conyza canadiensis</i>	Ethanol extract
Conifereae	<i>J. virginiana</i> , <i>J. chinensis</i>	Podophyllotoxin (lignans)
Cucurbitaceae	<i>Cucurbita moschata</i> , <i>Mormodica charantia</i>	Proteins
Cupressaceae	<i>Chamaecyparis lawsonianna</i> , <i>Thujopsis dolabrata</i>	Alkaloids
Ericaceae	<i>Vaccinium macrocarpon</i> , <i>V. smallii</i>	Triterpenes, flavonol, glycosides
Eriocaulaceae	<i>Paepalanthus latipes</i>	Naphthoquinones
Euphorbiaceae	<i>Emblica officinalis</i>	Aqueous extracts, Norsesquiterpenoid, glycosides
	<i>Euphorbia amygdaloides</i> , <i>E. helioscopic</i> , <i>E. lathyris</i> , <i>E. mongolica</i> , <i>E. pubescens</i>	Jatrophone diterpenoids
	<i>Jatrophia curcas</i> , <i>J. macrorhiza</i>	Jatrophone diterpenoids and triterpenoids
	<i>Mallotus phillipinensis</i>	Rottlerin, phloroglucinol

		derivatives
	<i>Phyllanthus acuminatus, P. amarus, P. emblica, P. urinaria</i>	Glycosides
Fabaceae	<i>Ononis sicula, Ononis hirta</i>	Aqueous methanol extract
Flacourtiaceae	<i>Caesaria sylvestris</i>	Clerodane diterpenes
Guttiferae	<i>Garcinia hunburyi</i>	Processed extract
Hernandiaceae	<i>Hernandia</i> sp.	Lignans
Hyacinthaceae	<i>Scilla scilloides, S. peruviana</i>	Glycosides
Hypericaceae	<i>Hypericum perforatum, H. drummondii</i>	Polycyclic diones
Iridaceae	<i>Crocus sativus</i>	Carotenoids
Lamiaceae	<i>Hyptis martiusii, H. verticillata</i>	Diterpenes, lignans
	<i>Origanum vulgare, O. majorana</i>	Quinines, quinine glycosides
	<i>Rabdosia ternifolia, R. trichocarpa, R. macrophylla</i>	Diterpenoids, lactones
	<i>Salvia sclarea</i>	Lectins, Ursolic acid (carboxylic acids)
	<i>Salvia pinardi</i>	Aqueous methanol extract
	<i>Salvia przewalskii</i>	Quinones
	<i>Salvia dominica</i>	Ethanol extract
	<i>Scutellaria barbata, S. lateriflora, S. baicalensis</i>	Flavonoids, flavones
Lauraceae	<i>Cinnanomum camphora</i>	Cinnamaldehydes
Leguminosae	<i>Acacia catechu, A. victoriae, A. confuse, A. auriculiformis A. Cunn. and A. nilotica</i>	Proteins
	<i>Bauhinia racemosa</i>	Methanol extracts
	<i>Cassia acutifolia, C. angustifolia, C. torosa, C. leptophylla</i>	Anathracenes, polysaccharides, piperidin (alkaloids)
	<i>C. pudibunda Glycyrrhiza glabra, G. uralensis, G. inflata</i>	Anthraquinones Glycyrrhizic acid, glycyrrhetinic acid, flavonoids and
	<i>Tamarindus indica</i>	Triterpenoids, Polysaccharides
Liliaceae	<i>Colchicum autumnale, C. speciosum Crinum asiaticum var. toxicarium</i>	Alkaloids
Linaceae	<i>Linum usitatissimum</i>	Lignans
	<i>Viscum cruciatum</i>	Hexanoic acid extracts
Magnoliaceae	<i>M. officinalis, M. grandiflora, M. virginiana</i>	Neolignans
Malvaceae	<i>G. herbaceum G., indicum</i>	Cathechin (phenolics)

Meliaceae	<i>Azadirachta indica</i>	Limonoids (triterpenes)
	<i>Melia azedarach</i>	Limonoids (triterpenes)
Myrtaceae	<i>Eugenia jambos</i>	Hydrolyzable tannins
Ochnaceae	<i>Ouratea hexasperma</i> , <i>O. semiserrata</i>	Biflavonoids
Pinaceae	<i>Pseudolarix kaempferi</i>	Triterpene lactones, diterpenes
Polygalaceae	<i>Polygala vulgaris</i>	Xanthones
Polygonaceae	<i>Polygonum cuspidatum</i>	Flavonoids, anthraquinones
Ranunculaceae	<i>Nigella sativa</i>	Quinones, fatty acids
	<i>Pulsatilla koreana</i>	Lignans, saponins
Rosaceae	<i>Agrimonia pilosa</i>	Tannins, methanolic extracts
Rubiaceae	<i>Nauclea orientalis</i>	Angustine alkaloids
	<i>Psychotria. rubra</i> , <i>P. forsteriana</i>	Napthoquinones, alkaloids
	<i>Rubia cordifolia</i> , <i>R. akane</i>	Cyclic hexapeptides Naphthoquinones, anthraquinones
Rutaceae	<i>Aegle marmelos</i> Correa Fagara <i>macrophylla</i>	Hydroalcoholic extract Alkaloids
	<i>Acronychia oblonga</i> , <i>A. porteri</i> , <i>A. pedunculata</i> , <i>A. Baueri</i>	Flavonols, alkaloids
	<i>Zieridium pseudobtusifolium</i>	Flavonols
Sapindaceae	<i>Koelreuteria henryi</i>	Anthraquinone, stilbene, and flavonoids
Simaroubaceae	<i>Brucea dysenterica</i> , <i>B. javanica</i>	Quassinoid glycosides, Alkaloids
	<i>Eurycoma harmadiana</i>	Alkaloids, quassinoids
	<i>Hannoa chlorantha</i> , <i>H. kleineana</i>	Quassinoids, chaparrinones
Solanaceae	<i>Solanum pseudocapsicum</i>	Alkaloids
Scrophulariaceae	<i>Verbascum sinaiticum</i>	Aqueous methanol extract
Theaceae	<i>Camellia sinensis</i>	Polyphenols
Thymelaceae	<i>Stellera chasmaejasme</i>	Diterpenes
Tropaeolaceae	<i>Wikstroemia foetida</i> , <i>W. uvaurisi</i> , <i>W. indica</i> , <i>Tropaeloum majus</i>	Polysaccharides, Aromatic plant hormones
Umbellifereae	<i>Angelica archangelica</i> <i>A. keiskei</i> , <i>A. sinensis</i> , <i>A. gigas</i> , <i>A. acutiloba</i> , <i>A. radix</i> , <i>A. japonika</i> , <i>A. edulis</i>	Pyranocoumarins, chalcones, polysaccharides
Urticaceae	<i>Ficus carica</i>	Lectins
Verbenaceae	<i>Vitex rotundifolia</i>	Flavonoids
Violaceae	<i>Viola odorata</i>	Nucleotides
References: (Kintzios, 2006), (Abu-Dahab and Afifi, 2007), (Talib and Mahasneh, 2010)		

Table 2. Plants with potential anticancer activity

6. Plants as a source of antimicrobial agents

The discovery of antibiotics has decreased the spread and severity of a wide variety of inferior diseases. However, and as a result of their uncontrolled use, the efficiency of many antibiotics is being threatened by the emergence of microbial resistance to existing chemotherapeutic agents (Cowan, 1999). While bioactive natural compounds have been isolated mainly from cultivable microbial strains, an untapped biologically active metabolites of different resources including plants remains to be investigated (Quiroga et al., 2001) to alleviate or help responding to current health care situations; such situations include but not limited to unmet clinical needs, increasing cost of chemotherapy, mycobacterial reemergence, and the emergence of antibiotic resistant microbial strains such as MRSA (Alanis et al., 2005) .

Microbial resistance occurs mainly in three general mechanisms: prevention of interaction of the drug with target; direct destruction or modification of the drug; and efflux of the drug from the cell (Alviano & Alviano, 2009). These mechanisms were used by different microorganisms and led to the emergence of many pathogenic bacterial strains (Alanis et al., 2005). With pathogenic fungi, the situation is not so bright also, where Amphotericin B was for many years the only treatment available for fungal infections. In late 1980s fluconazole and itraconazole was developed as additional therapeutic options (Ficker et al., 2002). Recently, azole derivatives are most widely used antifungal agents, although resistance for these drugs is emerging (Groll et al., 1998). All the available antifungal drugs used to date are not ideal in efficiency, safety, and antifungal spectrum (Di Domenico, 1998). Combination antifungal therapy was also used to increase the efficiency but there is a real demand for a next generation of safer and more powerful antifungal agents (Bartoli et al., 1998). Knowing that modifying known antimicrobial compounds is increasingly difficult created an urgent and very pressing need for isolation and identification of new bioactive chemicals from new sources including plants (Barker, 2006).

Plant derived natural products represent an attractive source of antimicrobial agents since they are natural, have manageable side effects and available at affordable prices (Ghosh et al., 2008). Also plants derived agents may have different mechanisms than conventional drugs, and could be of clinical importance in health care improvement (Ellof, 1998).

There are two main classes of plant derived agents. 1) phytoalexins which are low molecular weight compounds produced in response to microbial, herbivorous, or environmental stimuli (Van Etten et al., 1994). Phytoalexins include simple phenylpropanoid derivatives, flavonoids, isoflavonoids, terpenes and polyketides (Grayer & Harborne, 1994). 2) Phytoanticipins which are produced in plants before infection or from pre-existing compounds after infection (Van Etten et al., 1994). Phytoanticipins include: glycosides, glucosinolates and saponins that are normally stored in the vacuoles of plant cells (Osborn, 1996). The antimicrobial potential of plant derived natural products is well documented. Schelz and colleagues reported the potential of menthol isolated from peppermint oil to eliminate the resistance plasmids of bacteria (Schelz et al., 2006). In another study carbazole alkaloids isolated from *Clausena anisata* stem bark showed high antibacterial and antifungal activities (Chakraborty et al., 2003).

Thousands of other phytochemicals having *in vitro* antimicrobial activities were also screened. Such screening programs are essential for validating the traditional use of

medicinal plants and for providing leads in the search for new antimicrobial agents (Alviano & Alviano, 2009). A number of studies of the bioactivity of plant extracts have been conducted and many of these studies showed promising results in developing new biologically active agents. The methanolic extract of clove *Caryophyllus aromaticus* showed antibacterial activity against many bacterial genera and the highest activity was against *Staphylococcus aureus* (Ushimaru et al., 2007). Association of clove extract and antibiotics showed synergistic antibacterial activity against antibiotic resistant *Pseudomonas aeruginosa* (Nascimento et al., 2000). Out of the 50 plants used in Indian traditional medicine, 72% showed antimicrobial activity including nine plants showed antifungal activity (Srivnivasan et al., 2001). When some Palestinian plants were tested for bioactivity, out of fifteen used in traditional medicine, only eight showed antibacterial activity against eight different bacterial strains (Essawi & Srour, 2000). Butanol extracts of *Rosa damascene*, *Narcissus tazetta*, and *Inula viscose* exhibited potent antimicrobial activities against different microorganism including Methicillin-resistant *Staphylococcus aureus* and *Candida albicans* (Talib & Mahasneh, 2010b).

During the past decade there is an increase in the number of immuno-compromised patients. This is probably due to the alteration of the immune system caused by human immunodeficiency virus (HIV), cancer chemotherapy, and organ and bone marrow transplantation in addition to the use of immune suppressors to treat many diseases (Alexander & Perfect, 1997). The compromised immune system facilitates microbial infections including systemic mycosis. This leads to extensive use of Amphotericin B and azole derivatives as antifungal agents. Unfortunately, the wide range use of these antifungal agents leads to the emergence of drug resistant pathogenic fungi (Alexander & Perfect, 1997). *Candida albicans* is opportunistic yeast that can cause vaginal, oral, and lung infections in addition to systemic tissue damage in AIDS patients (Madigan & Martinko, 2006). This yeast was the target of many researchers to develop new antifungal agents. Out of twenty four medicinal plants used in traditional medicine in South Africa, two showed high potential to treat candidiasis (Motsei et al., 2003). Also some indigenous plants of Lebanon showed antimicrobial activity against *Candida albicans* and other tested microorganisms (Barbour et al., 2004).

Successive isolation of antimicrobial compounds from plants depends upon the type of solvent used in extraction procedure (Parekh & Chanda, 2007). Literature reported the use of different solvents to extract antimicrobial agents. The ethanol/methanol extracts of thirty four medicinal plants were more active than aqueous extracts against bacterial strains belonging to *Enterobacteriaceae* (Parekh & Chanda, 2007). Methanolic extract of *Terminallia chebula* showed higher antibacterial potential compared with aqueous extract (Ghosh et al., 2008). Mahasneh found that butanol extracts of several plants including: *Lotus halophilus*, *Pulicaria gnaphaloides*, *Capparis spinosa*, *Medicago laciniata*, *Limonium axillare* to exhibit superior antimicrobial activity compared with ethanol and aqueous extracts (Mahasneh, 2002). The ethanolic extract of 11 plant species from Argentina showed high antimicrobial activity against a list of microorganisms including methicillin, oxacillin, and gentamicin resistant *Staphylococcus* (Zampini et al., 2009). Of 16 plants studied, the methanolic and aqueous extracts of 10 Yemeni plants exhibited significant antimicrobial activity against three gram positive, two gram negative bacteria, and one fungus (Mothana et al., 2009).

Jordanian plants were the focus of many researchers for their antimicrobial activities. Butanol, ethanol, petroleum ether, and aqueous extracts were prepared from nine Jordanian plants. Butanol extract showed superior antimicrobial activity compared with other extracts (Mahasneh & El-Oqlah, 1999). Of 27 ethanol extracts prepared from indigenous Jordanian plants, six plants showed promising antimicrobial activity against different test microorganisms (Al-Bakri & Afifi, 2007). In addition to their broad antimicrobial activities some Jordanian plants like *Sonchus oleraceus* and *Laurus nobilis* exhibited high anti-quorum sensing activities (Al-Hussaini & Mahasneh, 2009). Other Jordanian plants like *Crupina crupinastrum* and *Achillea biebersteinii* showed high antimicrobial activity against bacteria and fungi (Khalil et al., 2009). Additionally the methanolic extracts of two Jordanian plants *Artemisia herba-alba* and *Artemisia arborescens* showed high antibacterial activity against 32 isolates of *Mycoplasma* species (Al-Momani et al., 2007).

7. Conclusions

The potential isolation and use of new and novel bioactive products from plant origins is still very productive playground for the development of new drugs to improve health care in certain medical fields. It is essential to emphasize that extensive *in vitro* and *in vivo* tests must be conducted to assure the selection of active and nontoxic anticancer and antimicrobial phytochemicals.

8. References

- Abu-Dahab, R. & Afifi, F. (2007). Antiproliferative activity of selected medicinal plants of Jordan against a breast adenocarcinoma cell line (MCF7). *Scientia Pharmaceutica*, 75, 121-136.
- Abu-Irmaileh B. & Afifi F. (2003). Herbal medicine in Jordan with special emphasis on commonly used herbs. *Journal of Ethnopharmacology*, 89, 193-197.
- Aburjai, T., Mohammad, H., Rabab, T., Mohammed, Y., & Maher, Q. (2007). Ethnopharmacological survey of medicinal herbs in Jordan, the Ajloun heights region. *Journal of Ethnopharmacology*, 110, 294-304.
- Alanis, L. (2005) Resistance to antibiotics: are we in the post-antibiotic era?. *Archives in Medical Research*, 36, 697-705.
- Al-Bakri, A. & Afifi, F. (2007). Evaluation of antimicrobial activity of selected plant extracts by rapid XTT colorimetry and bacterial enumeration. *Journal of Microbiological Methods*, 68 (1), 19-25
- Alexander, B. & Perfect, J. (1997). Antifungal resistance trends toward the year 2000, implication for therapy and new approaches. *Drugs*, 54, 657-678.
- AL-Hussaini, R. & Mahasneh A. (2009). Microbial growth and quorum sensing antagonist activities of herbal plant extracts. *Molecules*, 14, 3425-3435.
- Al-Momani, W., Abu-Basha, E., Janakat, S., Nicholas, R. & Ayling, R. (2007). In vitro antimycoplasmal activity of six Jordanian medicinal plants against three *Mycoplasma* species. *Tropical Animal Health and Production*, 39, (7), 515-519
- Al-Qura'n, S. (2009). Ethnopharmacological survey of wild medicinal plants in Showbak, Jordan. *Journal of Ethnopharmacology*, 123, 45-50.

- Alviano, D. & Alviano, A. (2009). Plant extracts: search for new alternative to treat microbial diseases. *Current Pharmaceutical Biotechnology*, 10, 106-121.
- Azaizeh, H., Saad, B., Khalil, K. & Said, O. (2006). The state of the art of traditional Arab herbal medicine in the Eastern region of the Mediterranean: A Review. *Evidence-based Complementary and Alternative Medicine*, 3 (2), 229-235.
- Azaizeh, H., Fulder, S., Khalil, K., & Said, O. (2003). Ethnobotanical survey of local practitioners of the Middle Eastern region: the status of traditional Arabic medicine. *Fitoterapia*, 74, 98-108.
- Barbour, K., Al Sharif, M., Sagherian V., Habre, A., Talhouk, R. & Talhouk, S. (2004). Screening of selected indigenous plants of Lebanon for antimicrobial activity. *Journal of Ethnopharmacology*, 93, 1-7.
- Barker, J. (2006). Antibacterial drug discovery and structure-based design. *Drug Discovery Today*, 11, 391-404
- Bartoli J., Turmo, E., Alguero, M., Boncompte, E., Vericat, M., Conte, L., Ramis, J., Merlos, M., Garcia-Rafanell, J. & Forn, J. (1998). New azole antifungals. 3. Synthesis and antifungal activity of 3-substituted-4(3H)-quinazolinone derivatives of 3-amino-2-aryl-1-azoly-2-butanol. *Journal of Medicinal Chemistry*, 41, 1869-1882.
- Boik, J. (2001). *Natural Compounds in Cancer Therapy* (first edition), Oregon Medical Press, USA
- Brown, J., & Wilson, W. (2004). Exploiting tumor hypoxia in cancer treatment. *Nature Review/Cancer*, 4, 437-447.
- Chakrabarty, M. (2003). Microorganisms and cancer: quest for therapy. *Journal of Bacteriology*, 185 (9), 2683-2686
- Chang, K., Kung, M., Chow, N., & Su, S. (2004). Genistein arrests hepatoma cells at G2/M phase: involvement of ATM activation and upregulation of p21waf/cip1 and wee1. *Biochemical Pharmacology*, 67, 717-726
- Chen, S., Robey, R., Belinsky, M., Shchaveleva, I., Ren, X., Sugimoto, Y., Ross, D., Bates, S., & Kruh, G. (2003). Transport of methotrexate, methotrexate polyglutamates, and 17beta-estradiol 17-(beta-D-glucuronide) by ABCG2: effects of acquired mutations at R482 on methotrexate transport. *Cancer Research*, 63, 4048-4054.
- Chitme, H., Chandra, R., & Kaushik, S. (2003). Studies on anti-diarrheal activity of *Calotropis Gigantea* R. Br. in experimental animals. *Journal of Pharmacy and Pharmaceutical sciences*, 7, 70-75.
- Cordell, G. (2002). Natural products in drug discovery- creating a new vision, *Phytochemistry Reviews*, 1 (3), 261-273.
- Cowan, M. (1999), Plant products as antimicrobial agents. *Clinical Microbiology Reviews*, 12, 564-582.
- Dai, Y. & Grant, S. (2003). Cyclin-dependent kinase inhibitors. *Current Opinion in Pharmacology*, 3, 362-370.
- Dang, H., Bettegowda, C., Huso, D., Kinzler, K., & Vogelstein, B. (2001). Combination bacteriolytic therapy for the treatment of experimental tumors. *Proceeding of the National Academy of Sciences of the United States of America*, 98, 15155-15160.
- Di Domenico, B. (1999). Novel antifungal drugs. *Current Opinion in Microbiology*, 2, 509-515.

- Dixon, R. & Paiva, N. (1995). Stress-induced phenylpropanoid metabolism. *Plant Cell*, 7, 1085-1097.
- Duflos, A., Kruczynski, A., & Barret, J. (2002). Novel aspects of natural and modified vinka alkaloids. *Current Medicinal Chemistry - Anti-Cancer Agents*, 2(1), 55-70.
- Eloff, J. (1998). Which extractant should be used for the screening and isolation of antimicrobial components from plants?, *Journal of Ethnopharmacology*, 60, 1-8.
- Essawi, T. & Srouf, M. (2000). Screening of some Palestinian medicinal plants for antibacterial activity, *Journal of Ethnopharmacology*, 70, 343-349.
- Fabricant, D. & Farnsworth, N. (2001). The value of plants used in traditional medicine for drug discovery. *Environmental Health Perspectives*, 109 (1), 69-75.
- Facchini, P. (2001). Alkaloids biosynthesis in plants: biochemistry, cell biology, molecular regulation, and metabolic engineering application. *Annual Review of Plant Physiology and Plant Molecular Biology*, 52, 29-66.
- Ficker, E., Arnason, T., Vindas, S., Alvarez, P., Akpagana, K., & Gbeassor, M. (2005). Inhibition of human pathogenic fungi by ethnobotanically selected plant extracts. *Mycoses*, 46, 29-37.
- Ghosh, A., Das, B., Roy, A., Mandal, B. & Chandra, G. (2008). Antibacterial activity of some medicinal plant extracts. *Journal of Natural Medicines*, 62, 259-262.
- Grayer, R. & Harborne, J. (1994). A survey of antifungal compounds from plants, 1982-1993. *Phytochemistry*, 37, 19-42.
- Griggs, J., Hesketh, R. Smith, G., Brindle, K., Metcalfe, J., Thomas, G., & Williams, E. (2001). Combretastatin-A4 disrupts neovascular development in non-neoplastic tissue. *British Journal of Cancer*, 84(6), 832-835
- Groll, A., Piscitelli, S., & Walsh, T. (1998). Clinical pharmacology of systemic antifungal agents: a comprehensive review of agents in clinical use, current investigational compounds, and putative targets for antifungal drug development. *Advances in Pharmacology*, 44, 343-500.
- Hirano, K., Budiyo, E., Swastika, N., & Fujii, K. (1995). Population dynamics of the whitefly, *Bemisia tabaci*(Gennadius) (Homoptera: Aleyrodidae), in Java, Indonesia, with special reference to spatio-temporal changes in the quantify of food Resources. *Ecological Research*, 10, 75-85.
- Horwitz, S., Cohen, D., Rao, S., Ringel, I., Shen, H., & Yang, C. (1993). Taxol: mechanisms of action and resistance. *Journal of the National Cancer Institute Monographs*, 15, 55-61.
- Joshi, B., Li, L., Taffe, B., Zhu, Z., Wahl, S., Tian, H., Ben-Josef, E., Taylor, J., Porter, A., & Tang, D. (1999). Apoptosis induction by a novel anti-prostate cancer compound, BMD188 (a fatty acid containing hydroxamic acid), requires the mitochondrial respiratory chain. *Cancer Research*, 59, 4343-4355.
- Karp G. (1999). *Cell and Molecular Biology* (second edition), Wiley, USA
- Khalil, A., Dababneh, B., & Al-Gabbiesh, A. (2009). Antimicrobial activity against pathogenic microorganisms by extracts from herbal Jordanian plants. *Journal of Food, Agriculture and Environment*, 7 (2), 103-106.
- Kim, H. (2005). Do not put too much value on conventional medicines. *Journal of Ethnopharmacology*, 100, 37-39.

- Kintzios, E. (2006). Terrestrial plant-derived anticancer agents and plant species used in anticancer research. *Critical Reviews in Plant Science*, 25, 79–113.
- Kozawa, M., Baba, K., & Matsuyama, Y. (1982). Components of the root of *Anthriscus sylvestris* Hoffm.2. insecticidal activity. *Chemical and Pharmaceutical Bulletin*, 30, 2885–2888.
- Lee, K. (1999). Anticancer drug design based on plant-derived natural products. *Journal of Biomedical Science*, 6, 236–350.
- Leong, O., Muhammad, T. & Sulaiman, S. (2011). Cytotoxic activities of *Physalis minima* L. chloroform extract on human lung adenocarcinoma NCI-H23 cell lines by induction of apoptosis. *Evidence-based Complementary and Alternative Medicine*, doi:10.1093/ecam/nep057
- Li, W., Chan S., Guo, D., Chung M., Leung, T., & Yu, P. (2009). Water extract of *Rheum officinale* Baill. induces apoptosis in human lung adenocarcinoma A549 and human breast cancer MCF-7 cell lines. *Journal of Ethnopharmacology*, 124, 251–256.
- Littlewood, A., Maguire, H., & Bulmahn J. (2002). Byzantine Garden Culture, *Speculum*, 79 (1), 232–235.
- Madhuri, S., & Pandey, G. (2009). some anticancer medicinal plants of foreign origin. *Current Science*, 96 (6), 779–782.
- Madigan, T., Martinko, J., & Parker, J. (2006). *Biology of Microorganisms* (eleventh Edition), Pearson, USA.
- Mahasneh, A. (2002). Screening of some indigenous Qatari medicinal plants for antimicrobial activity. *Phytotherapy Research*, 16(8), 751–3.
- Mahasneh, A. & El-Oqlah, A. (1999). Antimicrobial activity of extracts of herbal plants used in the traditional medicine in Jordan. *Journal of Ethnopharmacology*, 64, 271–276.
- Mahoney, B., Baggett, B. & Gillies, R. (2003). Tumor acidity, ion trapping and chemotherapeutics. I. Acid pH affects the distribution of chemotherapeutic agents *in vitro*. *Biochemical Pharmacology*, 66, 1207–1118.
- Mesquita, M., dePaula, J., Pessoa, C., Moraes, M., Costa-lotufo, L., Grougnet, R., Michel, S., Tillequin, F., & Espindola, L. (2009). Cytotoxic activity of Brazilian Cerrado plants used in traditional medicine against cancer cell lines. *Journal of Ethnopharmacology*, 123, 439–445.
- Mothana, R., Lindequist, U., Gruenert, R., & Bednarski, P. (2009). Studies of the *in vitro* anticancer, antimicrobial and antioxidant potentials of selected Yemeni medicinal plants from the island Soqatra. *BMC Complementary and Alternative Medicine*, 9, record number 7.
- Motsei, M., Lindsey, J., van Staden & Jager, A. (2003). Screening of Traditionally used South African Plants for Antifungal Activity against *Candida albicans*. *Journal of Ethnopharmacology*, (86), 235–241.
- Mukherjee, A., Basu, S., Sarkar, N., & Ghosh, A. (2001). Advances in cancer therapy with plant based natural products, *Current medicinal chemistry*, 8, 1467–1486
- Nagamine, M., Silva, T., Matsuzaki, P., Pinello, K., Cogliati, B., Pizzo, C., Akisue, G., Haraguchi, M., Górniak, S., Sinhörini, I., Rao, K., Barbuto J. & Dagli, M. (2009). Cytotoxic effects of butanolic extract from *Pfaffia paniculata* (Brazilian Ginseng) on

- cultured human breast cancer cell line MCF-7. *Experimental and Toxicologic Pathology*, 61, 75-82.
- Nascimento F., Gilson, J., Paulo, F. & Giuliana, S. (2000). Antibacterial activity of plant extracts and phytochemicals on antibiotic-resistant bacteria. *Brazilian Journal of Microbiology*, 31, 247-256.
- Neergheen, V., Bahorun, T., Taylor, E., Jen, L., & Aruoma, O. (2009). Targeting specific cell signaling transduction pathways by dietary and medicinal phytochemicals in cancer chemoprevention, *Toxicology*, 278, 229-241
- Newman, D., Cragg, G., & Snader, K. (2003). Natural products as sources of new drugs over the period 1981- 2002. *Journal of Natural Products*, 66 (7), 1022-1037.
- Nobili, S., Lippi, D., Witort, E., Donnini, M., Bausi, L., Mini, E., & Capaccioli, S. (2009). Natural compounds for cancer treatment and prevention. *Pharmacological Research*, 59, 365-378.
- Onetto, N., Canetta, R., Winograd, B., Catane, R., Dougan, M., Grechko, J., Burroughs, J., & Rozencweig, M. (1993). Overview of Taxol safety. *Journal of the National Cancer Institute Monographs*, 15,131-9.
- Osbourn, A. (1996). Preformed antimicrobial compounds and plant defense against fungal attack. *The Plant Cell*, 8, 1821-1831.
- Ozmen, A., Madlener, S., Bauer, S., Krasteva, S., Vonach, C., Giessrigl, B., Gridling, M., Viola, K., Stark, N., Saiko, P., Michel, B., Fritzer-Szekeres, M., Szekeres, T., Askin-Celik, T., Krenn, L. & Krupitza, G. (2010). *In vitro* anti-leukemic activity of the ethnopharmacological plant *Scutellaria orientalis* ssp. *carica* endemic to western Turkey. *Phytomedicine*, 17, 55-62.
- Palombo, E. (2009). Traditional medicinal plant extracts and natural products with activity against oral bacteria: potential application in the prevention and treatment of oral diseases. *Evidence-based Complementary and Alternative Medicine*, doi:10.1093/ecam/nep067.
- Pareke, J. & Chanda, S. (2007). *In vitro* screening of antibacterial activity of aqueous and alcoholic extracts of various Indian plant species against selected pathogens from Enterobacteriaceae. *African Journal of Microbiology Research*, 6, 92-99.
- Quiroga, E., Sampietro, A. & Vattuone, A. (2001). Screening antifungal activities of selected medicinal plants, *Journal of Ethnopharmacology*, 74, 89-96.
- Raskin, I., Ribnicky, D., Komarnytsky, S., Ilic, N., Poulev, A., Borisjuk, N., Brinker, A., Moreno, A., Ripoll, C., Yakoby, N., Cornwell, T., Pastor, I. & Fridlender, B. (2002). Plants and human health in the twenty-first century, *Trends In Biotechnology*, 20 (12), 522-531.
- Reed, J. & Pellecchia, M. (2005). Apoptosis-based therapies for hematologic malignancies, *Blood*, 106, 408-418.
- Saad, B., Azaizeh, H. & Said, O. (2006). Tradition and perspectives of Arab herbal medicine: A Review. *Evidence-based Complementary and Alternative Medicine*, 2, 475-479.
- Said, O., Khalil, K., Fulder, S., & Azaizeh, H. (2002). Ethnopharmacological survey of medicinal herbs in Israel, the Golan Heights and the West Bank region. *Journal of Ethnopharmacology*, 83, 251-165.

- Sakarkar, D. & Deshmukh, V. (2011). Ethnopharmacological review of traditional medicinal plants for anticancer activity. *International Journal of Pharma Tech Research*, 3, 298-308.
- Schelz, Z., Molnar, J., & Hohmann, J. (2006). Antimicrobial and antiplasmodic activities of essential oils, *Fitoterapia*, 77(4):279-285.
- Souza, G., Hass, A., Poser, G., Schapoval, E., & Elisabetsky, E. (2004). Ethnopharmacological studies of antimicrobial remedies in the south of Brazil. *Journal of Ethnopharmacology*, 90 (1), 135-143.
- Srinivasan, D., Sangeetha, N., Suresh, T. & Perumalsamy, P. (2001). Antimicrobial activity of certain Indian medicinal plants used in folkloric medicine. *Journal of Ethnopharmacology*, 74, 217-220.
- Talib, W. & Mahasneh A. (2010a). Antiproliferative activity of plant extracts used against cancer in traditional medicine. *Scientia Pharmaceutica*, 78: 33-45
- Talib, W. & Mahasneh, A. (2010b). Antimicrobial, Cytotoxicity and Phytochemical Screening of Jordanian Plants Used in Traditional Medicine. *Molecules*, 15, 1811-1824.
- Tannock, F. (1998). Conventional cancer therapy: promise broken or promise delayed?. *Lancet*, 352, 9-16.
- Taraphdar, A., Madhumita, R. & Bhattacharya, R. (2001). Natural products as inducers of apoptosis: Implication for cancer therapy and prevention. *Current Science*, 80 (11), 1387-1397.
- Trapp, S. & Croteau, R. (2001). Defensive resin biosynthesis in conifers. *Annual Review of Plant Physiology and Plant Molecular Biology*, 52, 689-724.
- Ushimaru, P., da Silva, M., Di Stasi, L., Barbosa, L. & Junior, A. (2007). Antibacterial activity of medicinal plant extracts, *Brazilian Journal of Microbiology*, 38:717-719
- Van Etten, H., Mansfield, W., Bailey, J. & Farmer, E. (1994). Two classes of plant antibiotics: phytoalexins versus phytoanticipins. *Plant Cell*, 6, 1191-1192.
- Verpoorte, R. (2000). Pharmacognosy in the new millenium: leadfinding and biotechnology. *Journal of Pharmacy and Pharmacology*, 52, 253-262.
- Wall, M., Wani, M., Cook, C., Palmer, K., McPhail, A. & Sim, G. (1966). Plant antitumor agents. 1. The isolation and structure of camptothecin, a novel alkaloidal leukemia and tumor inhibitor from *Camptotheca acuminata*. *Journal of the American Chemical Society*, 88, 3888-3890.
- Wani, M., Taylor, H., Wall, M., Coggan, P & McPhail, A. (1971). Plant antitumor agents. VI. the isolation and structure of Taxol, a novel antileukemic and antitumor agent from *Taxus brevifolia*. *Journal of the American Chemical Society*, 93, 2325-2327.
- Wood-Sheldon, J., Balick, M. & Laird, S. (1997). *Medicinal Plants: Can Utilization and Conservation Coexist?*. The New York Botanical Garden. USA.
- Xu, J., Liu, X., Zhou, S. & Wei, M. (2009). Combination of immunotherapy with anaerobic bacteria for immunogene therapy of solid tumours. *Gene Therapy and Molecular Biology*, 13, 36-52.
- Young, S. & Chaplin, D. (2004). Combretastatin A4 phosphate: background and current clinical status. *Expert Opinion on Investigational Drugs*, 13(9), 1171-1182.
- Yun-San Yip, A., Yuen-Yuen Ong, E. & Chow, C. (2008). Vinflunine: clinical perspectives of an emerging anticancer agent, *Expert Opinion on Investigational Drugs*, 17, 583-591.

Zampini, I., Cuello, S., Alberto, M., Ordoñez, R., Almeida, R., Solorzano, E. & Isla, M. (2009). Antimicrobial activity of selected plant species from "the Argentine Puna" against sensitive and multi-resistant bacteria. *Journal of Ethnopharmacology*, 124(3):499-505.

IntechOpen

IntechOpen

Phytochemicals - Bioactivities and Impact on Health

Edited by Prof. Iraj Rasooli

ISBN 978-953-307-424-5

Hard cover, 388 pages

Publisher InTech

Published online 22, December, 2011

Published in print edition December, 2011

Among the thousands of naturally occurring constituents so far identified in plants and exhibiting a long history of safe use, there are none that pose - or reasonably might be expected to pose - a significant risk to human health at current low levels of intake when used as flavoring substances. Due to their natural origin, environmental and genetic factors will influence the chemical composition of the plant essential oils. Factors such as species and subspecies, geographical location, harvest time, plant part used and method of isolation all affect chemical composition of the crude material separated from the plant. The screening of plant extracts and natural products for antioxidative and antimicrobial activity has revealed the potential of higher plants as a source of new agents, to serve the processing of natural products.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Wamidh Hadi Talib (2011). Anticancer and Antimicrobial Potential of Plant-Derived Natural Products, *Phytochemicals - Bioactivities and Impact on Health*, Prof. Iraj Rasooli (Ed.), ISBN: 978-953-307-424-5, InTech, Available from: <http://www.intechopen.com/books/phytochemicals-bioactivities-and-impact-on-health/anticancer-and-antimicrobial-potential-of-plant-derived-natural-products>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2011 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](https://creativecommons.org/licenses/by/3.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen