

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

185,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.  
For more information visit [www.intechopen.com](http://www.intechopen.com)



## EMG in People with Different Heel Height Condition

Xiaoxiang Su<sup>1</sup> and Yaodong Gu<sup>2</sup>

<sup>1</sup>*School of Engineering, Liverpool John Moores University, Liverpool*

<sup>2</sup>*Faculty of Sport Science, Ningbo University, Ningbo*

<sup>1</sup>UK

<sup>2</sup>China

### 1. Introduction

It has been reported that a prevalence proportion of foot problems in women were associated with wearing high-heeled shoes [1]. Women's walking pattern and the biomechanical behavior of the foot could be altered significantly by raising the heel height [2, 3]. Biomechanical studies showed that walking in high-heeled shoes may alter lower-extremity joint function [4], raise the peak pressure in the forefoot [5], and alter the load distribution on the media foot region [6]. Some of these biomechanical changes can be detrimental to the foot structure, such as increasing the metabolic cost of gait, accelerate on of muscle fatigue [7], high inner stress level of metatarsal bones [8]. In Latin dancing, partners perform tightly choreographed routines in which they must project energy and passion. The pair's appearance, including costuming and shoes, is important to the success of the routine. A detailed biomechanical study of different types of human movement such as walking, dancing, etc. would help to improve the understanding of the biomechanics and provide important data for design optimization to reduce the adverse effects associated with high heeled shoes.

The detailed diagnosis and correction of pathologic states of human mobility is mainly limited by an incomplete knowledge of the musculoskeletal system and the detailed and quantitative description of the activity of the individual muscles in the synergistic patterns which combine to produce different human movement. Commonly used techniques which employ myoelectric, force plate and kinematic gait pattern acquisition only provide a representation of the external manifestations of the underlying muscle physiology. At present, there is no exist research equipment to study the actions of the involved muscles non-invasively. A model of the musculoskeletal system could, in principle, given kinematic data and with assumptions to mitigate muscle redundancy, predict individual muscle behavior. Surface electromyography (SEMG), has been used in both research and clinical applications for non-invasive neuromuscular assessment in several different fields such as sport science, neurophysiology and rehabilitation. Different variables, related to different aims, can be monitored: muscle activation intervals are useful to evaluate motor coordination and treatment efficacy [9]; myoelectrical manifestations of fatigue [10] can be used to assess EMG signal

modifications in pathologies such as exercise [11], back pain [12] and neurological diseases [13]. Previous EMG study in high heel gait found out the low back EMG increased significantly while walking with high-heeled shoes, which successfully explained the reason why high heel wearers always complained of lumber back pain [14]. But there is very limited study of the high-heeled state of muscle activity in a high intensity movement such as dancing. Female dancers have to wear high-heeled shoes during routine exercise and competitions, while passively raised heel could greatly increased the foot injury probability [5]. The objective of this study is to investigate the lower limb muscles' surface SEMG activity during Latin dancing with different heel height shoes.

## 2. Methods

The study involved 10 professional female dancers to participate in the biomechanical tests and data collection. The mean age of the subjects was 22.8years (S.D. 2.3 years), the mean weight of the subjects was 49.5kg (S.D. 5.2 kg), and the mean height of the subject was 163.6cm (S.D. 2.5 cm). Each participant has more than five years' dancing history. The subjects' feet/lower limb was clinically checked to ensure that there was no recent injury phenomenon or medical conditions that may affect their performances. The shoes used in this study have a similar structure apart from the heel height. Typical dancing conditions included four shoes with a heel height of 1.0cm, 4.5cm, 7.5cm and 10cm, designated as flat, low heel, medium high heel and high heel shoes, respectively.



Fig. 1. The structure of typical dancing shoes.

Typical dancing shoes are made of leather, women's Latin dance shoes are lightweight, flexible and have suede soles, which provide the right blend of grip and slide while moving across the floor (Fig.1). In the study, shoes of different heel heights (1.0cm, 4.5cm, 7.5cm and 10cm) were randomly assigned to the participants. Each participant walked 5 minutes before the test in order to habituate to each heel height shoes, and took a 20-min rest in-between dance to prevent body fatigue. All the subjects were required to perform a similar warm up exercise before the test, and the Latin basic motion *visk* step (basic

crossing step, Fig.2) was selected as experiment motion, concomitance with the dancing music.



Fig. 2. Illustration of *visk* (crossing) step in the latin dancing: RFM (right foot move); LFM (left foot move).

SEMG activities of the tibialis anterior (TA), medial and lateral sides of gastrocnemius (MG and LG), soleus (SO) and biceps femoris (BF) were obtained using self-adhesive pairs of disposable Ag/AgCl surface electrode pairs with an electrode diameter of 10mm and an inter-electrode spacing of 20mm. Low impedance at the skin-electrode interface was obtained by skin preparation procedure of shaving, light abrasion and cleaning. For electrode placement, the SENIAM [15] recommendations for SEMG were followed (Fig.3). Furthermore, the electrode placement was confirmed by palpations of muscle bulk during brief maximal isometric contraction. The raw SEMG signal was recorded at the sampling rate of 1000Hz using a device (ME6000, Mega Electronics, Finland). The average EMG (aEMG) for the five muscles were calculated which using the rectified signal with a 512ms window, and used for further analysis.

Statistical data analysis was conducted with the statistical software package SPSS (version 13.0). Analysis of variance was employed to study the effects of heel height, and Turkey's HSD test was used for post hoc comparison. The level of significance was set at the 5% level.

### 3. Results

Fig.4 illustrated the comparison of aEMG data in five lower limb muscles during dancing under different heel height. It was clearly shown that all the muscles aEMG values except BF were significantly increased in heightened heel shoes comparing to wearing the flat heel height dancing ( $p < 0.05$ ). The largest increasing value of aEMG between flat and low heel height was appeared in LG, which was up to 36%. Differences were not found in LG, MG and SO among low, medium and high heel height.

Meanwhile, it was interesting to find that the aEMG value was quite close between the low and medium heel height in LG. Also, in BF the muscle activity was not changed a lot between the medium and high heel height. Although the mean value of aEMG data increased from low to high heel height, the significant difference only appeared at TA in which the high heel height was larger than the value in both low and medium heel height.




Fig. 3. Electrode placement methods.




Fig. 4. Comparison of aEMG data in five muscles of lower limb during different heel heights dancing (Significant change \* $p < 0.05$ ).

Another interesting finding in these dancing testing subjects, in which 70% are found have a certain extent deformation in the first metatarsal-phalangeal joint (Fig.5, should to further think about the muscles activity to cause joint deformation).



Fig. 5. The certain extent deformation in the first metatarsal-phalangeal joint of the testing subject.

#### 4. Discussion

The results of this study showed that increasing heel height caused increases in SEMG activity in lower limb muscles during dancing. Although previous studies in high heel gait reported similar trend [16, 17], to the authors' knowledge, no studies of heeled dancing movement have evaluated at lower limb muscles activity. It's interesting to find the SEMG



activity of the TA was seemed as the most vulnerable part to the heel height increasing. While the heel height increased to 10cm, the aEMG data of TA significantly higher than both 4.5cm and 7.5cm heel height. Due to the function of TA was to adjust dorsiflexion of the foot, acting as a stabilizer of the ankle, this implies that dancing with high heel shoes would consume more energy to control the balance. The trend of activity of LG increased more versus that of MG in heel height increased from flat to low. This different response of the two sides of the gastrocnemius under different heel height was close to one other study conducted by Gefen et al. [18]. The increased muscle activity would apparently cause higher loads on the shoe's front part and accelerate its damage or worn out that correlates well with failure analysis of dancing shoes after a long time of training (Fig. 6).




Fig. 6. Damage part in shoe's front part after a long time of training.

There are limitations to the current study, which could be developed further. The lack of kinematic and kinetic data means that confounding variables may be present. In addition, all participants who took part in this study were experienced high heel wearers. Muscle response to heel height may difference between regular and non-regular wearers of high-heeled shoes.

## 5. References

- [1] Lee KH, Shieh JC, Matteliano A et al. (1990) Electromyographic changes of leg muscles with heel lifts in women: therapeutic implications. *Arch Phys Med Rehabil* 71: 31-33.
- [2] Rong M, Wang YX, Gu YD (2009) Plantar pressure distribution character in the Latin dance. *Footwear Science* 1: 36-37.
- [3] Speksnijder CM, Munckhof RJ, Moonen SA et al. (2005) The higher the heel the higher the forefoot pressure in ten healthy women. *The foot* 15: 17-21.
- [4] Esenyel M, Walsh K, Walden JG et al. (2003) Kinetics of high-heeled gait. *J Am Podiatr Med Assoc* 93: 27-32.
- [5] Mandato MG, Nester E (1999) The effects of increasing heel height on forefoot peak pressure. *J Am Podiatr Med Assoc* 89: 75-80.
- [6] Eisengardt JR, Cook D, Pregler I et al (1996) Changes in temporal gait characteristics and pressure distribution for bare feet versus heel various heel height. *Gait Posture* 12: 280-286.
- [7] Snow RE, Williams KR, Holmes GB (1992) The effects of wearing high heeled shoes on pedal pressure in women. *Foot Ankle* 13: 85-92.
- [8] Gu YD, Li JS (2005) Finite element analysis of the instep fatigue trauma in the high-heeled gait. *WJMS* 1: 117-122.
- [9] Benedetti M (2001) Muscle activation intervals and EMG envelope in clinical gait analysis. *IEEE Eng Med Biol Mag* 20: 33-34.
- [10] Merletti R, Knaflitz M, De Luca CJ (1990) Myoelectric manifestations of fatigue in voluntary and electrically elicited contractions. *J Appl Physiol* 69:1810-1820.
- [11] Casale R, Rainoldi A, Nilsson J et al (2003) Can continuous physical training counteract aging effect on myoelectric fatigue? A surface electromyography study application. *Arch Phys Med Rehabil* 84:513-517.
- [12] Mannion AF, Dumas GA, Stevenson JM et al (1998) The influence of muscle fiber size and type distribution on electromyographic measures of back muscle fatigability. *Spine* 23:576-584.
- [13] Zwarts MJ, Drost G, Stegeman DF (2000) Recent progress in the diagnostic use of surface EMG for neurological diseases. *J Electromyogr Kinesiol* 10:287-291.
- [14] Lee CM, Jeong EH, Freivalds A (2001) Biomechanical effects of wearing high-heeled shoes *Int J Ind Ergon* 28:321-326.
- [15] Hermens HJ, Freriks B, Merletti R. et al. (1999) *Seniam 8: European recommendations for surface electromyography*. Roessingh Research and Development, Netherlands
- [16] Stefanyshyn DJ, Nigg BM, Fisher V et al. (2000) The influence of high heeled shoes on kinematics, kinetics and muscle EMG of normal female gait. *J Appl Biomech* 16: 309-319.
- [17] Lee CM, Jeong EH, Frievalds A (2001) Biomechanical effects of wearing high-heeled shoes. *Int J Ind Ergon* 28: 321-326.



- [18] Gefen A, Megido-Ravid M, Itzhak Y et al. (2002) Analysis of muscular fatigue and foot stability during high heeled gait. *Gait & Posture* 15: 56-63.

IntechOpen

IntechOpen



## **Applications of EMG in Clinical and Sports Medicine**

Edited by Dr. Catriona Steele

ISBN 978-953-307-798-7

Hard cover, 396 pages

**Publisher** InTech

**Published online** 11, January, 2012

**Published in print edition** January, 2012

This second of two volumes on EMG (Electromyography) covers a wide range of clinical applications, as a complement to the methods discussed in volume 1. Topics range from gait and vibration analysis, through posture and falls prevention, to biofeedback in the treatment of neurologic swallowing impairment. The volume includes sections on back care, sports and performance medicine, gynecology/urology and orofacial function. Authors describe the procedures for their experimental studies with detailed and clear illustrations and references to the literature. The limitations of SEMG measures and methods for careful analysis are discussed. This broad compilation of articles discussing the use of EMG in both clinical and research applications demonstrates the utility of the method as a tool in a wide variety of disciplines and clinical fields.

### **How to reference**

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Xiaoxiang Su and Yaodong Gu (2012). EMG in People with Different Heel Height Condition, Applications of EMG in Clinical and Sports Medicine, Dr. Catriona Steele (Ed.), ISBN: 978-953-307-798-7, InTech, Available from: <http://www.intechopen.com/books/applications-of-emg-in-clinical-and-sports-medicine/emg-in-people-with-different-heel-height-condition>

**INTECH**  
open science | open minds

### **InTech Europe**

University Campus STeP Ri  
Slavka Krautzeka 83/A  
51000 Rijeka, Croatia  
Phone: +385 (51) 770 447  
Fax: +385 (51) 686 166  
[www.intechopen.com](http://www.intechopen.com)

### **InTech China**

Unit 405, Office Block, Hotel Equatorial Shanghai  
No.65, Yan An Road (West), Shanghai, 200040, China  
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元  
Phone: +86-21-62489820  
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](https://creativecommons.org/licenses/by/3.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen