We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Skeletogenesis and the Hematopoietic Niche

Elizabeth Sweeney and Olena Jacenko
University of Pennsylvania
 USA

1. Introduction

The reciprocal regulation of the skeletal and the immune systems has been clinically appreciated for years. In particular, factors produced by immune cells during homeostasis and activation markedly affect the skeleton, which in turn affects the marrow niche environments (as reviewed in (Compston 2002)). This relationship also extends to an interdependence between bone and hematopoiesis during immune cell development, however the critical cell types and extracellular matrix components involved in establishing and maintaining hematopoietic niches within the bone marrow are only recently beginning to be defined. Indeed, some immuno-osseous disorders with hematopoietic defects such as bone marrow failure and immune dysfunction, as well as certain cancers, may result from a defective hematopoietic niche (Spranger et al. 1991; Kuijpers et al. 2004; Hermanns et al. 2005; Walkley et al. 2007; Walkley et al. 2007; Raaijmakers et al. 2010). Likewise during aging, a progressive decline in cell replacement and repair manifests in both the skeletal and hematopoietic systems with reduced bone mass and diminished blood cell formation respectively (as reviewed in (Rossi et al. 2008) and (Gruver et al. 2007)). Further, this altered hematopoiesis due to aging leads to deficient immune function and increased incidence of malignancies (Rossi et al. 2005; Janzen et al. 2006; Mayack et al. 2010). Thus, the dynamic relationship between skeletal and hematopoietic maintenance throughout life suggests that these clinical outcomes may ensue from cell signaling deficiencies or from defects in the structural environment supporting hematopoiesis. This chapter provides an overview of our current understanding of how hematopoietic niches may be established, how they promote hematopoiesis, and how the skeletal status may modulate niche function.

2. Coordinate skeletal and hematopoietic development

The vertebrate skeleton develops by one of two essential processes, endochondral (EO) and intramembranous (IO) ossification mechanisms (as reviewed in (Chan, D. and Jacenko 1998)). The direct differentiation of ectomesenchymal cells to osteoblasts in IO represents the rudimentary mechanism through which many skull bones and all periosteal bones form. The IO-derived bone is referred to as “dense”, “compact” or “cortical”, and as the names imply, is a solid bone with primary functions relating to weight bearing and protection (**Fig. 1C**). In contrast, EO relies on the generation of a cartilaginous skeletal blueprint that is gradually replaced by a “trabecular”, “spongy”, “cancellous” bone and a marrow capable of sustaining hematopoiesis (Chan, D. and Jacenko 1998; Mackie et al. 2008) (**Fig. 1C**). This replacement mechanism of EO is responsible for the formation of the vertebrate axial and appendicular skeleton, as well as certain cranial bones (Jacenko et al. 1991; Chan, D. and Jacenko 1998).

As EO initiates during embryogenesis, its distinctive feature is the emergence of hypertrophic cartilage, which is present in all skeletal elements that will develop a marrow cavity, e.g. long bones, hips, vertebrae, ribs, certain skull bones. The eventual replacement of cartilage by bone and marrow via EO relies on the sequential maturation of chondrocytes from resting, to proliferating, to hypertrophic (**Fig. 1A**). Chondrocyte hypertrophy manifests with a dramatic increase in cell size, cessation of proliferation, and synthesis of a new repertoire of differentiation-specific gene products (Godman and Porter 1960; Chan, D. and Jacenko 1998; Alvarez et al. 2001; James et al. 2010). Among these is the matrix protein collagen X, which represents the predominant biosynthetic product of hypertrophic cartilage (Gibson and Flint 1985; Schmid and Linsenmayer 1985). Concomitant with hypertrophy is a transformation from a non-calcified avascular cartilage matrix, to a calcifiable one that is permissive to vascular invasion. Morphometric analysis suggests that before vascular invasion, the terminal hypertrophic chondrocytes undergo either autophagy (Srinivas and Shapiro 2006; Bohensky et al. 2007) or apoptosis (Farnum and Wilsman 1989), the rate of which controls longitudinal growth of the skeletal element, as well as the transition from cartilage to trabecular bone and marrow (Farnum and Wilsman 1989).

Subsequent vascular entry into hypertrophic cartilage is critical to skeleto-hematopoietic development, since it leads to an influx of mesenchymal cells, hematopoietic precursors, and chondro/osteoclasts. This influx of cells, together with growth factors, cytokines and hormones, establishes the primary center of ossification and the marrow environment where hematopoiesis ensues (**Fig. 1**). Specifically, while chondro/osteoclasts degrade hypertrophic cartilage, multipotent stromal cells, including mesenchymal and perivascular reticular cells, form the marrow stroma, a meshwork of non-hematopoietic cells supporting hematopoiesis by providing structural scaffolding and producing hematopoietic factors (Taichman et al. 1996; Bianco et al. 1999). As hypertrophic cartilage continues to be degraded, matrix remnants serve as scaffolds upon which differentiating osteoblasts deposit bone matrix, thus forming trabecular bony spicules with hypertrophic cartilage cores (**Fig. 1B**) (Chan, D. and Jacenko 1998). Of note, the origin of the trabecular bone osteoblasts at the junction between marrow and the hypertrophic cartilage, termed the chondro-osseous junction, is still debated (Roach 1992; Roach et al. 1995; Roach and Erenpreisa 1996; Nakamura et al. 2006; Hilton et al. 2007; Maes et al. 2010). Following the formation of the primary ossification zones in the central or diaphyseal regions of skeletal elements, the establishment of secondary ossification centers at outer epiphyseal ends of bones defines the growth plate regions at the metaphysis (**Fig. 1A**). The growth plates occupy the narrow space that separates the marrow of the primary and secondary ossification centers, and are composed of a gradient of differentiating chondrocytes culminating in a zone of hypertrophic chondrocytes (**Fig. 1A & B**) (as reviewed in (Lefebvre and Smits 2005). The continual replacement of the hypertrophic chondrocytes by trabecular bone and marrow allows for longitudinal skeletal growth, robust hematopoiesis, and the progression of EO without consumption of the skeletal model until maturity, when in most non-rodent vertebrates EO ceases and growth plates close (**Fig. 1C**) (Kilborn et al. 2002). Thus, the end result of EO is a porous network of primary trabecular bone, consisting of a hybrid hypertrophic cartilage-bone matrix, and engulfed by a hematopoietic marrow (**Fig. 1B & C**). Subsequent bone remodeling gradually leads to a complete replacement of the hybrid primary bone by mature secondary bone, and is coincident with a gradual decline in lymphopoiesis and the onset of immunosenescence (**Fig. 1C**) (as reviewed in (Compston 2002; Gruver et al. 2007).

Fig. 1. Architecture of the endochondral bone. A) A schematic of a developing long bone illustrating its architecture. The epiphysis, or the bulbous end, lined by articular cartilage and containing the secondary ossification center with marrow, is supported by the flared metaphysis, which in turn rests upon the slender cylindrical shaft of the diaphysis. The growth plate separates the primary and secondary ossification centers, and consists of a gradient of differentiating chondrocyte zones; the proliferative cartilage (PC) and hypertrophic cartilage (HC) zones are marked, as well as the hybrid trabecular bone (TB) protruding into the marrow. The locations of the two-layered periosteal membrane surrounding the diaphysis and the inner endosteal network are marked. B) A longitudinal tibial section from a week-3 wild type mouse stained with safranin-orange, hematoxylin & eosin (H&E) and counterstained with fast green. Using these stains, the negatively charged cartilaginous matrix appears orange while the bone stains light blue-green; mature erythrocytes stain green, while other marrow elements stain pink-purple with H&E. The boxed inset is a high magnification of the chondro-osseous junction containing hypertrophic chondrocytes, bone and marrow with vascular. The hybrid nature of the trabecular bone can

be appreciated by the orange staining of the cartilaginous core, with green-blue bone matrix deposited on the surface (magnification, 10x). C) The inorganic mineralized matrix of a mature zebra bone illustrates the structural differences between the EO-derived trabecular /spongy/cancellous bone and the IO-derived compact/dense/cortical bone. Boxed is a high magnification of the EO- and IO-derived bone tissues. Note the mesh-like structure of trabecular bone for hematopoietic cell support.

Taken together, the proper differentiation of chondrocytes, vascular invasion and the gradual replacement of the cartilaginous anlagen by trabecular bone and marrow through EO, underscore the intricate orchestration of skeleto-hematopoietic development. Moreover, the coincident establishment and localization of trabecular bone within the site of active hematopoiesis likely reflects a critical hematopoietic niche in the chondro-osseous region (**Fig. 1B boxed**) (Jacenko et al. 1993; Nilsson et al. 1997; Gress and Jacenko 2000; Nilsson et al. 2001; Jacenko et al. 2002; Yoshimoto et al. 2003; Arai, F. et al. 2004; Balduino et al. 2005; Sweeney et al. 2008; Kohler et al. 2009; Lo Celso et al. 2009; Xie et al. 2009; Sweeney et al. 2010). This skeleto-hematopoietic link is strongly supported by several animal models where alterations in process of EO leads to hematopoietic defects (**Table 1**), including mouse models with altered: collagen X (Jacenko et al. 1993; Gress and Jacenko 2000; Jacenko et al. 2002; Sweeney et al. 2008; Sweeney et al. 2010), parathyroid hormone related protein (PTHrP) receptor in osteoblasts (Calvi et al. 2001; Calvi et al. 2003; Kuznetsov et al. 2004; Wu et al. 2008), osteoblast numbers (Visnjic et al. 2001; Visnjic et al. 2004; Zhu et al. 2007), bone morphogenic protein (BMP) receptor type 1A in marrow cells (Zhang, J. et al. 2003), osteoclast function (Blin-Wakkach et al. 2004; Mansour et al. 2011), retinoic acid receptor gamma (Purton et al. 2006; Walkley et al. 2007), $G_s\alpha$ in osteoblasts (Wu et al. 2008), Dicer in osteoblasts (Raaijmakers et al. 2010), glypican-3 (Viviano et al. 2005), and perlecan (Rodgers et al. 2008). **Table 1** presents a list of mouse models with defects in hematopoiesis due to alterations in a component within the niche environment. Only those mouse models are summarized that were proven, by and large, via bone marrow transplantation experiments to have an aberrant niche environment, since wild type marrow cells could not rescue the disease phenotype of the host.

3. Overview of the hematopoietic niche

Hematopoiesis is the process by which hematopoietic stem cells (HSCs) generate and replenish progenitors that develop into fully mature blood and immune cells, and populate the periphery. During vertebrate ontogeny, hematopoiesis is established sequentially in several different anatomic sites (see Development of Hematopoietic Stem Cells chapter in this book for review). Coincident with the onset of EO (approximately the last third of embryonic development), hematopoiesis shifts from the fetal liver and spleen to the EO-derived marrow, which represents the predominant site of blood cell production after birth (Aguila and Rowe 2005; Cumano and Godin 2007). Therefore, the marrow has become a tissue of study for hematopoietic cell biology post parturition. Additionally, due to the ease of marrow cell isolation in combination with the extensive list of cell markers identifying HSCs at different stages of differentiation (as summarized in (Morrison and Spradling 2008)), stem cell niche biology has also utilized the marrow environment for study.

Year	Protein/ cell	Description	Hematopoietic highlights	References
1993	Collagen X	Collagen X is a short chain collagen secreted by hypertrophic chondrocytes in the growth plate of endochondrally developing bones. In these mice, collagen X is either knocked-out or mutated to cause dominant interference.	-Altered chondro-osseous junction -Decreased trabecular bone, -No change in HSPCs, -Decreased B lymphocytes throughout life, -Diminished immunity in vitro and in vivo, -Altered hematopoietic/lymphopoietic cytokines, e.g. decreased SCF, CXCL-12, IL-7, -Disease phenotype retained after transfer of wild type HSPCs into collagen X mice	Jacenko, 1993; Jacenko, 1996; Gress, 2000; Jacenko 2001; Jacenko 2002; Sweeney, 2008; Sweeney, 2010
2001	PTH/PTHrP receptor	Parathyroid hormone (PTH) and the PTH related protein (PTHrP) receptor (PPR) are involved in calcium homeostasis and activation of osteoblasts, thus indirectly osteoclasts. In these mice, a constitutively active form of PPR is expressed in mesenchymal cells under the 2.3kb promotor of collagen I.	-Delayed hematopoiesis due to delay in the bone to marrow transition, -With aging, increased trabecular bone and osteoblast numbers, -Increased HSPCs, -Increase in Notch signaling, -Increase in IL-6, CXCL-12, SCF, -Decrease in B cells at all stages	Calvi, 2001; Calvi, 2003; Kuznetsov, 2004; Wu 2008
2001	Osteoblast deletion	Osteoblasts have been implicated as a niche cell where increased in osteoblasts resulted in increases in HSPCs. These mice were generated with the herpes thymidine kinase gene under the 2.3kb promotor of collagen I expressing mesenchymal cells after ganciclovir treatment.	-Loss of bone lining cells and trabecular bone elements, -Decrease in marrow cellularity, -Decreases in HSPCs, -Decreases in lymphoid, erythroid, and myeloid progenitor cells, -Decrease in osteoclasts, -Increased extramedullary hematopoiesis, -Recovery from disease phenotype after ganciclovire removed	Visnjic, 2001; Visnjic, 2004; Zhu, 2007
2003	BMPRI1A	The bone morphogenic protein receptor, type 1A (BMPRI1A) is a receptor for BMPs, which have been shown to influence hematopoiesis. These mice were generated with a PolyI:C-inducible Mx-1-Cre to delete BMPRI1A in hematopoietic and stromal cells.	-Increases in spindle-shaped N-cadherin+ CD45- osteoblasts (SNO), -Increases in HSPCs, -LT-HSC observed attached to SNO cells, -No block in HPC differentiation to the lymphoid or myeloid lineages	Zhang, 2003

Continued on facing page

Table 1. Mouse models with altered hematopoetic niche enviroments.

Year	Protein/ cell	Description	Hematopoietic highlights	References
2004	Tcirg 1	The T cell immune regulator 1 (Tcirg1) is a subunit of the vacuolar protein pump (V-ATPase) involved in osteoclast resorption of bone. In these mice there is a mutation of Tcirg1 rendering osteoclasts ineffective at bone resorption.	-Increases in myelomonocytic differentiation, -Reduced medullary cavity size, -Defective B cell differentiation leading to reduced numbers, -Reduced interferon-γ secretion from T cells, -Decreased IL-7 secretion from bone marrow cells, -Rescue of phenotype with restoration of marrow environment after wild type cell transfer	Blin-Wakkach, 2004; Mansour, 2011
2005	Osteopontin	Osteopontin is an extracellular matrix glycoprotein made by several cell types, including osteoblasts, fibroblasts, chondrocytes, etc. In these mice, osteopontin is knocked-out, and in 1998 Rittling et.al. showed no alterations to bone mophology, but increased osteoclastogenesis.	-Increased number of HSPCs, -No change in lymphoid or myeloid cell production, -Decreased HSPC apoptosis, -Increased Jagged 1 and Angiopoietin 1 stromal expression, -Disease phenotype retained after transfer of wild type HSPCs to Oprn-/- mice	Rittling, 1998; Stier, 2005; Nilsson, 2005
2005	p27Kip1 and MAD1	Cyclin-dependent kinase inhibitors MAD1 and p27Kip1 are negative regulators of cell cycle. In these two studies, mice had either p27Kip1 knocked-out (Chien, 2006) or p27Kip1 and MAD1 knocked-out (Walkley, 2005).	-Hyperplasia of hematopoietic organs, -Increased myeloid and erythroid colony forming cells, -Increase in LT-HSCs, -Disease phenotype retained after transfer of wild type HSPCs into p27Kip1-/- mice	Walkley, 2005; Chien, 2006
2005	gp130	Glycoprotein 103 (gp130) is a subunit of the cytokine receptors for the IL-6 family. These mice were generated with the Tere mouse for excision of gp103 in hematopoietic and endothelial cells.	-Hypocellular marrow, -Impairment in erythro-and thrombopoiesis, -Reduction in T lymphocytes in the thymus, -Reduction of B lymphopoiesis in the marrow, -Extramedullary hematopoiesis, -Disease phenotype retained after transplant of wild type HSPCs into gp130 mice	Yao, 2005
Continued on facing page				

Table 1. continued

Year	Protein/ cell	Description	Hematopoietic highlights	References
2006	RAR gamma	Retinoic acid receptors (RAR) are nuclear hormone receptors that act as ligand-dependent transcriptional regulators. In these mice the RARgamma is knocked-out.	-Decrease in HSPCs, -Increased granulopoiesis, -Myeloproliferative-like disease with excessive extramedullary hematopoiesis, -Loss of trabecular bone by 12 weeks of age, -Disease phenotype retained after transfer of wild type HSPCs into RARγ ^{-/-} mice	Purton, 2006; Walkley, 2007
2007	Rb	Retinoblastoma protein (Rb) is a central regulator of the cell cycle and several downstream regulators have been shown to have affects on hematopoiesis. These mice were generated using an inducible deletion construct in conjunction with Mx-1 Cre for deletion of Rb in hematopoietic and stromal cells.	-Myeloproliferative disease phenotype, -Increased HSPC differentiation, -Egress of HSPCs from marrow to extramedullary sites, -Loss of Rb was found to be necessary from both the myeloid-derived cells and the environment for presentation of the disease phenotype	Walkley, 2007
2008	Gsa	Gsa is a heterotrimeric G protein subunit that activates the cAMP-dependent pathway by activating adenylate cyclase and is part of the parathyroid hormone (PTH) and the PTH related protein (PTHrP) receptor. In these mice, Gsa was ablated in cells expressing ostromix, e.g. early osteoprogenitors and chondrocytes.	-Decreases in B cell precursors in the marrow and in the periphery, -No negative affect on other hematopoietic lineages, -Decreases in IL-7 expression from osteoblasts, -Decreases in trabecular bone, -Disease phenotype not transferable to wild type mice with marrow transplant	Wu, 2008
2008	Bis (BAG-3 or CAIR-1)	Bcl-2 interacting cell death suppressor (Bis) is a protein involved in antiapoptotic and antistress pathways. In this mouse, Bis was truncated for loss of function.	-Reduced lymphoid tissues, -Perturbed vasculature with defects in endothelial cells, -Loss of HSPCs, -Defect in B lymphopoiesis, -Decreased splenic hematopoietic cell numbers, -Defects in stromal progenitor cells, -Loss of stromal cells expressing CXCL-12 and IL-7, -Osteoblast lineage unaffected, -Disease phenotype retained after transplant of wild type cells into Bis ^{-/-} mice	Youn, 2008; Kwon, 2010

Continued on facing page

Table 1. continued

Year	Protein/ cell	Description	Hematopoietic highlights	References
2008	Hf2/merlin	Neurofibromin 2 (Nf2)/moesin-ezrin-radixin-like (merlin) is a cytoskeletal scaffolding protein involved in cell-cell communications. These mice were generated with a PolyI:C-inducible Mx-1-Cre to delete Nf2/merlin in hematopoietic and stromal cells.	-Egress of HSPCs, -Hypocellular marrow, -Increases in marrow vascularity, -Hematopoietic lineages unaffected, -After time, increases in trabecular bone and osteoblast numbers accompanied by restoration in marrow cellularity, -Disease phenotype retained after transplant of wild type cells into Nf2/merlin mice	Larsson, 2008
2010	Dicer	Dicer is an endoribonuclease that cleaves double-stranded RNA and pre-miRNA into small interfering RNA. In these mice, Dicer was ablated in cells expressing ostrix, e.g. early osteoprogenitors and chondrocytes.	-Decreases in leukocytes, platelets and red blood cells, -Decreases in B cell number with increases in myeloid cells, -No change in HSPC number, -Extramedullary hematopoiesis, -Slight decrease in osteoblast number with impaired differentiation, -Bone volume unchanged, but altered bone texture, -Disease phenotype retained after transplant of wild type cells into Dicer mice	Raaijmakers, 2010
2010	CAR cells	CXC chemokine ligand (CXCL) 12 abundant reticular (CAR) cells have been implicated as a hematopoietic niche cell in the marrow. These mice were designed to have inducible selective ablation of CAR cells with the diphtheria toxin receptor.	-Reduced cycling of lymphoid and erythroid progenitor cells, -Reduced HSPC numbers, -HSPCs more quiescent, -HSPCs express more myeloid genes, -No changes in osteoblasts or endothelial cells, -Impaired production of SCF and CXCL-12	Omatsu, 2010
2010	Nestin+ cells	Nestin is an intermediate filament protein expressed in nerve cells and reported here in rare non-hematopoietic cells with mesenchymal progenitor cell qualities. These mice were generated using the inducible diphtheria toxin with Nes-creERT2 for selective depletion of nestin expressing cells.	-Rapid reduction of HSPCs in marrow and increase in HSPCs in spleen, -Reduction of adrenergic nerve fibers, -Reduction of wild type HSPC homing to marrow of Nestin mice	Mendez-Ferrer, 2010
Continued on facing page				

Table 1. continued

Year	Protein/ cell	Description	Hematopoietic highlights	References
2010	Ebf2	Early B cell factor 2 (Ebf2) is a transcription factor expressed in neurons, immature osteoblasts and adipocytes. In these mice, Ebf2 is knocked-out.	-Impaired lymphopoiesis, -Reduced numbers of HSPCs, -Mice were smaller than wild type cohorts, -Disease phenotype rescued after transplant of Ebf2-/- cells into wild type mice	Corradi, 2003; Kieslinger, 2010
2011	Agrin	Agrin is a heparan sulfate proteoglycan matrix protein expressed on MSCs and trabecular osteoblasts in the niche. In these mice, agrin expression is deficient with low levels of muscle specific kinase (MuSK) expression for postnatal survival.	-Marrow hypoplasia with decreases in myeloid and lymphoid cells, -Reduced CD45+ cell numbers in marrow, spleen and marrow, -Decreases in thymus T-lineage populations, -Decreases in ST-HSC in marrow, but normal levels in fetal liver, -Disease phenotype is not transferred to wild type mice after transplant with MuSK-L: Agrin-/- hematopoietic cells	Mazzon, 2011

Table 1. continued

The idea of a unique tissue environment, or stem cell niche, as a tissue setting that can direct progenitor cell behavior, e.g. quiescence, proliferation, differentiation, etc. was proposed over four decades ago (Wolf and Trentin 1968; Trentin 1971; Schofield 1978; Wolf 1979). Hematopoietic niches, or hematopoietic microenvironments (HME), are defined by the association of particular cell types, their secreted matrix products and their soluble hematopoietic factors (Yin and Li 2006; Rodgers et al. 2008). The identity of the cellular, matrix and soluble components that influence HSCs, including the long-term populating (LT-HSC), short-term populating (ST-HSC), or the more differentiated hematopoietic progenitor cells (HPC), as well as the lymphoid and myeloid lineages, remains an active topic of investigation. However, at least two hematopoietic niches have been described, an osteoblastic (or endosteal) niche, ascribed to osteoblasts residing on bone surfaces, and a vascular niche, ascribed to endothelial cells and subendothelial MSCs or pericytes lining marrow sinusoids. Many have argued that more quiescent LT-HSCs and ST-HSCs are located in the osteoblast niche, while differentiating HPCs are located in the vascular niche for mobilization to the periphery (Lord et al. 1975; Shackney et al. 1975; Gong 1978; Nilsson et al. 2001; Heissig et al. 2002; Arai, F. et al. 2004; Balduino et al. 2005; Jang and Sharkis 2007; Bourke et al. 2009). However, these regions in the marrow are so close in proximity (Arai, F. et al. 2004; Kiel et al. 2005; Lo Celso et al. 2009; Xie et al. 2009), that the osteoblast and vascular niches may be the same, or perhaps interchangeable to some degree. Additionally, recent work has identified other cells involved in hematopoiesis that do not fully comply with these proposed niche regions, such as CXC chemokine ligand (CXCL)-12 expressing reticular cells that are scattered throughout the marrow (Tokoyoda et al. 2004; Sugiyama et al. 2006; Omatsu et al. 2010). Below we will discuss the different cellular, matrix and soluble components within the marrow that have been shown to influence hematopoiesis.

3.1 Cells of the niche

Experiments designed to identify the cells of the HME date back over fifty years (Pfeiffer 1948; Tavassoli and Crosby 1968; Tavassoli and Weiss 1971; Meck et al. 1973; Friedenstein et al. 1974; Tavassoli and Khademi 1980; Friedenstein et al. 1982; Patt et al. 1982; Tavassoli 1984; Friedenstein et al. 1987; Gurevitch and Fabian 1993; Kawai et al. 1994; Kuznetsov et al. 1997; Hara et al. 2003; Akintoye et al. 2006; Mankani et al. 2007; Sacchetti et al. 2007; Chan, C.K. et al. 2008; Mankani et al. 2008; Song et al. 2010). In these studies, ectopic bone with a functional HME was generated in host mice using various bone marrow derived osteoprogenitor seed cells. More recently, different osteoprogenitor pools have been isolated that can either generate EO-like bone with active hematopoiesis or compact IO-like bone without an HME (Akintoye et al. 2006; Chan, C.K. et al. 2008). Collectively, these studies have shown that functional HMEs form through the progression of EO with contributions from cartilage, bone, vasculature, and marrow stromal cells. Further, similar conclusions about the necessity of EO-derived components, e.g. cells and matrix molecules, were obtained through analyses of several mouse models with skeleto-hematopoietic defects (**Table 1**), including mice with disrupted collagen X function in the HME (Jacenko et al. 1993; Gress and Jacenko 2000; Jacenko et al. 2002; Sweeney et al. 2008; Sweeney et al. 2010). Discussed below are the data describing which cell type(s), associated matrix and soluble factors are necessary for blood cell development in the marrow, including: multipotent stromal cells (fibroblasts, pericytes, reticular cells, and adipocytes), osteoblasts, chondrocytes, endothelial cells, and cells of hematopoietic origin (hematopoietic stem/progenitor cells (HSPC), osteoclasts and macrophages).

3.1.1 Fibroblasts and perivascular cells

As early as the 1970's, in vitro studies with marrow stromal adherent colonies showed that this pool of cells is able to support hematopoiesis (Friedenstein et al. 1970; Dexter et al. 1973; Friedenstein et al. 1974; Friedenstein et al. 1976; Dexter et al. 1977). These plastic adherent colonies have been thought to contain mesenchymal progenitor cells, and likely mesodermal progenitor cells as well (Petrini et al. 2009). This would account for recent data indicating that marrow derived progenitor cells generate adipocytes, chondrocytes and osteoblasts, traditional mesenchymal cell types, as well as fibroblasts, smooth muscle cells, endothelial cells, and pericytes/subendothelial cells (Bentley and Foidart 1980; Muguruma et al. 2006; Sacchetti et al. 2007; Crisan et al. 2008; Kalajzic et al. 2008; Augello et al. 2010; Mendez-Ferrer et al. 2010). Thus, isolated marrow stromal cells will be referred to as multipotent stromal cells (MSC) throughout (Horwitz et al. 2005). The MSCs isolated for ectopic bone assays have been described as osteoprogenitor cells that generate bone with a HME able to support host-derived hematopoiesis (Kuznetsov et al. 1997; Akintoye et al. 2006; Sacchetti et al. 2007; Chan, C.K. et al. 2008; Morikawa et al. 2009). Of note, many of these osteoprogenitor cells have been shown to have stem-like qualities, such as self-replication and the ability to differentiate into several different cell types. For example, Sacchetti et al. isolated human-derived MCAM/CD146-expressing subendothelial cells, which can self-replicate as well as give rise to osteoblasts, chondrocytes and reticular cells in ectopic HMEs (Sacchetti et al. 2007). More recently, these data were replicated in the mouse by Morikawa et al. who also identified a perivascular cell type that has the ability to self-replicate and give rise to adipocytes, osteoblasts, chondrocytes and endothelial cells (Morikawa et al. 2009). An

additional perivascular cell has also been identified as a HME cell type, the nestin-expressing MSCs (Mendez-Ferrer et al. 2010). These, nestin⁺ MSCs have been shown to be spatially associate with HSPCs in the marrow and to express several HSPC maintenance genes, e.g. CXCL-12, stem cell factor (SCF)/kit ligand, angiopoietin (Ang)-1, interleukin (IL)-7, vascular cell adhesion molecule (VCAM)-1, and osteopontin (Mendez-Ferrer et al. 2010). Further, numbers of HSPC were rapidly reduced in the marrow of mice that were selectively depleted of nestin⁺ MSCs (**Table 1**) (Mendez-Ferrer et al. 2010). These mouse models also revealed a necessity for marrow nestin⁺ MSCs for homing of transferred HSPCs (Mendez-Ferrer et al. 2010). Together, these data suggest that MSCs and associated daughter cells not only make up the physical structure of the HME, but also provide maintenance and differentiation signals to HSPCs.

3.1.2 CXCL-12 abundant reticular cells

Reticular cells are of mesodermal origin and are a type of fibroblast cell localized to the intertrabecular region of the marrow near both the osteoblast and vascular niches (Weiss 1976; Rouleau et al. 1990). Recently, a sub-set of marrow reticular cells has been shown to express high levels of CXCL-12 (or stromal derived factor (SDF)-1) and have been termed CXCL-12 abundant reticular (CAR) cells (Tokoyoda et al. 2004; Sugiyama et al. 2006). CXCL-12 is reportedly involved in several aspects of hematopoiesis, including HSPC homing and maintenance, as well as B cell development (Nagasawa et al. 1994; Nagasawa et al. 1996; Ara et al. 2003; Broxmeyer et al. 2005; Jung et al. 2006; Sugiyama et al. 2006). Using CXCL-12/GFP knock-in mice, HSPCs, early lineage B cells and plasma B cells have been shown to spatially associate with CAR cells (Tokoyoda et al. 2004), suggesting CAR cells are a HME cell type. To address the importance of CXCL-12 expressing cells in the HME, Omatsu et al. designed a mouse model with selective ablation of CAR cells (**Table 1**) (Omatsu et al. 2010). These assays showed no change in the osteoblast or vascular niches, but impaired production of SCF and CXCL-12, combined with marked reduction in cycling lymphoid and erythroid progenitors. Further, the HSPC population in these mice was more quiescent, diminished in numbers and expressed myeloid selector genes. Finally, CAR cells can give rise to adipocytes and osteoblasts (Bianco et al. 1988; Balduino et al. 2005; Sipkins et al. 2005; Omatsu et al. 2010). Thus, these data combined with the ectopic bone assays and the nestin⁺ reticular cell studies discussed above, raise the possibility that the CAR, osteoprogenitor and nestin⁺ cells are from a similar cell pool, sharing differentiation capabilities and roles in hematopoietic support.

3.1.3 Adipocytes

Within the young marrow there are few adipocytes, however this phenomenon is reversed with aging and after marrow insult, such as post irradiation (Burkhardt et al. 1987; Verma et al. 2002). Although adipocytes have been described as having a positive influence on hematopoiesis via growth factors secretion (Lanotte et al. 1982), other studies have reported that these growth factors are in too low a concentration to influence HSPCs and that adipocytes secrete anti-hematopoietic factors as well (Hotamisligil et al. 1993; Zhang, Y. et al. 1995; Yokota et al. 2000; Corre et al. 2006; Belaid-Choucair et al. 2008; Miharada et al. 2008). Further, an increase in marrow adiposity has been negatively correlated with hematopoiesis in vivo (Touw and Lowenberg 1983; Naveiras et al. 2009). Interestingly,

Naveiras et al. have shown that in the adult mouse spine there is a proximal to distal gradient of marrow adiposity, with thoracic vertebrae being virtually free of adipocytes. This provides an *in vivo* model to study the affects of adipocytes on hematopoietic cells under homeostatic conditions. These studies showed that the number, frequency and cycling capacity of HSPCs was reduced as the number of adipocytes was increased (Naveiras et al. 2009). In support, after irradiation and marrow transplantation of mice genetically incapable of forming adipocytes, or in wild type mice treated with an inhibitor of adipogenesis, there was enhanced HSPC expansion compared to non-treated wild type cohorts (Naveiras et al. 2009). Of note, in these models of reduced/abrogated adipogenesis, as well as in a model where the fatty marrow is surgically removed, the increase in hematopoiesis is concomitant with an increase in bone formation (Tavassoli et al. 1974; Naveiras et al. 2009). These data suggest that after marrow insult and in the absence of adipocytes, there are signals enhancing osteoblast activity and bone formation, which may contribute to the enhanced hematopoiesis measured, as discussed below. This is supported by the clinical observations that aged patients have an increase in adiposity in the marrow, which is correlated with a decrease in bone formation and decreased hematopoiesis (Verma et al. 2002; Rossi et al. 2005; Mayack et al. 2010).

3.1.4 Osteoblasts

With the identification of osteoblast-like cells in stromal cultures (Friedenstein et al. 1987; Benayahu et al. 1991; Benayahu et al. 1992) and osteoprogenitors within marrow preparations for ectopic bone assays (Kuznetsov et al. 1997; Akintoye et al. 2006; Sacchetti et al. 2007; Chan, C.K. et al. 2008; Morikawa et al. 2009), Emerson and Taichman designed *in vitro* assays to assess the ability of isolated osteoblasts to support hematopoiesis (Taichman and Emerson 1994; Taichman et al. 1996; Taichman et al. 1997; Jung et al. 2005; Jung et al. 2006; Zhu et al. 2007). These assays showed that osteoblasts can support hematopoiesis through the secretion of pro-hematopoietic cytokines, e.g. granulocyte colony-stimulating factor (G-CSF), granulocyte-macrophage colony-stimulating factor (GM-CSF), CXCL-12, IL-6, and IL-7, and that cell-cell contact is necessary for support via integrin binding (very late antigen (VLA)-4/5 and VCAM/ICAM). Further, these assays confirmed a connection between osteoblasts and B cell development. To assess the contribution of osteoblasts to hematopoiesis *in vivo*, several different mouse models have been generated that either increase osteoblasts (Calvi et al. 2001; Calvi et al. 2003; Zhang, J. et al. 2003), decrease osteoblasts (Visnjic et al. 2001; Visnjic et al. 2004), disrupt EO-based trabecular bone formation (Jacenko et al. 1993; Gress and Jacenko 2000; Jacenko et al. 2001; Jacenko et al. 2002; Sweeney et al. 2008; Sweeney et al. 2010), alter osteoblast signaling (Wu et al. 2008), or modify osteoblast RNA processing (**Table 1**) (Raaijmakers et al. 2010). These *in vivo* models have confirmed that osteoblasts can support hematopoiesis, are involved in B lymphopoiesis, make pro-hematopoietic cytokines, make cell-cell contact with HSPCs, and moreover, have implicated osterix expressing osteoprogenitors as mediators of hematopoiesis.

The above studies, as well as many imaging studies of HSPC in bone, support the osteoblast hematopoietic niche theory and suggest that the osteoblast niche may additionally encompass the B lymphopoietic niche (Nilsson et al. 1997; Nilsson et al. 2001; Yoshimoto et al. 2003; Arai, F. et al. 2004; Balduino et al. 2005; Kohler et al. 2009; Lo Celso et al. 2009; Xie et

al. 2009). For instance, Xie et al. showed that GFP⁺ HSPCs home to the trabecular bone surface in the marrow, and others have reported early developing B lymphocytes at the endosteal region of the marrow (Hermans et al. 1989; Jacobsen and Osmond 1990; Osmond 1990; Xie et al. 2009). This zone is the chondro-osseous region where hypertrophic chondrocytes, trabecular osteoblasts and marrow cells are juxtaposed (**Fig. 1**). Hypertrophic chondrocytes and their matrix components are also essential for trabecular bone formation, and are proposed to be part of the osteoblast/lymphopoietic niche (Jacenko et al. 2002; Rodgers et al. 2008; Sweeney et al. 2008; Sweeney et al. 2010). Interestingly, hypertrophic chondrocytes have been described to trans-differentiate into osteoblasts (Roach 1992; Galotto et al. 1994; Roach et al. 1995; Roach and Erenpreisa 1996), and express osteoblast-like markers, e.g. osterix, osteocalcin, osteonectin, osteopontin and collagen I (Roach 1992; Yagi et al. 2003), suggesting similarities in the cells of the chondro-osseous niche. Indeed, it has been suggested that chondro-osteoprogenitor cells that expresses the chondrocyte-like marker collagen II contribute to both the perichondrial and trabecular osteoblast populations (Nakamura et al. 2006; Hilton et al. 2007). In contrast, however Maes et al. do not report any contribution from the collagen II labeled hypertrophic chondrocytes to trabecular bone (Maes et al. 2010). Taken together, these findings are reminiscent of the reports indicating that progenitor cells in the marrow can give rise to different cell types with many similarities, e.g. osteoprogenitors, CAR and nestin⁺ cells not only share gene expression profiles, but are all located in the chondro-osseous environment (Weiss 1976; Rouleau et al. 1990; Sugiyama et al. 2006). The possible overlap between the cells of the chondro-osseous region can also be appreciated when comparing osteoblasts and reticular cells that can support B lymphopoiesis and both express VCAM-1 and IL-7 (Ryan et al. 1991; Funk et al. 1995; Zhu et al. 2007). Moreover, using an osteoblast lineage tracer mouse generated with an osterix-LacZ construct, cells of the perichondrium, trabecular bone, cartilage and marrow stroma, some intimately associated with blood vessels in a pericyte-like fashion, were all positive for osterix expression (Maes et al. 2010), again confirming an overlap of cell phenotypes in the chondro-osseous HME.

3.1.5 Chondrocytes

As previously discussed, chondrocytes provide the blueprint for future bone with a marrow cavity during EO, and are adjacent to the postulated osteoblast and vascular niches (**Fig. 1B boxed**) (Arai, F. et al. 2004; Kiel et al. 2005; Kohler et al. 2009; Xie et al. 2009). Indeed, growth plate chondrocytes, as well as osteoblasts and vascular cells, express leukemia inhibitory factor (LIF), which can synergize with growth factors to promote the proliferation of HSPCs (Keller et al. 1996; Grimaud et al. 2002). Additionally, Wei et al. recently showed that hypertrophic chondrocytes express CXCR-4, the receptor for CXCL-12 made by stromal cells and osteoblasts (Peled et al. 1999; Kortessidis et al. 2005; Dar et al. 2006; Jung et al. 2006; Sacchetti et al. 2007; Wei et al. 2010). These data begin to reveal the cross talk between the hypertrophic chondrocytes and the cells of the chondro-osseous environment that are players within the hematopoietic niche, e.g. osteoblasts, stromal and hematopoietic cells.

To date, no imaging studies have attempted to localize HSPCs to hypertrophic chondrocytes. However, the contribution of various matrix components, in particular the heparan sulfate proteoglycans (HSPG), in establishing reservoirs of soluble factors for cell signaling and/or retention of HSPCs has been well established (as reviewed in (Rodgers et

al. 2008) also see (Gordon et al. 1988; Roberts et al. 1988; Siczkowski et al. 1992; Verfaillie 1993; Allouche and Bikfalvi 1995; Bruno et al. 1995; Klein et al. 1995; Gupta et al. 1996; Gupta et al. 1998; Borghesi et al. 1999; Siebertz et al. 1999; Zweegman et al. 2004; Rodgers et al. 2008; Spiegel et al. 2008). In support, our laboratory has shown altered localization of both hyaluronan and HSPGs within the hypertrophic cartilage zone of the growth plates in the collagen X mouse models that display altered hematopoiesis (Jacenko et al. 2001), which directly links EO and the hypertrophic cartilage matrix to hematopoiesis (Jacenko et al. 1993; Gress and Jacenko 2000; Jacenko et al. 2001; Jacenko et al. 2002; Sweeney et al. 2008; Sweeney et al. 2010). Briefly, collagen X is a short chain, network forming collagen that is the major secreted matrix protein of hypertrophic chondrocytes and is localized to the hypertrophic cartilage/chondro-osseous region (Campbell et al. 2004), where it is proposed to form a hexagonal lattice-like network in the matrix (Jacenko et al. 1991; Chan, D. and Jacenko 1998). Affinity co-electrophoresis studies demonstrated that collagen X and heparin, a structural analog of heparan sulfate, can endogenously bind (Sweeney, unpublished). We thus proposed that the hypertrophic chondrocyte derived matrix, made up of the collagen X network that is likely stabilized by associating with the HSPGs, is enriched with hematopoietic factors and is a vital component of the HME (Jacenko et al. 2001; Rodgers et al. 2008). In accord, mouse models where the function of collagen X was altered via transgenesis or targeted gene knock-out have both an altered HME structure, as well as aberrant hematopoiesis. Specifically, alterations within the EO-derived chondro-osseous junction include aberrant growth plate histomorphometry, collapsed hypertrophic chondrocyte matrix network, diminished and altered localization for HSPG within hypertrophic cartilage and trabecular bone, and decreased trabecular bone. The hematopoietic changes include diminished B lymphopoiesis throughout life, perinatal lethality within a sub-set of mice due to opportunistic infections by the third week of life, decreased responses to concanavalin A by splenocytes from mice at all ages, and the succumbing of all collagen X mice to non-virulent pathogen challenge (**Table 1**) (Jacenko et al. 1993; Rosati et al. 1994; Jacenko et al. 1996; Kwan et al. 1997; Gress and Jacenko 2000; Jacenko et al. 2001; Jacenko et al. 2002; Sweeney et al. 2008; Sweeney et al. 2010). Additionally, altered levels of hematopoietic cytokines have been measured from the collagen X mouse derived hypertrophic chondrocytes and trabecular osteoblasts when compared to wild type cohorts (Sweeney, 2011 Ann N Y Acad. Sci. in press). The altered cytokine availability may negatively affect hematopoiesis, which may be even further amplified by the altered chondro-osseous matrix in the collagen X mice, e.g. loss of functional collagen X and diminished HSPGs at the chondro-ossous HME (Jacenko et al. 2001). In support of the notion that matrix/cytokine signaling can affect cell differentiation in the HME, fate switching was sited as the cause of decreased osteoblast progenitors and bone formation in a knock-out mouse for the critical transforming growth factor (TGF)- β binding proteoglycans, biglycan and decorin (Bi et al. 2005). Overall, the collagen X mouse models have highlighted the contribution of EO-derived cells and matrix components to HMEs and to hematopoietic cell development.

3.1.6 Endothelial cells

Marrow arterioles and capillaries supply the sinusoids, which in turn supply the marrow with cells and nutrients (reviewed in (Kopp et al. 2005). These are the sites of the vascular niche, and examples of active vascular niches can be appreciated during development when

hematopoiesis takes place in the yolk sac, aorta-gonad-mesonephros region and placenta perivascularly (Cumano et al. 1996; Medvinsky and Dzierzak 1996; Sanchez et al. 1996; de Bruijn et al. 2002; North et al. 2002; Gekas et al. 2005; Ottersbach and Dzierzak 2005). Additionally, throughout life in some species, such as the zebra fish, hematopoiesis is not coincident with bone (Murayama et al. 2006). Further, the characteristics of the marrow sinusoids, e.g. chemokine and adhesion molecule expression, not only allow them to be conduits for hematopoietic cells to and from the circulation, but also to serve as an area for HSPC differentiation (Rafii et al. 1994; Rafii et al. 1995; Schweitzer et al. 1996; Naiyer et al. 1999; Abkowitz et al. 2003; Avezilla et al. 2004). In agreement, endothelial cells from several sources are able to support HSPC maintenance and differentiation toward lymphoid and myeloid lineages in culture (Rafii et al. 1995; Ohneda et al. 1998; Li et al. 2004; Wittig et al. 2009; Butler et al. 2010). Visualizing the vascular niche in vivo with a pure sub-set of HSCs also provided support for the vascular niche theory. These studies revealed approximately 60% of HSC residing in the chondro-osseous HME, and of that population, approximately 60% were proximal to the vasculature, where as 15% were near bone (Kiel et al. 2005). Moreover, one aspect of the osteoblast niche theory maintains that HSCs are bound to osteoblasts by a N-cadherin-mediated homophilic adhesion, however in a mouse model where HSC specific N-cadherin was depleted, hematopoiesis was fully functional (Zhang, J. et al. 2003; Kiel et al. 2009). More in vivo support is provided by the biglycan deficient mice, which present with decreased trabecular osteoblasts and bone formation, however show no defects in hematopoiesis, HSC frequency or function, and show HSC localization to the vasculature (Kiel et al. 2007). These data suggest an overlap in niche location with perhaps some ability for compensation between the osteoblast and vascular niches, albeit limited. In support, via histochemistry of the long bone, one can appreciate the spatial proximity of osteoblasts to the vasculature (**Fig. 1B boxed**).

3.1.7 Hematopoietic derived cells influence the niche

Discussed above are the data linking non-hematopoietic cells with hematopoiesis, however there are also data supporting reciprocal affects of hematopoietic derived cells influencing non-hematopoietic lineages. For example, HSPCs regulate MSC differentiation toward the osteoblast lineage via expression of BMP-2 and -6, suggesting that the HSPCs can actively maintain the osteoblast niche (Jung et al. 2008). Additionally, macrophages intercalated throughout bone have been described as osteoblast helper cells since they promote osteoblast mineralization in vitro and form a canopy over osteoblasts generating bone in vivo (Chang et al. 2008). An additional player in osteoblastogenesis is the megakaryocyte. In mouse models with increased numbers of megakaryocytes due to maturation arrest, increased osteoblast proliferation and bone mass were measured (Kacena et al. 2004). Additionally, megakaryocytes have been described as niche restoring cells post-irradiation since they migrate to the damaged bone surfaces and increase local concentrations of CXCL-12, platelet-derived growth factor (PDGF)- β and basic fibroblast growth factor (bFGF), which are associated with osteoblast proliferation (Kacena et al. 2006; Dominici et al. 2009).

A reciprocal balance between bone deposition by osteoblasts, bone resorption by osteoclasts and signaling by osteocytes is extensively noted in the literature, and is appreciated clinically. These coupled interactions will not be discussed here other than to acknowledge

that the function of one cell type highly depends upon and is affected by the actions of the other (Khosla 2003). Thereby, the continual signaling between cells of the mesenchymal and hematopoietic lineages underlies the tightly coupled process of bone remodeling, and its uncoupling can lead to skeletal disorders such as osteoporosis, osteopetrosis, as well as calcium homeostasis imbalances. Such examples are also seen in mouse models; for example, in one model where osteoclasts are depleted, instead of having increased bone mass due to lack of resorption, there is decreased bone mass compared to wild type cohorts (Kong et al. 1999). Collectively, the intricate cross talk between cells within the HME can both positively and negatively affect the niche environment as well as hematopoiesis, and is a vast area of research that remains to be adequately explored.

3.2 Soluble factors and the extracellular matrix in the niche

3.2.1 Cytokines, chemokines, growth factors, and neurotransmitters

As referred to above, the hematopoietic and non-hematopoietic cells within the HME are surrounded in all dimensions by matrix components and soluble factors. The building blocks of the extracellular matrices found within the chondro-osseous environment can include collagens, proteoglycans (PGs; including the HSPGs) and their glycosaminoglycan (GAG) constituents and glycoproteins. Collagens generally provide structural support for cells in the niche by forming supramolecular aggregates around cells (Jacenko et al. 1991), while the PGs, such as the HSPGs, can trap and store soluble factors for presentation to local cells (as reviewed in (Rodgers et al. 2008)). The amount and ratio of these molecules in the matrix also dictates the mechanical properties of the HME, which has recently become a topic of investigation in the stem cell field. For instance, matrix elasticity in the HME can influence fate choices of HSPCs (Holst et al. 2010), a phenomenon also reported with MSCs (as reviewed in (Discher et al. 2009)). Thus, the matrix provides structural integrity to the HME, acts as a substrate for cell migration and anchorage, and actively regulates cell morphology, development and metabolic function (Peerani and Zandstra 2010). The cells in the HME receive information for maintenance, development, differentiation, etc. via cell-cell interactions, cell-matrix interactions, and exposure to variable concentrations and combinations of soluble factors, e.g. cytokines, chemokines, hormones, and growth factors. Many soluble factors have been implicated in hematopoiesis, such as CXCL-12, SCF, Fms-related tyrosine kinase 3 ligand (Flt-3L), thrombopoietin (TPO), FGF, G-CSF, GM-CSF, LIF, Wnt, BMP-4, IL-3, 6, 7, 8, 11, 12, 14, 15 (Guba et al. 1992; Heinrich et al. 1993; Verfaillie 1993; Funk et al. 1995; Rafii et al. 1997; Taichman et al. 1997; Peled et al. 1999; Majumdar et al. 2000; Ponomaryov et al. 2000; Petit et al. 2002; Avecilla et al. 2004; Kortessidis et al. 2005; Dar et al. 2006; Jung et al. 2006; Spiegel et al. 2007; Wittig et al. 2009), or in maintenance and quiescence of HSCs, e.g. Ang-1 and TGF (Eaves et al. 1991; Fortunel et al. 2000; Arai, F. et al. 2004) (cytokine functions reviewed in (Zhang, C.C. and Lodish 2008)). The HME cell types discussed above are the primary sources of these soluble factors, however lymphocytes have also been shown to stimulate and suppress hematopoiesis through the release of different factors during both homeostasis and immune activation (Nathan et al. 1978; Bacigalupo et al. 1980; Mangan et al. 1982; Harada et al. 1985; Trinchieri et al. 1987; Crawford et al. 2010). Interestingly, many of these soluble factors are sequestered and presented by the matrix, specifically HSPGs, which have been described as key orchestrators of hematopoiesis (Bruno et al. 1995; Gupta et al. 1996; Gupta et al. 1998; Borghesi et al. 1999). This is particularly

relevant to the collagen X mouse models that present with a disrupted collagen X network coupled with a decreased HSPG staining intensity at the chondro-osseous HME and diminished cytokine levels, leading to diminished B lymphopoiesis (Gress and Jacenko 2000; Jacenko et al. 2001; Jacenko et al. 2002; Sweeney et al. 2008; Sweeney et al. 2010). These findings imply that the structural defects in the matrix may lead to changes in the cytokine reservoirs, which in turn would negatively affect hematopoietic cell development. By extension, human diseases associated with altered matrix components at the HME may have altered hematopoiesis due to changes in cytokine availability, such as with Simpson-Golabi Behmel syndrome where alterations in the HSPG glypican-3 results in skeletal and hematopoietic abnormalities (Pilia et al. 1996; Viviano et al. 2005).

There is increasing evidence that the nervous system can also affect the immune system through neurotransmitter signaling. The bone and marrow are supplied with autonomic efferent and afferent sensory innervations, specifically at the epiphysis and metaphysis of long bone, which includes the chondro-osseous HME (**Fig. 1A**) (reviewed in (Mignini et al. 2003). Catecholamines, acetylcholine and peptide transmitters of neural and non-neural origin are released in the HME, which contribute to neuro-immune modulations. For example, signaling from the nervous system can regulate HSPC egress and repopulation of the marrow (Katayama et al. 2006; Spiegel et al. 2007), which has been shown to be coupled to the circadian rhythm (Mendez-Ferrer et al. 2008). Indeed, HSPCs have receptors for several neurotransmitters, which can stimulate cell proliferation (Spiegel et al. 2008; Kalinkovich et al. 2009). Of note, beta-adrenergic agonists have been shown to stimulate osteoclast activity (Arai, M. et al. 2003), which could have an effect on the osteoblast niche via two methods, a) physically by decreasing bone lining cells and releasing HSPCs, and b) chemically via the release of soluble factor (Kollet et al. 2006; Mansour et al. 2011). Calcium is an example of one such soluble factor, which serves as an attractant to HSPCs encouraging homing to the osteoblast niche (Adams et al. 2006). Hematopoietic cell egress from the marrow has also been linked to many systemic causes, including exercise, inflammation, bleeding, cytotoxic drugs, and psychological anxiety (reviewed in (Lapid et al. 2008). These data serve as reminders that hematopoiesis and the HME can be influenced by factors outside of the local environment.

3.2.2 Cell and matrix influence

In the HME, cell-cell interactions influence cell fate decisions and mobility/homing; examples of such interactions include: Notch-1/Jagged-1, N-cadherin/N-cadherin and VLA-4/VCAM-1 (reviewed in (Coskun and Hirschi 2010). On the other hand, cell-matrix interactions influence not only cell behavior, but also cell anchorage to the niche. To date, the matrix proteins within the HME include: collagens (Types I, II, III, IV and X), glycoproteins (fibronectin, laminin, nidogen, tenascin C, thrombospondin, vitronectin), PGs (perlecan, decorin, agrin) and the GAG hyaluronan (Bentley and Foidart 1980; Bentley 1982; Spooncer et al. 1983; Zuckerman and Wicha 1983; Zuckerman et al. 1985; Klein 1995; Ohta et al. 1998; Campbell et al. 2004; Mazzon et al. 2011). These matrix constituents can signal to hematopoietic cells through cell receptors such as: integrins, immunoglobulin-like molecules, cadherins, selectins, and mucins (Teixido et al. 1992; Coulombel et al. 1997; Levesque and Simmons 1999; Zhang, J. et al. 2003; Merzaban et al. 2011). Notably, the HME matrix network is not static, but is continually remodeled by different enzymes including

metalloproteinases, neutrophil elastase and heparanase. Matrix turnover can thus assist in the release of hematopoietic cells from the niche (Levesque et al. 2001; Heissig et al. 2002; Petit et al. 2002; Spiegel et al. 2008), as well as liberate bound soluble hematopoietic factors (Heissig et al. 2002; Spiegel et al. 2008). An example of one such cell-matrix interaction in the HME is the VLA-4/fibronectin binding between HSPCs and the matrix, which provides the hematopoietic cell with anchorage as well as proliferation stimuli (Weinstein et al. 1989; Klein et al. 1998; Sagar et al. 2006). Hyaluronan also impacts HSPC maintenance, propagation, homing and homeostasis via CD44 binding (Avigdor et al. 2004; Matrosova et al. 2004; Haylock and Nilsson 2006). Most recently, Mazzon et al. have found the binding between agrin, expressed by trabecular osteoblasts and MSCs in the niche, and HSPCs leads to survival and proliferation signals (Mazzon et al. 2011). Finally, mature plasma B cells homing back to the marrow via CXCL-12 signals are anchored to their marrow niche via matrix-bound ligands produced by local myeloid cells (O'Connor et al. 2004; Crowley et al. 2005; Ingold et al. 2005; Moreaux et al. 2005; Nagasawa 2006; Schwaller et al. 2007; Huard et al. 2008; Moreaux et al. 2009). In fact, it has been shown that this interaction maintains long-lived antibody producing plasma B cells in the marrow by stimulating expression of anti-apoptotic genes in the lymphocytes (O'Connor et al. 2004; Huard et al. 2008). Thus, the cell-matrix interactions in the marrow serve to support hematopoietic maintenance and development, as well as support the persistence of mature hematopoietic cells that have returned to the marrow.

4. Summary and perspectives

All the specialized cells of the blood are generated through hematopoiesis via the directed differentiation of HSCs. The bone marrow, which is the predominant hematopoietic tissue after birth (Aguila and Rowe 2005; Cumano and Godin 2007), is formed through EO, where the cartilage anlage serves as a transient template for trabecular bone, and defines the environment of the marrow stroma. Thereby, either directly or indirectly, the process of EO establishes the hematopoietic niche by providing the niche with both the structure matrix constituents and the cellular components (Jacenko et al. 1993; Taichman and Emerson 1994; Taichman et al. 1996; Taichman et al. 1997; Gress and Jacenko 2000; Calvi et al. 2001; Visnjic et al. 2001; Jacenko et al. 2002; Calvi et al. 2003; Zhang, J. et al. 2003; Visnjic et al. 2004; Jung et al. 2005; Jung et al. 2006; Zhu et al. 2007; Chan, C.K. et al. 2008; Sweeney et al. 2008; Wu et al. 2008; Raaijmakers et al. 2010; Sweeney et al. 2010). Many cell types, matrix components and soluble factors contribute to the HME (**Fig. 2**). Through several different methods, HSPCs have been visualized in the chondro-osseous HME (Nilsson et al. 1997; Nilsson et al. 2001; Yoshimoto et al. 2003; Arai, F. et al. 2004; Balduino et al. 2005; Kiel et al. 2005; Kohler et al. 2009; Xie et al. 2009), which is comprised of vasculature sinusoids, sympathetic nerves, complex and diverse matrix regions, as well as osteoblasts, hypertrophic chondrocytes, endothelial cells, pericytes, CXCL-12 expressing cells, adipocytes, nestin⁺ cells, MSCs, macrophages intercalated in the endosteum, and cells of the immune system, both developing and recirculating. The cells of the HME express the cytokines, growth factors and chemokines utilized throughout hematopoiesis, as well as the matrix molecules that provides structural support, cell-matrix signaling and reservoirs of soluble factors (**Fig. 2**). We propose that the chondro-oseous HME is not static, but is continuously changing in response to various systemic influences (Lapid et al. 2008), as well as to remodeling of the hybrid trabecular bone-hypertrophic cartilage spicules into mature secondary bone (**Fig. 1C & 2**). During remodeling of the HME, both cells and

Fig. 2. The cells, matrix components and soluble factors within the chondro-osseous hematopoietic microenvironment. The putative hematopoietic niche has been localized to the chondro-osseous junction (COJ) of endochondrally developing bones, e.g. the juncture of the growth plate (GP) hypertrophic chondrocytes and the trabecular bone (TB) and marrow (M). Boxed, a cartoon in higher magnification represents the chondro-osseous hematopoietic environment, which includes: osteoblasts, hypertrophic chondrocytes, endothelial cells, pericytes, CXCL-12 abundant cells (CAR), adipocytes, endosteal intercalating macrophages, and other marrow cells, such as lymphocytes. Additionally, progenitor cells such as hematopoietic stem/progenitor cells (HSPC), multipotent stromal cells (MSC) and nestin expressing (nestin +) MSC are represented. All of these cells serve as a source of secreted factors, including cytokines, chemokines and growth factors, as well as neurotransmitters from the sympathetic nervous system (SNS). Also note the gradients of oxygen (O_2) and calcium (Ca^{++}) released from the vasculature or remodeled bone respectively. Finally, the extracellular matrix (ECM) in the marrow and in the core of trabecular bone, consisting of hypertrophic cartilage derived lattice-like collagen X and HSPGs, can serve as a substrate for cell anchorage, migration, and/or signaling through cell-ECM binding and as a reservoirs of secreted factors.

soluble factors are released into the milieu; for example matrix degrading enzymes release HSPCs for egress, and the combined activity of osteoclasts and enzymes release soluble factors from bone, including the trabecular bone with a collagen X/HSPG core (**Fig. 2**). Thus, the chondro-osseous HME is a continually active site with intrinsic and systemic signals influencing the cellular cross talk that ensures proper quiescence, maintenance, and differentiation of the HSPCs.

As itemized in **Table 1** and discussed above, various mouse models have revealed important cell and matrix players in the HME, while others have implicated the importance of proper cell cycle (Walkley et al. 2005; Walkley et al. 2007), transcription (Corradi et al. 2003; Purton et al. 2006; Walkley et al. 2007; Kieslinger et al. 2010), cell-cell communication (Larsson et al. 2008), and survival signals (Youn et al. 2008; Kwon et al. 2010) in the HME (**Table 1**). Still, the exact cellular make up and location of the HME is continually under debate, though most agree the cells of the osteoblast and vascular niches are important players in hematopoiesis. The possibility of osteoblast and vascular niches spatially overlapping or having the ability to provide some compensation for each other to some extent, is another intriguing theory. Evidence for this is observed with the biglycan deficient mouse model that presents with reduced bone, yet intact hematopoiesis (Kiel et al. 2007), as well as with the recently identified CAR cells that unite the osteoblast and vascular niches in the chondro-osseous HME by the observation that they are in contact with 90% of HSPCs throughout the trabecular bone region and the marrow sinusoidal region (Sugiyama et al. 2006; Omatsu et al. 2010). Additionally, there is similarity in the expression of hematopoietic soluble factors between the osteoblasts, endothelial cells, and CAR cells, such as CXCL-12 implicated in the homing, growth, development and maintenance of hematopoietic cells (Peled et al. 1999; Ponomaryov et al. 2000; Tokoyoda et al. 2004; Broxmeyer et al. 2005; Kortessidis et al. 2005; Dar et al. 2006; Sugiyama et al. 2006). Moreover, Medici et al. have confirmed an endothelial to osteoblast and chondrocyte transition in vivo, further supporting a cell interchange and overlap theory (Medici et al. 2010). In contrast, there is also evidence that the osteoblast and vascular niches provide different roles in hematopoiesis, e.g. a quiescence and maintenance role versus a differentiation and egress role (Lord et al. 1975; Shackney et al. 1975; Gong 1978; Nilsson et al. 2001; Heissig et al. 2002; Arai, F. et al. 2004; Balduino et al. 2005; Jang and Sharkis 2007; Bourke et al. 2009). Further, although these niches are proximal in the chondro-oseous region, mathematical modeling has predicted a layer of only two myeloid cells is sufficient to deplete most oxygen provided by a near by sinusoid (Chow et al. 2001). Thus, the local environment within each niche may differ significantly in chemical signals, such as with oxygen and calcium (**Fig. 2**).

Overall, the research generated in the hematopoietic niche field is beginning to shed light on many hematologic disorders, such as myelodysplasia, myeloproliferative syndromes and leukemias that seem to be influenced by the quality of the marrow environment (Walkley et al. 2007; Walkley et al. 2007; Raaijmakers et al. 2010). Further information on niche components, specifically the matrix molecules, will assist in generating bio-mimicking composites necessary for in vitro culture and expansion of patient specific hematopoietic tissues, for such clinical applications as autologous marrow transfers. The past sixty years of hematopoietic biology research has increased our understanding of marrow stromal cell types, as well as the three-dimensional regions that provide structure and organization of cell signaling for the maintenance and propagation of HSPCs. The study of the

hematopoietic niche will continue to provide details on necessary niche components, which may assist in the understanding of other stem cell niches, including the vascular, skin, hair, and neural niches, and provide therapeutic cues for immuno-osseous diseases that present with skeletal defects and altered hematopoiesis, such as McKusick type metaphyseal chondrodysplasia (cartilage-hair hypoplasia; CHH), Shwachmen-Diamond syndrome, Schimke dysplasia (Spranger et al. 1991; Kuijpers et al. 2004; Hermanns et al. 2005), and others.

5. References

- Abkowitz, J. L., A. E. Robinson, S. Kale, M. W. Long & J. Chen (2003). "Mobilization of hematopoietic stem cells during homeostasis and after cytokine exposure." *Blood* 102(4): 1249-1253
- Adams, G. B., K. T. Chabner, I. R. Alley, D. P. Olson, Z. M. Szczepiorkowski, M. C. Poznansky, C. H. Kos, M. R. Pollak, E. M. Brown & D. T. Scadden (2006). "Stem cell engraftment at the endosteal niche is specified by the calcium-sensing receptor." *Nature* 439(7076): 599-603
- Aguila, H. L. & D. W. Rowe (2005). "Skeletal development, bone remodeling, and hematopoiesis." *Immunol Rev* 208: 7-18
- Akintoye, S. O., T. Lam, S. Shi, J. Brahim, M. T. Collins & P. G. Robey (2006). "Skeletal site-specific characterization of orofacial and iliac crest human bone marrow stromal cells in same individuals." *Bone* 38(6): 758-768
- Allouche, M. & A. Bikfalvi (1995). "The role of fibroblast growth factor-2 (FGF-2) in hematopoiesis." *Prog Growth Factor Res* 6(1): 35-48
- Alvarez, J., M. Balbin, M. Fernandez & J. M. Lopez (2001). "Collagen metabolism is markedly altered in the hypertrophic cartilage of growth plates from rats with growth impairment secondary to chronic renal failure." *J Bone Miner Res* 16(3): 511-524
- Ara, T., K. Tokoyoda, T. Sugiyama, T. Egawa, K. Kawabata & T. Nagasawa (2003). "Long-term hematopoietic stem cells require stromal cell-derived factor-1 for colonizing bone marrow during ontogeny." *Immunity* 19(2): 257-267
- Arai, F., A. Hirao, M. Ohmura, H. Sato, S. Matsuoka, K. Takubo, K. Ito, G. Y. Koh & T. Suda (2004). "Tie2/angiopoietin-1 signaling regulates hematopoietic stem cell quiescence in the bone marrow niche." *Cell* 118(2): 149-161
- Arai, M., T. Nagasawa, Y. Koshihara, S. Yamamoto & A. Togari (2003). "Effects of beta-adrenergic agonists on bone-resorbing activity in human osteoclast-like cells." *Biochim Biophys Acta* 1640(2-3): 137-142
- Augello, A., T. B. Kurth & C. De Bari (2010). "Mesenchymal stem cells: a perspective from in vitro cultures to in vivo migration and niches." *Eur Cell Mater* 20: 121-133
- Avecilla, S. T., K. Hattori, B. Heissig, R. Tejada, F. Liao, K. Shido, D. K. Jin, S. Dias, F. Zhang, T. E. Hartman, N. R. Hackett, R. G. Crystal, L. Witte, D. J. Hicklin, P. Bohlen, D. Eaton, D. Lyden, F. de Sauvage & S. Rafii (2004). "Chemokine-mediated interaction of hematopoietic progenitors with the bone marrow vascular niche is required for thrombopoiesis." *Nat Med* 10(1): 64-71
- Avigdor, A., P. Goichberg, S. Shivtiel, A. Dar, A. Peled, S. Samira, O. Kollet, R. Hershkovich, R. Alon, I. Hardan, H. Ben-Hur, D. Naor, A. Nagler & T. Lapidot (2004). "CD44 and hyaluronin acid cooperate with SDF-1 in the trafficking of human CD34+ stem/progenitor cells to bone marrow." *Blood* 103(8): 2981-2989

- Bacigalupo, A., M. Podesta, M. C. Mingari, L. Moretta, M. T. Van Lint & A. Marmont (1980). "Immune suppression of hematopoiesis in aplastic anemia: activity of T-gamma lymphocytes." *J Immunol* 125(4): 1449-1453
- Balduino, A., S. P. Hurtado, P. Frazao, C. M. Takiya, L. M. Alves, L. E. Nasciutti, M. C. El-Cheikh & R. Borojevic (2005). "Bone marrow subendosteal microenvironment harbours functionally distinct haemosupportive stromal cell populations." *Cell Tissue Res* 319(2): 255-266
- Belaïd-Choucair, Z., Y. Lepelletier, G. Poncin, A. Thiry, C. Humblet, M. Maachi, A. Beaulieu, E. Schneider, A. Briquet, P. Mineur, C. Lambert, D. Mendes-Da-Cruz, M. L. Ahui, V. Asnafi, M. Dy, J. Boniver, B. V. Nusgens, O. Hermine & M. P. Defresne (2008). "Human bone marrow adipocytes block granulopoiesis through neuropilin-1-induced granulocyte colony-stimulating factor inhibition." *Stem Cells* 26(6): 1556-1564
- Benayahu, D., A. Fried, D. Zipori & S. Wientroub (1991). "Subpopulations of marrow stromal cells share a variety of osteoblastic markers." *Calcif Tissue Int* 49(3): 202-207
- Benayahu, D., M. Horowitz, D. Zipori & S. Wientroub (1992). "Hemopoietic functions of marrow-derived osteogenic cells." *Calcif Tissue Int* 51(3): 195-201
- Bentley, S. A. (1982). "Collagen synthesis by bone marrow stromal cells: a quantitative study." *Br J Haematol* 50(3): 491-497
- Bentley, S. A. & J. M. Foidart (1980). "Some properties of marrow derived adherent cells in tissue culture." *Blood* 56(6): 1006-1012
- Bi, Y., C. H. Stuelten, T. Kilts, S. Wadhwa, R. V. Iozzo, P. G. Robey, X. D. Chen & M. F. Young (2005). "Extracellular matrix proteoglycans control the fate of bone marrow stromal cells." *J Biol Chem* 280(34): 30481-30489
- Bianco, P., M. Costantini, L. C. Dearden & E. Bonucci (1988). "Alkaline phosphatase positive precursors of adipocytes in the human bone marrow." *Br J Haematol* 68(4): 401-403
- Bianco, P., M. Riminucci, S. Kuznetsov & P. G. Robey (1999). "Multipotential cells in the bone marrow stroma: regulation in the context of organ physiology." *Crit Rev Eukaryot Gene Expr* 9(2): 159-173
- Blin-Wakkach, C., A. Wakkach, P. M. Sexton, N. Rochet & G. F. Carle (2004). "Hematological defects in the oc/oc mouse, a model of infantile malignant osteopetrosis." *Leukemia* 18(9): 1505-1511
- Bohensky, J., I. M. Shapiro, S. Leshinsky, S. P. Terkhorn, C. S. Adams & V. Srinivas (2007). "HIF-1 regulation of chondrocyte apoptosis: induction of the autophagic pathway." *Autophagy* 3(3): 207-214
- Borghesi, L. A., Y. Yamashita & P. W. Kincade (1999). "Heparan sulfate proteoglycans mediate interleukin-7-dependent B lymphopoiesis." *Blood* 93(1): 140-148
- Bourke, V. A., C. J. Watchman, J. D. Reith, M. L. Jorgensen, A. Dieudonne & W. E. Bolch (2009). "Spatial gradients of blood vessels and hematopoietic stem and progenitor cells within the marrow cavities of the human skeleton." *Blood* 114(19): 4077-4080
- Broxmeyer, H. E., C. M. Orschell, D. W. Clapp, G. Hangoc, S. Cooper, P. A. Plett, W. C. Liles, X. Li, B. Graham-Evans, T. B. Campbell, G. Calandra, G. Bridger, D. C. Dale & E. F. Srour (2005). "Rapid mobilization of murine and human hematopoietic stem and progenitor cells with AMD3100, a CXCR4 antagonist." *J Exp Med* 201(8): 1307-1318

- Bruno, E., S. D. Luikart, M. W. Long & R. Hoffman (1995). "Marrow-derived heparan sulfate proteoglycan mediates the adhesion of hematopoietic progenitor cells to cytokines." *Exp Hematol* 23(11): 1212-1217
- Burkhardt, R., G. Kettner, W. Bohm, M. Schmidmeier, R. Schlag, B. Frisch, B. Mallmann, W. Eisenmenger & T. Gilg (1987). "Changes in trabecular bone, hematopoiesis and bone marrow vessels in aplastic anemia, primary osteoporosis, and old age: a comparative histomorphometric study." *Bone* 8(3): 157-164
- Butler, J. M., D. J. Nolan, E. L. Vertes, B. Varnum-Finney, H. Kobayashi, A. T. Hooper, M. Seandel, K. Shido, I. A. White, M. Kobayashi, L. Witte, C. May, C. Shawber, Y. Kimura, J. Kitajewski, Z. Rosenwaks, I. D. Bernstein & S. Rafii (2010). "Endothelial cells are essential for the self-renewal and repopulation of Notch-dependent hematopoietic stem cells." *Cell Stem Cell* 6(3): 251-264
- Calvi, L. M., G. B. Adams, K. W. Weibrecht, J. M. Weber, D. P. Olson, M. C. Knight, R. P. Martin, E. Schipani, P. Divieti, F. R. Bringhurst, L. A. Milner, H. M. Kronenberg & D. T. Scadden (2003). "Osteoblastic cells regulate the haematopoietic stem cell niche." *Nature* 425(6960): 841-846
- Calvi, L. M., N. A. Sims, J. L. Hunzelman, M. C. Knight, A. Giovannetti, J. M. Saxton, H. M. Kronenberg, R. Baron & E. Schipani (2001). "Activated parathyroid hormone/parathyroid hormone-related protein receptor in osteoblastic cells differentially affects cortical and trabecular bone." *J Clin Invest* 107(3): 277-286
- Campbell, M. R., C. J. Gress, E. H. Appleman & O. Jacenko (2004). "Chicken collagen X regulatory sequences restrict transgene expression to hypertrophic cartilage in mice." *Am J Pathol* 164(2): 487-499
- Chan, C. K., C. C. Chen, C. A. Luppen, J. B. Kim, A. T. Deboer, K. Wei, J. A. Helms, C. J. Kuo, D. L. Kraft & I. L. Weissman (2008). "Endochondral ossification is required for haematopoietic stem-cell niche formation." *Nature*
- Chan, D. & O. Jacenko (1998). "Phenotypic and biochemical consequences of collagen X mutations in mice and humans." *Matrix Biol* 17(3): 169-184
- Chang, M. K., L. J. Raggatt, K. A. Alexander, J. S. Kuliwaba, N. L. Fazzalari, K. Schroder, E. R. Maylin, V. M. Ripoll, D. A. Hume & A. R. Pettit (2008). "Osteal tissue macrophages are intercalated throughout human and mouse bone lining tissues and regulate osteoblast function in vitro and in vivo." *J Immunol* 181(2): 1232-1244
- Chow, D. C., L. A. Wenning, W. M. Miller & E. T. Papoutsakis (2001). "Modeling pO(2) distributions in the bone marrow hematopoietic compartment. I. Krogh's model." *Biophys J* 81(2): 675-684
- Compston, J. E. (2002). "Bone marrow and bone: a functional unit." *J Endocrinol* 173(3): 387-394
- Corradi, A., L. Croci, V. Broccoli, S. Zecchini, S. Previtali, W. Wurst, S. Amadio, R. Maggi, A. Quattrini & G. G. Consalez (2003). "Hypogonadotropic hypogonadism and peripheral neuropathy in Ebf2-null mice." *Development* 130(2): 401-410
- Corre, J., C. Barreau, B. Cousin, J. P. Chavoin, D. Caton, G. Fournial, L. Penicaud, L. Casteilla & P. Laharrague (2006). "Human subcutaneous adipose cells support complete differentiation but not self-renewal of hematopoietic progenitors." *J Cell Physiol* 208(2): 282-288

- Coskun, S. & K. K. Hirschi (2010). "Establishment and regulation of the HSC niche: Roles of osteoblastic and vascular compartments." *Birth Defects Res C Embryo Today* 90(4): 229-242
- Coulombel, L., I. Auffray, M. H. Gaugler & M. Roseblatt (1997). "Expression and function of integrins on hematopoietic progenitor cells." *Acta Haematol* 97(1-2): 13-21
- Crawford, L. J., R. Peake, S. Price, T. C. Morris & A. E. Irvine (2010). "Adiponectin is produced by lymphocytes and is a negative regulator of granulopoiesis." *J Leukoc Biol* 88(4): 807-811
- Crisan, M., S. Yap, L. Casteilla, C. W. Chen, M. Corselli, T. S. Park, G. Andriolo, B. Sun, B. Zheng, L. Zhang, C. Norotte, P. N. Teng, J. Traas, R. Schugar, B. M. Deasy, S. Badylak, H. J. Buhning, J. P. Giacobino, L. Lazzari, J. Huard & B. Peault (2008). "A perivascular origin for mesenchymal stem cells in multiple human organs." *Cell Stem Cell* 3(3): 301-313
- Crowley, J. E., L. S. Trembl, J. E. Stadanlick, E. Carpenter & M. P. Cancro (2005). "Homeostatic niche specification among naive and activated B cells: a growing role for the BLyS family of receptors and ligands." *Semin Immunol* 17(3): 193-199
- Cumano, A., F. Dieterlen-Lievre & I. Godin (1996). "Lymphoid potential, probed before circulation in mouse, is restricted to caudal intraembryonic splanchnopleura." *Cell* 86(6): 907-916
- Cumano, A. & I. Godin (2007). "Ontogeny of the hematopoietic system." *Annu Rev Immunol* 25: 745-785
- Dar, A., O. Kollet & T. Lapidot (2006). "Mutual, reciprocal SDF-1/CXCR4 interactions between hematopoietic and bone marrow stromal cells regulate human stem cell migration and development in NOD/SCID chimeric mice." *Exp Hematol* 34(8): 967-975
- de Bruijn, M. F., X. Ma, C. Robin, K. Ottersbach, M. J. Sanchez & E. Dzierzak (2002). "Hematopoietic stem cells localize to the endothelial cell layer in the midgestation mouse aorta." *Immunity* 16(5): 673-683
- Dexter, T. M., T. D. Allen & L. G. Lajtha (1977). "Conditions controlling the proliferation of haemopoietic stem cells in vitro." *J Cell Physiol* 91(3): 335-344
- Dexter, T. M., T. D. Allen, L. G. Lajtha, R. Schofield & B. I. Lord (1973). "Stimulation of differentiation and proliferation of haemopoietic cells in vitro." *J Cell Physiol* 82(3): 461-473
- Discher, D. E., D. J. Mooney & P. W. Zandstra (2009). "Growth factors, matrices, and forces combine and control stem cells." *Science* 324(5935): 1673-1677
- Dominici, M., V. Rasini, R. Bussolari, X. Chen, T. J. Hofmann, C. Spano, D. Bernabei, E. Veronesi, F. Bertoni, P. Paolucci, P. Conte & E. M. Horwitz (2009). "Restoration and reversible expansion of the osteoblastic hematopoietic stem cell niche after marrow radioablation." *Blood* 114(11): 2333-2343
- Eaves, C. J., J. D. Cashman, R. J. Kay, G. J. Dougherty, T. Otsuka, L. A. Gaboury, D. E. Hogge, P. M. Lansdorp, A. C. Eaves & R. K. Humphries (1991). "Mechanisms that regulate the cell cycle status of very primitive hematopoietic cells in long-term human marrow cultures. II. Analysis of positive and negative regulators produced by stromal cells within the adherent layer." *Blood* 78(1): 110-117

- Farnum, C. E. & N. J. Wilsman (1989). "Cellular turnover at the chondro-osseous junction of growth plate cartilage: analysis by serial sections at the light microscopical level." *J Orthop Res* 7(5): 654-666
- Fortunel, N., J. Hatzfeld, L. Aoustin, P. Batard, K. Ducos, M. N. Monier, A. Charpentier & A. Hatzfeld (2000). "Specific dose-response effects of TGF-beta1 on developmentally distinct hematopoietic stem/progenitor cells from human umbilical cord blood." *Hematol J* 1(2): 126-135
- Friedenstein, A. J., R. K. Chailakhyan & K. S. Lalykina (1970). "The development of fibroblast colonies in monolayer cultures of guinea-pig bone marrow and spleen cells." *Cell Tissue Kinet* 3(4): 393-403
- Friedenstein, A. J., R. K. Chailakhyan & U. V. Gerasimov (1987). "Bone marrow osteogenic stem cells: in vitro cultivation and transplantation in diffusion chambers." *Cell Tissue Kinet* 20(3): 263-272
- Friedenstein, A. J., R. K. Chailakhyan, N. V. Latsinik, A. F. Panasyuk & I. V. Keiliss-Borok (1974). "Stromal cells responsible for transferring the microenvironment of the hemopoietic tissues. Cloning in vitro and retransplantation in vivo." *Transplantation* 17(4): 331-340
- Friedenstein, A. J., J. F. Gorskaja & N. N. Kulagina (1976). "Fibroblast precursors in normal and irradiated mouse hematopoietic organs." *Exp Hematol* 4(5): 267-274
- Friedenstein, A. J., N. W. Latzinik, A. G. Grosheva & U. F. Gorskaya (1982). "Marrow microenvironment transfer by heterotopic transplantation of freshly isolated and cultured cells in porous sponges." *Exp Hematol* 10(2): 217-227
- Funk, P. E., R. P. Stephan & P. L. Witte (1995). "Vascular cell adhesion molecule 1-positive reticular cells express interleukin-7 and stem cell factor in the bone marrow." *Blood* 86(7): 2661-2671
- Galotto, M., G. Campanile, G. Robino, F. D. Cancedda, P. Bianco & R. Cancedda (1994). "Hypertrophic chondrocytes undergo further differentiation to osteoblast-like cells and participate in the initial bone formation in developing chick embryo." *J Bone Miner Res* 9(8): 1239-1249
- Gekas, C., F. Dieterlen-Lievre, S. H. Orkin & H. K. Mikkola (2005). "The placenta is a niche for hematopoietic stem cells." *Dev Cell* 8(3): 365-375
- Gibson, G. J. & M. H. Flint (1985). "Type X collagen synthesis by chick sternal cartilage and its relationship to endochondral development." *J Cell Biol* 101(1): 277-284
- Godman, G. C. & K. R. Porter (1960). "Chondrogenesis, studied with the electron microscope." *J Biophys Biochem Cytol* 8: 719-760
- Gong, J. K. (1978). "Endosteal marrow: a rich source of hematopoietic stem cells." *Science* 199(4336): 1443-1445
- Gordon, M. Y., G. P. Riley & D. Clarke (1988). "Heparan sulfate is necessary for adhesive interactions between human early hemopoietic progenitor cells and the extracellular matrix of the marrow microenvironment." *Leukemia* 2(12): 804-809
- Gress, C. J. & O. Jacenko (2000). "Growth plate compressions and altered hematopoiesis in collagen X null mice." *J Cell Biol* 149(4): 983-993
- Grimaud, E., F. Blanchard, C. Charrier, F. Gouin, F. Redini & D. Heymann (2002). "Leukaemia inhibitory factor (lif) is expressed in hypertrophic chondrocytes and vascular sprouts during osteogenesis." *Cytokine* 20(5): 224-230

- Gruver, A. L., L. L. Hudson & G. D. Sempowski (2007). "Immunosenescence of ageing." *J Pathol* 211(2): 144-156
- Guba, S. C., C. I. Sartor, L. R. Gottschalk, Y. H. Jing, T. Mulligan & S. G. Emerson (1992). "Bone marrow stromal fibroblasts secrete interleukin-6 and granulocyte-macrophage colony-stimulating factor in the absence of inflammatory stimulation: demonstration by serum-free bioassay, enzyme-linked immunosorbent assay, and reverse transcriptase polymerase chain reaction." *Blood* 80(5): 1190-1198
- Gupta, P., J. B. McCarthy & C. M. Verfaillie (1996). "Stromal fibroblast heparan sulfate is required for cytokine-mediated ex vivo maintenance of human long-term culture-initiating cells." *Blood* 87(8): 3229-3236
- Gupta, P., T. R. Oegema, Jr., J. J. Brazil, A. Z. Dudek, A. Slungaard & C. M. Verfaillie (1998). "Structurally specific heparan sulfates support primitive human hematopoiesis by formation of a multimolecular stem cell niche." *Blood* 92(12): 4641-4651
- Gurevitch, O. & I. Fabian (1993). "Ability of the hemopoietic microenvironment in the induced bone to maintain the proliferative potential of early hemopoietic precursors." *Stem Cells* 11(1): 56-61
- Hara, H., H. Ohdan, D. Tokita, T. Onoe, W. Zhou & T. Asahara (2003). "Construction of ectopic xenogeneic bone marrow structure associated with persistent multi-lineage mixed chimerism by engraftment of rat bone marrow plugs into mouse kidney capsules." *Xenotransplantation* 10(3): 259-266
- Harada, M., K. Odaka, K. Kondo, S. Nakao, M. Ueda, K. Matsue, T. Mori & T. Matsuda (1985). "Effect of activated lymphocytes on the regulation of hematopoiesis: suppression of in vitro granulopoiesis and erythropoiesis by OKT8+ Ia- T cells induced by concanavalin-A stimulation." *Exp Hematol* 13(9): 963-967
- Haylock, D. N. & S. K. Nilsson (2006). "The role of hyaluronic acid in hemopoietic stem cell biology." *Regen Med* 1(4): 437-445
- Heinrich, M. C., D. C. Dooley, A. C. Freed, L. Band, M. E. Hoatlin, W. W. Keeble, S. T. Peters, K. V. Silvey, F. S. Ey, D. Kabat & et al. (1993). "Constitutive expression of steel factor gene by human stromal cells." *Blood* 82(3): 771-783
- Heissig, B., K. Hattori, S. Dias, M. Friedrich, B. Ferris, N. R. Hackett, R. G. Crystal, P. Besmer, D. Lyden, M. A. Moore, Z. Werb & S. Rafii (2002). "Recruitment of stem and progenitor cells from the bone marrow niche requires MMP-9 mediated release of kit-ligand." *Cell* 109(5): 625-637
- Hermanns, P., A. A. Bertuch, T. K. Bertin, B. Dawson, M. E. Schmitt, C. Shaw, B. Zabel & B. Lee (2005). "Consequences of mutations in the non-coding RMRP RNA in cartilage-hair hypoplasia." *Hum Mol Genet* 14(23): 3723-3740
- Hermans, M. H., H. Hartsuiker & D. Opstelten (1989). "An in situ study of B-lymphocytopoiesis in rat bone marrow. Topographical arrangement of terminal deoxynucleotidyl transferase-positive cells and pre-B cells." *J Immunol* 142(1): 67-73
- Hilton, M. J., X. Tu & F. Long (2007). "Tamoxifen-inducible gene deletion reveals a distinct cell type associated with trabecular bone, and direct regulation of PTHrP expression and chondrocyte morphology by Ihh in growth region cartilage." *Dev Biol* 308(1): 93-105
- Holst, J., S. Watson, M. S. Lord, S. S. Eamegdool, D. V. Bax, L. B. Nivison-Smith, A. Kondyurin, L. Ma, A. F. Oberhauser, A. S. Weiss & J. E. Rasko (2010). "Substrate

- elasticity provides mechanical signals for the expansion of hemopoietic stem and progenitor cells." *Nat Biotechnol* 28(10): 1123-1128
- Horwitz, E. M., K. Le Blanc, M. Dominici, I. Mueller, I. Slaper-Cortenbach, F. C. Marini, R. J. Deans, D. S. Krause & A. Keating (2005). "Clarification of the nomenclature for MSC: The International Society for Cellular Therapy position statement." *Cytotherapy* 7(5): 393-395
- Hotamisligil, G. S., N. S. Shargill & B. M. Spiegelman (1993). "Adipose expression of tumor necrosis factor- α : direct role in obesity-linked insulin resistance." *Science* 259(5091): 87-91
- Huard, B., T. McKee, C. Bosshard, S. Durual, T. Matthes, S. Myit, O. Donze, C. Frossard, C. Chizzolini, C. Favre, R. Zubler, J. P. Guyot, P. Schneider & E. Roosnek (2008). "APRIL secreted by neutrophils binds to heparan sulfate proteoglycans to create plasma cell niches in human mucosa." *J Clin Invest* 118(8): 2887-2895
- Ingold, K., A. Zumsteg, A. Tardivel, B. Huard, Q. G. Steiner, T. G. Cachero, F. Qiang, L. Gorelik, S. L. Kalled, H. Acha-Orbea, P. D. Rennert, J. Tschopp & P. Schneider (2005). "Identification of proteoglycans as the APRIL-specific binding partners." *J Exp Med* 201(9): 1375-1383
- Jacenko, O., D. Chan, A. Franklin, S. Ito, C. B. Underhill, J. F. Bateman & M. R. Campbell (2001). "A dominant interference collagen X mutation disrupts hypertrophic chondrocyte pericellular matrix and glycosaminoglycan and proteoglycan distribution in transgenic mice." *Am J Pathol* 159(6): 2257-2269
- Jacenko, O., S. Ito & B. R. Olsen (1996). "Skeletal and hematopoietic defects in mice transgenic for collagen X." *Ann N Y Acad Sci* 785: 278-280
- Jacenko, O., P. A. LuValle & B. R. Olsen (1993). "Spondylometaphyseal dysplasia in mice carrying a dominant negative mutation in a matrix protein specific for cartilage-to-bone transition." *Nature* 365(6441): 56-61
- Jacenko, O., B. R. Olsen & P. LuValle (1991). "Organization and regulation of collagen genes." *Crit Rev Eukaryot Gene Expr* 1(4): 327-353
- Jacenko, O., D. W. Roberts, M. R. Campbell, P. M. McManus, C. J. Gress & Z. Tao (2002). "Linking hematopoiesis to endochondral skeletogenesis through analysis of mice transgenic for collagen X." *Am J Pathol* 160(6): 2019-2034
- Jacobsen, K. & D. G. Osmond (1990). "Microenvironmental organization and stromal cell associations of B lymphocyte precursor cells in mouse bone marrow." *Eur J Immunol* 20(11): 2395-2404
- James, C. G., L. A. Stanton, H. Agoston, V. Ulici, T. M. Underhill & F. Beier (2010). "Genome-wide analyses of gene expression during mouse endochondral ossification." *PLoS One* 5(1): e8693
- Jang, Y. Y. & S. J. Sharkis (2007). "A low level of reactive oxygen species selects for primitive hematopoietic stem cells that may reside in the low-oxygenic niche." *Blood* 110(8): 3056-3063
- Janzen, V., R. Forkert, H. E. Fleming, Y. Saito, M. T. Waring, D. M. Dombkowski, T. Cheng, R. A. DePinho, N. E. Sharpless & D. T. Scadden (2006). "Stem-cell ageing modified by the cyclin-dependent kinase inhibitor p16INK4a." *Nature* 443(7110): 421-426
- Jung, Y., J. Song, Y. Shiozawa, J. Wang, Z. Wang, B. Williams, A. Havens, A. Schneider, C. Ge, R. T. Franceschi, L. K. McCauley, P. H. Krebsbach & R. S. Taichman (2008). "Hematopoietic stem cells regulate mesenchymal stromal cell induction into

- osteoblasts thereby participating in the formation of the stem cell niche." *Stem Cells* 26(8): 2042-2051
- Jung, Y., J. Wang, A. Havens, Y. Sun, J. Wang, T. Jin & R. S. Taichman (2005). "Cell-to-cell contact is critical for the survival of hematopoietic progenitor cells on osteoblasts." *Cytokine* 32(3-4): 155-162
- Jung, Y., J. Wang, A. Schneider, Y. X. Sun, A. J. Koh-Paige, N. I. Osman, L. K. McCauley & R. S. Taichman (2006). "Regulation of SDF-1 (CXCL12) production by osteoblasts; a possible mechanism for stem cell homing." *Bone* 38(4): 497-508
- Kacena, M. A., C. M. Gundberg & M. C. Horowitz (2006). "A reciprocal regulatory interaction between megakaryocytes, bone cells, and hematopoietic stem cells." *Bone* 39(5): 978-984
- Kacena, M. A., R. A. Shivdasani, K. Wilson, Y. Xi, N. Troiano, A. Nazarian, C. M. Gundberg, M. L. Bouxsein, J. A. Lorenzo & M. C. Horowitz (2004). "Megakaryocyte-osteoblast interaction revealed in mice deficient in transcription factors GATA-1 and NF-E2." *J Bone Miner Res* 19(4): 652-660
- Kalajzic, Z., H. Li, L. P. Wang, X. Jiang, K. Lamothe, D. J. Adams, H. L. Aguila, D. W. Rowe & I. Kalajzic (2008). "Use of an alpha-smooth muscle actin GFP reporter to identify an osteoprogenitor population." *Bone* 43(3): 501-510
- Kalinkovich, A., A. Spiegel, S. Shvitiel, O. Kollet, N. Jordaney, W. Piacibello & T. Lapidot (2009). "Blood-forming stem cells are nervous: direct and indirect regulation of immature human CD34+ cells by the nervous system." *Brain Behav Immun* 23(8): 1059-1065
- Katayama, Y., M. Battista, W. M. Kao, A. Hidalgo, A. J. Peired, S. A. Thomas & P. S. Frenette (2006). "Signals from the sympathetic nervous system regulate hematopoietic stem cell egress from bone marrow." *Cell* 124(2): 407-421
- Kawai, M., H. Hattori, K. Yasue, H. Mizutani, M. Ueda, T. Kaneda & T. Hoshino (1994). "Development of hemopoietic bone marrow within the ectopic bone induced by bone morphogenetic protein." *Blood Cells* 20(1): 191-199; discussion 200-191
- Keller, J. R., J. M. Gooya & F. W. Ruscetti (1996). "Direct synergistic effects of leukemia inhibitory factor on hematopoietic progenitor cell growth: comparison with other hematopoietins that use the gp130 receptor subunit." *Blood* 88(3): 863-869
- Khosla, S. (2003). "Parathyroid hormone plus alendronate--a combination that does not add up." *N Engl J Med* 349(13): 1277-1279
- Kiel, M. J., M. Acar, G. L. Radice & S. J. Morrison (2009). "Hematopoietic stem cells do not depend on N-cadherin to regulate their maintenance." *Cell Stem Cell* 4(2): 170-179
- Kiel, M. J., G. L. Radice & S. J. Morrison (2007). "Lack of evidence that hematopoietic stem cells depend on N-cadherin-mediated adhesion to osteoblasts for their maintenance." *Cell Stem Cell* 1(2): 204-217
- Kiel, M. J., O. H. Yilmaz, T. Iwashita, O. H. Yilmaz, C. Terhorst & S. J. Morrison (2005). "SLAM family receptors distinguish hematopoietic stem and progenitor cells and reveal endothelial niches for stem cells." *Cell* 121(7): 1109-1121
- Kieslinger, M., S. Hiechinger, G. Dobrev, G. G. Consalez & R. Grosschedl (2010). "Early B cell factor 2 regulates hematopoietic stem cell homeostasis in a cell-nonautonomous manner." *Cell Stem Cell* 7(4): 496-507

- Kilborn, S. H., G. Trudel & H. Uthoff (2002). "Review of growth plate closure compared with age at sexual maturity and lifespan in laboratory animals." *Contemp Top Lab Anim Sci* 41(5): 21-26
- Klein, G. (1995). "The extracellular matrix of the hematopoietic microenvironment." *Experientia* 51(9-10): 914-926
- Klein, G., S. Conzelmann, S. Beck, R. Timpl & C. A. Muller (1995). "Perlecan in human bone marrow: a growth-factor-presenting, but anti-adhesive, extracellular matrix component for hematopoietic cells." *Matrix Biol* 14(6): 457-465
- Klein, G., C. Kibler, F. Schermutzki, J. Brown, C. A. Muller & R. Timpl (1998). "Cell binding properties of collagen type XIV for human hematopoietic cells." *Matrix Biol* 16(6): 307-317
- Kohler, A., V. Schmithorst, M. D. Filippi, M. A. Ryan, D. Daria, M. Gunzer & H. Geiger (2009). "Altered cellular dynamics and endosteal location of aged early hematopoietic progenitor cells revealed by time-lapse intravital imaging in long bones." *Blood* 114(2): 290-298
- Kollet, O., A. Dar, S. Shivtiel, A. Kalinkovich, K. Lapid, Y. Sztainberg, M. Tesio, R. M. Samstein, P. Goichberg, A. Spiegel, A. Elson & T. Lapidot (2006). "Osteoclasts degrade endosteal components and promote mobilization of hematopoietic progenitor cells." *Nat Med* 12(6): 657-664
- Kong, Y. Y., H. Yoshida, I. Sarosi, H. L. Tan, E. Timms, C. Capparelli, S. Morony, A. J. Oliveira-dos-Santos, G. Van, A. Itie, W. Khoo, A. Wakeham, C. R. Dunstan, D. L. Lacey, T. W. Mak, W. J. Boyle & J. M. Penninger (1999). "OPGL is a key regulator of osteoclastogenesis, lymphocyte development and lymph-node organogenesis." *Nature* 397(6717): 315-323
- Kopp, H. G., S. T. Avezilla, A. T. Hooper & S. Rafii (2005). "The bone marrow vascular niche: home of HSC differentiation and mobilization." *Physiology (Bethesda)* 20: 349-356
- Kortesidis, A., A. Zannettino, S. Isenmann, S. Shi, T. Lapidot & S. Gronthos (2005). "Stromal-derived factor-1 promotes the growth, survival, and development of human bone marrow stromal stem cells." *Blood* 105(10): 3793-3801
- Kuijpers, T. W., E. Nannenberg, M. Alders, R. Bredius & R. C. Hennekam (2004). "Congenital aplastic anemia caused by mutations in the SBDS gene: a rare presentation of Shwachman-Diamond syndrome." *Pediatrics* 114(3): e387-391
- Kuznetsov, S. A., P. H. Krebsbach, K. Satomura, J. Kerr, M. Riminucci, D. Benayahu & P. G. Robey (1997). "Single-colony derived strains of human marrow stromal fibroblasts form bone after transplantation in vivo." *J Bone Miner Res* 12(9): 1335-1347
- Kuznetsov, S. A., M. Riminucci, N. Ziran, T. W. Tsutsui, A. Corsi, L. Calvi, H. M. Kronenberg, E. Schipani, P. G. Robey & P. Bianco (2004). "The interplay of osteogenesis and hematopoiesis: expression of a constitutively active PTH/PTHrP receptor in osteogenic cells perturbs the establishment of hematopoiesis in bone and of skeletal stem cells in the bone marrow." *J Cell Biol* 167(6): 1113-1122
- Kwan, K. M., M. K. Pang, S. Zhou, S. K. Cowan, R. Y. Kong, T. Pfordte, B. R. Olsen, D. O. Sillence, P. P. Tam & K. S. Cheah (1997). "Abnormal compartmentalization of cartilage matrix components in mice lacking collagen X: implications for function." *J Cell Biol* 136(2): 459-471

- Kwon, K. R., J. Y. Ahn, M. S. Kim, J. Y. Jung, J. H. Lee & I. H. Oh (2010). "Disruption of *bis* leads to the deterioration of the vascular niche for hematopoietic stem cells." *Stem Cells* 28(2): 268-278
- Lanotte, M., D. Metcalf & T. M. Dexter (1982). "Production of monocyte/macrophage colony-stimulating factor by preadipocyte cell lines derived from murine marrow stroma." *J Cell Physiol* 112(1): 123-127
- Lapid, K., Y. Vagima, O. Kollet & T. Lapidot (2008). "Egress and mobilization of hematopoietic stem and progenitor cells."
- Larsson, J., M. Ohishi, B. Garrison, M. Aspling, V. Janzen, G. B. Adams, M. Curto, A. I. McClatchey, E. Schipani & D. T. Scadden (2008). "Nf2/merlin regulates hematopoietic stem cell behavior by altering microenvironmental architecture." *Cell Stem Cell* 3(2): 221-227
- Lefebvre, V. & P. Smits (2005). "Transcriptional control of chondrocyte fate and differentiation." *Birth Defects Res Part C: Embryo Today: Reviews* 75(3): 200-212
- Levesque, J. P. & P. J. Simmons (1999). "Cytoskeleton and integrin-mediated adhesion signaling in human CD34+ hemopoietic progenitor cells." *Exp Hematol* 27(4): 579-586
- Levesque, J. P., Y. Takamatsu, S. K. Nilsson, D. N. Haylock & P. J. Simmons (2001). "Vascular cell adhesion molecule-1 (CD106) is cleaved by neutrophil proteases in the bone marrow following hematopoietic progenitor cell mobilization by granulocyte colony-stimulating factor." *Blood* 98(5): 1289-1297
- Li, W., S. A. Johnson, W. C. Shelley & M. C. Yoder (2004). "Hematopoietic stem cell repopulating ability can be maintained in vitro by some primary endothelial cells." *Exp Hematol* 32(12): 1226-1237
- Lo Celso, C., H. E. Fleming, J. W. Wu, C. X. Zhao, S. Miake-Lye, J. Fujisaki, D. Cote, D. W. Rowe, C. P. Lin & D. T. Scadden (2009). "Live-animal tracking of individual haematopoietic stem/progenitor cells in their niche." *Nature* 457(7225): 92-96
- Lord, B. I., N. G. Testa & J. H. Hendry (1975). "The relative spatial distributions of CFUs and CFUc in the normal mouse femur." *Blood* 46(1): 65-72
- Mackie, E. J., Y. A. Ahmed, L. Tatarczuch, K. S. Chen & M. Mirams (2008). "Endochondral ossification: how cartilage is converted into bone in the developing skeleton." *Int J Biochem Cell Biol* 40(1): 46-62
- Maes, C., T. Kobayashi, M. K. Selig, S. Torrekens, S. I. Roth, S. Mackem, G. Carmeliet & H. M. Kronenberg (2010). "Osteoblast precursors, but not mature osteoblasts, move into developing and fractured bones along with invading blood vessels." *Dev Cell* 19(2): 329-344
- Majumdar, M. K., M. A. Thiede, S. E. Haynesworth, S. P. Bruder & S. L. Gerson (2000). "Human marrow-derived mesenchymal stem cells (MSCs) express hematopoietic cytokines and support long-term hematopoiesis when differentiated toward stromal and osteogenic lineages." *J Hematother Stem Cell Res* 9(6): 841-848
- Mangan, K. F., G. Chikkappa, L. Z. Bieler, W. B. Scharfman & D. R. Parkinson (1982). "Regulation of human blood erythroid burst-forming unit (BFU-E) proliferation by T-lymphocyte subpopulations defined by Fc receptors and monoclonal antibodies." *Blood* 59(5): 990-996

- Mankani, M. H., S. A. Kuznetsov, G. W. Marshall & P. G. Robey (2008). "Creation of new bone by the percutaneous injection of human bone marrow stromal cell and HA/TCP suspensions." *Tissue Eng Part A* 14(12): 1949-1958
- Mankani, M. H., S. A. Kuznetsov & P. G. Robey (2007). "Formation of hematopoietic territories and bone by transplanted human bone marrow stromal cells requires a critical cell density." *Exp Hematol* 35(6): 995-1004
- Mansour, A., A. Anginot, S. J. Mancini, C. Schiff, G. F. Carle, A. Wakkach & C. Blin-Wakkach (2011). "Osteoclast activity modulates B-cell development in the bone marrow." *Cell Res* 21(7): 1102-1115
- Matrosova, V. Y., I. A. Orlovskaya, N. Serobyanyan & S. K. Khaldoyanidi (2004). "Hyaluronic acid facilitates the recovery of hematopoiesis following 5-fluorouracil administration." *Stem Cells* 22(4): 544-555
- Mayack, S. R., J. L. Shadrach, F. S. Kim & A. J. Wagers (2010). "Systemic signals regulate ageing and rejuvenation of blood stem cell niches." *Nature* 463(7280): 495-500
- Mazzon, C., A. Anselmo, J. Cibella, C. Soldani, A. Destro, N. Kim, M. Roncalli, S. J. Burden, M. L. Dustin, A. Sarukhan & A. Viola (2011). "The critical role of agrin in the hematopoietic stem cell niche." *Blood* 118(10): 2733-2742
- Meck, R. A., J. E. Haley & G. Brecher (1973). "Hematopoiesis versus osteogenesis in ectopic bone marrow transplants." *Blood* 42(5): 661-669
- Medici, D., E. M. Shore, V. Y. Lounev, F. S. Kaplan, R. Kalluri & B. R. Olsen (2010). "Conversion of vascular endothelial cells into multipotent stem-like cells." *Nat Med* 16(12): 1400-1406
- Medvinsky, A. & E. Dzierzak (1996). "Definitive hematopoiesis is autonomously initiated by the AGM region." *Cell* 86(6): 897-906
- Mendez-Ferrer, S., D. Lucas, M. Battista & P. S. Frenette (2008). "Haematopoietic stem cell release is regulated by circadian oscillations." *Nature* 452(7186): 442-447
- Mendez-Ferrer, S., T. V. Michurina, F. Ferraro, A. R. Mazloom, B. D. Macarthur, S. A. Lira, D. T. Scadden, A. Ma'ayan, G. N. Enikolopov & P. S. Frenette (2010). "Mesenchymal and haematopoietic stem cells form a unique bone marrow niche." *Nature* 466(7308): 829-834
- Merzaban, J. S., M. M. Burdick, S. Z. Gadhoun, N. M. Dagia, J. T. Chu, R. C. Fuhlbrigge & R. Sackstein (2011). "Analysis of glycoprotein E-selectin ligands on human and mouse marrow cells enriched for hematopoietic stem/progenitor cells." *Blood* 118(7): 1774-1783
- Mignini, F., V. Streccioni & F. Amenta (2003). "Autonomic innervation of immune organs and neuroimmune modulation." *Auton Autacoid Pharmacol* 23(1): 1-25
- Miharada, K., T. Hiroyama, K. Sudo, I. Danjo, T. Nagasawa & Y. Nakamura (2008). "Lipocalin 2-mediated growth suppression is evident in human erythroid and monocyte/macrophage lineage cells." *J Cell Physiol* 215(2): 526-537
- Moreaux, J., F. W. Cremer, T. Reme, M. Raab, K. Mahtouk, P. Kaukel, V. Pantesco, J. De Vos, E. Jourdan, A. Jauch, E. Legouffe, M. Moos, G. Fiol, H. Goldschmidt, J. F. Rossi, D. Hose & B. Klein (2005). "The level of TACI gene expression in myeloma cells is associated with a signature of microenvironment dependence versus a plasmablastic signature." *Blood* 106(3): 1021-1030
- Moreaux, J., A. C. Sprynski, S. R. Dillon, K. Mahtouk, M. Jourdan, A. Ythier, P. Moine, N. Robert, E. Jourdan, J. F. Rossi & B. Klein (2009). "APRIL and TACI interact with

- syndecan-1 on the surface of multiple myeloma cells to form an essential survival loop." *Eur J Haematol* 83(2): 119-129
- Morikawa, S., Y. Mabuchi, Y. Kubota, Y. Nagai, K. Niiibe, E. Hiratsu, S. Suzuki, C. Miyauchi-Hara, N. Nagoshi, T. Sunabori, S. Shimmura, A. Miyawaki, T. Nakagawa, T. Suda, H. Okano & Y. Matsuzaki (2009). "Prospective identification, isolation, and systemic transplantation of multipotent mesenchymal stem cells in murine bone marrow." *J Exp Med* 206(11): 2483-2496
- Morrison, S. J. & A. C. Spradling (2008). "Stem cells and niches: mechanisms that promote stem cell maintenance throughout life." *Cell* 132(4): 598-611
- Muguruma, Y., T. Yahata, H. Miyatake, T. Sato, T. Uno, J. Itoh, S. Kato, M. Ito, T. Hotta & K. Ando (2006). "Reconstitution of the functional human hematopoietic microenvironment derived from human mesenchymal stem cells in the murine bone marrow compartment." *Blood* 107(5): 1878-1887
- Murayama, E., K. Kissa, A. Zapata, E. Mordelet, V. Briolat, H. F. Lin, R. I. Handin & P. Herbomel (2006). "Tracing hematopoietic precursor migration to successive hematopoietic organs during zebrafish development." *Immunity* 25(6): 963-975
- Nagasawa, T. (2006). "Microenvironmental niches in the bone marrow required for B-cell development." *Nat Rev Immunol* 6(2): 107-116
- Nagasawa, T., S. Hirota, K. Tachibana, N. Takakura, S. Nishikawa, Y. Kitamura, N. Yoshida, H. Kikutani & T. Kishimoto (1996). "Defects of B-cell lymphopoiesis and bone-marrow myelopoiesis in mice lacking the CXC chemokine PBSF/SDF-1." *Nature* 382(6592): 635-638
- Nagasawa, T., H. Kikutani & T. Kishimoto (1994). "Molecular cloning and structure of a pre-B-cell growth-stimulating factor." *Proc Natl Acad Sci U S A* 91(6): 2305-2309
- Naiyer, A. J., D. Y. Jo, J. Ahn, R. Mohle, M. Peichev, G. Lam, R. L. Silverstein, M. A. Moore & S. Rafii (1999). "Stromal derived factor-1-induced chemokinesis of cord blood CD34(+) cells (long-term culture-initiating cells) through endothelial cells is mediated by E-selectin." *Blood* 94(12): 4011-4019
- Nakamura, E., M. T. Nguyen & S. Mackem (2006). "Kinetics of tamoxifen-regulated Cre activity in mice using a cartilage-specific CreER(T) to assay temporal activity windows along the proximodistal limb skeleton." *Dev Dyn* 235(9): 2603-2612
- Nathan, D. G., L. Chess, D. G. Hillman, B. Clarke, J. Breard, E. Merler & D. E. Housman (1978). "Human erythroid burst-forming unit: T-cell requirement for proliferation in vitro." *J Exp Med* 147(2): 324-339
- Naveiras, O., V. Nardi, P. L. Wenzel, P. V. Hauschka, F. Fahey & G. Q. Daley (2009). "Bone-marrow adipocytes as negative regulators of the haematopoietic microenvironment." *Nature* 460(7252): 259-263
- Nilsson, S. K., M. S. Dooner, C. Y. Tiarks, H. U. Weier & P. J. Quesenberry (1997). "Potential and distribution of transplanted hematopoietic stem cells in a nonablated mouse model." *Blood* 89(11): 4013-4020
- Nilsson, S. K., H. M. Johnston & J. A. Coverdale (2001). "Spatial localization of transplanted hemopoietic stem cells: inferences for the localization of stem cell niches." *Blood* 97(8): 2293-2299
- North, T. E., M. F. de Bruijn, T. Stacy, L. Talebian, E. Lind, C. Robin, M. Binder, E. Dzierzak & N. A. Speck (2002). "Runx1 expression marks long-term repopulating

- hematopoietic stem cells in the midgestation mouse embryo." *Immunity* 16(5): 661-672
- O'Connor, B. P., V. S. Raman, L. D. Erickson, W. J. Cook, L. K. Weaver, C. Ahonen, L. L. Lin, G. T. Mantchev, R. J. Bram & R. J. Noelle (2004). "BCMA is essential for the survival of long-lived bone marrow plasma cells." *J Exp Med* 199(1): 91-98
- Ohneda, O., C. Fennie, Z. Zheng, C. Donahue, H. La, R. Villacorta, B. Cairns & L. A. Lasky (1998). "Hematopoietic stem cell maintenance and differentiation are supported by embryonic aorta-gonad-mesonephros region-derived endothelium." *Blood* 92(3): 908-919
- Ohta, M., T. Sakai, Y. Saga, S. Aizawa & M. Saito (1998). "Suppression of hematopoietic activity in tenascin-C-deficient mice." *Blood* 91(11): 4074-4083
- Omatsu, Y., T. Sugiyama, H. Kohara, G. Kondoh, N. Fujii, K. Kohno & T. Nagasawa (2010). "The essential functions of adipo-osteogenic progenitors as the hematopoietic stem and progenitor cell niche." *Immunity* 33(3): 387-399
- Osmond, D. G. (1990). "B cell development in the bone marrow." *Semin Immunol* 2(3): 173-180
- Ottersbach, K. & E. Dzierzak (2005). "The murine placenta contains hematopoietic stem cells within the vascular labyrinth region." *Dev Cell* 8(3): 377-387
- Patt, H. M., M. A. Maloney & M. L. Flannery (1982). "Hematopoietic microenvironment transfer by stromal fibroblasts derived from bone marrow varying in cellularity." *Exp Hematol* 10(9): 738-742
- Peerani, R. & P. W. Zandstra (2010). "Enabling stem cell therapies through synthetic stem cell-niche engineering." *J Clin Invest* 120(1): 60-70
- Peled, A., V. Grabovsky, L. Habler, J. Sandbank, F. Arenzana-Seisdedos, I. Petit, H. Ben-Hur, T. Lapidot & R. Alon (1999). "The chemokine SDF-1 stimulates integrin-mediated arrest of CD34(+) cells on vascular endothelium under shear flow." *J Clin Invest* 104(9): 1199-1211
- Petit, I., M. Szyper-Kravitz, A. Nagler, M. Lahav, A. Peled, L. Habler, T. Ponomaryov, R. S. Taichman, F. Arenzana-Seisdedos, N. Fujii, J. Sandbank, D. Zipori & T. Lapidot (2002). "G-CSF induces stem cell mobilization by decreasing bone marrow SDF-1 and up-regulating CXCR4." *Nat Immunol* 3(7): 687-694
- Petrini, M., S. Pacini, L. Trombi, R. Fazzi, M. Montali, S. Ikehara & N. G. Abraham (2009). "Identification and purification of mesodermal progenitor cells from human adult bone marrow." *Stem Cells Dev* 18(6): 857-866
- Pfeiffer, C. A. (1948). "Development of bone from transplanted marrow in mice." *Anat Rec* 102(2): 225-243
- Pilia, G., R. M. Hughes-Benzie, A. MacKenzie, P. Baybayan, E. Y. Chen, R. Huber, G. Neri, A. Cao, A. Forabosco & D. Schlessinger (1996). "Mutations in GPC3, a glypican gene, cause the Simpson-Golabi-Behmel overgrowth syndrome." *Nat Genet* 12(3): 241-247
- Ponomaryov, T., A. Peled, I. Petit, R. S. Taichman, L. Habler, J. Sandbank, F. Arenzana-Seisdedos, A. Magerus, A. Caruz, N. Fujii, A. Nagler, M. Lahav, M. Szyper-Kravitz, D. Zipori & T. Lapidot (2000). "Induction of the chemokine stromal-derived factor-1 following DNA damage improves human stem cell function." *J Clin Invest* 106(11): 1331-1339

- Purton, L. E., S. Dworkin, G. H. Olsen, C. R. Walkley, S. A. Fabb, S. J. Collins & P. Chambon (2006). "RARgamma is critical for maintaining a balance between hematopoietic stem cell self-renewal and differentiation." *J Exp Med* 203(5): 1283-1293
- Raaijmakers, M. H., S. Mukherjee, S. Guo, S. Zhang, T. Kobayashi, J. A. Schoonmaker, B. L. Ebert, F. Al-Shahrour, R. P. Hasserjian, E. O. Scadden, Z. Aung, M. Matza, M. Merckenschlager, C. Lin, J. M. Rommens & D. T. Scadden (2010). "Bone progenitor dysfunction induces myelodysplasia and secondary leukaemia." *Nature* 464(7290): 852-857
- Rafii, S., R. Mohle, F. Shapiro, B. M. Frey & M. A. Moore (1997). "Regulation of hematopoiesis by microvascular endothelium." *Leuk Lymphoma* 27(5-6): 375-386
- Rafii, S., F. Shapiro, R. Pettengell, B. Ferris, R. L. Nachman, M. A. Moore & A. S. Asch (1995). "Human bone marrow microvascular endothelial cells support long-term proliferation and differentiation of myeloid and megakaryocytic progenitors." *Blood* 86(9): 3353-3363
- Rafii, S., F. Shapiro, J. Rimarachin, R. L. Nachman, B. Ferris, B. Weksler, M. A. Moore & A. S. Asch (1994). "Isolation and characterization of human bone marrow microvascular endothelial cells: hematopoietic progenitor cell adhesion." *Blood* 84(1): 10-19
- Roach, H. I. (1992). "Trans-differentiation of hypertrophic chondrocytes into cells capable of producing a mineralized bone matrix." *Bone Miner* 19(1): 1-20
- Roach, H. I. & J. Erenpreisa (1996). "The phenotypic switch from chondrocytes to bone-forming cells involves asymmetric cell division and apoptosis." *Connect Tissue Res* 35(1-4): 85-91
- Roach, H. I., J. Erenpreisa & T. Aigner (1995). "Osteogenic differentiation of hypertrophic chondrocytes involves asymmetric cell divisions and apoptosis." *J Cell Biol* 131(2): 483-494
- Roberts, R., J. Gallagher, E. Spooncer, T. D. Allen, F. Bloomfield & T. M. Dexter (1988). "Heparan sulphate bound growth factors: a mechanism for stromal cell mediated haemopoiesis." *Nature* 332(6162): 376-378
- Rodgers, K. D., J. D. San Antonio & O. Jacenko (2008). "Heparan sulfate proteoglycans: A GAGgle of skeletal-hematopoietic regulators." *Dev Dyn* 237(10): 2622-2642
- Rosati, R., G. S. Horan, G. J. Pinero, S. Garofalo, D. R. Keene, W. A. Horton, E. Vuorio, B. de Crombrughe & R. R. Behringer (1994). "Normal long bone growth and development in type X collagen-null mice." *Nat Genet* 8(2): 129-135
- Rossi, D. J., D. Bryder, J. M. Zahn, H. Ahlenius, R. Sonu, A. J. Wagers & I. L. Weissman (2005). "Cell intrinsic alterations underlie hematopoietic stem cell aging." *Proc Natl Acad Sci U S A* 102(26): 9194-9199
- Rossi, D. J., C. H. Jamieson & I. L. Weissman (2008). "Stems cells and the pathways to aging and cancer." *Cell* 132(4): 681-696
- Rouleau, M. F., J. Mitchell & D. Goltzman (1990). "Characterization of the major parathyroid hormone target cell in the endosteal metaphysis of rat long bones." *J Bone Miner Res* 5(10): 1043-1053
- Ryan, D. H., B. L. Nuccie, C. N. Abboud & J. M. Winslow (1991). "Vascular cell adhesion molecule-1 and the integrin VLA-4 mediate adhesion of human B cell precursors to cultured bone marrow adherent cells." *J Clin Invest* 88(3): 995-1004
- Sacchetti, B., A. Funari, S. Michienzi, S. Di Cesare, S. Piersanti, I. Saggio, E. Tagliafico, S. Ferrari, P. G. Robey, M. Riminucci & P. Bianco (2007). "Self-renewing

- osteoprogenitors in bone marrow sinusoids can organize a hematopoietic microenvironment." *Cell* 131(2): 324-336
- Sagar, B. M., S. Rentala, P. N. Gopal, S. Sharma & A. Mukhopadhyay (2006). "Fibronectin and laminin enhance engraftability of cultured hematopoietic stem cells." *Biochem Biophys Res Commun* 350(4): 1000-1005
- Sanchez, M. J., A. Holmes, C. Miles & E. Dzierzak (1996). "Characterization of the first definitive hematopoietic stem cells in the AGM and liver of the mouse embryo." *Immunity* 5(6): 513-525
- Schmid, T. M. & T. F. Linsenmayer (1985). "Immunohistochemical localization of short chain cartilage collagen (type X) in avian tissues." *J Cell Biol* 100(2): 598-605
- Schofield, R. (1978). "The relationship between the spleen colony-forming cell and the haemopoietic stem cell." *Blood Cells* 4(1-2): 7-25
- Schwaller, J., P. Schneider, P. Mhaweche-Fauceglia, T. McKee, S. Myit, T. Matthes, J. Tschopp, O. Donze, F. A. Le Gal & B. Huard (2007). "Neutrophil-derived APRIL concentrated in tumor lesions by proteoglycans correlates with human B-cell lymphoma aggressiveness." *Blood* 109(1): 331-338
- Schweitzer, K. M., A. M. Drager, P. van der Valk, S. F. Thijsen, A. Zevenbergen, A. P. Theijssmeijer, C. E. van der Schoot & M. M. Langenhuijsen (1996). "Constitutive expression of E-selectin and vascular cell adhesion molecule-1 on endothelial cells of hematopoietic tissues." *Am J Pathol* 148(1): 165-175
- Shackney, S. E., S. S. Ford & A. B. Wittig (1975). "Kinetic-microarchitectural correlations in the bone marrow of the mouse." *Cell Tissue Kinet* 8(6): 505-516
- Siczkowski, M., D. Clarke & M. Y. Gordon (1992). "Binding of primitive hematopoietic progenitor cells to marrow stromal cells involves heparan sulfate." *Blood* 80(4): 912-919
- Siebertz, B., G. Stocker, Z. Drzeniek, S. Handt, U. Just & H. D. Haubeck (1999). "Expression of glypican-4 in haematopoietic-progenitor and bone-marrow-stromal cells." *Biochem J* 344 (Pt 3): 937-943
- Sipkins, D. A., X. Wei, J. W. Wu, J. M. Runnels, D. Cote, T. K. Means, A. D. Luster, D. T. Scadden & C. P. Lin (2005). "In vivo imaging of specialized bone marrow endothelial microdomains for tumour engraftment." *Nature* 435(7044): 969-973
- Song, J., M. J. Kiel, Z. Wang, J. Wang, R. S. Taichman, S. J. Morrison & P. H. Krebsbach (2010). "An in vivo model to study and manipulate the hematopoietic stem cell niche." *Blood* 115(13): 2592-2600
- Spiegel, A., A. Kalinkovich, S. Shvitiel, O. Kollet & T. Lapidot (2008). "Stem cell regulation via dynamic interactions of the nervous and immune systems with the microenvironment." *Cell Stem Cell* 3(5): 484-492
- Spiegel, A., S. Shvitiel, A. Kalinkovich, A. Ludin, N. Netzer, P. Goichberg, Y. Azaria, I. Resnick, I. Hardan, H. Ben-Hur, A. Nagler, M. Rubinstein & T. Lapidot (2007). "Catecholaminergic neurotransmitters regulate migration and repopulation of immature human CD34+ cells through Wnt signaling." *Nat Immunol* 8(10): 1123-1131
- Spiegel, A., E. Zcharia, Y. Vagima, T. Itkin, A. Kalinkovich, A. Dar, O. Kollet, N. Netzer, K. Golan, I. Shafat, N. Ilan, A. Nagler, I. Vlodavsky & T. Lapidot (2008). "Heparanase regulates retention and proliferation of primitive Sca-1+/c-Kit+/Lin- cells via modulation of the bone marrow microenvironment." *Blood* 111(10): 4934-4943

- Spooncer, E., J. T. Gallagher, F. Krizsa & T. M. Dexter (1983). "Regulation of haemopoiesis in long-term bone marrow cultures. IV. Glycosaminoglycan synthesis and the stimulation of haemopoiesis by beta-D-xylosides." *J Cell Biol* 96(2): 510-514
- Spranger, J., G. K. Hinkel, H. Stoss, W. Thoenes, D. Wargowski & F. Zepp (1991). "Schimke immuno-osseous dysplasia: a newly recognized multisystem disease." *J Pediatr* 119(1 (Pt 1)): 64-72
- Srinivas, V. & I. M. Shapiro (2006). "Chondrocytes embedded in the epiphyseal growth plates of long bones undergo autophagy prior to the induction of osteogenesis." *Autophagy* 2(3): 215-216
- Sugiyama, T., H. Kohara, M. Noda & T. Nagasawa (2006). "Maintenance of the hematopoietic stem cell pool by CXCL12-CXCR4 chemokine signaling in bone marrow stromal cell niches." *Immunity* 25(6): 977-988
- Sweeney, E., M. Campbell, K. Watkins, C. A. Hunter & O. Jacenko (2008). "Altered endochondral ossification in collagen X mouse models leads to impaired immune responses." *Dev Dyn* 237(10): 2693-2704
- Sweeney, E., D. Roberts, T. Corbo & O. Jacenko (2010). "Congenic mice confirm that collagen X is required for proper hematopoietic development." *PLoS One* 5(3): e9518
- Taichman, R. S. & S. G. Emerson (1994). "Human osteoblasts support hematopoiesis through the production of granulocyte colony-stimulating factor." *J Exp Med* 179(5): 1677-1682
- Taichman, R. S., M. J. Reilly & S. G. Emerson (1996). "Human osteoblasts support human hematopoietic progenitor cells in vitro bone marrow cultures." *Blood* 87(2): 518-524
- Taichman, R. S., M. J. Reilly, R. S. Verma & S. G. Emerson (1997). "Augmented production of interleukin-6 by normal human osteoblasts in response to CD34+ hematopoietic bone marrow cells in vitro." *Blood* 89(4): 1165-1172
- Tavassoli, M. (1984). "Hemopoiesis in ectopically implanted bone marrow." *Kroc Found Ser* 18: 31-54
- Tavassoli, M. & W. H. Crosby (1968). "Transplantation of marrow to extramedullary sites." *Science* 161(836): 54-56
- Tavassoli, M. & R. Khademi (1980). "The origin of hemopoietic cells in ectopic implants of spleen and marrow." *Experientia* 36(9): 1126-1127
- Tavassoli, M., A. Maniatis & W. H. Crosby (1974). "Induction of sustained hemopoiesis in fatty marrow." *Blood* 43(1): 33-38
- Tavassoli, M. & L. Weiss (1971). "The structure of developing bone marrow sinuses in extramedullary autotransplant of the marrow in rats." *Anat Rec* 171(4): 477-494
- Teixido, J., M. E. Hemler, J. S. Greenberger & P. Anklesaria (1992). "Role of beta 1 and beta 2 integrins in the adhesion of human CD34hi stem cells to bone marrow stroma." *J Clin Invest* 90(2): 358-367
- Tokoyoda, K., T. Egawa, T. Sugiyama, B. I. Choi & T. Nagasawa (2004). "Cellular niches controlling B lymphocyte behavior within bone marrow during development." *Immunity* 20(6): 707-718
- Touw, I. & B. Lowenberg (1983). "No stimulative effect of adipocytes on hematopoiesis in long-term human bone marrow cultures." *Blood* 61(4): 770-774
- Trentin, J. J. (1971). "Determination of bone marrow stem cell differentiation by stromal hemopoietic inductive microenvironments (HIM)." *Am J Pathol* 65(3): 621-628

- Trinchieri, G., M. Murphy & B. Perussia (1987). "Regulation of hematopoiesis by T lymphocytes and natural killer cells." *Crit Rev Oncol Hematol* 7(3): 219-265
- Verfaillie, C. M. (1993). "Soluble factor(s) produced by human bone marrow stroma increase cytokine-induced proliferation and maturation of primitive hematopoietic progenitors while preventing their terminal differentiation." *Blood* 82(7): 2045-2053
- Verma, S., J. H. Rajaratnam, J. Denton, J. A. Hoyland & R. J. Byers (2002). "Adipocytic proportion of bone marrow is inversely related to bone formation in osteoporosis." *J Clin Pathol* 55(9): 693-698
- Visnjic, D., I. Kalajzic, G. Gronowicz, H. L. Aguila, S. H. Clark, A. C. Lichtler & D. W. Rowe (2001). "Conditional ablation of the osteoblast lineage in Col2.3delta^{tk} transgenic mice." *J Bone Miner Res* 16(12): 2222-2231
- Visnjic, D., Z. Kalajzic, D. W. Rowe, V. Katavic, J. Lorenzo & H. L. Aguila (2004). "Hematopoiesis is severely altered in mice with an induced osteoblast deficiency." *Blood* 103(9): 3258-3264
- Viviano, B. L., L. Silverstein, C. Pflederer, S. Paine-Saunders, K. Mills & S. Saunders (2005). "Altered hematopoiesis in glypican-3-deficient mice results in decreased osteoclast differentiation and a delay in endochondral ossification." *Dev Biol* 282(1): 152-162
- Walkley, C. R., M. L. Fero, W. M. Chien, L. E. Purton & G. A. McArthur (2005). "Negative cell-cycle regulators cooperatively control self-renewal and differentiation of haematopoietic stem cells." *Nat Cell Biol* 7(2): 172-178
- Walkley, C. R., G. H. Olsen, S. Dworkin, S. A. Fabb, J. Swann, G. A. McArthur, S. V. Westmoreland, P. Chambon, D. T. Scadden & L. E. Purton (2007). "A microenvironment-induced myeloproliferative syndrome caused by retinoic acid receptor gamma deficiency." *Cell* 129(6): 1097-1110
- Walkley, C. R., J. M. Shea, N. A. Sims, L. E. Purton & S. H. Orkin (2007). "Rb regulates interactions between hematopoietic stem cells and their bone marrow microenvironment." *Cell* 129(6): 1081-1095
- Wei, L., K. Kanbe, M. Lee, X. Wei, M. Pei, X. Sun, R. Terek & Q. Chen (2010). "Stimulation of chondrocyte hypertrophy by chemokine stromal cell-derived factor 1 in the chondro-osseous junction during endochondral bone formation." *Dev Biol* 341(1): 236-245
- Weinstein, R., M. A. Riordan, K. Wenc, S. Kreczko, M. Zhou & N. Dainiak (1989). "Dual role of fibronectin in hematopoietic differentiation." *Blood* 73(1): 111-116
- Weiss, L. (1976). "The hematopoietic microenvironment of the bone marrow: an ultrastructural study of the stroma in rats." *Anat Rec* 186(2): 161-184
- Wittig, O., J. Paez-Cortez & J. Cardier (2009). "Liver Sinusoidal Endothelial Cells Promote B Lymphopoiesis from Primitive Hematopoietic Cells." *Stem Cells Dev*
- Wolf, N. S. (1979). "The haemopoietic microenvironment." *Clin Haematol* 8(2): 469-500
- Wolf, N. S. & J. J. Trentin (1968). "Hemopoietic colony studies. V. Effect of hemopoietic organ stroma on differentiation of pluripotent stem cells." *J Exp Med* 127(1): 205-214
- Wu, J. Y., L. E. Purton, S. J. Rodda, M. Chen, L. S. Weinstein, A. P. McMahon, D. T. Scadden & H. M. Kronenberg (2008). "Osteoblastic regulation of B lymphopoiesis is mediated by Gs{alpha}-dependent signaling pathways." *Proc Natl Acad Sci U S A* 105(44): 16976-16981
- Xie, Y., T. Yin, W. Wiegraabe, X. C. He, D. Miller, D. Stark, K. Perko, R. Alexander, J. Schwartz, J. C. Grindley, J. Park, J. S. Haug, J. P. Wunderlich, H. Li, S. Zhang, T.

- Johnson, R. A. Feldman & L. Li (2009). "Detection of functional haematopoietic stem cell niche using real-time imaging." *Nature* 457(7225): 97-101
- Yagi, K., K. Tsuji, A. Nifuji, K. Shinomiya, K. Nakashima, B. DeCrombrugghe & M. Noda (2003). "Bone morphogenetic protein-2 enhances osterix gene expression in chondrocytes." *J Cell Biochem* 88(6): 1077-1083
- Yin, T. & L. Li (2006). "The stem cell niches in bone." *J Clin Invest* 116(5): 1195-1201
- Yokota, T., K. Oritani, I. Takahashi, J. Ishikawa, A. Matsuyama, N. Ouchi, S. Kihara, T. Funahashi, A. J. Tenner, Y. Tomiyama & Y. Matsuzawa (2000). "Adiponectin, a new member of the family of soluble defense collagens, negatively regulates the growth of myelomonocytic progenitors and the functions of macrophages." *Blood* 96(5): 1723-1732
- Yoshimoto, M., T. Shinohara, T. Heike, M. Shiota, M. Kanatsu-Shinohara & T. Nakahata (2003). "Direct visualization of transplanted hematopoietic cell reconstitution in intact mouse organs indicates the presence of a niche." *Exp Hematol* 31(8): 733-740
- Youn, D. Y., D. H. Lee, M. H. Lim, J. S. Yoon, J. H. Lim, S. E. Jung, C. E. Yeum, C. W. Park, H. J. Youn, J. S. Lee, S. B. Lee, M. Ikawa, M. Okabe, Y. Tsujimoto & J. H. Lee (2008). "Bis deficiency results in early lethality with metabolic deterioration and involution of spleen and thymus." *Am J Physiol Endocrinol Metab* 295(6): E1349-1357
- Zhang, C. C. & H. F. Lodish (2008). "Cytokines regulating hematopoietic stem cell function." *Curr Opin Hematol* 15(4): 307-311
- Zhang, J., C. Niu, L. Ye, H. Huang, X. He, W. G. Tong, J. Ross, J. Haug, T. Johnson, J. Q. Feng, S. Harris, L. M. Wiedemann, Y. Mishina & L. Li (2003). "Identification of the haematopoietic stem cell niche and control of the niche size." *Nature* 425(6960): 836-841
- Zhang, Y., A. Harada, H. Bluethmann, J. B. Wang, S. Nakao, N. Mukaida & K. Matsushima (1995). "Tumor necrosis factor (TNF) is a physiologic regulator of hematopoietic progenitor cells: increase of early hematopoietic progenitor cells in TNF receptor p55-deficient mice in vivo and potent inhibition of progenitor cell proliferation by TNF alpha in vitro." *Blood* 86(8): 2930-2937
- Zhu, J., R. Garrett, Y. Jung, Y. Zhang, N. Kim, J. Wang, G. J. Joe, E. Hexner, Y. Choi, R. S. Taichman & S. G. Emerson (2007). "Osteoblasts support B-lymphocyte commitment and differentiation from hematopoietic stem cells." *Blood* 109(9): 3706-3712
- Zuckerman, K. S., R. K. Rhodes, D. D. Goodrum, V. R. Patel, B. Sparks, J. Wells, M. S. Wicha & L. A. Mayo (1985). "Inhibition of collagen deposition in the extracellular matrix prevents the establishment of a stroma supportive of hematopoiesis in long-term murine bone marrow cultures." *J Clin Invest* 75(3): 970-975
- Zuckerman, K. S. & M. S. Wicha (1983). "Extracellular matrix production by the adherent cells of long-term murine bone marrow cultures." *Blood* 61(3): 540-547
- Zweegman, S., J. Van Den Born, A. M. Mus, F. L. Kessler, J. J. Janssen, T. Netelenbos, P. C. Huijgens & A. M. Drager (2004). "Bone marrow stromal proteoglycans regulate megakaryocytic differentiation of human progenitor cells." *Exp Cell Res* 299(2): 383-392

Advances in Hematopoietic Stem Cell Research

Edited by Dr. Rosana Pelayo

ISBN 978-953-307-930-1

Hard cover, 464 pages

Publisher InTech

Published online 27, January, 2012

Published in print edition January, 2012

This book provides a comprehensive overview in our understanding of the biology and therapeutic potential of hematopoietic stem cells, and is aimed at those engaged in stem cell research: undergraduate and postgraduate science students, investigators and clinicians. Starting from fundamental principles in hematopoiesis, *Advances in Hematopoietic Stem Cell Research* assemble a wealth of information relevant to central mechanisms that may regulate differentiation, and expansion of hematopoietic stem cells in normal conditions and during disease.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Elizabeth Sweeney and Olena Jacenko (2012). Skeletogenesis and the Hematopoietic Niche, *Advances in Hematopoietic Stem Cell Research*, Dr. Rosana Pelayo (Ed.), ISBN: 978-953-307-930-1, InTech, Available from: <http://www.intechopen.com/books/advances-in-hematopoietic-stem-cell-research/skeletogenesis-and-the-hematopoietic-niche>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](https://creativecommons.org/licenses/by/3.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen