We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Adipose Tissue Metabolism and Effect of Postmenopausal Hormone Therapy on Change of Body Composition

Kyong Wook Yi¹ and Seung Yup Ku²

¹*Department of Obstetrics and Gynecology,
College of Medicine, Korea University,*

²*Department of Obstetrics and Gynecology,
College of Medicine, Seoul National University, Seoul,
Korea*

1. Introduction

Obesity is a process by which excess energy accumulates and results in increasing fat deposition in various parts of the body. Obesity is associated with economic, social, and lifestyle factors, and is commonly induced by imbalanced energy intake and consumption of high calorie foods or low physical activity. Obesity is a worldwide issue in public health that significantly increases the risk for type 2 diabetes, metabolic syndrome, atherosclerosis, and cardiovascular disease.

Body weight increases with aging in both genders, irrespective of the baseline weight in normal and obese individuals [1]. This increase in body weight is attributed to a reduction in energy expenditure with decreased physical activity. The global prevalence of obesity has been reported to be higher in females than males [2]. A US population survey estimated that approximately two thirds of women 40 - 60 years of age are overweight or obese [3].

In addition to aging, the menopause is considered an important factor for contributing to altered adiposity in women. Menopause is defined as a decline in endogenous estrogen production from the ovaries, and clinically represents cessation of menstruation and loss of fertility. Estrogen loss is the most significant event impacting a variety of physiologic and psychological changes in women. In the peri- or postmenopausal period, a change in adiposity has been well described. Weight gain during the menopausal transition has been scrutinized as a critical factor to midlife body weight in women [1]. Several observational studies have shown increased weight gain during the menopausal transition [4, 5]. A number of clinical trials have demonstrated a significant association between menopausal status with changes in anthropometry, blood pressure, lipid profile, and glucose/insulin metabolism [6-11], which can be linked to increased cardiovascular morbidity and mortality during the postmenopausal period [12].

Given the collective evidence on the change in adiposity across the menopausal transition, the roles of sex hormones, especially estrogen, are of increased interest in understanding the regulation of adiposity. This review discusses the association between estrogen and adiposity, and the interaction of estrogen with other biological metabolites and substances

involved in obesity. In addition we summarize evidence for the effect of postmenopausal hormone therapy (HT) on changes in body composition.

2. Adipose tissue

Adipose tissue has a vital role in the lives of mammals. The primary function of adipose tissue is to store excess energy within the body in the form of free fatty acids (FFAs) and heat production. However, adipocytes are currently regarded as an independent endocrine organ since the metabolic and endocrine actions have been revealed.

In mammals, two types of adipose tissue exist (white adipose tissue [WAT] and brown adipose tissue [BAT]) [13]. WAT represents the major component of adipose tissue and provides most of the total body fat [13]. Moreover, WAT is the main source of FFAs that are available as energy substrates for generation through oxidative phosphorylation of adenosine triphosphate (ATP) high-energy bonds [13, 14]. Excess WAT in the upper parts of the body (android type obesity) represents a strong risk factor for some inflammatory pathologies [15]. In contrast, accumulation of WAT in lower body parts (gynecoid type obesity) is not associated with metabolic complications [13, 15].

In contrast to WAT, BAT participates in energy expenditure from non-oxidative phosphorylation in the form of heat for cold adaption [16]. The uncoupling of phosphorylation in BAT is attributed to the activity of uncoupling protein-1, which is expressed on the mitochondrial membrane, by creating a proton leak that depletes the electrochemical gradient needed for oxidative phosphorylation [14, 16]. BAT represents a smaller number of fat cells that have a rich vascular supply, which can respond more rapidly to sympathetic nervous system stimulation, and elicits heat production, rather than ATP production from nonshivering cold adaptive thermogenesis [14, 16]. In humans, BAT helps to maintain body temperature in newborns, but BAT regresses with increasing age and is completely lost in adulthood [17, 18]. Recently, Virtanen et al. studied BAT deposition in healthy adults using the glucose analogue, ^{18}F -fluorodeoxyglucose, uptake by PET and computed tomography [18]. Metabolically-active BAT depots in paracervical and supraclavicular adipose tissue, which can be induced in response to cold and sympathetic nervous system activation, has been reported [14, 16, 18]. The presence of BAT in human adults is of potential interest in understanding the mechanism of obesity, and may provide a rationale for pharmacologic and gene expression manipulation to combat human obesity [14, 18, 19].

3. Deposition and distribution of adipose tissue

Deposition of the fat mass has a different pattern between the genders. Young males have little subcutaneous fat and do not show a central-peripheral difference [20], whereas women of reproductive age have more subcutaneous fat than men at all measured subcutaneous regions [21]. In addition, with an increasing severity of obesity, adipose tissue thickness is higher in the central regions in men, but women exhibit a peripheral deposition [20]. This pattern of fat deposition is altered in women across the menopausal transition; specifically, the total amount of body fat increases, and the peripheral fat shifts around the abdomen. Furthermore, the change in visceral adipose tissue has been ascribed to both chronologic aging and menopause [22, 23]. The results from a clinical study with 156 healthy women

during 4 years of follow-up showed an increase in visceral adipose tissue and total body fat, and a 32% reduction in fat oxidation during the menopausal transition [4]. In the study, the subcutaneous adipose tissue increased in accordance with age independent of menopausal status, while the findings of increased visceral adipose tissue and total body fat were noted only in postmenopausal women. This distinctive physiologic change in amount and distribution of fat in women is noteworthy because the central fat deposition has a more deleterious effect on the development of cardiovascular and metabolic disease [24, 25].

Although the exact mechanism regarding fat redistribution after menopause remains unclear, the phenomenon with declining estrogen level may be due to alterations in adipose tissue metabolism [4]. Several studies have shown that estrogen directly promotes subcutaneous fat accumulation [26], and the loss of estrogen by menopause is associated with an increase in central fat. Several longitudinal studies lend support in suggesting that estrogen plays an important role in regulating body fat distribution. Postmenopausal women who receive estrogen replacement have significantly lower waist-to-hip ratios and less visceral adipose tissue than women who have never received estrogen replacement therapy [27, 28].

In experimental studies, 17 β -estradiol (E2) has been shown to regulate adipose tissue by increasing the number of adipocytes through effects on proliferation and differentiation [29, 30]. The number and size of adipocytes is a determining factor for adiposity, and the adipocyte size is balanced by lipogenesis and the lipolysis pathway. Palin et al. have found a direct regulatory effect of E2 on the expression of lipoprotein lipase (LPL) and hormone-sensitive lipase (HSL) in human subcutaneous abdominal adipose tissue [31]. LPL is a major modulator of lipid deposition as triglycerides into adipocytes, and HSL is the rate-limiting enzyme involved in the process of lipolysis. Therefore, the direct effect of estrogen on these enzymes might lead to fat redistribution in postmenopausal women.

As other mechanisms suggest, the role of the estrogen receptor (ER) is focused on estrogen-related action on the regulation of adiposity. Adipocytes express two main subtypes of ERs (alpha [ER α] and beta [ER β]). ER α was discovered first, and the biological effects of ER α on adiposity have been thoroughly described. ER α is considered to be essential for genomic actions of E2 on the regulation of body fat [32]. Because the hypothalamic nuclei that regulate energy homeostasis express ER α , E2 action could affect adiposity [33-35]. An animal study conducted by Heine et al. demonstrated that glucose intolerance, hypertrophy, and hyperplasia of adipocytes are induced in ER α knockout mice [36], thus supporting a critical role of ER α in determining adiposity.

In contrast, the biological implications of the more recently discovered ER β have been less revealing than ER α . The binding affinity of estrogen to ER α and ER β is known to be similar, but the two subtypes of ER only have 56% identity in the ligand binding domain [26, 37, 38]. Therefore, different or competitive roles between both ERs have been repeatedly suggested. Naaz et al. studied the role of ER β in adipose tissue [39] in mice with ER α KO. When compared to the results generated in ER α KO mice, it was shown that removing E2/ER β signaling induced a decrease in body weight, the amount of fat, and adipocyte size. Therefore, the authors suggested a potential role for ER β in regulating adiposity, as well as ER α , but with opposing actions. Thus, the roles for ERs in adipocytes might be an interesting target to further elucidate the estrogen effects on regulating adiposity.

4. Proinflammatory cytokines production

Obesity is a low grade systemic inflammatory state; several studies have addressed adipocyte-related mechanisms involved in regulating proinflammatory [15, 40] and adipocyte-specific cytokines. Systemic chronic inflammation is due to an inflammatory response in adipose tissues that are under quick expansion, and in which macrophages and adipocytes are activated and stimulate the production of proinflammatory cytokines and adipokines [41]. Visceral WAT is considered a main source of inflammation related to obesity. Obese subjects with higher visceral fat exhibit monocyte-chemotactic protein (MCP)-1 expression and infiltration of macrophages in visceral fat compared to subcutaneous fat [42]. Also, several *in vivo* studies have shown higher levels of plasminogen activator inhibitor (PAI)-1, IL-6, and TNF- α in visceral obesity [43, 44].

Recent studies have shown the inhibitory role of estrogen to the inflammatory response in adipose tissue, and cardiovascular and nervous systems [35]. ER α is located in cytokine-producing cells, such as macrophages and microglia, and several *in vitro* studies have reported that E2/ ER α decreases the number of pro-inflammatory cytokines [35, 45, 46]. Another *in vivo* study has demonstrated that both ER α and ER β regulate proinflammatory cytokine and chemokine production through E2-dependent and independent mechanisms [47].

Because energy expenditure is related to inflammation, a role for nuclear receptor kappa B (NF- κ B) has been suggested. NF- κ B is a protein complex that controls transcription factor, and is known to have a crucial role in regulating immune responses to infection or stress. NF- κ B induces the transcription of inflammatory cytokines, such as TNF- α , IL-1, IL-6, and MCP-1 [41]. In the classical pathway, NF- κ B is mediated through IKK β -induced phosphorylation and proteasome-mediated degradation of I κ B α [48]. The other pathway is the activation by hypoxia in the absence of I κ B α degradation, a pathway in adipocytes and macrophages which contributes to chronic inflammation in the adipose tissues of obese subjects [41, 49]. Limited evidence has proposed an inhibitory effect of E2 via ER α on NF- κ B [50, 51], which partially explains the anti-inflammatory properties of estrogen.

5. Adipokine expression/secretion

Adipokines are a family of cytokines which include leptin, resistin, adiponectin, and TNF- α and are primarily secreted from adipocytes. These adipokines are released from different tissues and organs, and are not exclusively produced by WAT [14, 52].

Leptin, the product of the *ob* gene, has been shown to be a key metabolic hormone in regulating appetite body weight and energy homeostasis [53, 54]. In humans, circulating leptin levels have a parallel correlation with the amount of body fat. In addition, serum levels of leptin have been shown to be higher in women than men [55]. This finding has been hypothesized to be due to a different pattern in fat deposition between the genders and/or the effects of a different hormonal milieu [56]. With respect to the change in leptin levels according to menopausal status, data are inconsistent; some data have shown no change in leptin levels between the pre- and post-menopause [57], while other studies have demonstrated a decrease in leptin levels during the menopausal transition [58, 59]. Some investigators have suggested that the low amount of visceral fat compared with subcutaneous fat in the genders [60, 61] makes it unlikely that the leptin secretion rates from these two depots differ, and may therefore be the cause for the sexual dimorphism in leptin concentrations, suggesting an important role for sex steroid hormones [53]. According to

recent data derived from healthy pre- and post-menopausal women, postmenopausal women had increased levels of tissue plasminogen activator antigen (tPA), MCP-1, and adiponectin [24]. Furthermore, an increase in intraabdominal fat was correlated with C-reactive protein, tPA, and leptin, and negatively with adiponectin levels. The results imply that during the menopausal transition, women have adverse changes in inflammatory markers and adipokines which correlate with increased visceral obesity.

Several animal studies have indicated a stimulatory effect of estrogen on leptin expression and secretion in rat adipose tissues [59, 62]. Machinal et al. reported that there are regional differences in leptin expression between subcutaneous and deep fat tissues in rats, and that leptin secretion increased in the deep fat tissue tissues [63]. Based on one study involving human adipocytes, estrogen is likely to stimulate leptin expression [53]. In a previous study, the association between ER and adipokine expression in 3T3-L1 adipocytes was investigated [64]. The results showed that ER α has a stimulatory effect on leptin expression, while the expression of ER β is inversely correlated with leptin expression. Therefore, discordant findings regarding the estrogen effect on leptin expression/secretion from other studies can be explained by the different expression of the two ER subtypes, which have opposite actions on the expression of leptin. In addition, if there are regulating factors (genetic or environmental) for ER α or ER β expression in adipocytes, this may explain how the different expression of ER in adipose tissue can affect the diverse obesity phenotypes.

Adiponectin is a 30 kDa protein secreted abundantly from adipocytes [65, 66], and functions to exert anti-diabetic and anti-atherogenic properties. The serum adiponectin levels are decreased in obese individuals, metabolic syndrome, and type 2 DM [67, 68]. Similar to leptin, circulating adiponectin levels show a sexual dimorphism with higher levels in women than men [69]. Although the estrogen effects for adiponectin expression are limited, some *in vitro* studies have reported no direct regulatory effect in humans and mouse adipocytes [64, 70].

Resistin is a cysteine-rich protein that was originally described as an adipose-derived protein in rodents that links obesity to insulin resistance [71]. However, in human data, the relationship of resistin with obesity and insulin metabolism is still debated. A widely reported biological role of resistin is the regulatory effect involved in inflammatory processes. In addition to adipocytes, resistin is induced by lipopolysaccharides and TNF- α in macrophages [72]. The other data showed that resistin induces or is induced by IL-6 and TNF- α via NF- κ B in human monocytes [73, 74]. Data on estrogen effects for resistin expression are limited. In a mouse adipocyte model, the regulatory effect of 17 β -E2 on resistin expression is discordant [64, 75].

Recently described novel adipokines, such as visfatin, retinol binding protein-4, and omentin, have been shown to exert some metabolic properties, but their biological actions linked to obesity and interactions with estrogen need to be elucidated.

6. Postmenopausal hormone therapy and change in body composition

Hormone therapy (HT) is widely used for the treatment of menopausal symptoms and preventing bone loss in postmenopausal women. Estrogen replacement, when combined with various progestogens for endometrial protection, has been established as a conventional formulation. The benefit of HT is known to reduce vasomotor symptoms,

prevent osteoporotic fractures, and improve well-being and quality of life during the menopausal period. In contrast, the risks for breast cancer and thromboembolic disease may increase, and cardiovascular effects remain controversial.

As previously mentioned, aging and menopause in women are related to an increase in body fat and redistribution of fat mass to the central portion of the body. These changes are linked to an increase in metabolic and cardiovascular disease after the menopause. Therefore, it has been continuously questioned and theorized that HT may have a favorable effect on change in body composition and anthropometries in postmenopausal women. With respect to this issue, data are still debated. In a meta-analysis of 24 RCTs, no effect of ET or HT on body weight was described [76]. However, subsequent studies suggested some beneficial effects of HT on body composition. A sub-study of the estrogen plus progestin trial of the Women's Health Initiative (WHI) investigated whether or not postmenopausal HT affects age-related changes in body composition [77]. The WHI study was originally designed to evaluate the risks and benefits of HT (EPT/ET) with an enrollment of 16,608 postmenopausal women between 1993 and 1998 [78]. The sub-analysis included 835 women who had whole-body dual-energy X-ray absorptiometry scans for measurement of body composition at baseline and year 3. Based on the results of the study, women who received EPT lost less lean soft tissue mass (-0.04 kg) than women who received placebo (-0.44 kg) 3 year after intervention. Furthermore, less upper-body fat distribution was noted in the EPT group than the placebo group (ratio of trunk to leg fat mass, -0.025 for the EPT group and 0.004 for the placebo group, $P = 0.003$). The investigators concluded that EPT has a favorable effect by reducing central fat deposition, but the real health benefits of this effect remains to be confirmed due to the small size of the effect. In a randomized, single-blind study, the effects of HT on body fat composition were studied in 59 postmenopausal women (mean age 49.9 ± 3.8 years) [79]. The participants were assigned into the following three groups according to the type of HT: transdermal estradiol (E2)/norethisterone acetate (NETA); transdermal continuous E2/ oral medroxyprogesterone acetate (MPA); and oral continuous E2/NETA. The results showed that all types of HT caused a significant decrease in WC, subcutaneous fat, and WHR. Thus, HT reduced fat deposition in the central part of the body, and such an effect was more marked in women with a WC ≥ 88 cm and subcutaneous fat ≥ 33 cm. Another placebo-controlled study investigated the effects of oral continuous E2/NETA on anthropometric changes and serum leptin levels in postmenopausal women [80]. In agreement with a previous study, WC and HC decreased significantly in the E2/NETA group, while the body weight (BW) increased in the placebo group. With this reduction in central fat deposition, the serum leptin levels were positively related to the changes in subcutaneous fat tissue. The authors advocated that HT may have a protective effect on CVD via a slimming effect on the central region in postmenopausal women.

Although the effect of postmenopausal HT on the change of body composition is not clearly understood, the limited data have suggest that HT has a small, but significant effect on preventing an android fat shift (central fat depot) regardless of the body weight.

Tibolone is a synthetic steroid hormone that exerts estrogenic, androgenic, and progestogenic properties. The biological actions of tibolone are mediated through the metabolites of tibolone (3α -OH-tibolone, 3β -OH-tibolone, and Δ^4 -isomer) by binding estrogen or progesterone receptors in multiple target tissues and organs (brain, bone, breast, and endometrium). Based on several randomized trials, tibolone has shown comparable

effects with traditional estrogen therapy (ET)/HT on relieving vasomotor symptoms/genital atrophy, improving the quality of life, and preventing bone loss in postmenopausal women [81-84]. Currently 4 studies have shown the effects of tibolone on BW or body composition in postmenopausal women. In a 2-year follow up study comparing three regimens of HRT, tibolone had a stable effect on body fat and lean mass [85]. However, the fat mass increased (+3.6%) and the lean body mass decreased (-1.7%) in the control group. Another study demonstrated that tibolone significantly increased fat-free mass by 0.85 kg and total body water by 0.78 liter during a 1-year observation period [86]. The authors reported an effect of tibolone on preventing a decline in lean body mass, but mentioned the need for a long-term observational study.

Arabi et al. compared the effects of tibolone with EPT (E2/NETA) on changes in body composition and bone densitometry in postmenopausal women [87]. Both EPT and tibolone increased lean body mass, whereas the android fat and android obesity index decreased.

Tibolone might provide further benefits in increasing lean body mass and decreasing the fat component. Steroid hormones exert their action by binding to an intra-nuclear receptor [88]. Therefore, like estrogen, tibolone might have a direct effect on skeletal muscle by binding to ER α or ER β expressed within human skeletal muscle [88-90]. In addition, tibolone increases the serum IGF-1 levels [91], which promotes muscle protein synthesis, and increases the number of myogenic satellite cells and the proliferation of myogenic satellite cells. The anabolic effect of tibolone on muscle has been suggested to be mediated in part by local IGF-1, independent of the serum IGF-1 level [92].

7. Summary

Adipose tissue is the largest endocrine organ in the human body. The amount and distribution of adipose tissue reflects energy balance. Adipose tissue also releases a variety of biologically active molecules or cytokines. Adipocyte metabolism and physiology have been extensively studied over recent years, but the exact mechanism and effect of sex steroid hormones involved in regulating adiposity remain to be defined.

Current data give evidence that estrogen appears to have direct effects on cell proliferation or differentiation for adipocytes, and in regulating key enzymes involved in fat deposition. Another possible mechanism involves the hormonal or paracrine effects by secretion of various adipokines and cytokines, in which interactions with estrogen are promising and need additional studies.

At present, it is not known whether or not HT provides a significant effect on modulating fat mass or preventing fat redistribution. Furthermore, in recent years the HT formulations have been changed to include lower doses of estrogen (< 50 μ g) in combination with new progestins. The effects of the currently used HT regimens on changes in body fat composition are limited and require more data. Tibolone includes estrogenic, androgenic, and progestogenic properties which may affect the body composition (adipose tissue or lean mass) differently than conventional HT.

The respective roles of sex steroid hormones and their receptors (ER subtypes and PR) on body fat distribution could be an interesting target for understanding the estrogen effect on adiposity, and may provide selective therapeutic approaches, such as hormonal manipulation for adiposity related to estrogen change throughout the menopausal period.

8. References

- [1] H. N. Polotsky, and A. J. Polotsky, "Metabolic implications of menopause," *Semin Reprod Med*, vol. 28, no. 5, pp. 426-434.
- [2] "Obesity: preventing and managing the global epidemic. Report of a WHO consultation," *World Health Organization Technical Report Series*, vol. 894, pp. i-xii, 1-253, 2000.
- [3] C. L. Ogden, M. D. Carroll, L. R. Curtin et al., "Prevalence of overweight and obesity in the United States, 1999-2004," *JAMA*, vol. 295, no. 13, pp. 1549-1555, 2006.
- [4] J. C. Lovejoy, C. M. Champagne, L. de Jonge et al., "Increased visceral fat and decreased energy expenditure during the menopausal transition," *Int J Obes (Lond)*, vol. 32, no. 6, pp. 949-958, 2008.
- [5] R. R. Wing, K. A. Matthews, L. H. Kuller et al., "Weight gain at the time of menopause," *Archives of Internal Medicine*, vol. 151, no. 1, pp. 97-102, 1991.
- [6] J. F. Owens, C. M. Stoney, and K. A. Matthews, "Menopausal status influences ambulatory blood pressure levels and blood pressure changes during mental stress," *Circulation*, vol. 88, no. 6, pp. 2794-2802, 1993.
- [7] J. H. Shin, J. Y. Hur, H. S. Seo et al., "The ratio of estrogen receptor alpha to estrogen receptor beta in adipose tissue is associated with leptin production and obesity," *Steroids*, vol. 72, no. 6-7, pp. 592-599, 2007.
- [8] M. P. Brincat, Y. M. Baron, and R. Galea, "Estrogens and the skin," *Climacteric*, vol. 8, no. 2, pp. 110-123, 2005.
- [9] F. A. Tremollieres, J. M. Pouilles, C. Cauneille, and C. Ribot, "Coronary heart disease risk factors and menopause: a study in 1684 French women," *Atherosclerosis*, vol. 142, no. 2, pp. 415-423, 1999.
- [10] K. Fukami, K. Koike, K. Hirota et al., "Perimenopausal changes in serum lipids and lipoproteins: a 7-year longitudinal study," *Maturitas*, vol. 22, no. 3, pp. 193-197, 1995.
- [11] M. E. Piche, S. J. Weisnagel, L. Corneau et al., "Contribution of abdominal visceral obesity and insulin resistance to the cardiovascular risk profile of postmenopausal women," *Diabetes*, vol. 54, no. 3, pp. 770-777, 2005.
- [12] K. A. Matthews, E. Meilahn, L. H. Kuller et al., "Menopause and risk factors for coronary heart disease," *New England Journal of Medicine*, vol. 321, no. 10, pp. 641-646, 1989.
- [13] S. Gesta, Y. H. Tseng, and C. R. Kahn, "Developmental origin of fat: tracking obesity to its source," *Cell*, vol. 131, no. 2, pp. 242-256, 2007.
- [14] C. R. Balistreri, C. Caruso, and G. Candore, "The role of adipose tissue and adipokines in obesity-related inflammatory diseases," *Mediators of Inflammation*, vol. 2010, p. 802078.
- [15] R. Cencello, and K. Clement, "Is obesity an inflammatory illness? Role of low-grade inflammation and macrophage infiltration in human white adipose tissue," *BJOG*, vol. 113, no. 10, pp. 1141-1147, 2006.
- [16] R. N. Redinger, "Fat storage and the biology of energy expenditure," *Transl Res*, vol. 154, no. 2, pp. 52-60, 2009.
- [17] M. E. Lean, "Brown adipose tissue in humans," *Proceedings of the Nutrition Society*, vol. 48, no. 2, pp. 243-256, 1989.
- [18] K. A. Virtanen, M. E. Lidell, J. Orava et al., "Functional brown adipose tissue in healthy adults," *New England Journal of Medicine*, vol. 360, no. 15, pp. 1518-1525, 2009.

- [19] G. Fruhbeck, S. Becerril, N. Sainz et al., "BAT: a new target for human obesity?," *Trends in Pharmacological Sciences*, vol. 30, no. 8, pp. 387-396, 2009.
- [20] P. Bjorntorp, "Adipose tissue distribution and function," *International Journal of Obesity*, vol. 15 Suppl 2, pp. 67-81, 1991.
- [21] L. Sjostrom, U. Smith, M. Krotkiewski, and P. Bjorntorp, "Cellularity in different regions of adipose tissue in young men and women," *Metabolism: Clinical and Experimental*, vol. 21, no. 12, pp. 1143-1153, 1972.
- [22] C. J. Ley, B. Lees, and J. C. Stevenson, "Sex- and menopause-associated changes in body-fat distribution," *American Journal of Clinical Nutrition*, vol. 55, no. 5, pp. 950-954, 1992.
- [23] H. Shimokata, R. Andres, P. J. Coon et al., "Studies in the distribution of body fat. II. Longitudinal effects of change in weight," *International Journal of Obesity*, vol. 13, no. 4, pp. 455-464, 1989.
- [24] C. G. Lee, M. C. Carr, S. J. Murdoch et al., "Adipokines, inflammation, and visceral adiposity across the menopausal transition: a prospective study," *Journal of Clinical Endocrinology and Metabolism*, vol. 94, no. 4, pp. 1104-1110, 2009.
- [25] W. B. Kannel, L. A. Cupples, R. Ramaswami et al., "Regional obesity and risk of cardiovascular disease; the Framingham Study," *Journal of Clinical Epidemiology*, vol. 44, no. 2, pp. 183-190, 1991.
- [26] M. Krotkiewski, P. Bjorntorp, L. Sjostrom, and U. Smith, "Impact of obesity on metabolism in men and women. Importance of regional adipose tissue distribution," *Journal of Clinical Investigation*, vol. 72, no. 3, pp. 1150-1162, 1983.
- [27] J. Munoz, A. Derstine, and B. A. Gower, "Fat distribution and insulin sensitivity in postmenopausal women: influence of hormone replacement," *Obesity Research*, vol. 10, no. 6, pp. 424-431, 2002.
- [28] M. A. Espeland, M. L. Stefanick, D. Kritz-Silverstein et al., "Effect of postmenopausal hormone therapy on body weight and waist and hip girths. Postmenopausal Estrogen-Progestin Interventions Study Investigators," *Journal of Clinical Endocrinology and Metabolism*, vol. 82, no. 5, pp. 1549-1556, 1997.
- [29] M. N. Dieudonne, R. Pecquery, M. C. Leneuve, and Y. Giudicelli, "Opposite effects of androgens and estrogens on adipogenesis in rat preadipocytes: evidence for sex and site-related specificities and possible involvement of insulin-like growth factor 1 receptor and peroxisome proliferator-activated receptor gamma2," *Endocrinology*, vol. 141, no. 2, pp. 649-656, 2000.
- [30] D. A. Roncari, and R. L. Van, "Promotion of human adipocyte precursor replication by 17beta-estradiol in culture," *Journal of Clinical Investigation*, vol. 62, no. 3, pp. 503-508, 1978.
- [31] S. L. Palin, P. G. McTernan, L. A. Anderson et al., "17Beta-estradiol and anti-estrogen ICI:compound 182,780 regulate expression of lipoprotein lipase and hormone-sensitive lipase in isolated subcutaneous abdominal adipocytes," *Metabolism: Clinical and Experimental*, vol. 52, no. 4, pp. 383-388, 2003.
- [32] S. Musatov, W. Chen, D. W. Pfaff et al., "Silencing of estrogen receptor alpha in the ventromedial nucleus of hypothalamus leads to metabolic syndrome," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 104, no. 7, pp. 2501-2506, 2007.

- [33] I. Merchenthaler, M. V. Lane, S. Numan, and T. L. Dellovade, "Distribution of estrogen receptor alpha and beta in the mouse central nervous system: in vivo autoradiographic and immunocytochemical analyses," *Journal of Comparative Neurology*, vol. 473, no. 2, pp. 270-291, 2004.
- [34] D. L. Voisin, S. X. Simonian, and A. E. Herbison, "Identification of estrogen receptor-containing neurons projecting to the rat supraoptic nucleus," *Neuroscience*, vol. 78, no. 1, pp. 215-228, 1997.
- [35] L. M. Brown, and D. J. Clegg, "Central effects of estradiol in the regulation of food intake, body weight, and adiposity," *Journal of Steroid Biochemistry and Molecular Biology*, vol. 122, no. 1-3, pp. 65-73.
- [36] P. A. Heine, J. A. Taylor, G. A. Iwamoto et al., "Increased adipose tissue in male and female estrogen receptor-alpha knockout mice," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 97, no. 23, pp. 12729-12734, 2000.
- [37] S. Mosselman, J. Polman, and R. Dijkema, "ER beta: identification and characterization of a novel human estrogen receptor," *FEBS Letters*, vol. 392, no. 1, pp. 49-53, 1996.
- [38] G. G. Kuiper, E. Enmark, M. Peltö-Huikko et al., "Cloning of a novel receptor expressed in rat prostate and ovary," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 93, no. 12, pp. 5925-5930, 1996.
- [39] A. Naaz, M. Zakroczyński, P. Heine et al., "Effect of ovariectomy on adipose tissue of mice in the absence of estrogen receptor alpha (ERalpha): a potential role for estrogen receptor beta (ERbeta)," *Hormone and Metabolic Research*, vol. 34, no. 11-12, pp. 758-763, 2002.
- [40] C. E. Juge-Aubry, E. Henrichot, and C. A. Meier, "Adipose tissue: a regulator of inflammation," *Best Pract Res Clin Endocrinol Metab*, vol. 19, no. 4, pp. 547-566, 2005.
- [41] J. Ye, and J. N. Keller, "Regulation of energy metabolism by inflammation: a feedback response in obesity and calorie restriction," *Aging (Albany NY)*, vol. 2, no. 6, pp. 361-368.
- [42] I. Harman-Boehm, M. Bluher, H. Redel et al., "Macrophage infiltration into omental versus subcutaneous fat across different populations: effect of regional adiposity and the comorbidities of obesity," *Journal of Clinical Endocrinology and Metabolism*, vol. 92, no. 6, pp. 2240-2247, 2007.
- [43] G. Winkler, S. Kiss, L. Keszthelyi et al., "Expression of tumor necrosis factor (TNF)-alpha protein in the subcutaneous and visceral adipose tissue in correlation with adipocyte cell volume, serum TNF-alpha, soluble serum TNF-receptor-2 concentrations and C-peptide level," *European Journal of Endocrinology / European Federation of Endocrine Societies*, vol. 149, no. 2, pp. 129-135, 2003.
- [44] G. He, S. B. Pedersen, J. M. Bruun et al., "Differences in plasminogen activator inhibitor 1 in subcutaneous versus omental adipose tissue in non-obese and obese subjects," *Hormone and Metabolic Research*, vol. 35, no. 3, pp. 178-182, 2003.
- [45] E. Vegeto, S. Belcredito, S. Etteri et al., "Estrogen receptor-alpha mediates the brain antiinflammatory activity of estradiol," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 100, no. 16, pp. 9614-9619, 2003.

- [46] E. Vegeto, C. Bonincontro, G. Pollio et al., "Estrogen prevents the lipopolysaccharide-induced inflammatory response in microglia," *Journal of Neuroscience*, vol. 21, no. 6, pp. 1809-1818, 2001.
- [47] C. M. Brown, T. A. Mulcahey, N. C. Filipek, and P. M. Wise, "Production of proinflammatory cytokines and chemokines during neuroinflammation: novel roles for estrogen receptors alpha and beta," *Endocrinology*, vol. 151, no. 10, pp. 4916-4925.
- [48] M. Karin, and Y. Ben-Neriah, "Phosphorylation meets ubiquitination: the control of NF-[kappa]B activity," *Annual Review of Immunology*, vol. 18, pp. 621-663, 2000.
- [49] J. Ye, Z. Gao, J. Yin, and Q. He, "Hypoxia is a potential risk factor for chronic inflammation and adiponectin reduction in adipose tissue of ob/ob and dietary obese mice," *Am J Physiol Endocrinol Metab*, vol. 293, no. 4, pp. E1118-1128, 2007.
- [50] S. Ghisletti, C. Meda, A. Maggi, and E. Vegeto, "17beta-estradiol inhibits inflammatory gene expression by controlling NF-kappaB intracellular localization," *Molecular and Cellular Biology*, vol. 25, no. 8, pp. 2957-2968, 2005.
- [51] B. Stein, and M. X. Yang, "Repression of the interleukin-6 promoter by estrogen receptor is mediated by NF-kappa B and C/EBP beta," *Molecular and Cellular Biology*, vol. 15, no. 9, pp. 4971-4979, 1995.
- [52] F. Lago, R. Gomez, J. J. Gomez-Reino et al., "Adipokines as novel modulators of lipid metabolism," *Trends in Biochemical Sciences*, vol. 34, no. 10, pp. 500-510, 2009.
- [53] C. Di Carlo, G. A. Tommaselli, and C. Nappi, "Effects of sex steroid hormones and menopause on serum leptin concentrations," *Gynecological Endocrinology*, vol. 16, no. 6, pp. 479-491, 2002.
- [54] Y. Zhang, R. Proenca, M. Maffei et al., "Positional cloning of the mouse obese gene and its human homologue," *Nature*, vol. 372, no. 6505, pp. 425-432, 1994.
- [55] M. Maffei, J. Halaas, E. Ravussin et al., "Leptin levels in human and rodent: measurement of plasma leptin and ob RNA in obese and weight-reduced subjects," *Nature Medicine*, vol. 1, no. 11, pp. 1155-1161, 1995.
- [56] C. Di Carlo, G. A. Tommaselli, A. Sammartino et al., "Serum leptin levels and body composition in postmenopausal women: effects of hormone therapy," *Menopause*, vol. 11, no. 4, pp. 466-473, 2004.
- [57] P. J. Havel, S. Kasim-Karakas, G. R. Dubuc et al., "Gender differences in plasma leptin concentrations," *Nature Medicine*, vol. 2, no. 9, pp. 949-950, 1996.
- [58] M. Rosenbaum, M. Nicolson, J. Hirsch et al., "Effects of gender, body composition, and menopause on plasma concentrations of leptin," *Journal of Clinical Endocrinology and Metabolism*, vol. 81, no. 9, pp. 3424-3427, 1996.
- [59] H. Shimizu, Y. Shimomura, Y. Nakanishi et al., "Estrogen increases in vivo leptin production in rats and human subjects," *Journal of Endocrinology*, vol. 154, no. 2, pp. 285-292, 1997.
- [60] K. Kotani, K. Tokunaga, S. Fujioka et al., "Sexual dimorphism of age-related changes in whole-body fat distribution in the obese," *International Journal of Obesity and Related Metabolic Disorders*, vol. 18, no. 4, pp. 207-202, 1994.
- [61] R. Leenen, K. van der Kooy, P. Deurenberg et al., "Visceral fat accumulation in obese subjects: relation to energy expenditure and response to weight loss," *American Journal of Physiology*, vol. 263, no. 5 Pt 1, pp. E913-919, 1992.

- [62] T. Murakami, M. Iida, and K. Shima, "Dexamethasone regulates obese expression in isolated rat adipocytes," *Biochemical and Biophysical Research Communications*, vol. 214, no. 3, pp. 1260-1267, 1995.
- [63] F. Machinal, M. N. Dieudonne, M. C. Leneuve et al., "In vivo and in vitro ob gene expression and leptin secretion in rat adipocytes: evidence for a regional specific regulation by sex steroid hormones," *Endocrinology*, vol. 140, no. 4, pp. 1567-1574, 1999.
- [64] K. W. Yi, J. H. Shin, H. S. Seo et al., "Role of estrogen receptor-alpha and -beta in regulating leptin expression in 3T3-L1 adipocytes," *Obesity (Silver Spring)*, vol. 16, no. 11, pp. 2393-2399, 2008.
- [65] E. Hu, P. Liang, and B. M. Spiegelman, "AdipoQ is a novel adipose-specific gene dysregulated in obesity," *Journal of Biological Chemistry*, vol. 271, no. 18, pp. 10697-10703, 1996.
- [66] P. E. Scherer, S. Williams, M. Fogliano et al., "A novel serum protein similar to C1q, produced exclusively in adipocytes," *Journal of Biological Chemistry*, vol. 270, no. 45, pp. 26746-26749, 1995.
- [67] K. Hotta, T. Funahashi, Y. Arita et al., "Plasma concentrations of a novel, adipose-specific protein, adiponectin, in type 2 diabetic patients," *Arteriosclerosis, Thrombosis, and Vascular Biology*, vol. 20, no. 6, pp. 1595-1599, 2000.
- [68] Y. Arita, S. Kihara, N. Ouchi et al., "Paradoxical decrease of an adipose-specific protein, adiponectin, in obesity," *Biochemical and Biophysical Research Communications*, vol. 257, no. 1, pp. 79-83, 1999.
- [69] H. Nishizawa, I. Shimomura, K. Kishida et al., "Androgens decrease plasma adiponectin, an insulin-sensitizing adipocyte-derived protein," *Diabetes*, vol. 51, no. 9, pp. 2734-2741, 2002.
- [70] S. Horenburg, P. Fischer-Posovszky, K. M. Debatin, and M. Wabitsch, "Influence of sex hormones on adiponectin expression in human adipocytes," *Hormone and Metabolic Research*, vol. 40, no. 11, pp. 779-786, 2008.
- [71] C. M. Steppan, S. T. Bailey, S. Bhat et al., "The hormone resistin links obesity to diabetes," *Nature*, vol. 409, no. 6818, pp. 307-312, 2001.
- [72] P. D. Anderson, N. N. Mehta, M. L. Wolfe et al., "Innate immunity modulates adipokines in humans," *Journal of Clinical Endocrinology and Metabolism*, vol. 92, no. 6, pp. 2272-2279, 2007.
- [73] S. S. Pang, and Y. Y. Le, "Role of resistin in inflammation and inflammation-related diseases," *Cell Mol Immunol*, vol. 3, no. 1, pp. 29-34, 2006.
- [74] M. Bokarewa, I. Nagaev, L. Dahlberg et al., "Resistin, an adipokine with potent proinflammatory properties," *Journal of Immunology*, vol. 174, no. 9, pp. 5789-5795, 2005.
- [75] Y. H. Chen, M. J. Lee, H. H. Chang et al., "17 beta-estradiol stimulates resistin gene expression in 3T3-L1 adipocytes via the estrogen receptor, extracellularly regulated kinase, and CCAAT/enhancer binding protein-alpha pathways," *Endocrinology*, vol. 147, no. 9, pp. 4496-4504, 2006.
- [76] R. J. Norman, I. H. Flight, and M. C. Rees, "Oestrogen and progestogen hormone replacement therapy for peri-menopausal and post-menopausal women: weight and body fat distribution," *Cochrane Database Syst Rev*, no. 2, p. CD001018, 2000.

- [77] Z. Chen, T. Bassford, S. B. Green et al., "Postmenopausal hormone therapy and body composition--a substudy of the estrogen plus progestin trial of the Women's Health Initiative," *American Journal of Clinical Nutrition*, vol. 82, no. 3, pp. 651-656, 2005.
- [78] J. E. Rossouw, G. L. Anderson, R. L. Prentice et al., "Risks and benefits of estrogen plus progestin in healthy postmenopausal women: principal results From the Women's Health Initiative randomized controlled trial," *JAMA*, vol. 288, no. 3, pp. 321-333, 2002.
- [79] H. Yuksel, A. R. Odabasi, S. Demircan et al., "Effects of postmenopausal hormone replacement therapy on body fat composition," *Gynecological Endocrinology*, vol. 23, no. 2, pp. 99-104, 2007.
- [80] H. Yuksel, A. R. Odabasi, S. Demircan et al., "Effects of oral continuous 17beta-estradiol plus norethisterone acetate replacement therapy on abdominal subcutaneous fat, serum leptin levels and body composition," *Gynecological Endocrinology*, vol. 22, no. 7, pp. 381-387, 2006.
- [81] S. R. Cummings, B. Ettinger, P. D. Delmas et al., "The effects of tibolone in older postmenopausal women," *New England Journal of Medicine*, vol. 359, no. 7, pp. 697-708, 2008.
- [82] E. A. Nijland, W. C. Weijmar Schultz, and S. R. Davis, "Effects of tibolone and raloxifene on health-related quality of life and sexual function," *Maturitas*, vol. 58, no. 2, pp. 164-173, 2007.
- [83] M. L. Hammar, P. van de Weijer, H. R. Franke et al., "Tibolone and low-dose continuous combined hormone treatment: vaginal bleeding pattern, efficacy and tolerability," *BJOG*, vol. 114, no. 12, pp. 1522-1529, 2007.
- [84] M. L. Bots, G. W. Evans, W. Riley et al., "The effect of tibolone and continuous combined conjugated equine oestrogens plus medroxyprogesterone acetate on progression of carotid intima-media thickness: the Osteoporosis Prevention and Arterial effects of tiboLone (OPAL) study," *European Heart Journal*, vol. 27, no. 6, pp. 746-755, 2006.
- [85] W. Hanggi, K. Lippuner, P. Jaeger et al., "Differential impact of conventional oral or transdermal hormone replacement therapy or tibolone on body composition in postmenopausal women," *Clinical Endocrinology*, vol. 48, no. 6, pp. 691-699, 1998.
- [86] I. B. Meeuwssen, M. M. Samson, S. A. Duursma, and H. J. Verhaar, "The effect of tibolone on fat mass, fat-free mass, and total body water in postmenopausal women," *Endocrinology*, vol. 142, no. 11, pp. 4813-4817, 2001.
- [87] A. Arabi, P. Garnero, R. Porcher et al., "Changes in body composition during postmenopausal hormone therapy: a 2 year prospective study," *Human Reproduction*, vol. 18, no. 8, pp. 1747-1752, 2003.
- [88] D. E. Jacobsen, M. M. Samson, S. Kezic, and H. J. Verhaar, "Postmenopausal HRT and tibolone in relation to muscle strength and body composition," *Maturitas*, vol. 58, no. 1, pp. 7-18, 2007.
- [89] A. Wiik, M. Ekman, G. Morgan et al., "Oestrogen receptor beta is present in both muscle fibres and endothelial cells within human skeletal muscle tissue," *Histochemistry and Cell Biology*, vol. 124, no. 2, pp. 161-165, 2005.

- [90] S. Lemoine, P. Granier, C. Tiffocche et al., "Estrogen receptor alpha mRNA in human skeletal muscles," *Medicine and Science in Sports and Exercise*, vol. 35, no. 3, pp. 439-443, 2003.
- [91] A. Porcile, E. Gallardo, P. Duarte, and S. Aedo, "[Differential effects on serum IGF-1 of tibolone (5 mg/day) vs combined continuous estrogen/progestagen in post menopausal women]," *Revista Medica de Chile*, vol. 131, no. 10, pp. 1151-1156, 2003.
- [92] M. I. Lewis, G. D. Horvitz, D. R. Clemmons, and M. Fournier, "Role of IGF-I and IGF-binding proteins within diaphragm muscle in modulating the effects of nandrolone," *Am J Physiol Endocrinol Metab*, vol. 282, no. 2, pp. E483-490, 2002.

Sex Steroids

Edited by Dr. Scott M. Kahn

ISBN 978-953-307-857-1

Hard cover, 330 pages

Publisher InTech

Published online 27, January, 2012

Published in print edition January, 2012

This book, entitled "Sex Steroids", features a valuable collection of reviews and research articles written by experts in signal transduction, cellular biology, diseases and disorders. "Sex Steroids" is comprised of four sections, "The Biology of Sex Steroids", "Sex Steroids, Memory, and the Brain", "Sex Steroids and the Immune Response", and "Therapy"; individual chapters address a broad range of recognized and predicted functions and applications of sex steroids. "Sex Steroids" is intended to provide seasoned veterans as well as newcomers to this area of research with informative, resourceful, and provocative insights. Readers of "Sex Steroids" should emerge with an appreciation and understanding of the multitude and complexity of biologic processes attributed to these important hormones, and possible future directions of research in this fascinating and ever evolving field.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Kyong Wook Yi and Seung Yup Ku (2012). Adipose Tissue Metabolism and Effect of Postmenopausal Hormone Therapy on Change of Body Composition, Sex Steroids, Dr. Scott M. Kahn (Ed.), ISBN: 978-953-307-857-1, InTech, Available from: <http://www.intechopen.com/books/sex-steroids/effect-of-postmenopausal-hormone-therapy-on-adipose-tissue>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](https://creativecommons.org/licenses/by/3.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen