We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Tumor Infiltrating Lymphocytes as Prognostic Factor of Early Recurrence and Poor Prognosis of Colorectal Cancer After Radical Surgical Treatment

Vaclav Liska¹, Ondrej Daum², Petr Novak¹, Vladislav Treska¹,
Ondrej Vycital¹, Jan Bruha¹, Pavel Pitule¹ and Lubos Holubec³

¹Department of Surgery,

²Institute of Pathology,

³Department of Oncology, Charles University Prague,
Medical School and Teaching Hospital Pilsen,
Czech Republic

1. Introduction

Sixty percent of patients with colorectal cancer (CRC) are afflicted with distant metastases (liver or lung metastatic process) or a local relapse of malignancy (Bird et al., 2006). The possibilities of surgical and oncological treatment of this disease offer us a large spectrum of treatments including the combination of surgical procedures and consecutive oncological treatments. In the case of radical surgical therapy we can consider the curative access. The main medical problem of CRC is the high rate of recurrences after radically performed surgical therapy. The operability of recurrence is only about 30% in the case of local relapse and 20% in the case of distant metastases (Coleman et al., 2008; Kobayashi et al., 2007). The second dominant problem is the early recurrence of CRC after radical surgical treatment, when the patients undergo a difficult and exhausting procedure with a high risk of perioperative complications without any significant differences in overall survival against modern palliative therapy (Van den Eynde & Hendlisch, 2009).

Contemporary clinical and histopathological prognostic factors (staging, grading, etc.) used for the detection of patients with a high risk of relapse and a short overall survival rate and for the indication of adjuvant oncological treatment after radical surgery are not sufficient. Tumor infiltrating lymphocytes (TIL) were described as a good prognostic factor for patients with a high risk of relapse. They are critical indicators of efficient antitumor immunological response. Their number, type and morphology of TIL cells determine resulting tumor prognosis (Atreya & Neurath, 2008; Galon et al., 2006). They could be connected also with the suppression of micrometastatic disease after radical surgery (Gajewski et al., 2006; Pages et al., 2005). We can recognize either the type of immune cells or distinguish their morphological aspects (infiltration of any part of tumor or surrounding of tumor or tributary lymph nodes) (Talmadge et al., 2007).

We detail only short overview of their types and function. We recommend the readers with deeper interest in these problems to find comprehensive reviews in the cited papers (Jochems et al., 2011; Ohtani 2007). From this view we find CD8+ and CD4+ T lymphocytes (Fig. 1a & Fig. 1b), natural killer cells (Fig. 1c), dendritic cells (Fig. 1d), macrophages, etc. The exact function of these cells is under current discussion. We only know that they play main role in controlling tumor development and growth. CD8+ T lymphocytes within cancer cell nests of colorectal cancer have significant impact on the survival of patients. They contain the cytolytic enzyme granzyme-B. In case of increased proliferating activity of CD8+ T lymphocytes we observe their activated and cytotoxic phenotype that is significantly associated with the absence of early metastatic events (vascular emboli, lymphatic invasion or perineural invasion of tumor cells) and with a decreased rate of cancer recurrence (Atreya & Neurath, 2008, Pages et al., 2010). A high density of memory T lymphocytes within colorectal cancer tissue was more frequently observed in patients without early detectable signs of metastatic events and was associated with both improved disease free interval (DFI) and prolonged overall survival (OS) (Galon et al., 2006; Pages et al., 2005).

Natural killer cells (NK cells) mediate an effective lysis of cancer cells but the mechanism of detection of cancer cells is different from CD8+ T lymphocytes (Cooper et al., 2009). NK cells are mainly involved in the innate immune response and do not recognize specific tumor associated antigen on the surface of cancer cells as CD8+ T lymphocytes. NK cells lyse the cancer cells that are opsonized by surface antibody. NK cells also respond to other signals as cytokines produced by antigen presenting cells, which allow them to mediate early host responses against pathogen (Moretta et al., 2006). Decreased preoperative number of NK cells was associated with increased frequency of postoperative recurrence of colorectal cancer (Atreya & Neurath, 2008; Cooper et al., 2001). Their crucial role in the elimination of haematological malignancies, primary and secondary tumors has been recognized (Lucas et al., 2007; Ljunggren & Malmberg, 2007, Stojanovic & Cerwenka, 2011). In the last year there are some signs that NK-cells have the capacity for memory-like responses, a property that was previously thought to be limited to adaptive immunity, but in this view the discussion still continues (Cooper et al., 2009).

Dendritic cells are considered to be most potent antigen presenting cells. They play key role in activation, stimulation and recruitment of T lymphocytes. They can also induce antigen-specific unresponsiveness or immune tolerance. Immature dendritic cells enter tumor tissue, uptake and process its antigens. Then after they migrate to lymph nodes, undergo maturation and interact with T-lymphocytes that are able to recognize presented antigen and so T-lymphocytes play effector role of this tumor-specific immunity (Atreya & Neurath, 2008; Pages et al., 2005; Sandel et al., 2005; Steinman et al., 2003).

Macrophages are important producers of different factors that have function during tumor progression and also during tumor progression control. Their function is not fully understand, but it was described that the number of tumor infiltrating macrophages correlates with overall survival of colorectal cancer patients (Atreya & Neurath, 2008; Pollard, 2004; Forssell et al., 2007). It seems that several types of tumor infiltrating macrophages influence the balance between pro- and anti-tumor properties of immune system (Forssell et al., 2007).

From the morphologic view we can observe TIL in the specific portions of tumor and so we detect lymphocytic infiltration intratumoral (ITL – intratumoral lymphocytes) (Fig.

2a), intrastromal (ISL – intrastromal lymphocytes) (Fig. 2b), peritumoral (PTL – peritumoral lymphocytes) (Fig. 2c) and Crohn-like reaction (Crohn-like PTL)(Fig. 2d). We can also describe reactive histological changes in tributary lymph nodes (LN reactions). It means follicular hyperplasia (LN-FH) (Fig. 3a.), sinus histiocytosis (LN-SH) (Fig. 3b.) and the presence of granulomas (LN-GR) (Fig. 3c) (Ogino et al, 2009; Pages et al., 2005).

The aim of this study was to analyze the relationship of contemporary clinical and histopathological factors and TIL to determine patients with a high risk of poor overall survival and tendency to early recurrence of malignancy with shortened disease free interval (DFI) after radical surgery for CRC.

Fig. 1. Examples of tumor infiltration by immune cells: a) CD8+ T lymphocytes; b) CD4+ T lymphocytes; c) Natural killer cell (CD 57 staining) and d) Dendritic cells (S 100 staining).

Fig. 2. Different localisation of TIL within the tumor tissue: a) intratumoral lymphocytes; b) intrastromal lymphocytes; c) peritumoral lymphocytes and d) Crohn-like peritumoral lymphocytes. All sections stained with hematoxylin-eosin.

Fig. 3. Reactive histological changes in tributary lymph nodes: a) follicular hyperplasia; b) sinus histiocytosis and c) presence of granulomas. All sections stained with hematoxylin-eosin.

2. Methods

We analyzed 150 patients who underwent radical surgical procedure for CRC between the years 2004-2007 at the Department of Surgery, Medical School and Teaching Hospital in Pilsen, Charles University in Prague. We selected only patients who were operated on electively – our aim was to decrease the risk of inflammation that is often connected with the acute operation of CRC and does not depend on the immune reaction against a tumor but could be evoked by the distension of the bowel. We also excluded patients who had risk of understaging (for example low number of analysed lymph nodes) and patients with a synchronous metastatic process. The inclusion standard was also the entire follow-up of patients during the whole postoperative period to increase the number of patients with a diagnosed early recurrence of CRC.

The following clinical parameters were statistically analysed in relation to the disease free interval (DFI) and the overall survival (OS): staging, grading, preoperative leukocytosis, type of surgical procedure (radical vs. palliative), postoperative complications and postoperative oncological treatment.

2.1 Histology

We examined three different sections of each tumor and also sections of all found lymph nodes. Tissue for light microscopy was fixed in 4% formaldehyde and embedded in paraffin using routine procedures. Five micrometer-thick sections were cut from the tissue blocks and stained with hematoxylin-eosin.

The type and grade of all tumors were determined according to WHO 2000 guidelines. The stage of tumors was established according to UICC 2002 guidelines. We evaluated endovascular (VI), endolymphatic (LI) and perineural infiltration (PI) by cancer cells (0 – none, 1 – yes). Lymphocytic infiltration was detected as intratumoral (ITL – intratumoral lymphocytes), intrastromal (ISL – intrastromal lymphocytes), peritumoral (PTL – peritumoral lymphocytes) and Crohn-like reaction (Crohn-like PTL), and scaled as none (0), mild (1), moderate (2) and severe (3). Reactive histological changes in lymph nodes (LN reactions) were detected as follicular hyperplasia (LN-FH), sinus histiocytosis (LN-SH) and the presence of granulomas (LN-GR), and all these parameters were quantified in the same manner as lymphocytic infiltration.

2.2 Immunohistochemistry

For immunohistochemical investigations the following primary antibodies were used: CD4 (clone 4B12, 1:50, Vector Laboratories, Burlingame, CA, USA) and CD8 (clone C8/144B, 1:50, Dako, Glostrup, Denmark). Microwave pretreatment was used in both cases. The primary antibodies were visualized using the supersensitive streptavidin-biotin-peroxidase complex (Biogenex, San Ramon, CA). Appropriate positive and negative control slides were employed. The density of intratumoral infiltration by lymphocytes was evaluated in five High power microscopical fields (HPF) and expressed as the number of immunopositive cells per HPF.

2.3 Statistical evaluation

Statistical analysis was processed by the statistical software Statistica 9.0. The mean, median, standard deviation (SD), minimum, maximum, quartiles, frequencies and other basic statistical measurements were computed in given groups and subgroups of patients corresponding to studied clinical and histopathological parameters.

The relationships between the variables were described by Spearman rank correlation coefficients.

The analyses of Overall survival (OS) and Disease free interval (DFI) were performed by Kaplan-Meier’s survival functions. The influence of given covariates (clinical and histopathological factors) was tested by the Log-Rank test and Wilcoxon test. The Cox regression hazard model, hazard ratio (HR) and 95% confidence interval (CI) for HR were computed for the evaluation of given clinical and histopathological factors to OS or DFI. Multivariate analysis was performed by the use of classification and regression trees (CART). The Cox regression hazard model (stepwise regression) was applied to find the predictors in CART.

3. Results

The statistical analysis of the studied cohort of patients after surgical treatment for colorectal cancer demonstrated an acceptable distribution of basic statistical description parameters (gender ratio 93:57 (male vs. female)). 1, 3 and 5 years overall survival was 92.2%, 76.5% and 70.2% and 1, 3 and 5 year DFI was 85.3%, 64.3% and 49.4%.

The Spearman rank correlation coefficient did not prove any stronger correlation than a moderate correlation at endolymphatic invasion (LI) and lymph node infiltration by metastatic process (Spearman rank correlation coefficient 0.56, $p<0,05$). All the other studied factors were independent factors or factors with a low correlation.

Statistical analysis proved lymph node infiltration by metastatic process as statistically significant for the prognosis of overall survival ($p<0.05$) and N2 status of lymph nodes increased the risk of shorter overall survival 9.3x (Fig. 4.).

Fig. 4. Statistical analysis proved the lymph node infiltration by a metastatic process as statistically significant for the prognosis of overall survival ($p<0.05$), and the N2 status of lymph nodes increased the risk of a shorter overall survival 9.3x.

Endovascular infiltration (VI) was proved as a negative prognostic factor of shorter overall survival (Fig.5.). Patients with positive histopathological findings of VI have 3.1x increased risk for shorter overall survival. The presence of peritumoral lymphocytes (PTL) (Fig.6.) and of Crohn-like PTL (Fig.7.) was proved as a positive prognostic factor of OS. Patients with a positive histopathological finding of PTL and Crohn-like PTL have a decreased risk for shorter overall survival (2.3x and 2.3x respectively). Lymph node follicular hyperplasia (LN-FH) was verified as a positive prognostic factor for longer overall survival (Fig.8.). The statistical significance of LN-FH increased also with the raised density of infiltration. LN-FH positivity decreased the risk of shorter overall survival 3.3times. The severity of CD8+ lymphocytic infiltration was proved by the Cox regression hazard model as a positive prognostic factor enlarging overall survival (cut off 30 cells/HPF). The severity of CD4+ lymphocytic infiltration was proved as a significant factor for the prognosis of overall survival (cut off 4cells/HPF) with 2.5x increased hazard ratio in patients over the cut off (Fig.9.). Statistical analysis did not confirm the statistical significance of CD8/CD4 ratio.

Fig. 5. Endovascular infiltration by cancer cells (VI) was proved as a statistically significant factor for the prognosis of overall survival ($p<0.05$). The patients with a positive histopathological finding of VI have 3.1x higher risk ratio for shorter overall survival.

Fig. 6. Presence of peritumoral lymphocytes (PTL) was proved as a statistically significant positive factor for the prognosis of overall survival ($p<0.05$). The patients with a positive histopathological finding of PTL have 2.3x lower risk ratio for a shorter overall survival.

Fig. 7. Presence of Crohn-like PTL was proved as a statistically significant positive factor for the prognosis of overall survival ($p<0.05$). The patients with a positive histopathological finding of PTL have 2.3x lower risk ratio for a shorter overall survival.

Fig. 8. Lymph node follicular hyperplasia (LN-FH) was verified as a positive prognostic factor for a longer overall survival ($p<0.05$). The statistical significance of LN-FH increased also with the raised density of infiltration. LN-FH positivity decreased the risk of a shorter overall survival 3.3x.

Fig. 9. Severity of CD4+ lymphocytic intratumoral infiltration was proved as a significant factor for the prognosis of overall survival (cut off 4cells/HPF) with a 2.5x increased hazard ratio in patients over the cut off ($p<0.05$).

The Multivariate Cox Regression Hazard Model proved the combination of the severity of lymph node infiltration by metastatic process and LN-FH as the best prognostic factors for the prediction of the risk of shorter overall survival. This situation is demonstrated in the Classification and Regression Tree (CART)($p<0.05$) (Fig.10.). All other studied parameters were not proved as statistically significant for the prognosis of overall survival.

Classification and Regression Tree

Overall Survival

Fig. 10. Multivariate Cox Regression Hazard Model proved the combination of the severity of lymph node infiltration by a metastatic process and LN-FH as the best prognostic factors for the prediction of risk of a shorter overall survival ($p<0.05$). This situation is demonstrated in the Classification and Regression Tree (CART).

Perineural infiltration (PI) was proved as a negative prognostic factor of an earlier recurrence (Fig.11.). Patients with a positive histopathological finding of PI have 3.8x increased risk for shorter DFI.

The severity of CD8+ lymphocytic infiltration was proved by the Cox regression hazard model as a positive prognostic factor enlarging DFI (cut off 30cells/HPF) (Fig.12.). Patients over the cut off have 2.2x increased risk of an early recurrence. The severity of CD4+ lymphocytic infiltration was not proved as a significant factor for the prognosis of DFI. Statistical analysis did not confirm the statistical significance of the CD8/CD4 ratio.

Fig. 11. Perineural infiltration (PI) was proved as a negative prognostic factor of an earlier recurrence ($p<0.05$). Patients with a positive histopathological finding of PI have a 3.8x increased risk for shorter DFI.

Statistical analysis proved lymph node infiltration by a metastatic process as statistically significant for the prognosis of DFI and N2 status of lymph nodes increased the risk of shorter DFI 5x (Fig. 13.).

Fig. 12. Severity of CD8+ lymphocytic intratumoral infiltration was proved as a positive prognostic factor enlarging DFI (cut off 30 cells/HPF) by the Cox regression hazard model ($p<0.05$). Patients over the cut off have a 2.2x increased risk of an early recurrence.

Fig. 13. Statistical analysis proved lymph node infiltration by a metastatic process as statistically significant for the prognosis of DFI, and a N2 status of lymph nodes 5x increased the risk of a shorter DFI ($p<0.05$).

The Multivariate Cox Regression Hazard Model proved the combination of the severity of the lymph node infiltration by a metastatic process and the severity of CD8 positivity of infiltrating lymphocytes as the best prognostic factors for the prediction of risk of early recurrence ($p<0.05$). This situation is demonstrated in the Classification and Regression Tree (CART) (Fig. 14). All other studied parameters were not proved as statistically significant for the prognosis of DFI.

4. Discussion

The role of the adaptive immunological response in controlling the growth and relapse of CRC remains controversial and contemporary studies have not answered all the questions about the prognosis of patients after radical surgical treatment of CRC (Galon et al., 2006; Ohtani, 2007; Van den Eynde & Hendlisz, 2009). We analysed our large cohort of patients of CRC with consideration to detect the negative and also positive prognostic factors of early recurrence of the disease and the poor overall survival after radical surgery. It was stimulated by the unsatisfactory situation and some dilemmas in the indication of surgical and oncological treatment, when early recurrence depreciates our effort to radical surgery with a high risk of complications and the long time of the decreased quality of life of our patients.

In the presented clinico-pathological study we demonstrated that lymph node infiltration by a metastatic process, N2 status of lymph nodes, VI, and extent of CD4+ lymphocytic intratumoral infiltration as negative prognostic factors of OS. In contrary PTL, Crohn-like PTL, LN-FH, and severity of CD8+ lymphocytic intratumoral infiltration were proved as positive prognostic factors of the overall survival.

Fig. 14. Multivariate Cox Regression Hazard Model proved the combination of the severity of lymph node infiltration by a metastatic process and the severity of CD8 positivity as the best prognostic factors for the prediction of risk of an early recurrence ($p<0.05$). This situation is demonstrated in the Classification and Regression Tree (CART).

The combination of the severity of the lymph node infiltration by a metastatic process and LN-FH were proved as the best prognostic factors for the prediction of risk of a shorter overall survival by the Multivariate Cox Regression Hazard Model.

We also demonstrated PI, lymph node metastatic infiltration and the N2 status of lymph nodes as negative prognostic factors of an earlier recurrence, and the severity of CD8+ lymphocytic intratumoral infiltration as a positive prognostic factor enlarging DFI. The combination of the severity of the lymph node infiltration by a metastatic process and the severity of CD8+ lymphocytic intratumoral infiltration were proved as the best prognostic factors for the prediction of the risk of an early recurrence by the Multivariate Cox Regression Hazard Model.

Our results support the hypothesis that the adaptive immunological response in tumor tissue and its reaction in regional lymph nodes can influence the behavior of CRC and so affect the prognosis of patients (Atreya & Neurath, 2008; Galon et al., 2006). CD4 and CD8 positivity of ITL was demonstrated as a key histopathological sign of tumor-specific immune response that could reflect the contemporary clinical situation and a tendency to relapse (CD4+) or the larger overall survival (CD8+) (Chiba et al. 2004; Koch et al., 2006, Pages et al., 2010).

We assessed several types of tumor infiltrating lymphocytes and clearly identified their relationships to relapse and the overall survival as positive or negative prognostic factors in contrary to previous publications that evaluated only the infiltration of the tumor but not the quality and type of infiltration (Ogino et al., 2009).

Tumor infiltration by lymphocytes seems to be a promising prognostic factor reflecting the risk of patients to early recurrence or poor overall survival. Future work has to be focused on the molecular-biological background of tumor infiltration by lymphocytes to understand their pathophysiological functions (Pages et al., 2005, Zbar, 2004).

5. Conclusion

Tumor infiltrating lymphocytes seem to be promising prognostic factors that could find their use in colorectal surgery and consecutive oncological treatment as an indicator of the type or combinations of therapies reflecting the risk of patients to early recurrence or poor overall survival. The TIL status corresponds to immune control of cancer progression.

6. Acknowledgment

This article was supported by research project MSM 0021620819 (Replacement of and support to some vital organs) and grant IGA MZ CR 10230 and IGA MZ CR 12025 and specific students research grant of Charles University SVV-2011- 262 806.

7. References

- Atreya, I., & Neurath, M. F. (2008). Immune cells in colorectal cancer: prognostic relevance and therapeutic strategies. *Expert Review of Anticancer Therapy*, Vol.8, No.4, (April 2008), pp. 561-572, ISSN 1473-7140
- Bird, N. C., Mangnall, D., & Majeed, A. W. (2006). Biology of colorectal liver metastases: a review. *Journal of Surgical Oncology*, Vol.94, No.1, (July 2006), pp. 68-80, ISSN 0022-4790
- Chiba, T., Ohtani, H., Mizoi, T., Naito, Y., Sato, E., Nagura, H., Ohuchi, A., Ohuchi, K., Shiiba, K., Kurokawa, Y. & Satomi, S. (2004). Intraepithelial CD8+ T-cell-count becomes a prognostic factor after a longer follow-up period in human colorectal carcinoma: possible association with suppression of micrometastasis. *British Journal of Cancer*, Vol.91, No.9, (November 2004), pp. 1711-1717, ISSN 0007-0920
- Coleman, M. P., Quaresma, M., Berrino, F., Lutz, J. M., De Angelis, R., Capocaccia, R., Baili, P., Rachet, B., Gatta, G., Hakulinen, T., Micheli, A., Sant, M., Weir, H. K., Elwood, J. M., Tsukuma, H., Koifman, S., Silva, E., Francisci, S., Santaquilani, M., Verdecchia, A., Storm, H. H., Young, J. L. & CONCORD Working Group. (2008). Cancer survival in five continents: a worldwide population-based study (CONCORD). *The Lancet Oncology*, Vol.9, No.8, (August 2008), pp. 730-756, ISSN 1470-2045
- Cooper, M. A., Fehniger, T. A., & Caligiuri, M. A. (2001). The biology of human natural killer – cell subsets. *Trends in Immunology*, Vol.22, No.11, (November 2001), pp. 633-640, ISSN 1471-4906
- Cooper, M. A., Colonna, M., & Yokoyama, W. M. (2009). Hidden talents of natural killers: NK cells in innate and adaptive immunity. *Embo reports*, Vol.10, No.10, (October 2009), pp. 1103-1110, ISSN 1469-221X

- Forssell, J., Oberg, A., Henriksson, M. L., Stenling, R., Jung, A., & Palmquist, R. (2007). High macrophage infiltration along the tumor front correlates with improved survival in colon cancer. *Clinical Cancer research*, Vol.1, No.13, (March 2007), pp. 1472-1479, ISSN 1078-0432.
- Gajewski, T. F., Meng, Y., Harlin, H. (2006). Immune suppression in the tumor microenvironment. *Journal of Immunotherapy*, Vol.29, No.3, (May-June 2006), pp. 233-240, ISSN 1524-9557
- Galon, J., Costes, A., Sanchez-Cabo, F., Kirilovsky, A., Mlecnik, B., Lagorce-Pagès, C., Tosolini, M., Camus, M., Berger, A., Wind, P., Zinzindohoué, F., Bruneval, P., Cugnenc, P. H., Trajanoski, Z., Fridman, W. H. & Pagès, F. (2006). Type, density, and location of immune cells within human colorectal tumors predict clinical outcome. *Science*, Vol.313, No.5795, (September 2006), pp. 1960-1964, ISSN 0036-8075
- Jochems C, Schlom J. (2011). Tumor-infiltrating immune cells and prognosis: the potential link between conventional cancer therapy and immunity. *Exp Biol Med (Maywood)*. Vol. 236, No.5 (May 2011), pp. 567-79, ISSN 1535-3702
- Kobayashi, H., Mochizuki, H., Sugihara, K., Sugihara, K., Morita, T., Kotake, K., Teramoto, T., Kameoka, S., Saito, Y., Takahashi, K., Hase, K., Oya, M., Maeda, K., Hirai, T., Kameyama, M., Shirouzu, K. & Muto T. (2007). Characteristics of recurrence and surveillance tools after curative resection for colorectal cancer: a multicenter study. *Surgery*, Vol.141, No.1, (January 2007), pp. 67-75, ISSN 0039-6060
- Koch, M., & Bechthove, P. (2006). Op den Winkel J et al. Tumor infiltrating T lymphocytes in colorectal cancer: Tumor-selective activation and cytotoxic activity in situ. *Annals of Surgery*, Vol.244, No.6, (December 2006), pp. 986-992, ISSN 0003-4932
- Ljunggren, H.G., & Malmberg, K.J. (2007). Prospects for the use of NK cells in immunotherapy of human cancer. *Nature Reviews. Immunology*, Vol.7, No.5, (May 2007), pp. 329-339, ISSN 1474-1733
- Lucas, M., Schachterle, W., Oberle, K., Aichele, P., & Diefenbach, A. (2007). Dendritic cells prime natural killer cells by trans-presenting interleukin 15. *Immunity*, Vol.26, No.4, (April 2007), pp. 503-517, ISSN 1074-7613
- Moretta, L., Ferlazzo, G., Bottino, C., Vitale, M., Pende, D., Mingari M. C., & Moretta, A. (2006). Effector and regulatory events during natural killer-dendritic cell interaction. *Immunological Reviews*, Vol.214, No.1, (December 2006), pp. 219-228, ISSN 0105-2896
- Ogino, S., Nosho, K., Irahia, N., Meyerhardt, J. A., & Baba, Y. (2009). Lymphocytic reaction to colorectal cancer is associated with longer survival, independent of lymph node count, microsatellite instability, and CpG island methylation phenotype. *Clinical Cancer Research*, Vol. 15, No. 20, (October 2009), pp. 6412-6420, ISSN 1078-0432
- Ohtani, H. (2007). Focus on TILs: prognostic significance of tumor infiltrating lymphocytes in human colorectal cancer, *Cancer immunity*, Vol. 7, (February 2007), pp. 4, ISSN 1424-9634
- Pagès, F., Berger, A., Camus, M., Costes, A., Molitor, R., Mlecnik, B., Kirilovsky, A., Nilsson, M., Damotte, D., Meatchi, T., Bruneval, P., Cugnenc, P. H., Trajanoski, Z., Fridman, W. H., & Galon, J. (2005). Effector memory T cells, early metastasis, and survival in colorectal cancer. *The New England Journal of Medicine*, Vol.353, No.25, (December 2005), pp. 2654-2666, ISSN 0028-4793

- Pagès, F., Galon, J., Die-Nosjeanu, M. C., Tartour, E., & Sautes-Fridman, C. (2010). Immune infiltration in human tumors: a prognostic factor that should not be ignored. *Oncogene*, Vol.29, No.8, (February 2010), pp. 1093-1102, ISSN 0950-9232
- Pollard, J. W. (2004). Tumor-educated macrophages promote tumour progression and metastasis. *Nature Reviews. Cancer*, Vol.4, No.1, (January 2004), pp. 71-78, ISSN 1474-175X
- Sandel, M. H., Dadabayev, A. R., Menon, A. G., Morreau, H., Melief, C. J., Offringa, R., van der Burg, S. H., Janssen-van Rhijn, C. M., Ensink, N. G., Tollenaar, R. A., van de Velde, C. J., & Kuppen, P. J. (2005). Prognostic value of tumor-infiltrating dendritic cells in colorectal cancer: role of maturation status and intratumoral localization. *Clinical Cancer Research*, Vol.11, No.7, (April 2005), pp. 2576-2582, ISSN 1078-0432
- Steinman, R. M., Hawiger, D., & Nussenzweig, M. C. (2003). Tolerogenic dendritic cells. *Annual review of immunology*, Vol.21, pp. 685-711, ISSN 0732-0582
- Stojanovic, A., & Cerwenka, A. (2011). Natural Killer cells and solid tumors. In: *Journal of Innate Immunity*, 10.6.2011, Available from: <<http://content.karger.com/produktedb/produkte.asp?doi=325465>>.
- Talmadge, J. E., Donkor, M., & Scholar, E. (2007). Inflammatory cell infiltration of tumors: Jekyll or Hyde. *Cancer and Metastasis Reviews*. Vol.26, No.3-4, (December 2007), pp. 373-400, ISSN 0167-7659
- Van den Eynde, M., & Hendlisz, A. (2009) Treatment of colorectal liver metastases: a review. *Reviews on Recent Clinical Trials*, Vol.4, No.1, (January 2009), pp. 56-62, ISSN 1574-8871
- Zbar, A. P. (2004) The immunology of colorectal cancer. *Surgical Oncology*, Vol.13, No.2-3, (August-November 2004), pp. 45-53, ISSN 0960-7404

IntechOpen

Colorectal Cancer Biology - From Genes to Tumor

Edited by Dr. Rajunor Ettarh

ISBN 978-953-51-0062-1

Hard cover, 446 pages

Publisher InTech

Published online 10, February, 2012

Published in print edition February, 2012

Colorectal cancer is a common disease, affecting millions worldwide and represents a global health problem. Effective therapeutic solutions and control measures for the disease will come from the collective research efforts of clinicians and scientists worldwide. This book presents the current status of the strides being made to understand the fundamental scientific basis of colorectal cancer. It provides contributions from scientists, clinicians and investigators from 20 different countries. The four sections of this volume examine the evidence and data in relation to genes and various polymorphisms, tumor microenvironment and infections associated with colorectal cancer. An increasingly better appreciation of the complex inter-connected basic biology of colorectal cancer will translate into effective measures for management and treatment of the disease. Research scientists and investigators as well as clinicians searching for a good understanding of the disease will find this book useful.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Vaclav Liska, Ondrej Daum, Petr Novak, Vladislav Treska, Ondrej Vycital, Jan Bruha, Pavel Pitule and Lubos Holubec (2012). Tumor Infiltrating Lymphocytes as Prognostic Factor of Early Recurrence and Poor Prognosis of Colorectal Cancer After Radical Surgical Treatment, *Colorectal Cancer Biology - From Genes to Tumor*, Dr. Rajunor Ettarh (Ed.), ISBN: 978-953-51-0062-1, InTech, Available from:
<http://www.intechopen.com/books/colorectal-cancer-biology-from-genes-to-tumor/tumor-infiltrating-lymphocytes-as-prognostic-factor-of-early-recurrence-and-poor-prognosis-of-colore>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](https://creativecommons.org/licenses/by/3.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen