We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Fragile X Syndrome: From Pathophysiology to New Therapeutic Perspectives

Simona D'Antoni¹, Michela Spatuzza¹, Carmela M. Bonaccorso²,
Elisabetta Aloisi^{1,3}, Sebastiano Musumeci² and Maria Vincenza Catania^{1,2}

¹*Institute of Neurological Sciences, CNR, Catania,*

²*Oasi Maria SS Troina (EN),*

³*PhD Program in Neuropsychopharmacology, University of Catania,
Italy*

1. Introduction

In the present chapter we will provide an overview of recent literature regarding new therapeutic perspectives in Fragile X syndrome (FXS), which are based on a rational approach well-grounded on a deeper understanding of the disease pathophysiology. FXS represents a paradigmatic example of how research can be translated into therapy targeting dysfunctional mechanisms rather than symptoms. Several clinical trials using these new strategies are underway. Here, we will mainly describe the basic mechanisms and the animal studies which suggest the use of these innovative pharmacological approaches. In addition, an emerging concept is that developmental pathologies with intellectual disability (ID) presenting common features such as autism, behavioural disturbances and epilepsy might share dysregulation of the same biochemical pathways. The identification of common altered pathways in ID might help to develop new therapeutic strategies helpful for apparently diverse pathologies.

2. Fragile X syndrome

2.1 FXS: Clinic and genetics

ID, also referred with the term Mental Retardation, is the most common developmental disorder, with a prevalence of 1-3%, and includes a highly diverse group of cognitive disorders. It is defined, according to the American Psychiatric Association, by an intelligence quotient (IQ) of 70 or below, and deficits in at least two behaviours related to adaptive functioning diagnosed by 18 years of age. Gene defects account for about half of all patients and mutations have been identified in more than 400 genes, of which 97 are positioned in the X chromosome (reviewed in Kaufman et al., 2010).

FXS is an X-linked developmental disorder which represents the most common form of inherited ID, affecting approximately 1 in 2500-6000 males and 1 in 4000-8000 females. ID ranges from severe to mild and may be associated with Attention Deficit and/or Hyperactive Disorder (ADHD), autism, behavioral disturbances, hyperactivity, seizures and hypersensitivity to sensory stimuli. People with FXS may also exhibit facial dysmorphic

features including a long face with prominent ears and arched palate, hyperextensible joints, mitral valve prolapse and macroorchidism (R.J. Hagerman, 2002).

The most common genetic defect in FXS is a CGG trinucleotide repeat expansion of >200 repeats in the 5' untranslated region of the FMR1 (fragile X mental retardation 1) gene, located on the long arm of the X chromosome at position 27.3 (Verkerk et al., 1991). This triplet amplification is associated to methylation of the FMR1 promoter region and transcriptional silencing of the FMR1 gene with consequent loss or significant reduction of the FMR1 encoded protein FMRP (fragile X mental retardation protein) (Devys et al., 1993; O'Donnel & Warren, 2002). Expansions of CGG repeats are instable during meiosis, increasing in length from one generation to the next. In carriers of the premutation, the expansion is between 55–200 repeats (normal is <45), and does not result in FMR1 methylation and loss of FMRP expression, but gives rise to two independent pathologies such as fragile X-associated tremor/ataxia syndrome (FXTAS) and premature ovarian failure, primarily in males and females respectively (reviewed in Berry-Kravis et al., 2007; P.J. Hagerman et al., 2008; Toniolo, 2006). It has been hypothesised that these conditions are caused by a gain of function toxic effect of increased levels of CGG repeat-containing FMR1 mRNA reviewed in (reviewed in Berry-Kravis et al., 2007), although decreased levels of FMRP may also play a role (Qin et al., 2011).

2.2 FXS: Alterations of dendritic spine morphology

Microscopic analysis of brain material from both patients with FXS and mouse models of the disease reveals no gross morphological abnormalities (Bakker et al., 1994; Reyniers et al., 1999). However, in certain brain areas such as cortex and hippocampus, long and thin dendritic spines have been observed, consistent with an immature spine phenotype (Comery et al., 1997; Irwin et al., 2001, 2002; Nimchinsky et al., 2001).

Dendritic spines are protrusions of dendritic membrane and serve as the postsynaptic component for the vast majority of central nervous system (CNS) excitatory synapses. Spines are dynamic structures that can regulate many neurochemical events related to synaptic transmission and modulate synaptic efficacy. The tip of the spine contains an electrondense region, the "postsynaptic density" (PSD), that is a protein dense specialization and consists of receptors, channels, and signaling proteins involved in synaptic transmission. Spines are highly motile structures, their density varies across areas of different brain regions but also within individual dendritic trees; spine morphology changes with development and requires actin cytoskeleton remodelling and local protein translation in response to synaptic activity. Notably, spines are equipped with translational machinery and protein synthesis may occur in response to receptor activation. The structural modifications of spines are correlated with synaptic plasticity (see below); Long Term Depression (LTD) is generally associated with a shrinkage of spines, whereas Long Term Potentiation (LTP) causes formation of new spines and enlargements of existing spines (Tada & Sheng, 2006).

Abnormalities in dendrites and spines have been implicated in several psychiatric disorders and have been associated with cognitive impairment and mental retardation disorders (Tuberous Sclerosis Type I, Fetal alcohol syndrome, Down syndrome, Rett syndrome, autism and FXS) (Nimchinsky et al., 2002), but the causes of these malformations are not yet well understood.

2.3 FMRP: Expression, structure and interacting proteins

FMRP is an RNA binding protein involved in the regulation of target mRNA translation and transport. It belongs to a small family of highly conserved RNA binding proteins referred to as the fragile X-related (FXR) proteins; it is expressed in several tissues and organs and has been found to be most abundant in the brain and testis. FMRP is highly expressed in neurons and is associated with translating polyribosomes and ribonucleoprotein complexes (mRNP) in the cytoplasm, in dendrites and dendritic spines where it is believed to regulate mRNA translation (De Diego Otero et al., 2002). Recent data also suggest that FMRP is present in axons and pre-synaptic terminals (Christie et al., 2008).

The analysis of the structure of FMRP has revealed the presence of different functional motifs and has contributed to elucidate the function of the protein. FMRP contains three different RNA binding domains: two hnRNP K-protein homology (KH) domains and an Arg-Gly-Gly (RGG) box (Siomi et al., 1993), which bind sequence-specific elements such as the U-rich sequences called FMRP kissing complex and G-quartet, respectively (Darnell et al., 2001, 2005). Interestingly, a missense mutation in the second hnRNP KH binding domain (I304N) abolishes FMRP association with polyribosomes and causes FXS. The presence within FMRP of a nuclear localization signal (NLS) and a nuclear export signal (NES) suggests that FMRP is a shuttle protein and that it travels between the nucleus and the cytoplasm (Darnell et al., 2001, 2005; Eberhart et al., 1996). In the nucleus, FMRP binds to RNAs and proteins to form the mRNP particle and is then exported to the cytoplasm where it could associate with translating ribosomes (Corbin et al., 1997; Eberhart et al., 1996; Feng et al., 1997a; Khandjian et al., 1996). The mRNP complex can stay in the neuronal cell body or it can move to the dendritic spines via the microtubule structures present in the dendrites. In this way, FMRP can control the local protein synthesis at the synapses, influencing synaptic function, structure and plasticity (Bardoni et al., 2006; Feng et al., 1997b; Miyashiro et al., 2003; Zukin et al., 2009).

The structure of FMRP presents also two coiled coil (CC) domains involved in protein-protein interactions. Using immunoprecipitation two-hybrid screens or large mass spectrometry analysis several FMRP interacting proteins have been identified including its two close paralogs, FXR1P and FXR2P (Fragile X Related Protein 1/2), NUFIP1 (Nuclear FMRP Interacting Protein 1), 82-FIP (82 kDa-FMRP Interacting Protein) and the two closely related proteins CYFIP1 and CYFIP2 (Cytoplasmic FMRP Interacting Protein 1/2). The role and importance of these interacting proteins in the function of FMRP is not clear; it is possible that the interaction with these proteins might modulate the function of FMRP in different cellular compartments (reviewed by Bardoni et al., 2006).

FXR1P and FXR2P show a similar structure to that of FMRP, being characterized by the presence of two KH and one RGG box RNA binding domains and nuclear localization and export signals (NLS and NES). In the absence of FMRP there is not a compensatory increase in levels of FXR1P and FXR2P, which would suggest functional redundancy. However, the precise role of the two FMRP paralogs and their reciprocal interaction is still under investigation.

CYFIP1 and CYFIP2 are highly homologous to each other; CYFIP2 interacts with all members of the FXR family, while CYFIP1 is specific for FMRP (Schenck et al., 2003). CYFIP1 and 2 are localized at synapses and CYFIP1 also interacts with activated Rac1 (Kobayashi et al., 1998; Schenck et al., 2003), a small RhoGTPase involved in maturation and

maintenance of dendritic spines (Govek et al., 2005), suggesting that FMRP might influence cytoskeleton remodelling through Rho/Rac GTPase (Schenck et al., 2003). The interaction between FMRP and CYFIP1 has been proposed to mediate the inhibition of translation initiation by sequestering the cap-binding protein eIF4E (Napoli et al., 2008).

2.4 FMRP: Regulation of target mRNA translation and transport

There is a general consensus that FMRP acts mainly as a negative regulator of translation although the underlying mechanisms are not clear. Several mechanisms have been proposed and they may not be mutually exclusive. The majority of co-sedimentation studies has found an association of FMRP with polyribosomes and suggests that FMRP acts by repressing elongation (reviewed by Bardoni et al., 2006), although other studies suggest that FMRP is associated with BC1 (a non translatable RNA), a complex which will block the initiation step through an interaction with eIF-4E-BP and CYFIP1 (Napoli et al., 2008). FMRP has been found also associated to high-density granules, which represent ribonucleic aggregates where mRNA translation is stalled (Aschrafi et al., 2005). A recent work supports a model in which FMRP acts to stall ribosomal translocation during elongation; although the exact mechanism by which FMRP stalls ribosomes remains to be determined, authors suggest that it is a dynamic and reversible mechanism related with plastic changes occurring both in the cytoplasm and at synapses (Darnell et al., 2011). Another mechanism by which FMRP might control expression levels of proteins is through the regulation of transcript stability, such as that of microRNA-124a (miRNA-124a) and PSD-95 (Xu et al., 2008; Zalfa et al., 2007). A further element of complexity is added by recent data suggesting that FMRP may also promote translation of target mRNAs, such as Trailer-Hitch and Superoxide Dismutase 1 (SOD1) transcripts (Bechara et al., 2009; Monzo et al., 2006). Thus, the translation and expression of FMRP targets can be either positively or negatively affected by FMRP expression, indicating that the potential role of FMRP as a translational regulator is much more complex than it was originally believed.

In addition to its role as a regulator of translation FMRP has been involved in the regulation of RNA transport along dendrites. A number of putative RNA targets have been found to be abundantly expressed in dendrites, although no major changes have been detected in the steady-state distribution and expression levels in the absence of FMRP (Bassel & Warren, 2008). FMRP traffics in the form of motile "RNA granules", structures different in size and composition containing translationally repressed mRNP complexes which travel on microtubules to the dendrites. mRNAs, once localized to the appropriate sites, are released from granules and translated in response to appropriate stimuli (reviewed in Bassel & Warren, 2008). FMRP trafficking is regulated in response to activation of group-I metabotropic glutamate (mGlu) receptors. Application of 3,5-Dihydroxyphenylglycine (DHPG), a selective agonist of group-I mGlu receptors, enhances the dendritic transport of several FMRP target mRNAs, including those encoding FMRP, Map1b, CaMKII in hippocampal cultured neurons (Antar et al., 2004; Dictenberg et al., 2008; Ferrari et al., 2007). Dictenberg shows that FMRP, upon DHPG stimulation, interacts more efficiently with the kinesin light chain and this mGlu-receptor mediated transport is markedly attenuated in the absence of FMRP. These data suggest that FMRP is involved in promoting the activity-dependent localization of bound mRNAs, but not in the constitutive transport of mRNAs in dendrites.

It is clear that, as a consequence of the lack of FMRP, levels of several synaptic and non-synaptic proteins are altered and key biochemical pathways might be dysregulated in FXS.

The *in vivo* evidence that an overall increase of protein synthesis in several brain regions occurs in FXS has been provided by quantitative autoradiographic studies using radioactively labelled aminoacid L-[1-14C]leucine, which showed an increase in several regions of Fmr1 knock out (KO) mice compared to wild type (WT) (Qin et al., 2005). Accordingly, Dölen and collaborators have shown a 20% increase protein synthesis in hippocampal slices of Fmr1 KO mice compared to WT using 35S-methionine/cystine labelling (Dölen et al., 2007). These studies corroborate the view that FMRP acts mainly as inhibitor of protein synthesis in the brain, although do not exclude the possibility that certain proteins might be downregulated in a direct or an indirect way as a result of dysregulated pathways.

The identification of target mRNAs has been object of intense research during the last years, using a variety of *in vitro* assays. A recent work has identified 842 FMRP mRNA targets using a stringent high-throughput sequencing-cross-linking immunoprecipitation (HITS-CLIP) method (Darnell et al., 2011). An overlap has been found with a list of FMRP mRNA targets previously identified with a co-immunoprecipitation method (181 mRNAs) (V. Brown et al., 2001), but a significant number of mRNAs are newly identified. Interestingly, this list includes several well-studied autism candidate genes such as NLGN3, NRXN1, SHANK3, PTEN, TSC2 and NF1 and components of pre- and post-synaptic compartments.

2.5 FXS animal models

A major advancement towards a better understanding of the molecular mechanisms implicated in FXS is represented by the development of FXS animal models, which have been also used for pre-clinical studies aimed at testing potential therapeutic interventions. Mouse and *Drosophila melanogaster* are the main genetic model organisms used to these purposes. The mouse Fmr1 gene and its two related genes Fxr1 and Fxr2 are well conserved relative to their human homologs FMR1, FXR1 and FXR2, respectively (Bakker et al., 1994; Bontekoe et al., 2002; Mientjes et al., 2004), whereas the fly model organism has a single FMR1 homolog (dFmr1) that is more functionally similar to human FMRP than to human FXR1 or FXR2 (Coffee et al., 2010). Both the fly and the mouse models present phenotypic abnormalities that are similar to those observed in humans such as: behavioural changes, altered axon morphology and connectivity, social, memory and learning deficits. The Fmr1 KO mouse shows macroorchidism, hyperactivity, a mild spatial learning impairment in the Morris water maze (Bakker et al., 1994), abnormalities in dendritic spines (Comery et al., 1997; Nimchinski et al., 2001) and altered synaptic plasticity (see below). Fmr1 KO mice have also an increased susceptibility to audiogenic seizures (AGS) (Musumeci et al., 2000), which is specifically reverted by the introduction of constructs codifying the human FMR1 gene (Musumeci et al., 2007). In addition, Fmr1 KO mice is currently considered one of the leading animal models of autism (Bernardet & Crusio, 2006).

To study the function of FXR2P and FXR1P and their possible implication in FXS, Fxr1 and Fxr2 KO mouse models have been generated. Homozygous Fxr1 KO neonates die shortly after birth for cardiac or respiratory failure; whereas a mouse model expressing very low levels of FXR1P displays a strongly reduced limb musculature and has a reduced life span, suggesting a role for FXR1P in muscle mRNA transport/translation control similar to that seen for FMRP in neuronal cells (Mientjes et al., 2004).

Fxr2 KO mice do not show gross abnormalities in brain or testis, but are hyperactive in the open-field test, have reduced levels of prepulse inhibition, display less contextual

conditioned fear and are less sensitive to a heat stimulus. Interestingly, Fxr2 KO mice present some behavioural phenotypes similar to those observed in Fmr1 KO mice (Bontekoe et al., 2002).

A double Fmr1/Fxr2 KO mouse has also been created. These mice have exaggerated behavioural phenotypes in open-field activity, prepulse inhibition of acoustic startle response and contextual fear conditioning when compared with Fmr1 KO mice, Fxr2 KO mice or WT (Spencer et al., 2006). This is in line with the hypothesis that Fmr1 and Fxr2 play a similar role in pathways controlling locomotor activity, sensorimotor gating and cognitive processes. In addition, Fmr1/Fxr2 double KO mice exhibit more severe electrophysiological alterations than either single KO model, which suggests that FMRP and FXR2P regulate synaptic plasticity both together and separately (J. Zhang et al., 2009).

2.6 Role of FMRP in the formation of neuronal network

Although FXS has traditionally been thought of as a disorder of the postsynaptic compartment, several evidences suggest a potential axonal or pre-synaptic role for FMRP. The first evidence that suggests a pre-synaptic role for FMRP was the observation that FMRP is present in growth cones of developing axons and distal segments of mature axons in hippocampal cell cultures (Antar et al., 2006). More recently, FMRP (but also FXR1P and FXR2P) have been detected in pre-synaptic terminals in discrete small structures defined as granules (Fragile X granules) by light and electron microscopy in brain slices (Christie et al., 2009). The expression of such pre-synaptic FMRP granules is regulated both developmentally and regionally in the brain, being maximal in the frontal cortex and hippocampal area CA3 in two-weeks-old mice but virtually non-existent in adult neocortex or in CA1 (Christie et al., 2009). A second line of evidence comes from studies in *Drosophila*, where mutations in the Fmr1 gene result in axonal defects. It has been demonstrated that in *Drosophila* loss of dFMRP causes defects in axonal targeting and arborization (Y.Q. Zhang et al., 2001), misregulated pre-synaptic structure (Michel et al., 2004), neuromuscular junction (NMJ) synapse overelaboration (overgrowth, overbranching, excess synaptic boutons), and altered neurotransmission (Gatto & Broadie, 2008). Two recent papers in *Drosophila* highlights the role of FMRP in activity-dependent axon pruning and in regulation of synaptic structure during development (Gatto & Broadie, 2008; Tessier & Broadie 2008). Using the *Drosophila* model these authors addressed the question whether FXS is mainly a disease of development, characterized by structural defects, or a disease of plasticity, or both. The establishment of neural circuits proceeds via a two-stages process: an early, activity-independent wiring to produce a rough map characterized by excessive synaptic connections and subsequent, use-dependent pruning to eliminate inappropriate connections and reinforce maintained synapses. dFMRP expression and function are maximal during late-stage periods of axon pruning, which requires both dFMRP and sensory input activity. dFMRP has a primary role in activity-dependent neural circuit refinement during late brain development (Tessier & Broadie, 2008). Gatto and Broadie (2008) observed that constitutive neuronal dFMRP expression rescues all NMJ synaptic structural defects, demonstrating a strictly pre-synaptic dFMRP requirement. By contrast, targeted pre-synaptic dFMRP expression does not rescue neurotransmission function in the null mutant, indicating a separable post-synaptic dFMRP requirement. Temporally, transient early-development expression of dFMRP strongly rescues synaptic architecture, demonstrating primarily an early role for dFMRP in establishing synapse morphology.

Interestingly, acute dFMRP expression at maturity weakly rescues synaptic structure defects, showing that late-stage intervention might only partially compensate for structural abnormalities established early during development. Thus, FMRP may play a double crucial role by regulating the structure of neural circuits during development and by regulating synaptic plasticity during maturity.

Recent data in the mouse model also suggest that FMRP might be involved in the establishment of neuronal connectivity, possibly through mechanisms which involve guidance and stabilization of axons during development (Bureau, 2009). Bureau et al. (2008) investigated the development of excitatory projections in the barrel cortex of *Fmr1* KO mice and they observed that projections are altered both functionally and morphologically, suggesting an important role for FMRP in this process. Dysregulated neuronal connectivity in the barrel cortex causes defective glutamatergic synapse maturation, delayed and aberrant formation of sensory maps, and altered synaptic plasticity during the critical period (Harlow et al., 2010). In general, the absence of FMRP could lead to altered network synchrony and hyperexcitable neuronal networks (Chuang et al., 2005; Gibson et al., 2008).

These data have a very strong implication for the therapeutic approach to FXS, but also to other developmental disorders characterized by altered neuronal connectivity. Interestingly, in a list of newly identified FMRP mRNA targets several transcripts encode for pre-synaptic proteins and are implicated in autism spectrum disorders (Darnell et al., 2011). It will be important in the future to establish whether a therapeutic intervention is able to rescue these early established abnormalities in neuronal circuitry.

3. Therapeutic strategies in FXS

Current therapeutic approach to patients with FXS is aimed at correcting symptoms or behavioural deficits, including hyperactivity and anxiety. Medications include stimulants, antipsychotics, anti-depressants and anticonvulsants. Patients with FXS also seem to benefit from behavioural intervention and special educational programs. As demonstrated in the FXS mouse model, an enriched environment can improve behaviour, and thus this therapy might also be beneficial for patients (Restivo et al., 2005).

In the last few years the amount of scientific publications in the field of neurobiology of FXS have exponentially increased and these efforts have led to important discoveries which are now partially translated in therapeutic perspectives. These include the use of drugs to correct the abnormal activity of the mGlu receptor- and GABA-pathways. In addition, novel therapeutic targets will be discussed based on other pathways, which have been found to be dysregulated in mouse models of FXS.

3.1 mGlu5 receptor: A key protein for synaptic plasticity

Glutamate, the major excitatory neurotransmitter in the mammalian CNS, exerts its action by interacting with ionotropic (iGlu) and mGlu receptors. iGlu receptors are multimeric ion channels responsible for fast synaptic transmission and are subdivided into three distinct subtypes: AMPA (α -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid), kainate, and NMDA (N-methyl-D-aspartate) receptors. mGlu receptors are members of a G-protein-coupled receptor superfamily that includes GABA_B, Ca²⁺ sensing, some taste and pheromone receptors (Bockaert & Pin, 1999).

The family of mGlu receptors comprises eight subtypes (mGlu1-mGlu8) that are divided into three distinct groups on the basis of sequence similarities and different pharmacological responses. Group-I includes mGlu1 and mGlu5 receptor subtypes which are coupled to G_q/G_{11} proteins and whose activation stimulates polyphosphoinositide (PI) hydrolysis and an increase in intracellular Ca^{2+} release as a result of a PKC-mediated receptor phosphorylation (Kawabata et al., 1996). Activation of group-I mGlu receptors also stimulates the ERK1/2 MAP kinase and the phosphatidylinositol-3-kinase (PI3K) pathways, which are involved in cell proliferation, differentiation, and survival, as well as in processes of activity-dependent synaptic plasticity (Ferraguti et al., 1999; Peavy & Conn, 1998; Rong et al., 2003). Activation of ERK in striatum and PI3K in hippocampus (Mao et al., 2005; Rong et al., 2003) requires the interaction of group-I mGlu receptors with Homer proteins, a class of scaffolding proteins cross-linking group-I mGlu receptors (mGlu1 and mGlu5) to inositol triphosphate (IP_3) receptors and to other proteins of the post synaptic density such as SHANK (Tu et al., 1998, 1999). Homer proteins also control several functions of group-I mGlu receptors such as constitutive activity (Ango et al., 2001), cell surface expression and trafficking (Ango et al., 2002; Coutinho et al., 2001), lateral mobility (Sergé et al., 2002) and coupling to ion channels of the cytoplasmic membrane (Kammermeier et al., 2000). Group-II and group-III include mGlu2/3 and mGlu4,6,7,8, respectively and are coupled to G_i/G_o proteins. While mGlu1 and mGlu5 receptors are generally found in postsynaptic densities and modulate postsynaptic efficacy, mGlu2, -3, -4, -7, and -8 receptors are mainly (but not exclusively) pre-synaptic and regulate neurotransmitter release (Luján et al., 1997; Schoepp, 2001). The pharmacology of mGlu receptors has expanded in the last years and ligands for mGlu receptors are now considered the most promising drugs in the treatment of neurological and psychiatric disorders (reviewed by Nicoletti et al., 2011). Here we will focus on group-I mGlu receptors, namely mGlu5, for their implication in the pathophysiology of FXS.

mGlu1 and mGlu5 receptors have a different temporal and regional expression pattern. While the transcript of mGlu1 receptors is low at birth and progressively increases during postnatal development, the transcript of mGlu5 receptors is highly expressed early after birth and progressively decreases afterwards (Catania et al., 1994). Expression of mGlu5 receptors is high and widespread in the first two weeks of postnatal life (reviewed in Catania et al., 2007) when the PI response to group-I mGlu receptor agonists in brain slices is substantial (Nicoletti et al., 1986a, 1986b). A much lower receptor response is detected in hippocampal, cortical or striatal slices of adult rats, where only agonists endowed with high intrinsic efficacy can stimulate PI hydrolysis (Nicoletti et al., 1986a, 1986b). The mGlu1a receptor protein is highly expressed in discrete regions of the adult brain including the cerebellum, olfactory bulbs, thalamus and pars compacta of the substantia nigra, and is barely detectable during early development (Lopez-Bendito et al., 2002). These expression studies suggest that mGlu5 receptors may have an important role in plastic changes occurring early during post-natal development (Catania et al., 2007).

Most of group-I mGlu receptors are located in dendritic spines (Baude et al., 1993), in an annulus that circumscribes the PSD, but some (probably mGlu5) are also distributed on glutamatergic nerve terminals (Cochilla & Alford, 1998; Gereau & Conn, 1995; Rodriguez-Moreno et al., 1998; Sistiaga et al., 1998). mGlu5 receptors are also expressed in non-neuronal cells, including astrocytes, oligodendrocytes and microglia, stem progenitor cells, and a variety of peripheral cells (Nicoletti et al., 2011).

mGlu5 receptors are involved in the regulation of synaptic plasticity, including the induction of LTP (important for retaining nascent synapses) and LTD (important for activity-guided synapse elimination), two electrophysiological substrates that, working in concert, contribute to learning and memory storage throughout postnatal life (Bear, 1998). LTP is a long term increase in synaptic efficacy and is associated with the strengthening of the connection between a pre-synaptic and post-synaptic neuron, whereas LTD is defined as the weakening of the synapse, and is mainly reflected by a reduced number of iGlu responsive AMPA receptors at the post-synaptic membrane (Collingridge et al., 2010). Activation of mGlu5 receptors is involved in both LTP and LTD. Mice lacking mGlu5 receptors show impaired learning and reduced LTP in the hippocampal CA1 region (Lu et al., 1997).

There are two forms of LTD: one is dependent on activation of post-synaptic NMDA receptors, the other requires activation of post-synaptic group-I mGlu receptors (Oliet et al., 1997) and also can be readily induced by the selective group-I mGlu receptors agonist DHPG (Huber et al., 2001; Palmer et al., 1997). Both types of LTD determine a decrease in the number of post-synaptic AMPA receptors by distinct mechanisms (Bear et al., 2004). One important distinction is that LTD triggered by mGlu receptor activation (mGlu-LTD), but not NMDA-receptor-dependent LTD, requires the activation of mGlu5 receptors and the rapid translation of pre-existing mRNAs in the post-synaptic dendrites through a mechanism that involves ERK phosphorylation (Gallagher et al., 2004).

3.2 mGlu5 receptor: A pharmacological target in FXS

The first indication for a link between mGlu receptors and FXS was the evidence that activation of group-I mGlu receptors in rat and mouse brain synaptoneurosomes stimulates the rapid translation of pre-existing mRNAs, including the FMRP mRNAs (Weiler et al., 1997, 2004). Since, a growing number of studies was carried out to support a role of group-I mGlu receptors in the pathophysiology of FXS. In particular, the finding that mGlu5-/protein synthesis-dependent forms of synaptic plasticity, namely mGlu5-dependent LTD, are increased in the mouse model of FXS led Bear and collaborators to formulate the “mGlu theory” of FXS, which postulates that in the absence of FMRP, which acts reducing the mGlu5-activated mRNA translation at synapse, levels of FMRP-regulated proteins are increased and, as a consequence, can be reduced by mGlu5 pharmacological antagonism (Bear et al., 2004). Other forms of synaptic plasticity, including the more classical NMDA-receptor dependent LTD, show no abnormalities in the hippocampus of *Fmr1* KO mice. Another important step towards the understanding of FXS physiopathology was represented by the finding that, while in WT mice mGlu5-dependent LTD is blocked by inhibitors of protein synthesis, this is not the case in *Fmr1* KO mice, suggesting that in the absence of FMRP LTD proteins are constitutively and highly expressed before LTD induction (Waung & Huber, 2009).

Thus, the absence of FMRP causes an abnormal expression of dendritic proteins leading to the amplification of mGlu-mediated long-term responses. The identification of these proteins, which may be critical for the pathophysiology of synaptic dysfunction in FXS, is crucial. Some proteins encoded by FMRP target mRNAs may play a role. For example, Map1b interacts with the GluR2 interacting protein and scaffold GRIP1 (Davidkova & Carroll, 2007; Seog, 2004). Other proteins which are rapidly synthesized after mGlu5 receptor activation and that are basally elevated in *Fmr1* KO mice include CaMKII (Zalfa et

al., 2003), amyloid precursor protein (APP), Arc/Arg3.1 (Park et al., 2008; Zalfa et al., 2003) which are all involved in mechanisms underlying synaptic plasticity. The list of FMRP mRNAs targets has recently grown with the discovery of 842 mRNA by using the high stringent CLIP method (Darnell et al., 2011). Further studies examining the expression levels of the encoded proteins in FXS and their regulation by mGlu receptors may corroborate the link between mGlu5 activation and protein synthesis of FMRP target mRNAs and its role in synaptic plasticity under physiological and pathological conditions. In addition, as a direct or indirect consequence of altered protein synthesis at synapses, several mGlu-mediated signalling pathways might be dysregulated. Interestingly, mGlu5 receptors in *Fmr1* KO mice are less tightly associated to Homer proteins (Giuffrida et al., 2005), which suggests either an increase of mGlu5 constitutive activity or an altered coupling of mGlu5 receptors with downstream signalling pathways. Accordingly, Ronesi and Huber (2008) reported that induction of PI3K-Akt-mTOR signalling by mGlu5 is impaired in *Fmr1* KO mice and, differently than in WT mice, mGlu5 dependent LTD is insensitive to disruption of mGlu5/Homer interaction. Further studies are needed to understand how the lack of FMRP affects mGlu5 mediated responses in FXS.

Several pharmacological studies have supported the “mGlu theory”, by demonstrating that phenotypic features of FXS can be corrected with the use of antagonists of mGlu5 such as 2-methyl-6-(phenylethynyl)-pyridine (MPEP) and fenobam. MPEP is a systemically active negative allosteric modulator of mGlu5 receptors and can also inhibit constitutive activity of mGlu5 acting as an inverse agonist (Yan et al., 2005). Fenobam, which had previously been investigated as an anxiolytic, has been identified as a highly potent, selective negative modulator of mGlu5 receptor (Porter et al., 2005). In particular, MPEP blocks audiogenic seizure susceptibility of *Fmr1* KO mice (Chuang et al., 2005) and both MPEP and fenobam restore dendritic spine morphology in hippocampal cell cultures from *Fmr1* KO mice (De Vrij et al., 2008).

A more direct evidence that the “mGluR theory” might be corrected has been provided using genetic interaction experiments (Dölen et al., 2007). In this study, *Fmr1* KO mice were crossed with heterozygous mGlu5 receptor KO mice generating double mutants of *Fmr1* and *Grm5* (the gene that encodes mGlu5 receptor) and multiple phenotypes relevant to the pathogenesis of FXS were examined. Reduction of mGlu5 expression by 50% in the *Fmr1* KO/*Grm5* heterozygote cross rescued altered ocular dominance plasticity, increased density of dendritic spines, increased basal protein synthesis, exaggeration of avoidance extinction and audiogenic seizure susceptibility, but not macroorchidism (Dölen et al., 2007). Moreover, no change in protein synthesis was detected in *Grm5* heterozygote, suggesting that a therapeutic dose of an mGlu5 receptor antagonist for FXS patients should not have negative side effects in unaffected individuals. These pre-clinical studies support the therapeutic utility in FXS patients. Interestingly, the potential use of mGlu5 antagonists is not restricted to FXS but is considered for a variety of human conditions including anxiety, convulsions, pain, depression, Parkinson's disease and gastroesophageal reflux disease (see Nicoletti et al., 2011).

An initial small pilot open label, single dose trial with fenobam in adults with FXS did not reveal any adverse effect and produced promising results showing an improvement of prepulse inhibition (Berry-Kravis et al., 2009). More recently, the Novartis compound AFQ056 has been used in a randomized, double-blind study in 30 male FXS patients aged 18-35 years. Although an initial assessment did not show any improvement after treatment,

when patients were divided into two groups on the basis of a full or partial methylation of the FMR1 promoter a significant improvement on stereotypic behaviour, hyperactivity and inappropriate speech were detected only in the full methylation group (Jacquemont et al., 2011). While this work confirms the clinical efficacy of mGlu5 pharmacological blockade in FXS, there is no clear explanation for the lack of improvement in patients with partial methylated FMR1 gene. More clinical studies in a higher number of patients are needed.

3.3 GABA system as target of viable pharmacological treatments in FXS

In addition to the mGlu receptors, several evidences suggest that gamma-aminobutyric acid (GABA) signalling is another molecular pathway involved in FXS. Expression and functional studies suggest that defects in GABA transmission might be region specific and might involve different components of the GABAergic system in different brain regions.

GABA is the major inhibitory neurotransmitter in the CNS and plays a key role in modulating neuronal activity, by maintaining the inhibitory tone and the physiological balance between inhibition and excitation at synapses. GABA mediates its action via two distinct receptor systems, the ionotropic GABA_A and metabotropic GABA_B receptors.

Ionotropic GABA_A receptors are heteropentameric complexes, formed by the assembly of various classes of at least 19 different subunits (α 1–6, β 1–3, γ 1–3, δ , ϵ , θ , π and ρ 1–3) (Simon et al., 2004) associated with channels permeable to Cl⁻ ions. In brain, a high diversity of GABA_A receptor subtypes having a spatio-temporal specific distribution in different regions has been found (Barnard et al., 1998; Kneussel, 2002; Korpi et al., 2002). The subunit combination confers highly different pharmacological and physiological properties to GABA_A receptors (Fritschy et al., 1995).

GABA_B receptors are heterodimeric G protein-linked receptors constituted by two different subunits. They have a pre- and post-synaptic distribution; at pre-synaptic level they can inhibit the release of neurotransmitters through a decrease of Ca²⁺ entry, whereas, at post-synaptic level they reduce neuronal excitability through an increase of K⁺ conductance. In general, they mediate a slower and more prolonged inhibitory signal than GABA_A receptors (Bormann, 2000; Chebib et al., 1999). Interestingly, GABA_B receptors agonists inhibit pre-synaptic glutamate release and consequently the post-synaptic glutamate responses (reviewed in Chalifoux & Carter, 2011).

An important indication that the GABAergic system might be involved in FXS was the evidence, obtained using the Antibody Positioned RNA Amplification (APRA) technique, that the mRNA of the δ subunit of the GABA_A receptor is directly bound to FMRP (Miyashiro et al., 2003). In Fmr1 KO mouse changes in levels of expression of both GABA_A and GABA_B receptors have been found by different authors. Several studies have revealed in different brain regions, all playing an important role in cognitive functions (behaviour, learning, memory and anxiety), as cortex, hippocampus, diencephalon and brainstem an under expression of many distinct GABA_A receptor subunits (α 1, α 3, α 4, α 5 β 1 and β 2 and γ 1 and γ 2 and δ) at the mRNA (Curia et al., 2009; D'Hulst et al., 2006; Gantois et al., 2006) and protein level (Adusei et al., 2010; El Idrissi et al., 2005).

Altered GABA transmission has been reported in different brain regions. An alteration of both GABAergic and cholinergic system, with a lower inhibitory effect mediate by GABA_A receptor in subiculum neurons has been detected by electrophysiology in brain slices of

Fmr1 KO mice (D'Antuono et al., 2003). More recently, other electrophysiological findings in subiculum have shown that tonic GABA_A currents were down regulated in Fmr1 KO mice, whereas no significant differences were observed in phasic currents (Curia et al., 2009). An increased GABA transmission has been found in the striatum (Centonze et al., 2008), whereas a robust reduction in the inhibitory transmission has been revealed in the amygdala, which results in hyper-excitability of principal neurons and is likely due to pre-synaptic defects such as decreases in GABA production and release (Olmos-Serrano et al., 2010). Accordingly, a reduction of GABA has been detected in Fmr1 KO mice using a metabolomic approach (Davidovic et al., in press).

Furthermore, cytoarchitectonic and morphological studies from somatosensory cortex highlighted a significant reorganization of neocortical inhibitory circuits of GABAergic interneurons in the Fmr1 KO mouse. In fact, this animal model showed a marked reduction of parvalbumin-positive neurons compared to the WT mice, whereas no difference was observed for calbindin- and calretinin-positive neurons (Selby et al., 2007).

Thus, most expression and functional data suggest that increasing GABAergic transmission might result in a beneficial effect, at least in certain regions. Accordingly, experiments from Fmr1 mutant *Drosophila* have shown that GABA treatment during development using GABA, nipecotic acid (a known GABA reuptake inhibitor) and creatinine (a potential activator of GABA_A receptor) rescued the lethality induced by glutamate toxicity of dFmr1 mutant flies, when they were reared on food containing increased levels of glutamate (Chang et al., 2008) and rescued many Fmr1 mutant phenotypes, such as *Futsch* overexpression, defects in mushroom bodies structure and altered male courtship behaviour (Chang et al., 2008). In addition, treatment of Fmr1 KO mice with the GABA_A receptor agonist taurine is reported to increase acquisition of a passive-avoidance task (El Idrissi et al., 2009). More recently, a treatment with the systemically active agonist acting at δ subunit-containing GABA_A receptors, 4,5,6,7-Tetrahydroisoxazolo[5,4-c]pyridin-3-ol hydrochloride (THIP hydrochloride), that is able to determine an augmentation of tonic inhibitory tone (Glykys & Mody, 2007), was shown to rescue neuronal hyperexcitability recorded from principal neurons of BL nucleus of amygdala in Fmr1 KO mice (Olmos-Serrano et al., 2010).

The involvement of GABA_B receptors is also under investigation in FXS. In fact, it has been observed a reduced expression of the GABA_B R1 subunits in the forebrain of Fmr1 KO mice, early during the development and in adulthood; whereas no significant differences have been observed in GABA_B R2 expression (Adusei et al., 2010; Pacey et al., 2011). Reduced functioning of GABA_B receptors might explain the increased susceptibility of Fmr1 KO mice to audiogenic seizures (Musumeci et al., 2000). Accordingly, stimulation of GABA_B receptors with agonist Baclofen reduces the rate of audiogenic seizures in Fmr1 KO mice (Pacey et al., 2009). These receptors play also a role in the pathophysiology of anxiety and depression, so GABA_B receptor agonist treatment might be used for reducing anxiety symptoms in patients with FXS (Cryan & Kaupmann, 2005).

3.4 Protein dysregulation and other biochemical pathways as potential targets of intervention

As soon as the list of validated FMRP-targeted mRNAs will grow, more pathways will be shown to be affected and more drugs will be proposed for the future therapy of FXS. In the next paragraph we will discuss recent advances concerning relevant pathways which may lead to treatment.

3.4.1 Oxidative stress and fragile X syndrome

Several evidences suggest a role of oxidative stress in FXS. FXS patients display an increase in adrenocortical activity and an altered hypothalamic–pituitary–adrenal (HPA) axis (Hessl et al., 2004); adrenal hormones have been involved in the induction of brain oxidative stress resulting in oxidation of molecules and depletion of antioxidants such as glutathione (Herman & Cullinan, 1997). In *Fmr1* null flies changes in the expression of proteins involved in redox reactions have been observed, suggesting a possible alteration in the oxidative balance (Y.Q. Zhang et al., 2005). In the brain of *Fmr1* KO mice higher levels of reactive oxygen species, nicotinamide adenine dinucleotide phosphate (NADPH)-oxidase activation, lipid and protein oxidation have been found, suggesting that a moderate increase in the oxidative stress in the brain may play a role in the pathophysiology of FXS (el Bekay et al., 2007). In addition, microarray identification has revealed altered mRNA translational profiles in the absence of FMRP, involving proteins which participate in homeostasis of the antioxidant status such as glutathione transferase and SOD1 (M.R. Brown et al., 2001; Miyashiro et al., 2003). Recently, a reduction of protein levels of SOD1 has been found in *Fmr1* null cells and brain (Bechara et al., 2009), suggesting that in the absence of FMRP the increase in brain oxidative stress might be due to the altered SOD1 expression. A comprehensive profiling of the metabolome of the *Fmr1*-deficient brain has revealed an increase in lipid-oxidized species at early age (Davidovic et al., in press), further corroborating the hypothesis that oxidative stress is indeed involved in FXS pathophysiology.

The therapeutic implication of these findings is that anti-oxidant agents may be useful in the treatment of FXS and are supported by recent results obtained in *Fmr1* KO mice after treatment with alpha-tocopherol and melatonin (de Diego-Otero et al., 2009; Romero-Zerbo et al., 2009). Chronic pharmacological treatment with alpha-tocopherol reverses pathophysiological hallmarks including free radical overproduction, oxidative stress, macroorchidism, and also behaviour and learning deficits (de Diego-Otero et al., 2009). Chronic administration of melatonin protects the *Fmr1* KO mouse from the oxidative stress in brain and testis, reverses several behavioural and learning deficits, normalizes several abnormalities observed in the *Fmr1* KO mouse, including biochemical hallmarks, such as free radical production in macrophage cells and brain slices, as well as carbonyl content in proteins and lipid peroxidation (Romero-Zerbo et al., 2009). Additionally, it also normalizes reduced glutathione levels in the brain and testis of *Fmr1* KO mice. The treatment controls corticosterone plasma levels, locomotion (hyperactivity), anxiety responses and fear learning deficits.

3.4.2 Matrix metallo-proteinase 9 and minocycline

Another example of protein dysregulated in the mouse model of FXS and considered a valuable target of a pharmacological treatment is the matrix metallo-proteinase 9 (MMP-9). MMP-9 is an extracellular endopeptidase that cleaves extracellular matrix proteins that impact synaptogenesis and spine morphology (Ethell & Ethell, 2007). MMP-9 could affect dendritic spine morphology by cleaving components of the extracellular matrix and/or cell surface proteins that participate in synaptogenesis and dendritic spine maturation (Ethell & Ethell, 2007). It has been shown that MMP-9s are elevated in the hippocampus of *Fmr1* KO mice and may be partially responsible for the immature dendritic spine profile of

hippocampal neurons and for synaptic instability (Bilousova et al., 2006). A treatment with minocycline, a tetracycline analogue that can inhibit matrix MMP-9 and reduce inflammation in the CNS, promotes the formation of mature dendritic spines and reduces dendritic spine abnormalities respectively in WT and Fmr1 KO hippocampal neurons. Indeed, it has been shown that excessive MMP-9 activity disrupts mature dendritic spines in hippocampal neurons. The beneficial effects of this drug on dendritic spine morphology are also accompanied by changes in the behavioural performance of 3-weeks-old Fmr1 KO mice (Bilousova et al., 2009).

Clinical trials have been started for patients with FXS and an open-label trial has been recently completed to study the effects of minocycline in patients with FXS (Utari et al., 2010). The results show that minocycline provides significant functional benefits to FXS patients, it is well-tolerated, and both adolescents and adults with FXS can benefit from minocycline treatment.

3.4.3 Phosphoinositide 3-kinase and FXS

It has been hypothesized that FMRP controls protein synthesis-dependent regulation of synaptic morphology and function through regulation of PI3K signalling. PI3K regulates different pathways. Deficiency of FMRP results in excess activity of PI3K; loss of FMRP leads to excess mRNA translation and synaptic protein expression of p110beta, a catalytic subunit of PI3K and a putative FMRP-target mRNA (Miyashiro et al., 2003). FMRP regulates the synthesis and synaptic localization of p110beta. In WT, mGlu receptor activation induces p110beta translation, p110beta protein expression, and PI3K activity; in contrast, both p110beta protein synthesis and PI3K activity are elevated and insensitive to mGlu receptor stimulation in Fmr1 KO mice. Excess of PI3K activity in the absence of FMRP can occur independently of mGlu receptors (Gross et al., 2010). PI3K is a downstream signalling molecule of many cell surface receptors; aberrant regulation of p110beta could provide a molecular explanation for dysregulation of D1 dopamine receptors (Wang et al., 2008), of Gq-proteins (Volk et al., 2007), and of Ras (Hu et al., 2008) observed in Fmr1 KO mice. Dysregulated PI3K signalling may also underlie the synaptic impairments in FXS. In support of this hypothesis, it has been observed that a treatment with LY294002 (PI3K antagonist) in Fmr1 KO neurons can rescue the enhanced AMPA receptor internalization and the increased spine density (Gross et al., 2010). Targeting excessive PI3K activity might thus be another therapeutic strategy for FXS.

4. Conclusion and future direction

A deeper understanding of the function of FMRP and the molecular mechanisms underlying FXS using animal models has recently led to propose new therapeutic approaches, which will prove to be corrected in the next future as soon as several ongoing clinical trials will be completed. As a consequence of altered protein expression both at pre- and post-synaptic levels it is possible that several interconnected biochemical pathways are altered in FXS. It will be important to identify these cascades. System biology approaches and bioinformatic tools may help to identify the metabolic consequences of dysregulated biochemical cascades in FXS and in other neurological disorders associated with intellectual disability and autism. Given the high number of proteins and pathways which are likely to be dysregulated in FXS

it will be also very important to establish which of them are involved in determining structural changes during development and which are more involved in plasticity defects.

It is possible that different therapeutic interventions might be used during development and in adult patients.

5. Acknowledgements

Authors are funded by Telethon (GGP07264, Italy), Ministry of Health (Italy), PRIN (Italy), Foundation Jérôme Lejeune (France), IRCCS Oasi Maria SS.Troina. (Italy).

6. References

- Adusei, D.C., Pacey, L.K., Chen, D. & Hampson, D.R. (2010). Early developmental alterations in GABAergic protein expression in fragile X knockout mice. *Neuropharmacology*, Vol.59, No.3, (September 2010), pp. 167–171, ISSN 0028-3908
- Ango, F., Prézeau, L., Muller, T., Tu, J.C., Xiao, B., Worley, P.F., Pin, J.P., Bockaert, J. & Fagni, L. (2001). Agonist-independent activation of metabotropic glutamate receptors by the intracellular protein Homer. *Nature*, Vol. 411, No.6840, (June 2001), pp. 962–965, ISSN 0028-0836
- Ango, F., Robbe, D., Tu, J.C., Xiao, B., Worley, P.F., Pin, J.P., Bockaert, J. & Fagni, L. (2002). Homer-dependent cell surface expression of metabotropic glutamate receptor type 5 in neurons. *Molecular and Cellular Neuroscience*, Vol. 20, No.2, (June 2002), pp. 323–329, ISSN 1044-7431
- Antar, L.N., Afroz, R., Dictenberg, J.B., Carroll, R.C. & Bassell G.J. (2004). Metabotropic glutamate receptor activation regulates fragile x mental retardation protein and FMR1 mRNA localization differentially in dendrites and at synapses, *Journal of Neuroscience*, Vol.24, No.11, (March 2004), pp. 2648-2655, ISSN 1529-2401
- Antar, L.N., Li, C., Zhang, H., Carroll, R.C. & Bassell, G.J. (2006). Local functions for FMRP in axon growth cone motility and activity-dependent regulation of filopodia and spine synapses. *Molecular and Cellular Neuroscience*, Vol.32, No.1-2, (May-June 2006), pp. 37–48, ISSN 1044-7431
- Aschrafi, A., Cunningham, B.A., Edelman, G.M. & Vanderklish, P.W. (2005). The fragile X mental retardation protein and group I metabotropic glutamate receptors regulate levels of mRNA granules in brain. *Proceedings of the National Academy of Sciences of the United States of America*, Vol.102, No.6, (February 2005), pp. 2180-5, ISSN 0027-8424
- Bakker, C. E., Verheij, C., Willemsen, R., van der Helm, R., Oerlemans, F., Vermey, M., Bygrave, A., Hoogeveen, A.T. & Oostra, B.A. (1994). Fmr1 knockout mice: a model to study fragile X mental retardation. *Cell*, Vol.78, No.1, (July 1994), pp. 23-33, ISSN 0092-8674
- Bardoni, B., Davidovic, L., Bensaid, M. & Khandjian, E.W. (2006). The fragile X syndrome: exploring its molecular basis and seeking a treatment. *Expert Reviews in Molecular Medicine*, Vol.8, No.8, (April 2006), pp. 1–16, ISSN 1462-3994
- Barnard, E.A., Skolnick, P., Olsen, R.W., Mohler, H., Sieghart, W., Biggio, G., Braestrup, C., Bateson, A.N. & Langer, S.Z. (1998). International Union of Pharmacology. XV. Subtypes of g-aminobutyric acidA receptors: classification on the basis of subunit

- structure and receptor function. *Pharmacological reviews*, Vol.50, No.2, (June 1998) pp. 291– 313, ISSN 0031-6997
- Bassell, G.J. & Warren, S.T. (2008). Fragile X syndrome: loss of local mRNA regulation alters synaptic development and function. *Neuron*, Vol.60, No.2, (October 2008), pp. 201–214, ISSN 0896-6273
- Baude, A., Nusser, Z., Roberts, J.D., Mulvihill, E., McIlhinney, R.A. & Somogyi, P.(1993). The metabotropic glutamate receptor (mGluR1 alpha) is concentrated at perisynaptic membrane of neuronal subpopulations as detected by immunogold reaction. *Neuron*, Vol.11, No.4, (October 1993), pp. 771–787, ISSN 0896-6273
- Bear, M.F. (1998). *The role of LTD and LTP in development and learning*. In *Mechanistic Relationships between Development and Learning* (Carew TJ, Menzel R, Shatz CJ, eds), Wiley, New York, pp. 205–225
- Bear, M.F., Huber, K.M. & Warren, S.T. (2004). The mGluR theory of fragile X mental retardation. *Trends in Neurosciences*, Vol.27, No.7, (July 2004), pp. 370–377, ISSN 0166-2236
- Bechara, E. G., Didiot, M. C., Melko, M., Davidovic, L., Bensaid, M., Martin, P., Castets, M., Pognonec, P., Khandjian, E.W., Moine, H. & Bardoni, B. (2009). A novel function for fragile X mental retardation protein in translational activation. *PLoS Biology*, Vol.7, No.1, (January 2009), pp. 133–145, ISSN 1544-9173
- Bernardet, M. & Crusio, W.E. (2006). Fmr1 KO mice as a possible model of autistic features. *The Scientific World Journal*, Vol.6, (September 2006), pp. 1164–76, ISSN 1537-744X
- Berry-Kravis, E., Abrams, L., Coffey, S.M., Hall, D.A., Greco, C., Gane, L.W., Grigsby, J., Bourgeois, J.A., Finucane, B., Jacquemont, S., Brunberg, J.A., Zhang, L., Lin, J., Tassone, F., Hagerman, P.J., Hagerman, R.J. & Leehey, M.A. (2007). Fragile X associated tremor/ataxia syndrome: clinical features, genetics, and testing guidelines. *Movement Disorders*, Vol.22, No.14, (October 2007), pp. 2018–30, ISSN 0885-3185
- Berry-Kravis, E., Hessel, D., Coffey, S., Hervey, C., Schneider, A., Yuhas, J., Hutchison, J., Snape, M., Tranfaglia, M., Nguyen, D.V. & Hagerman, R. (2009). A pilot open label, single dose trial of fenobam in adults with fragile X syndrome. *Journal of Medical Genetics*, Vol.46, No.4, (April 2009), pp. 266–271, ISSN 00222593
- Bilousova, T., Rusakov, D.A., Ethell, D.W. & Ethell, I.M. (2006). Matrix metalloproteinase-7 disrupts dendritic spines in hippocampal neurons through NMDA receptor activation. *Journal of Neurochemistry*, Vol.97, No.1, (April 2006), pp. 44–56, ISSN 0022-3042
- Bilousova, T.V., Dansie, L., Ngo, M., Aye, J., Charles, J.R., Ethell, D.W. & Ethell, I.M. (2009). Minocycline promotes dendritic spine maturation and improves behavioural performance in the fragile X mouse model. *Journal of Medical Genetics*, Vol.46, No.2, (February 2009), pp. 94–102, ISSN 00222593
- Bockaert, J. & Pin, J.P. (1999). Molecular tinkering of G protein-coupled receptors: an evolutionary success. *The EMBO Journal*, Vol.18, No.7, (April 1999), pp. 1723–9, ISSN 0261-4189
- Bontekoe, C.J., McIlwain, K.L., Nieuwenhuizen, I.M., Yuva-Paylor, L. A., Nellis, A., Willemsen, R., Fang, Z., Kirkpatrick, L., Bakker, C.E., McAninch, R., Cheng, N.C., Merriweather, M., Hoogeveen, A.T., Nelson, D., Paylor, R. & Oostra, B.A. (2002).

- Knockout mouse model for Fxr2: a model for mental retardation. *Human Molecular Genetics*, Vol.11, No.5, (March 2002), pp. 487-498, ISSN 0964-6906
- Bormann, J. (2000). The 'ABC' of GABA receptors. *Trends in Pharmacological Sciences*, Vol.21, No.1, (January 2000), pp. 16-19, ISSN 0165-6147
- Breyse, N., Baunez, C., Spooren, W., Gasparini, F. & Amalric, M. (2002). Chronic but not acute treatment with a metabotropic glutamate 5 receptor antagonist reverses the akinetic deficits in a rat model of parkinsonism. *The Journal of Neuroscience*, Vol.22, No.13, (July 2002), pp. 5669-5678, ISSN 0270-6474
- Brodkin, J., Bradbury, M., Busse, C., Warren, N., Bristow, L. J., & Varney, M. A. (2002). Reduced stress-induced hyperthermia in mGluR5 knockout mice. *European Journal of Neuroscience*, Vol.16, No.11, (December 2002), pp. 2241-2244, ISSN 0953-816X
- Brown, M.R., Kronengold, J., Gazula, V.R., Chen, Y., Strumbos, J.G., Sigworth, F.J., Navaratnam, D. & Kaczmarek, L.K. (2010). Fragile X mental retardation protein controls gating of the sodiumactivated potassium channel Slack. *Nature Neuroscience*, Vol.13, No.7, (July 2010), pp. 819-821, ISSN 1546-1726
- Brown, V., Jin, P., Ceman, S., Darnell, J.C., O'Donnell, W.T., Tenenbaum, S.A., Jin, X., Feng, Y., Wilkinson, K.D., Keene, J.D., Darnell, R.B. & Warren, S.T. (2001). Microarray identification of FMRP-associated brain mRNAs and altered mRNA translational profiles in fragile X syndrome. *Cell*, Vol.107, No.4, (November 2001), pp. 477-487, ISSN 0092-8674
- Bureau, I., Shepherd, G.M. & Svoboda, K. (2008). Circuit and plasticity defects in the developing somatosensory cortex of FMR1 knock-out mice. *The Journal of Neuroscience*, Vol.28, No.20, (May 2008), pp. 5178-5188, ISSN 0270-6474
- Bureau, I. (2009). The development of cortical columns: role of Fragile X mental retardation protein. *The Journal of Physiology*, Vol.587, No.9, (May 2009), pp. 1897-1901, ISSN 0022-3751
- Catania, M.V., Landwehrmeyer, G.B., Testa, C.M., Standaert, D.G., Penney, J.B. Jr. & Young, A.B. (1994). Metabotropic glutamate receptors are differentially regulated during development. *Neuroscience*, Vol.61, No.3, (August 1994), pp. 481-495, ISSN 0306-4522
- Catania, M.V., D'Antoni, S., Bonaccorso, C.M., Aronica, E., Bear, M.F. & Nicoletti, F. (2007). Group I metabotropic glutamate receptors: a role in neurodevelopmental disorders? *Molecular Neurobiology*, Vol.35, No.3, (June 2007), pp. 298-307, ISSN 0893-7648
- Centonze, D., Rossi, S., Mercaldo, V., Napoli, I., Ciotti, M.T., De Chiara, V., Musella, A., Prosperetti, C., Calabresi, P., Bernardi, G. & Bagni, C. (2008). Abnormal striatal GABA transmission in the mouse model for the fragile X syndrome. *Biological Psychiatry*, Vol.63, No.10, (May 2008), pp. 963-973, ISSN 0006-3223
- Chalifoux, J.R. & Carter, A.G. (2011). GABAB receptor modulation of synaptic function. *Current Opinion in Neurobiology* Vol.21, No.2, (April 2011), pp. 339-344, ISSN 1873-6882
- Chang, S., Bray, S.M., Li, Z., Zarnescu, D.C., He, C., Jin, P. & Warren, S.T. (2008). Identification of small molecules rescuing fragile X syndrome phenotypes in *Drosophila*. *Nature Chemical Biology*, Vol.4, No.4, (April 2008), pp. 256-263, ISSN 1552-4450

- Chebib, M. & Johnston, G.A. (1999). The 'ABC' of GABA receptors: a brief review. *Clinical and Experimental Pharmacology and Physiology*, Vol.26, No.11, (November 1999), pp. 937-940, ISSN0305-1870
- Christie, S.B., Akins, M.R., Schwob, J.E. & Fallon, J.R. (2009). The FXG: a presynaptic fragile X granule expressed in a subset of developing brain circuits. *The Journal of Neuroscience*, Vol.29, No.5, (February 2009), pp. 1514-24, ISSN 1529-2401
- Chuang, S.C., Zhao, W., Bauchwitz, R., Yan, Q., Bianchi, R. & Wong, R.K. (2005). Prolonged epileptiform discharges induced by altered group I metabotropic glutamate receptor-mediated synaptic responses in hippocampal slices of a fragile X mouse model. *The Journal of Neuroscience*, Vol.25, No.35, (August 2005), pp. 8048-8055, ISSN 1529-2401
- Cochilla, A.J., & Alford, S. (1998). Metabotropic glutamate receptor-mediated control of neurotransmitter release. *Neuron*, Vol.20, No.5, (May 1998) pp. 1007-1016, ISSN 0896-6273
- Coffee, L.R. Jr., Tessier, C.R., Woodruff, E.A. 3rd & Broadie, K. (2010). Fragile X mental retardation protein has a unique, evolutionarily conserved neuronal function not shared with FXR1P or FXR2P. *Disease models & mechanisms*, Vol.3, No.7-8, (July-August 2010), pp. 471-485, ISSN 1754-8403
- Collingridge, G.L., Collingridge, G.L., Peineau, S., Howland, J.G. & Wang, Y.T. (2010). Long-term depression in the CNS. *Nature reviews. Neuroscience*, Vol.11, No.7, (July 2010), pp. 459-473, ISSN 1471-003X
- Comery, T.A., Harris, J.B., Willems, P.J., Oostra, B.A., Irwin, S.A., Weiler, I.J. & Greenough, W.T. (1997). Abnormal dendritic spines in fragile X knockout mice: maturation and pruning deficits. *Proceedings of the National Academy of Sciences of the United States of America*, Vol.94, No.10, (May 1997), pp. 5401-5404, ISSN 0027-8424
- Corbin, F., Bouillon, M., Fortin, A., Morin, S., Rousseau, F. & Khandjian, E.W. (1997). The fragile X mental retardation protein is associated with poly(A)⁺ mRNA in actively translating polyribosomes. *Human Molecular Genetics*, Vol.6, No.9, (September 1997), pp. 1465-1472, ISSN 0964-6906
- Coutinho, V., Kavanagh, I., Sugiyama, H., Tonesm, M.A. & Henleym, J.M. (2001). Characterization of a metabotropic glutamate receptor type 5-green fluorescent protein chimera (mGluR5-GFP): pharmacology, surface expression, and differential effects of Homer-1a and Homer-1c. *Molecular and Cellular Neuroscience*, Vol.18, No.3, (September 2001), pp. 296 -306, ISSN 1044-7431
- Curia, G., Papouin, T., Seguela, P. & Avoli, M. (2009). Downregulation of Tonic GABAergic Inhibition in a Mouse Model of Fragile X Syndrome. *Cerebral Cortex*, Vol.19, No.7, (July 2009), pp. 1515-1520, ISSN 1047-3211
- D'Antuono, M., Merlo, D. & Avoli, M. (2003). Involvement of cholinergic and gabaergic systems in the fragile X knockout mice. *Neuroscience*, Vol.119, No.1, (June 2003), pp. 9-13, ISSN 0306-4522
- Cryan, J.F. & Kaupmann, K. (2005). Don't worry 'B' happy!: a role for GABA(B) receptors in anxiety and depression. *Trends in Pharmacological Sciences*, Vol.26, No.1, (January 2005), pp. 36-43, ISSN0165-6147
- Darnell, J.C., Jensen, K.B., Jin, P., Brown, V., Warren, S.T. & Darnell R.B. (2001). Fragile X mental retardation protein targets G-quartet mRNAs important for neuronal function. *Cell*, Vol.107, No.4, (November 2001), pp. 489-499, ISSN 0092-8674

- Darnell, J.C., Fraser, C.E., Mostovetsky, O., Stefani, G., Jones, T.A., Eddy, S.R. & Darnell R.B. (2005). Kissing complex RNAs mediate interaction between the Fragile-X mental retardation protein KH2 domain and brain polyribosomes. *Genes and Development*, Vol.19, No.8, (April 2005), pp. 903-918, ISSN 0890-9369
- Darnell, J.C., Van Driesche, S.J., Zhang, C., Hung, K.Y., Mele, A., Fraser, C.E., Stone, E.F., Chen, C., Fak, J.J., Chi, S.W., Licatalosi, D.D., Richter, J.D. & Darnell, R.B. (2011). FMRP stalls ribosomal translocation on mRNAs linked to synaptic function and autism. *Cell*, Vol.146, No.2, (July 2011), pp. 247-261, ISSN 1097-417
- Davidkova, G. & Carroll, R.C. (2007). Characterization of the role of microtubule-associated protein 1B in metabotropic glutamate receptor-mediated endocytosis of AMPA receptors in hippocampus. *The Journal of Neuroscience*, Vol.27, No.48, (November 2007), pp. 13273-13278, ISSN 0270-6474
- Davidovic, L., Navratil, V., Bonaccorso, C.M., Catania, M.V., Bardoni, B. & Dumas, M.E. (2011). A metabolomic and systems biology perspective on the brain of the Fragile X syndrome mouse model. *Genome Research*, (September 2011), Epub ahead of print
- de Diego-Otero, Y., Severijnen, L.A., Van Cappellen, G., Schrier, M., Oostra, B.A. & Willemsen, R. (2002). Transport of fragile X mental retardation protein via granules in neurites of PC12 cells. *Molecular and Cellular Biology*, Vol.22, No.23, (December 2002), pp. 8332-8341, ISSN 0898-7750
- de Diego-Otero, Y., Romero-Zerbo, Y., el Bekay, R., Decara, J., Sanchez, L., Rodriguez-de Fonseca, F. & del Arco-Herrera, I. (2009). Alpha-tocopherol protects against oxidative stress in the fragile X knockout mouse: an experimental therapeutic approach for the Fmr1 deficiency. *Neuropsychopharmacology*, Vol.34, No.4, (March 2009), pp. 1011-1026, ISSN 0893-133X
- de Vrij, F.M., Levenga, J., van der Linde, H.C., Koekkoek, S.K., De Zeeuw, C.I., Nelson, D.L., Oostra, B.A. & Willemsen, R. (2008). Rescue of behavioral phenotype and neuronal protrusion morphology in Fmr1 KO mice. *Neurobiology of Disease*, Vol.31, No.1, (July 2008), pp. 127-132, ISSN 0969-9961
- Devys, D., Lutz, Y., Rouyer, N., Bellocq, J.P. & Mandel, J.L. (1993). The FMR-1 protein is cytoplasmic, most abundant in neurons and appears normal in carriers of a fragile X premutation. *Nature Genetics*, Vol.4, No.4, (August 1993), pp. 335-340, ISSN 1061-40
- D'Hulst, C., De Geest, N., Reeve, S.P., Van Dam, D., De Deyn, P.P., Hassan, B.A. & Kooy, R.F. (2006). Decreased expression of the GABAA receptor in fragile X syndrome. *Brain Research*, Vol.1121, No.1, (November 2006), pp. 238-245, ISSN 0006-8993
- Dictenberg, J.B., Swanger, S.A., Antar, L.N., Singer, R.H. & Bassel, G.J. (2008). A direct role for FMRP in activity-dependent dendritic mRNA transport links filopodial-spine morphogenesis to fragile X syndrome. *Developmental Cell*, Vol.14, No.6, (June 2008), pp. 926-939, ISSN 1878-1551
- Dölen, G., Osterweil, E., Rao, B.S., Smith, G.B., Auerbach, B.D., Chattarji, S. & Bear, M.F. (2007). Correction of fragile X syndrome in mice. *Neuron*, Vol.56, No.6, (December 2007), pp. 955-962, ISSN 0896-6273
- Eberhart, D.E., Malter, H.E., Feng, Y. & Warren, S.T. (1996). The fragile X mental retardation protein is a ribonucleoprotein containing both nuclear localization and nuclear export signals. *Human Molecular Genetics*, Vol.5, No.8, (August 1996), pp. 1083-1091, ISSN 0964-6906

- el Bekay, R., Romero-Zerbo, Y., Decara, J., Sanchez-Salido, L., Del Arco-Herrera, I., Rodríguez-de Fonseca, F. & de Diego-Otero, Y. (2007). Enhanced markers of oxidative stress, altered antioxidants and NADPH-oxidase activation in brains from Fragile X mental retardation 1-deficient mice, a pathological model for Fragile X syndrome. *European Journal of Neuroscience*, Vol.26, No.11, (December 2007), pp. 3169-3180, ISSN 0953-816X
- El Idrissi, A., Ding, X.H., Scalia, J., Trenkner, E., Brown, W.T. & Dobkin, C. (2005). Decreased GABA(A) receptor expression in the seizure-prone fragile X mouse. *Neuroscience Letters*, Vol. 377, No.3, (April 2005), pp. 141-146, ISSN 0304-3940
- El Idrissi, A., Boukarrou, L., Dokin, C. & Brown, W.T. (2009). Taurine improves congestive functions in a mouse model of fragile X syndrome. *Advances in Experimental Medicine and Biology*, Vol. 643, pp. 191-198, ISSN 0065-2598
- Ethell, I.M. & Ethell, D.W. (2007). Matrix metalloproteinases in brain development and remodeling: synaptic functions and targets. *Journal of Neuroscience Research*, Vol.85, No. 13, (October 2007), pp. 2813-2823, ISSN 1097-4547
- Feng, Y., Absher, D., Eberhart, D.E., Brown, V., Malter, H.E. & Warren, S.T. (1997a). FMRP associates with polyribosomes as an mRNP, and the I304N mutation of severe fragile X syndrome abolishes this association. *Molecular Cell*, Vol.1, No.1, (December 1997), pp. 109-118, ISSN 1097-2765
- Feng, Y., Gutekunst, C.A., Eberhart, D.E., Yi, H., Warren, S.T. & Hersch, S.M. (1997b). Fragile X mental retardation protein: nucleocytoplasmic shuttling and association with somatodendritic ribosomes. *The Journal of Neuroscience*, Vol.17, No.5, (March 1997), pp. 1539-47, ISSN 0270-6474
- Ferraguti, F., Baldani-Guerra, B., Corsi, M., Nakanishi, S. & Corti, C. (1999). Activation of the extracellular signal-regulated kinase 2 by metabotropic glutamate receptors. *The European Journal of Neuroscience*, Vol.11, No.6, (June 1999), pp. 2073-2082, ISSN 0953-816X
- Ferrari, F., Mercaldo, V., Piccoli, G., Sala, C., Cannata, S., Achsel, T. & Bagni, C. (2007). The fragile X mental retardation protein-RNP granules show an mGluR-dependent localization in the post-synaptic spines. *Molecular and Cellular Neuroscience*, Vol.34, No.3, (March 2007), pp. 343-354, ISSN 1044-7431
- Frisby, C.L., Mattsson, J.P., Jensen, J.M., Lehmann, A., Dent, J. & Blackshaw, L.A. (2005). Inhibition of transient lower esophageal sphincter relaxation and gastroesophageal reflux by metabotropic glutamate receptor ligands. *Gastroenterology*, Vol.129, No.3, (September 2005), pp. 995-1004, ISSN 0016-5085
- Fritschy, J.M. & Mohler, H. (1995). GABAA-receptor heterogeneity in the adult rat brain: differential regional and cellular distribution of seven major subunits. *Journal of Comparative Neurology*, Vol. 359, No.1, (August 1995), pp. 154-194, ISSN 0021-9967
- Gallagher, S.M., Daly, C.A., Bear, M.F. & Huber, K.M. (2004). Extracellular signalregulated protein kinase activation is required for metabotropic glutamate receptor-dependent long-term depression in hippocampal area CA1. *The Journal of Neuroscience*, Vol.24, No.20, (May 2004), pp. 4859-4864, ISSN 0270-6474
- Gantois, I., Vandesompele, J., Speleman, F., Reyniers, E., D'Hooge, R., Severijnen, L.A., Willemsen, R., Tassone, F. & Kooy, R.F. (2006). Expression profiling suggests underexpression of the GABAA receptor subunit delta in the fragile X knockout

- mouse model. *Neurobiology of Disease*, Vol.21, No.2, (February 2006), pp. 346–357, ISSN 0969-9961
- Gatto, C.L. & Broadie, K. (2008). Temporal requirements of the fragile X mental retardation protein in the regulation of synaptic structure. *Development*, Vol.135, No.15, (August 2008), pp. 2637–2648, ISSN 1011-6370
- Gereau, R.W. & Conn, P.J. (1995). Multiple presynaptic metabotropic glutamate receptors modulate excitatory and inhibitory synaptic transmission in hippocampal area CA1. *The Journal of Neuroscience*, Vol.15, No.10, (October 1995), pp. 6879–6889, ISSN 0270-6474
- Gibson, J.R., Bartley, A.F., Hays, S.A. & Huber, K.M. (2008). Imbalance of neocortical excitation and inhibition and altered UP states reflect network hyperexcitability in the mouse model of fragile X syndrome. *Journal of Neurophysiology*, Vol.100, No.5, (November 2008), pp. 2615–2626, ISSN 0022-30
- Giuffrida, R., Musumeci, S., D'Antoni, S., Bonaccorso, C.M., Giuffrida-Stella, A.M., Oostra, B.A. & Catania, M.V. (2005). A reduced number of metabotropic glutamate subtype 5 receptors are associated with constitutive homer proteins in a mouse model of fragile X syndrome. *Journal of Neuroscience*, Vol.25, No.39, (September 2005), pp. 8909–8916, ISSN 1529-2401
- Glykys, J. & Mody, I. (2007). Activation of GABAA receptors: views from outside the synaptic cleft. *Neuron*, Vol.56, No.5, (December 2007), pp. 763–770, ISSN 0896-6273
- Govek, E.E., Newey, S.E. & Van Aelst, L. (2005). The role of the Rho GTPases in neuronal development. *Genes Development*, Vol.19, No.1, (January 2005), pp. 1–49, ISSN 0890-9369
- Gross, C., Nakamoto, M., Yao, X., Chan, C.B., Yim, S.Y., Ye, K., Warren, S.T. & Bassell, G.J. (2010). Excess phosphoinositide 3-kinase subunit synthesis and activity as a novel therapeutic target in fragile X syndrome. *The Journal of Neuroscience*, Vol.30, No.32, (August 2010), pp. 10624–10638, ISSN 0270-6474
- Gross, C., Yao, X., Pong, D.L., Jeromin, A. & Bassell, G.J. (2011). Fragile X mental retardation protein regulates protein expression and mRNA translation of the potassium channel Kv4.2. *The Journal of Neuroscience*, Vol.31, No.15, (April 2011), pp. 5693–5698, ISSN 1529-2401
- Hagerman, R.J. (2002). The physical and behavioural phenotype. In: *Fragile X Syndrome: Diagnosis, Treatment, and Research*. (Hagerman RJ, Hagerman PJ editors), The Johns Hopkins University Press, Baltimore, MD, pp. 3–109
- Hagerman, P. J. (2008). The fragile X prevalence paradox. *Journal of Medical Genetics*, Vol.45, No.8, (August 2008), pp. 498–499, ISSN 0022-2593
- Harlow, E.G., Till, S.M., Russell, T.A., Wijetunge, L.S., Kind, P. & Contractor, A. (2010). Critical period plasticity is disrupted in the barrel cortex of FMR1 knockout mice. *Neuron*, Vol.65, No.3, (February 2010), pp. 385–398, ISSN 1097-4199
- Herman, J.P. & Cullinan, W.E. (1997). Neurocircuitry of stress: central control of the hypothalamo-pituitary-adrenocortical axis. *Trends in Neurosciences*, Vol.20, No.2, (February 1997), pp. 78–84, ISSN 0166-2236
- Hessl, D., Rivera, S.M. & Reiss, A.L. (2004). The neuroanatomy and neuroendocrinology of fragile X syndrome. *Mental Retardation and Developmental Disabilities Research Reviews*, Vol.10, No.1, (2004), pp. 17–24, ISSN 1080-4013

- Hu, H., Qin, Y., Bochorishvili, G., Zhu, Y., van Aelst, L. & Zhu, J.J. (2008). Ras signaling mechanisms underlying impaired GluR1-dependent plasticity associated with fragile X syndrome. *The Journal of Neuroscience*, Vol.28, No.31, (July 2008), pp. 7847–7862, ISSN 0270-6474
- Huber, K.M., Gallagher, S.M. & Bear, M.F. (2001). Evidence for a novel signal transduction pathway in hippocampal mGluR-dependent LTD. Program number 388.7. In Abstract Viewer/Itinerary Planner, Society for Neuroscience
- Irwin, S.A., Patel, B., Idupulapati, M., Harris, J.B., Crisostomo, R.A., Larsen, B.P., Kooy, F., Willems, P.J., Cras, P., Kozlowski, P.B., Swain, R.A., Weiler, I.J. & Greenough, W.T. (2001). Abnormal dendritic spine characteristics in the temporal and visual cortices of patients with fragile-X syndrome: a quantitative examination. *American Journal of Medical Genetics*, Vol.98, No.2, (January 2001), pp. 161–167, ISSN 0148-7299
- Irwin, S.A., Idupulapati, M., Gilbert, M.E., Harris, J.B., Chakravarti, A.B., Rogers, E.J., Crisostomo, R.A., Larsen, B.P., Mehta, A., Alcantara, C.J., Patel, B., Swain, R.A., Weiler, I.J., Oostra, B.A. & Greenough, W.T. (2002). Dendritic spine and dendritic field characteristics of layer V pyramidal neurons in the visual cortex of fragile-X knockout mice. *American Journal of Medical Genetics*, Vol. 111, No.2, (August 2002), pp. 140–146, ISSN 0148-7299
- Jacquemont, S., Curie, A., des Portes, V., Torrioli, M.G., Berry-Kravis, E., Hagerman, R.J., Ramos, F.J., Cornish, K., He, Y., Paulding, C., Neri, G., Chen, F., Hadjikhani, N., Martinet, D., Meyer, J., Beckmann, J.S., Delange, K., Brun, A., Bussy, G., Gasparini, F., Hilse, T., Floesser, A., Branson, J., Bilbe, G., Johns, D. Gomez-Mancilla, B. (2011). Epigenetic Modification of the FMR1 Gene in Fragile X Syndrome Is Associated with Differential Response to the mGluR5 Antagonist AFQ056. *Science Translational Medicine*, Vol.3, No. 64, (January 2011), pp. 64ra1, ISSN 1946-6234
- Kammermeier, P.J., Xiao, B., Tu, J.C., Worley, P.F. & Ikeda, S.R. (2000). Homer proteins regulate coupling of group I metabotropic glutamate receptors to N-type calcium and M-type potassium channels. *The Journal of Neuroscience*, Vol.20, No.19, (October 2000), pp. 7238–7245, ISSN 0270-6474
- Kao, D.I., Aldridge, G.M., Weiler, I.J. & Greenough, W.T. (2010). Altered mRNA transport, docking, and protein translation in neurons lacking fragile X mental retardation protein. *Proceedings of the National Academy of Sciences of the United States of America*, Vol.107, No.35, (August 2010) pp. 15601–15606, ISSN 1091-6490
- Kaufman, L., Ayub, M. & Vincent, J.B. (2010). The genetic basis of non-syndromic intellectual disability: a review. *Journal of Neurodevelopmental Disorders*, Vol.2, No.4, (December 2010), pp. 182–209, ISSN 1866-1947
- Kawabata, S., Tsutsumi, R., Kohara, A., Yamaguchi, T., Nakanishi, S. & Okada, M. (1996). Control of calcium oscillations by phosphorylation of metabotropic glutamate receptors. *Nature*, Vol.383, No.6595, (September 1996), pp. 89–92, ISSN 0028-0836
- Khandjian, E.W., Corbin, F., Woerly, S. & Rousseau, F. (1996). The fragile X mental retardation protein is associated with ribosomes. *Nature Genetics*, Vol.12, No.1, (January 1996), pp. 91–93, ISSN 1061-40
- Kneussel, M. (2002). Dynamic regulation of GABA(A) receptors at synaptic sites. *Brain Research Review*, Vol.39, No.1, (June 2002), pp. 74–83, ISSN 0165-0173
- Kobayashi, K., Kuroda, S., Fukata, M., Nakamura, T., Nagase, T., Nomura, N., Matsuura, Y., Yoshida-Kubomura, N., Iwamatsu, A. & Kaibuchi, K. (1998). p140Sra-1 (specifically

- Rac1-associated protein) is a novel specific target for Rac1 small GTPase. *The Journal of Biological Chemistry*, Vol.273, No.1, (January 1998), pp. 291-295, ISSN 0021-9258
- Korpi, E.R., Grunder, G. & Luddens, H. (2002). Drug interactions at GABA(A) receptors. *Progress in Neurobiology*, Vol.67, No.2, (June 2002), pp. 113-159, ISSN 0301-0082
- López-Bendito, G., Shigemoto, R., Fairén, A. & Luján, R. (2002). Differential distribution of group I metabotropic glutamate receptors during rat cortical development. *Cerebellar Cortex*, Vol.12, No.6, (June 2002), pp. 625-38, ISSN 1047-3211
- Lu, Y.M., Jia, Z., Janus, C., Henderson, J.T., Gerlai, R., Wojtowicz, J.M. & Roder, J.C. (1997). Mice lacking metabotropic glutamate receptor 5 show impaired learning and reduced CA1 long-term potentiation (LTP) but normal CA3 LTP. *The Journal of Neuroscience*, Vol.17, No.13, (July 1997), pp. 5196-205, ISSN 0270-6474
- Luján, R., Roberts, J.D., Shigemoto, R., Ohishi, H. & Somogyi, P. (1997). Differential plasma membrane distribution of metabotropic glutamate receptors mGluR1 alpha, mGluR2 and mGluR5, relative to neurotransmitter release sites. *Journal of Chemical Neuroanatomy*, Vol.13, No.4, (October 1997), pp. 219-41, ISSN 0891-0618
- Mao, L., Yang, L., Tang, Q., Samdani, S., Zhang, G. & Wang, J.Q. (2005). The scaffold protein Homer1b/c links metabotropic glutamate receptor 5 to extracellular signal-regulated protein kinase cascades in neurons. *The Journal of Neuroscience*, Vol.25, No.10, (March 2005), pp. 2741-2752, ISSN 0270-6474
- Michel, C.I., Kraft, R. & Restifo, L.L. (2004). Defective neuronal development in the mushroom bodies of *Drosophila* fragile X mental retardation 1 mutants. *The Journal of Neuroscience*, Vol. 24, No.25, (June 2004), pp. 5798-5809, ISSN 0270-6474
- Mientjes, E.J., Willemsen, R., Kirkpatrick, L.L., Nieuwenhuizen, I.M., Hoogeveen-Westerveld, M., Verweij, M., Reis, S., Bardoni, B., Hoogeveen, A.T., Oostra, B.A. & Nelson, D.L. (2004). Fxr1 knockout mice show a striated muscle phenotype: implications for Fxr1p function in vivo. *Human Molecular Genetics*, Vol.13, No.13, (July 2004), pp. 1291-1302, ISSN 0964-6906
- Miyashiro, K.Y., Beckel-Mitchener, A., Purk, T.P., Becker, K.G., Barret, T., Liu, L., Carbonetto, S., Weiler, I.J., Greenough, W.T. & Eberwine, J. (2003). RNA cargoes associating with FMRP reveal deficits in cellular functioning in *Fmr1* null mice. *Neuron*, Vol.37, No.3, (February 2003), pp. 417-431, ISSN 0896-6273
- Moldrich, R.X., Chapman, A.G., De Sarro, G. & Meldrum, B.S. (2003). Glutamate metabotropic receptors as targets for drug therapy in epilepsy. *European Journal of Pharmacology*, Vol.476, No.1-2, (August 2003), pp. 3-16, ISSN 0014-2999
- Monzo, K., Papoulas, O., Cantin, G.T., Wang, Y., Yates, J.R. 3rd & Sisson, J.C. (2006). Fragile X mental retardation protein controls trailer hitch expression and cleavage furrow formation in *Drosophila* embryos. *Proceedings of the National Academy of Sciences of the United States of America*, Vol.103, No.48, (November 2006), pp. 18160-18165, ISSN 0027-8424
- Musumeci, S.A., Bosco, P., Calabrese, G., Bakker, C., De Sarro, G.B., Elia, M., Ferri, R. & Oostra, B.A. (2000). Audiogenic seizures susceptibility in transgenic mice with fragile X syndrome. *Epilepsia*, Vol.41, No.1, (January 2000), pp. 19-23, ISSN 0013-9580
- Musumeci, S.A., Calabrese, G., Bonaccorso, C.M., D'Antoni, S., Brouwer, J.R., Bakker, C., Elia, M., Ferri, R., Nelson, D.L., Oostra, B.A. & Catania, M.V. (2007). Audiogenic

- seizure susceptibility is reduced in fragile X knockout mice after introduction of FMR1 transgenes. *Experimental Neurology*, Vol.203, No.1, (January 2007), pp. 233-240, ISSN 0014-4886
- Napoli, I., Mercaldo, V., Boyl, P.P., Eleuteri, B., Zalfa, F., De Rubeis, S., Di Marino, D., Mohr, E., Massimi, M., Falconi, M., Witke, W., Costa-Mattioli, M., Sonenberg, N., Achsel, T. & Bagni, C. (2008). The fragile X syndrome protein represses activity-dependent translation through CYFIP1, a new 4E-BP. *Cell*, Vol.134, No.6, (September 2008), pp. 1042-54, ISSN 1097-4172
- Nicoletti, F., Iadarola, M.J., Wroblewski, J.T. & Costa, E. (1986a). Excitatory amino acid recognition sites coupled with inositol phospholipid metabolism: developmental changes and interaction with alpha 1-adrenoceptors. *Proceedings of the National Academy of Sciences of the United States of America*, Vol.83, No.6, (March 1986), pp. 1931-5, ISSN 0027-8424
- Nicoletti, F., Meek, J.L., Iadarola, M.J., Chuang, D.M., Roth, B.L. & Costa, E. (1986b). Coupling of inositol phospholipid metabolism with excitatory amino acid recognition sites in rat hippocampus. *Journal of Neurochemistry*, Vol.46, No.1, (January 1986), pp. 40-6, ISSN 0022-3042
- Nicoletti, F., Bockaert, J., Collingridge, G.L., Conn, P.J., Ferraguti, F., Schoepp, D.D., Wroblewski J.T. & Pin, J.P. (2011). Metabotropic glutamate receptors: From the workbench to the bedside. *Neuropharmacology*, Vol.60, (October 2010), pp. 1017-1041, ISSN 0028-3908
- Nimchinsky, E. A., Oberlander, A. M. & Svoboda, K. (2001). Abnormal development of dendritic spines in FMR1 knock-out mice. *The Journal of Neuroscience*, Vol.21, No.14, (July 2001), pp. 5139-5146, ISSN 0270-6474
- Nimchinsky, E. A., Sabatini, B.L. & Svoboda, K. (2002). Structure and function of dendritic spines. *Annual Review of Physiology*, Vol.64, (March 2002), pp. 313-53, ISSN 0066-4278
- Nordquist, R.E., Steckler, T., Wettstein, J.G., Mackie, C. & Spooren, W. (2008). Metabotropic glutamate receptormodulation, translationmethods, and biomarkers: Relationships with anxiety. *Psychopharmacology (Berl)*, Vol.199, No.3, (August 2008), pp. 389-402
- O'Donnel, W.T. & Warren, S.T. (2002). A decade of molecular studies of fragile X syndrome. *Annual review of Neuroscience*, Vol.25, (March 2002), pp. 315-338, ISSN 0147-006X
- Oliet, S.H, Malenka, R.C. & Nicoll, R.A. (1997). Two distinct forms of long-term depression coexist in CA1 hippocampal pyramidal cells. *Neuron*, Vol.18, No.6, (June 1997), pp. 969-982, ISSN 0896-6273
- Olmos-Serrano, J.L., Paluszkievicz, S.M., Martin, B.S., Kaufmann, W.E., Corbin, J.G. & Huntsman, M.M. (2010). Defective GABAergic neurotransmission and pharmacological rescue of neuronal hyperexcitability in the amygdala in a mouse model of fragile X syndrome. *The Journal of Neuroscience*, Vol.30, No.29, (July 2010), pp. 9929-38, ISSN 0270-6474
- Pacey, L.K., Heximer, S.P. & Hampson, D.R. (2009). Increased GABA(B) receptor-mediated signaling reduces the susceptibility of fragile X knockout mice to audiogenic seizures. *Molecular Pharmacology*, Vol.76, No.1, (July 2009), pp. 18-24, ISSN 0026-895X

- Pacey, L.K., Doss, L., Cifelli, C., der Kooy, D.V., Heximer, S.P. & Hampson, D.R. (2011). Genetic deletion of Regulator of G-protein Signaling 4 (RGS4) rescues a subset of fragile X related phenotypes in the FMR1 knockout mouse. *Molecular and Cellular Neuroscience*, Vol.46, No.3, (March 2011), pp. 563-572, ISSN 1044-7431
- Palmer, M.J., Irving, A.J., Seabrook, G.R., Jane, D.E. & Collingridge, G.L. (1997). The group I mGlu receptor agonist DHPG induces a novel form of LTD in the CA1 region of the hippocampus. *Neuropharmacology*, Vol.36, No.11-12, (November-December 1997), pp. 1517-1532, ISSN 0028-3908
- Park, S., Park, J.M., Kim, S., Kim, J.-A., Shepherd J.D., Smith-Hicks, C.L., Chowdhury, S., Kaufmann, W., Kuhl, D., Ryazanov, A.G., Huganir, R.L., Linden, D.J. & Worley, P.F. (2008). Elongation Factor 2 and Fragile X Mental Retardation Protein Control the Dynamic Translation of Arc/Arg3.1 Essential for mGluR-LTD. *Neuron*, Vol.59, No.1, (July 2008), pp. 70-83, ISSN 0896-6273
- Peavy, R.D. & Conn, P.J. (1998). Phosphorylation of mitogen-activated protein kinase in cultured rat cortical glia by stimulation of metabotropic glutamate receptors. *Journal of Neurochemistry*, Vol.71, No.2, (August 1998), pp. 603-12, ISSN 0022-3042.
- Porter, R.H., Jaeschke, G., Spooren, W., Ballard, T.M., Büttelmann, B., Kolczewski, S., Peters, J.U., Prinssen, E., Wichmann, J., Vieira, E., Mühlemann, A., Gatti, S., Mutel, V. & Malherbe, P. (2005). Fenobam: a clinically validated nonbenzodiazepine anxiolytic is a potent, selective, and noncompetitive mGlu5 receptor antagonist with inverse agonist activity. *Journal of Pharmacology and Experimental Therapeutics*, Vol.315, No.2, (November 2005), pp. 711-721, ISSN 0022-3565
- Qin, M., Kang, J., Burlin, T.V., Jiang, C. & Smith, C.B. (2005). Postadolescent changes in regional cerebral protein synthesis: an in vivo study in the FMR1 null mouse. *The Journal of Neuroscience*, Vol.25, No.20, (May 2005), pp. 5087-95, ISSN 1529-2401
- Qin, M., Entezam, A., Usdin, K., Huang, T., Liu, Z.H., Hoffman, G.E. & Smith, C.B. (2011). A mouse model of the fragile X premutation: effects on behavior, dendrite morphology, and regional rates of cerebral protein synthesis *Neurobiology of Disease*, Vol.42, No.1, (April 2011), pp. 85-98, ISSN 1095-953X
- Reyniers, E., Martin, J.J., Cras, P., Van Marck, E., Handig, I., Jorens, H.Z., Oostra, B.A., Kooy, R.F. & Willems P.J. (1999). Postmortem examination of two fragile X brothers with an FMR1 full mutation. *American Journal of Medical Genetics*, Vol.84, No.3, (May 1999), pp. 245-249, ISSN 0148-7299
- Rodriguez-Moreno, A., Sistiaga, A., Lerma, J. & Sanchez-Prieto, J. (1998). Switch from facilitation to inhibition of excitatory synaptic transmission by group I mGluR desensitization. *Neuron*, Vol.21, No.6, (December 1998), pp. 1477-1486, ISSN 0896-6273
- Romero-Zerbo, Y., Decara, J., el Bekay, R., Sanchez-Salido, L., Del Arco-Herrera, I., de Fonseca, F.R. & de Diego-Otero, Y. (2009). Protective effects of melatonin against oxidative stress in Fmr1 knockout mice: a therapeutic research model for the fragile X syndrome. *Journal of Pineal Research*, Vol.46, No.2, (March 2009), pp. 224-234, ISSN 0742-3098
- Ronesi, J.A. & Huber, K.M. (2008). Homer interactions are necessary for metabotropic glutamate receptor-induced long-term depression and translational activation. *Journal of Neuroscience*, Vol.28, No.2, (January 2008), pp. 543-547 ISSN 1529-2401

- Rong, R., Ahn, J.Y., Huang, H., Nagata, E., Kalman, D., Kapp, J.A., Tu, J., Worley, P.F., Snyder, S.H. & Ye, K. (2003). PI3 kinase enhancer-Homer complex couples mGluRI to PI3 kinase, preventing neuronal apoptosis. *Nature Neuroscience*, Vol.6, No.11, (November 2003), pp. 1153-61, ISSN 1097-6256
- Schenck, A., Bardoni, B., Langmann, C., Harden, N., Mandel, J.L. & Giangrande, A. (2003). CYFIP/Sra-1 controls neuronal connectivity in *Drosophila* and links the Rac1 GTPase pathway to the fragile X protein. *Neuron*, Vol.38, No.6, (June 2003), pp. 887-898, ISSN 0896-6273
- Schoepp, D.D. (2001). Unveiling the functions of presynaptic metabotropic glutamate receptors in the central nervous system. *The Journal of Pharmacology and Experimental Therapeutics*, Vol.299, No.1, (October 2001), pp. 12-20, ISSN 0022-3565
- Selby, L., Zhang, C. & Sun, Q.Q. (2007). Major defects in neocortical GABAergic inhibitory circuits in mice lacking the fragile X mental retardation protein. *Neuroscience Letters*, Vol.412, No.3, (February 2007), pp. 227-232, ISSN 0304-3940
- Seog, D.H.(2004). Glutamate receptor-interacting protein 1 protein binds to the microtubule-associated protein. *Bioscience, Biotechnology, and Biochemistry*, Vol.68, No.8, (August 2004), pp. 1808-1810, ISSN 0916-8451
- Sergé, A., Fourgeaud, L., He´mar, A. & Choquet, D. (2002). Receptor activation and Homer differentially control the lateral mobility of metabotropic glutamate receptor 5 in the neuronal membrane. *The Journal of Neuroscience*, Vol.22, No.10, (May 2002), pp. 3910 -3920, ISSN 0270-6474
- Simon, J., Wakimoto, H., Fujita, N., Lalande, M. & Barnard, E.A. (2004). Analysis of the set of GABA(A) receptor genes in the human genome. *The Journal of Biological Chemistry*, Vol.279, No.40, (October 2004), pp. 41422-41435, ISSN 0021-9258
- Siomi, H., Siomi, M.C., Nussbaum, R.L. & Dreyfuss, G. (1993). The protein product of the fragile X gene, FMR1, has characteristics of an RNA-binding protein. *Cell*, Vol.74, No.2, (July 1993), pp. 291-298, ISSN 0092-8674
- Sistiaga, A., Herrero, I., Conquet, F. & Sanchez-Prieto, J. (1998). The metabotropic glutamate receptor 1 is not involved in the facilitation of glutamate release in cerebrocortical nerve terminals. *Neuropharmacology*, Vol.37, No.12, (December 1998), pp. 1485-1492, ISSN 0028-3908
- Spencer, C.M., Serysheva, E., Yuva-Paylor, L.A., Oostra, B.A., Nelson, D.L. & Paylor, R. (2006). Exaggerated behavioral phenotypes in Fmr1/Fxr2 double knockout mice reveal a functional genetic interaction between Fragile X-related proteins. *Human Molecular Genetics*, Vol.15, No.12 (June 2006), pp. 1984-1994, ISSN 0964-6906
- Strumbos, J.G., Brown, M.R., Kronengold, J., Polley, D.B. & Kaczmarek, L.K. (2010). Fragile X mental retardation protein is required for rapid experience-dependent regulation of the potassium channel Kv3.1b. *The Journal of Neuroscience*, Vol.30, No.31, (August 2010), pp. 10263-10271, ISSN 1529-2401
- Tada, T. & Sheng, M. (2006) Molecular mechanism of dendritic spine morphogenesis. *Current Opinion in Neurobiology* Vol.16, No.1, (February 2006), pp. 95-101, ISSN 0959-4388
- Tessier, C.R. & Broadie, K. (2008). *Drosophila* fragile X mental retardation protein developmentally regulates activity-dependent axon pruning. *Development*, Vol.135, No.8, (Apr 2008) pp. 1547-1557, ISSN 1011-6370

- Toniolo, D. (2006). X-linked premature ovarian failure: a complex disease. *Current Opinion in Genetics & Development*, Vol.16, No.3, (June 2006), pp. 293-300, ISSN 0959-437X
- Tu, J.C., Xiao, B., Yuan, J.P., Lanahan, A.A., Leoffert, K., Li, M., Linden, D.J. & Worley, P.F. (1998). Homer binds a novel proline-rich motif and links group 1 metabotropic glutamate receptors with IP3 receptors. *Neuron*, Vol.21, No.4, (October 1998), pp. 717-726, ISSN 0896-6273
- Tu, J.C., Xiao, B., Naisbitt, S., Yuan, J.P., Petralia, R.S., Brakeman, P., Doan, A., Aakalu, V.K., Lanahan, A.A., Sheng, M. & Worley, P.F. (1999). Coupling of mGluR/Homer and PSD-95 complexes by the Shank family of postsynaptic density proteins. *Neuron*, Vol.23, No.3, (July 1999), pp. 583-592, ISSN 0896-6273
- Utari, A., Chonchaiya, W., Rivera, S.M., Schneider, A., Hagerman, R.J., Faradz, S.M., Ethell, I.M. & Nguyen, D.V. (2010). Side effects of minocycline treatment in patients with fragile X syndrome and exploration of outcome measures. *American journal on intellectual and developmental disabilities*, Vol.115, No.5, (September 2010) pp. 433-443, ISSN 1944-7515
- Verkerk, A.J., Pieretti, M., Sutcliffe, J.S., Fu, Y.H., Kuhl, D.P., Pizzuti, A., Reiner, O., Richards, S., Victoria, M.F., Zhang, F., Eussen, B.E., van Ommen, G.B., Blonden, L.A.J., Riggins, G.J., Chastain, J.L., Kunst, C.B., Galjaard, H., Caskey, C.T., Nelson D.L., Oostra B.A. & Warren S.T. (1991). Identification of a gene (FMR-1) containing a CGG repeat coincident with a breakpoint cluster region exhibiting length variation in fragile X syndrome. *Cell*, Vol.65, No.5, (May 1991), pp. 905-914, ISSN 0092-8674
- Volk, L.J., Pfeiffer, B.E., Gibson, J.R. & Huber, K.M. (2007). Multiple Gq-coupled receptors converge on a common protein synthesis-dependent long-term depression that is affected in fragile X syndrome mental retardation. *The Journal of Neuroscience*, Vol.27, No.43, (October 2007), pp. 11624-11634, ISSN 0270-6474
- Walker, K., Bowes, M., Panesar, M., Davis, A., Gentry, C., Kesingland, A., Gasparini, F., Spooren, W., Stoehr, N., Pagano, A., Flor, P.J., Vranesic, I., Lingenhoehl, K., Johnson, E.C., Varney, M., Urban, L. & Kuhn, R. (2001). Metabotropic glutamate receptor subtype 5 (mGlu5) and nociceptive function. I. Selective blockade of mGlu5 receptors in models of acute, persistent and chronic pain. *Neuropharmacology*, Vol.40, No.1, (June 2001). pp. 1-9, ISSN 0028-3908
- Wang, H., Wu, L.J., Kim, S.S., Lee, F.J., Gong, B., Toyoda, H., Ren, M., Shang, Y.Z., Xu, H., Liu, F., Zhao, M.G. & Zhuo M. (2008). FMRP acts as a key messenger for dopamine modulation in the forebrain. *Neuron*, Vol.59, No.4, (August 2008), pp. 634-647, ISSN 0896-6273
- Waung, M.W. & Huber, K.M. (2009). Protein translation in synaptic plasticity: mGluR-LTD, Fragile X. *Current Opinion in Neurobiology*, Vol.19, No.3, (June 2009), pp. 319-326, ISSN 0959-4388
- Weiler, I.J., Irwin, S.A., Klintsova, A.Y., Spencer, C.M., Brazelton, A.D., Miyashiro, K., Comery, T.A., Patel, B., Eberwine, J. & Greenough, W.T. (1997). Fragile X mental retardation protein is translated near synapses in response to neurotransmitter activation. *Proceedings of the National Academy of Sciences of the United States of America*, Vol.94, No.10, (May 1997), pp. 5395-5400, ISSN 0027-8424
- Weiler, I.J., Spangler, C.C., Klintsova, A.Y., Grossman, A.W., Kim, S.H., Bertaina-Anglade, V., Khaliq, H., de Vries, E., Lambers, F.A.E., Hatia, F., Base, C.K. & Greenough,

- W.T. (2004). Fragile X mental retardation protein is necessary for neurotransmitter-activated protein translation at synapses. *Proceedings of the National Academy of Sciences of the United States of America*, Vol.101, No.50, (December 2004), pp. 17504 – 17509, ISSN 0027-8424
- Xu, X.L., Li, Y., Wang, F. & Gao, F.B. (2008). The steady-state level of the nervous-system-specific microRNA-124a is regulated by dFMR1 in *Drosophila*. *The Journal of Neuroscience*, Vol.28, No.46, (November 2008), pp. 11883-11889, ISSN 0270-6474
- Yan, Q.J., Rammal, M., Tranfaglia, M. & Bauchwitz, R.P. (2005). Suppression of two major Fragile X Syndrome mouse model phenotypes by the mGluR5 antagonist MPEP. *Neuropharmacology*, Vol.49, No.7, (December 2005), pp. 1053-1066, ISSN 0028-3908
- Zalfa, F., Giorgi, M., Primerano, B., Moro, A., Di Penta, A., Reis, S., Oostra, B. & Bagni, C. (2003). The fragile X syndrome protein FMRP associates with BC1 RNA and regulates the translation of specific mRNAs at synapses. *Cell*, Vol.112, No.3, (February 2003), pp. 317-327, ISSN 0092-8674
- Zalfa, F., Eleuteri, B., Dickson, K.S., Mercaldo, V., De Rubeis, S., di Penta, A., Tabolacci, E., Chiurazzi, P., Neri, G., Grant, S.G. & Bagni, C. (2007). A new function for the fragile X mental retardation protein in regulation of PSD-95 mRNA stability. *Nature Neuroscience*, Vol.10, No.5, (May 2007), pp. 578-587, ISSN 1097-6256
- Zhang, J., Hou, L., Klann, E. & Nelson, D.L. (2009). Altered hippocampal synaptic plasticity in the FMR1 gene family knockout mouse models. *Journal of Neurophysiology*, Vol.101, No.5, (May 2009), pp. 2572-2580, ISSN 0022-3077
- Zhang, Y.Q., Bailey, A.M., Matthies, H.J., Renden, R.B., Smith, M.A., Speese, S.D., Rubin, G.M. & Broadie, K. (2001). *Drosophila* fragile X-related gene regulates the MAP1B homolog Futsch to control synaptic structure and function. *Cell*, Vol.107, No.5, (November 2001), pp. 591-603, ISSN 0092-8674
- Zhang, Y.Q., Friedman, D.B., Wang, Z., Woodruff, E. 3rd, Pan, L., O'Donnell, J. & Broadie, K. (2005). Protein expression profiling of the *drosophila* fragile X mutant brain reveals up-regulation of monoamine synthesis. *Molecular & Cellular Proteomics*, Vol.4, No.3, (March 2005), pp. 278-290, ISSN 1535-9476
- Zukin, R.S., Richter, J.D. & Bagni, C. (2009). Signals, synapses, and synthesis: how new proteins control plasticity. *Frontiers in Neural Circuits*, Vol.3, No.14, (October 2009), pp. 1-8, ISSN 1662-5110

IntechOpen

Latest Findings in Intellectual and Developmental Disabilities Research

Edited by Prof. Uner Tan

ISBN 978-953-307-865-6

Hard cover, 404 pages

Publisher InTech

Published online 15, February, 2012

Published in print edition February, 2012

Intellectual and Developmental Disabilities presents reports on a wide range of areas in the field of neurological and intellectual disability, including habitual human quadrupedal locomotion with associated cognitive disabilities, Fragile X syndrome, autism spectrum disorders, Down syndrome, and intellectual developmental disability among children in an African setting. Studies are presented from researchers around the world, looking at aspects as wide-ranging as the genetics behind the conditions to new and innovative therapeutic approaches.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Simona D'Antoni, Michela Spatuzza, Carmela M. Bonaccorso, Elisabetta Aloisi, Sebastiano Musumeci and Maria Vincenza Catania (2012). Fragile X Syndrome: From Pathophysiology to New Therapeutic Perspectives, Latest Findings in Intellectual and Developmental Disabilities Research, Prof. Uner Tan (Ed.), ISBN: 978-953-307-865-6, InTech, Available from: <http://www.intechopen.com/books/latest-findings-in-intellectual-and-developmental-disabilities-research/fragile-x-syndrome-from-pathophysiology-to-new-therapeutic-perspectives>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](https://creativecommons.org/licenses/by/3.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen