We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Noncoding RNAs in Neural Stem Cell Development

Shan Bian and Tao Sun

*Department of Cell and Developmental Biology
Cornell University Weill Medical College
USA*

1. Introduction

Neural stem cells and neural progenitors/precursors (NSCs/NPs) are identified in both embryonic and adult central nervous system (CNS). NSCs can self-renew and give rise to neurons and glia. The development of NSCs is controlled by precisely orchestrated gene expression regulation. Recently, emerging evidence has shown the importance of noncoding RNA regulation in NSC self-renewal, proliferation, survival and differentiation. In this chapter, we will present new research of noncoding RNA functions in NSC development. We will highlight the future directions of applying noncoding RNAs in stem cell-based therapy for neurological diseases.

2. Noncoding RNAs

Noncoding RNAs (ncRNAs) are functional RNA molecules that do not show protein translation capability. ncRNAs consist of ribosomal RNAs (rRNAs), transfer RNAs (tRNAs), piwi-interacting RNAs (piRNAs), microRNAs (miRNAs), and long noncoding RNAs (lncRNAs) and so on. ncRNAs have shown to play distinct but also conserved roles in normal development in invertebrates and vertebrates.

2.1 piRNAs

piRNAs are a group of small RNAs with size between 26-31 nucleotides (nt), which are only found in male and female germlines of invertebrates and within testes in mammals (Houwing et al., 2007; 2009; Lander et al., 2001; Lau et al., 2006; Seto et al., 2007). piRNAs interact with piwi proteins to form RNA-protein complexes (Das et al., 2008; Houwing et al., 2007). The piRNA-protein complexes have been shown to silence transcription, specifically transposons (Brennecke et al., 2008; Das et al., 2008). Since piRNAs are mainly expressed during the germline stem cell development, they will not be discussed further in this chapter.

2.2 miRNAs

miRNAs are ~22 nt highly conserved small noncoding RNAs found in almost all eukaryotic cells (Khraiweh et al., 2010) (Fig. 1). Like coding genes, miRNAs are mainly transcribed by the

RNA Polymerase II (Pol II) into long primary miRNA transcripts (pri-miRNAs). Pri-miRNAs are next cleaved by Drosha, a class 2 RNase III enzyme, to produce short hairpin stem-loop structures, known as precursor miRNAs (pre-miRNAs) (Ambros, 2008; Gregory et al., 2006). Pre-miRNAs are then processed into about 20-25 nt double-stranded mature miRNAs by RNase III enzyme Dicer (Lund and Dahlberg, 2006). The duplex undergoes unwinding and the strand with the weakest base pairing at its 5' terminus, together with Dicer and many associated proteins including Argonaute proteins, form an active RNA-induced silencing complex (RISC) (Neilson and Sharp, 2008; Rana, 2007). A miRNA recognizes and binds to the 3' untranslated region (3' UTR) of target messenger RNAs (mRNAs) by imperfect complementary sequence recognition (Neilson and Sharp, 2008; Wang et al., 2004). Once bound, the RISC can negatively regulate miRNA targets by degrading the mRNA or repressing its translation (Khraiwesh et al., 2010; Pratt and MacRae, 2009) (Fig. 1).

miRNAs are classified into intergenic and intragenic miRNAs, depending on their genomic location (Lagos-Quintana et al., 2001; Lau et al., 2001; Lee and Ambros, 2001). Intergenic miRNAs are located at intergenic regions (between genes) in the genome. These miRNAs are usually transcribed using their own promoters. Intragenic miRNAs normally lie in intronic regions of coding genes with the same orientation. They are often transcribed together with the host genes.

Fig. 1. A scheme of microRNA biogenesis. microRNAs silence target coding genes by binding to the 3' untranslated region (3' UTR).

2.3 lncRNAs

The majority of the human genome has previously been considered as “junk” DNA, since only about 1.5% of the human genome, which occupies over 3 billion DNA base pairs,

consists of protein-coding genes (Lander et al., 2001). A recent study of a large-scale complementary DNA (cDNA) sequencing project has shown that four fifths of transcripts of the human genome are RNA transcripts that don't encode proteins (Kapranov et al., 2007). These RNA transcripts are normally longer than 200 nt, thus they are called long noncoding RNAs. Except no open reading frame (ORF) found within lncRNAs, they share many features with coding mRNAs such as 5' capping. lncRNAs usually contain exons and introns (Carninci et al., 2005).

lncRNAs display different features of genomic location and orientation in the genome. Emerging evidence has demonstrated potential functions of lncRNAs in many biological events. lncRNAs have been shown to modulate gene transcription via different mechanisms. For example, some *cis*-antisense lncRNAs complementary to protein-coding transcripts regulate expression of the coding genes (Feng et al., 2006; Kotake et al., 2010; Pasmant et al., 2007; Tochitani and Hayashizaki, 2008; Yu et al., 2008); some lncRNAs modulate transcription factors by acting as co-regulators (Feng et al., 2006; Nguyen et al., 2001; Shamovsky et al., 2006; Wang et al., 2008; Willingham et al., 2005; Yang et al., 2001). Moreover, some studies have shown that lncRNAs are involved in epigenetic regulations such as chromatin and histone modification, and X-chromosome inactivation (Denisenko et al., 1998; Mancini-Dinardo et al., 2006; Mazo et al., 2007; Rinn et al., 2007; Sanchez-Elsner et al., 2006; Tsai et al., 2010; Wutz et al., 2002).

3. Noncoding RNAs and neural stem cell development

Noncoding RNAs such as miRNAs and lncRNAs participate gene expression regulation in many ways. The underlying mechanisms of noncoding RNA functions in normal development are beginning to be uncovered. In this book chapter, we will focus on reviewing functions of miRNAs and lncRNAs in neural stem cell development, such as NSC self-renewal, cell fate determination and survival.

3.1 miRNAs and self-renewal and proliferation of NSCs

The ability of self-renewal is essential for NSCs/NPs to perpetuate themselves to maintain an undifferentiated status during the embryonic stage and even in the adulthood (Gage, 2000; Shi et al., 2008; Temple, 2001). NSC proliferation and self-renewal are modulated by a complicated regulation network that consists of growth factors, epigenetic regulators, transcription factors and extrinsic signaling molecules from the NSC niche. Recent discovers have indicated that ncRNAs also play important roles in NSC self-renewal through a post-transcriptional regulation mechanism (Doe, 2008; Shi et al., 2008).

miRNAs have been shown to play essential roles in regulating NSC proliferation. Since Dicer is the key enzyme in miRNA processing, several studies have reported the global effects of miRNAs in NSC development by ablating *Dicer* and in turn blocking biogenesis of all miRNAs in the CNS using tissue specific Cre lines. Conditional deletion of *Dicer* from the mouse cerebral cortex using the *Emx1-Cre* line results in a significant reduction in cortical size and the cortical NP pool (De Pietri Tonelli et al., 2008; Kawase-Koga et al., 2010; Kawase-Koga et al., 2009). *Dicer* ablation from the mouse CNS using the *Emx1-Cre* and *Nestin-Cre* line causes a reduction of NSC numbers and abnormal differentiation (Andersson et al., 2010; Kawase-Koga et al., 2009) (Fig. 2A). *Dicer*-deficient NSCs display apoptosis when

mitogens are withdrawn from the culture medium (Fig. 2A). Because Dicer is also involved in maintaining the heterochromatin assembly, the defects of NSCs in *Dicer* knockout mice need to be carefully interpreted (Fukagawa et al., 2004; Kanellopoulou et al., 2005). Examining functions of individual miRNAs will help reveal precise roles of miRNAs in the NSC self-renewal and proliferation (Fig. 3).

Fig. 2. A. *Dicer*-deficient (*Dicer-Ko*) neural stem cells (NSCs) did not survive well in a differentiation culture medium without mitogens. Most *Dicer-Ko* neurospheres died after 48 hours in culture. Many differentiated cells (arrows) migrated away from the control neurosphere but not from the *Dicer-Ko* neurospheres. B. Under the differentiation condition without mitogens, passaged (p-1) *Dicer-Ko* NSCs gave rise to cells expressing neuronal (Tuj1⁺) and glial (GFAP⁺ and O4⁺) markers. However, their morphology was abnormal, as shown with shorter neurites and processes than controls (Kawase-Koga et al., 2010).

let-7, the first identified miRNA (Reinhart et al., 2000), has been shown to regulate NSC proliferation and differentiation by targeting the nuclear receptor TLX and the cell cycle regulator cyclin D1 (Zhao et al., 2010a) (Fig. 3). Overexpression of *let-7b* inhibits NSC proliferation and enhances differentiation, while knockdown of *let-7b* promotes NSC proliferation (Zhao et al., 2010a). It appears that the expression levels of let-7 in NSCs are controlled by a feedback regulation of Lin-28, a pluripotency factor that controls miRNA processing in NSCs (Rybak et al., 2008). Lin-28 binds to the *let-7* precursor and inhibits its processing by Dicer. On the other hand, the expression of Lin-28 is repressed by let-7 and miR-125, allowing the maturation of let-7 (Fig. 4A). This feedback loop reveals an autoregulation between miRNA let-7 and miR-125, and transcription factor Lin-28 during NSC development (Rybak et al., 2008).

miR-124 is identified as a CNS-enriched miRNA and its expression is upregulated during neuronal differentiation (Lagos-Quintana et al., 2002) (Fig. 3). In the adult brain, NSCs are identified in the subventricular zone (SVZ). In cultured adult NSCs derived from the SVZ and in the SVZ *in vivo*, knocking down *miR-124* results in an increase of NSC proliferation and a decrease of differentiation, while overexpressing *miR-124* reduces the number of dividing precursors and enhances neuronal differentiation (Cheng et al., 2009). Moreover, miR-124 modulates NSC proliferation and differentiation by suppressing Sox9 expression in adult NSCs (Cheng et al., 2009). A recent study has shown that miR-124 regulates neuronal differentiation through a mutual inhibition mechanism of Ephrin-B1 (Arvanitis et al., 2010). In

addition, miR-124 promotes differentiation of NPs by modulating a network of nervous system-specific alternative splicing through suppressing expression of PTBP1, which encodes a global repressor of alternative pre-mRNA splicing (Makeyev et al., 2007). Together, miR-124 plays a general role in promoting differentiation of embryonic and adult NSCs and NPs. It appears that miR-124 executes its function through repressing various targets.

Fig. 3. Many miRNAs are involved in neural stem cell (NSC) self-renewal and differentiation into neurons, astrocytes and oligodendrocyte precursor cells (OPCs) and oligodendrocytes.

Fig. 4. Feedback loop regulation of miRNAs and their target genes. A. Let-7 processing is inhibited by Lin-28, and the 3' untranslated region (3' UTR) of Lin-28 has binding sites for Let-7. B. TLX inhibits miR-9 expression, while miR-9 displays silencing effects on TLX.

miR-9 is another CNS-enriched miRNA. miR-9 is shown to inhibit NSC proliferation but promote differentiation through a feedback regulation of a nuclear receptor TLX (Zhao et al., 2009) (Fig. 4B). In human embryonic stem cell (ESC) derived NPs, miR-9 is shown to have a positive effect on proliferation but a negative effect on migration by directly targeting *Stmn1*, which increases microtubule instability (Delaloy et al., 2010). The opposite effect of miR-9 on proliferation is perhaps caused by differential physical contacts of miR-9 with target genes and the different culture systems.

In the CNS of *Xenopus*, miR-9 knockdown promotes the proliferation of NPs in the hindbrain, leads to an increased expression of *cyclin D1* and a downregulation of *p27Xic1* (Bonev et al., 2011). miR-9 targets *Hairy1* and regulates proliferation of NPs (Bonev et al., 2011). In zebrafish, miR-9 promotes differentiation of NPs that give rise to neurons at the midbrain-hindbrain domain and controls the organization of the midbrain-hindbrain boundary by targeting several genes in the Fibroblast growth factor (Fgf) signaling, such as *fgf8-1* and *fgfr1* (Leucht et al., 2008). In the chick spinal cord, miR-9 specifies a subtype of motor neurons that project axons to the axial muscles from motor neuron progenitors by specifically targeting transcription factor FoxP1 (Otaegi et al., 2011).

In the mouse brain, miR-9 function is demonstrated by the generation of *miR-9-2* and *miR-9-3* double knockout mice. *miR-9* double mutants show reduced cortical layers, disordered migration of interneurons, and misrouted thalamocortical axons and cortical axon projections, suggesting an important role of miR-9 in NP proliferation, differentiation and migration during brain development (Shibata et al., 2011). Moreover, it appears that miR-9 regulates multiple target genes, including Foxg1, Pax6 and Gsh2, which have shown to be essential in cortical development (Shibata et al., 2011). Therefore, miR-9 plays an important role in controlling differentiation of NSCs/NPs in different regions in the CNS (Fig. 3).

The major role of let-7, miR-124 and miR-9 is to inhibit NSC/NP proliferation and to induce their differentiation into specific cell types. miRNAs that promote proliferation of NSCs and NPs have also been identified (Fig. 3). miR-134 plays a role in enhancing proliferation of cortical NPs by targeting doublecortin (*Dcx*) and/or Chordin-like 1 (*Chrdl1*) (Gauthwin et al., 2011). miR-25 is shown to be a major player in the miR-106-25 cluster in neural development. Overexpression of *miR-25* but not *miR-106b* and *miR-93* promotes adult NP proliferation (Brett et al., 2011). Interestingly, the expression of the miR-106-25 cluster is regulated by FoxO3, a transcription factor maintaining the NSC population (Renault et al., 2009).

During the retina development, *otx2* and *vsx1* genes are shown to control the division of retinal precursors and differentiation into bipolar retina neurons. In early retinal precursors, the expression of *otx2* and *vsx1* is inhibited, accompanied with a rapid precursor division. miR-129, miR-155, miR-214, and miR-222, which are highly expressed in the embryonic retina, have been identified to target and repress translation of *otx2* and *vsx1*, by which they promote proliferation of retinal precursors (Decembrini et al., 2009).

miRNA expression is also controlled by epigenetic regulators in the NSC development. The expression of miR-137 is regulated by DNA methyl-CpG-binding protein (MeCP2) and transcription factor Sox2. miR-137 modulates adult NSC proliferation and cell fate determination by targeting *Ezh2*, a histone methyltransferase and polycomb group protein (Szulwach et al., 2010). Ectopic expression of *miR-137* in adult NSCs enhances proliferation, while knockdown of *miR-137* promotes differentiation of adult NSCs (Szulwach et al., 2010).

In addition, miR-184 expression is suppressed by methyl-CpG binding protein 1 (MBD1) and miR-184 promotes adult NSC proliferation by repressing the expression of Numb-like (Numbl) (Liu et al., 2010).

3.2 lncRNAs and proliferation of NSCs

The lncRNAs may also play a role in controlling NSC proliferation, even though studies of lncRNAs in NSC development are still sparse. Sox2 is a transcription factor and plays a key role in the maintenance of the undifferentiating state of embryonic and adult NSCs (Pevny and Placzek, 2005). *Sox2 overlapping transcript (Sox2OT)* is a lncRNA containing *Sox2* gene and shares the same transcriptional orientation with *Sox2* (Fig. 5A). Similar to *Sox2*, *Sox2OT* is stably expressed in mouse embryonic stem cells and down-regulated during differentiation. *Sox2OT* is expressed in the neurogenic regions of the adult mouse brain including olfactory bulb (OB), rostral migratory stream (RMS) and SVZ, and is dynamically regulated during vertebrate CNS development, implying its role in regulating NSC self-renewal and neurogenesis (Amaral et al., 2009; Mercer et al., 2008).

Fig. 5. Genomic location and potential functions of long noncoding RNAs (lncRNAs) (Bian and Sun, 2011). A. *Sox2 overlapping transcript Sox2OT* is a lncRNA containing *Sox2* gene and shares the same transcriptional orientation with *Sox2*. B. *Evf2* is transcribed from the intergenic region between the *Dlx-5* and *Dlx-6* loci, and is overlapped with *Dlx-5/6* enhancer i (ei) and enhancer ii (eii) sequences. *Evf2* acts as a transcriptional co-activator of *Dlx-2* and activates the *Dlx5/6* enhancer. C. *Nkx2.2 antisense (Nkx2.2as)* is an antisense lncRNA to *Nkx2.2* gene and promotes *Nkx2-2* expression.

3.3 Summary

Taken together, self-renewal and differentiation of NSCs and NPs are controlled by complex gene regulation networks that consist of both protein coding genes and noncoding miRNAs. During proliferation and differentiation of NSCs and NPs, one miRNA can have multiple target genes and features a feedback regulation with their targets (Figs. 3 and 4). The availability of physical contacts and the binding affinity of a miRNA and its targets perhaps determine interactions of the miRNA with the specific targets. The interactions of miRNAs

and their target genes eventually produce proper protein output of key factors that directly control self-renewal, proliferation and differentiation of NSCs/NPs.

3.4 NSC survival controlled by noncoding RNAs

Several reports have shown that miRNAs play a general role in controlling cell survival. Conditional deletion of *Dicer* from neural crest cells using *Wnt1-Cre* mouse line results in an increased apoptosis of neural crest-derived cells (Zehir et al., 2010). Ablation of *Dicer* from postmitotic neurons in the cortex and the hippocampus using *calmodulin kinase II* (*CaMKII*) promoter-driven *Cre* transgenic mice results in smaller cortex, enhanced cortical cell death (Davis et al., 2008). *Emx1-Cre Dicer* conditional knockout mice have shown an increased apoptosis, especially in the ventricular zone (VZ) and SVZ (De Pietri Tonelli et al., 2008). Our own work of cortical NSCs of *Emx1-Cre Dicer* conditional knockout mice using proteomic analysis by mass spectrometry and bioinformatic assays has indicated that *Dicer* deletion results in an increase of pro-cell-death and a decrease of pro-survival proteins in *Dicer*-deficient NSCs (Kawase-Koga et al., 2010) (Fig. 2A). Interestingly, an upregulation of fragile X mental retardation protein (FMRP), a proven target for miR-124, and Caspase3, a key cell apoptosis molecule, are observed in *Dicer*-deficient NSCs. On the other hand, proteins such as transforming growth factor-beta receptor type II (TGFβR2) and SOD1 are downregulated in *Dicer*-deficient NSCs (Kawase-Koga et al., 2010). These observations suggest that miRNAs perhaps control survival of NSCs by modulating the balance of protein output of genes regulating apoptosis and survival.

Neurotrophins and their receptors play important roles in the NSC proliferation, survival and differentiation. miR-128 is shown to target the truncated non-catalytic form of the human neurotrophin-3 receptor (NTRK3), which affects membrane remodeling and cytoskeletal reorganization. Overexpression of *miR-128* in neuroblastoma cells leads to round cell body and shorter neurites, which is similar to knockdown of truncated NTRK3. miR-128 overexpression causes altered expression of genes involved in cell proliferation and apoptosis such as antiapoptotic factor Bcl-2, suggesting an important role of miR-128 on cell survival (Guidi et al., 2010). Moreover, miR-134 is shown to be required for inhibiting apoptosis initiated by Chrdl-1 in cortical progenitors (Gaughwin et al., 2011). Studies on an ethanol teratogenic culture model by exposing embryonic cortex-derived NPs in ethanol have revealed different roles of miRNAs during this pathological process (Sathyan et al., 2007). In NP cultures, miR-21 is suppressed by the ethanol exposure and the reduction of miR-21 causes cell apoptosis, suggesting an anti-apoptotic effect of miR-21 (Sathyan et al., 2007).

The BH3-only family is a group of pro-apoptotic regulators, including Bim, Hrk, Bmf, Puma and N-Bak, which induce cytochrome c release from mitochondria (Giam et al., 2008). Overexpression of miR-29b in neurons inhibits endogenous BH3-only proteins Bim, Puma and Bmf, and promotes neuronal survival (Giam et al., 2008; Kole et al., 2011). In the brain of the calorie-restricted mice, expression of three miRNAs, miR-181a-1*, miR-30e and miR-34a, is significantly downregulated with a corresponding upregulation of their target gene Bcl-2, a decrease of pro-apoptotic factor Bax and cleavage of Caspases (Khanna et al., 2011). Overexpressing these three genes results in an increased cell apoptosis, accompanied with a decrease in Bcl-2 expression (Khanna et al., 2011).

miRNAs also play an important role in neural tissue growth and organ development by regulating cell survival. In *Drosophila*, the Hippo pathway together with Yorkie transcriptional activator contribute to the regulation of tissue growth by stimulating cell proliferation and inhibiting apoptosis (Saucedo and Edgar, 2007). Recent studies have shown that Yorkie not only activates cyclin E and apoptosis inhibitor DIAP1, but also triggers the expression of *bantam* miRNA to promote proliferation and cell survival (Huang et al., 2005; Thompson and Cohen, 2006). A downregulation of *bantam* miRNA is found in dying Rim cells at the eye margin, and restoration of *bantam* miRNA to higher levels prevents apoptosis of these cells, suggesting a role of *bantam* miRNA in enhancing cell survival in eye development (Thompson and Cohen, 2006). In addition, as the largest miRNA family in *Drosophila*, miR-2/6/11/13/308 are required for inhibiting embryonic apoptosis by suppressing pro-apoptotic factors *hid*, *grim*, *reaper* and *sickle* (Leaman et al., 2005).

In the forebrain of *Xenopus*, miR-9 deletion results in apoptosis of NPs due to increased expression of p53 (Bonev et al., 2011). In addition, miR-24a is expressed in the retina of *Xenopus* (Walker and Harland, 2009). Overexpression of *miR-24a* in retinal cells prevents cells from death, while knockdown of *miR-24a* causes a reduction in eye size due to an increased apoptosis (Walker and Harland, 2009). miR-24a controls cell survival by a negative regulation of pro-apoptotic factors caspase9 and *apaf1*.

In summary, miRNAs play critical roles in regulating survival of both NSCs/NPs and postmitotic neurons. miRNAs either promote cell survival or lead to apoptosis, depending on functions of their target genes.

3.5 NSC differentiation and cell fate determination mediated by noncoding RNAs

In the mammalian CNS, different neural cell types arise and migrate in a precise temporospatial manner. In the developing mouse brain, neurons arise first by embryonic day 12 (E12), neurogenesis peaks at E14 and ceases by E18. Astrocytes appear around E18, with their numbers peaking in the postnatal period. Oligodendrocytes are generated after birth when the neurogenesis is almost complete. Studies have shown that ncRNAs play an important role in regulating both neurogenesis and gliogenesis.

3.5.1 Cell fate determination controlled by miRNAs

miRNAs play essential roles in NSC differentiation and the cell fate switch between neurons and glia (Cuellar et al., 2008; Hebert et al., 2010; Zheng et al., 2010). We have found that *Dicer*-deficient NSCs display abnormal differentiation, with shorter neurites in neurons and fewer processes in glial cells (Kawase-Koga et al., 2010) (Fig. 2B). Conditional deletion of *Dicer* from the mouse forebrain neurons using *CamKII-Cre* line results in neuronal degeneration and an increase in glial fibrillary acidic protein (GFAP)-positive astrocytes (Hebert et al., 2010). *Dicer* ablation in the dopaminergic neurons in the basal ganglia using a *dopamine receptor-1 (DR-1)-Cre* line leads to astroglialogenesis, but not neurodegeneration (Cuellar et al., 2008). Interestingly, in the mouse spinal cord, conditional deletion of *Dicer* using *Olig1-Cre* line disrupts production of both oligodendrocytes and astrocytes (Zheng et al., 2010). These observations suggest that global loss of miRNAs in specific precursor cells affects production of distinct cell types.

miRNA expression profiling studies have shown that some miRNAs are preferentially expressed in neurons or glia. For example, miR-124 and miR-128 are highly expressed in neurons, while miR-23 is restrictively expressed in astrocytes. miR-26 and miR-29 display higher expression in astrocytes than in neurons; and miR-9 and miR-125 are evenly expressed in neurons and astrocytes (Smirnova et al., 2005). Overexpressing *miR-124* in cultured NSCs and in embryonic cortical NPs using lenti-virus and *in utero* electroporation, respectively, promotes neurogenesis and stimulates cortical progenitor migration (Maiorano and Mallamaci, 2009). In cultured adult NSCs, overexpressing *miR-124* enhances neuronal differentiation (Cheng et al., 2009). Ectopic expression of miR-124a and miR-9 in the embryonic stem cell-derived NPs results in a great reduction of GFAP-positive astrocytes compared to the control groups, while knockdown of miR-9, but not miR-124a, switches differentiation of NPs from neurogenesis to astrogliogenesis (Krichevsky et al., 2006). miR-124 and miR-9 promotes neurogenesis by targeting phospholated signal transducer and activator of transcription 3 (STAT3), a transcription factor normally initiating astrogliogenesis (Bonni et al., 1997; Krichevsky et al., 2006).

miR-200 family members, including miR-200a, miR-200b, miR-200c, miR-141 and miR-429, are highly expressed in the developing olfactory bulb. Loss of function of the *miR-200* family prevents normal differentiation of olfactory precursors into mature neurons (Choi et al., 2008). *Foxg1*, *Zfhx1* and *Lfng* have been identified as the targets of the miR-200 family that affect neurogenesis of the olfactory bulb.

Specific miRNAs that promote gliogenesis have also been identified. Brain-enriched miR-125b is up-regulated in cultured interleukin-6 (IL-6)-induced human astrocytes. Loss of function of *miR-125b* causes an impaired proliferation of astrocytes, accompanied by an upregulation of a miR-125b target cyclin-dependent kinase inhibitor 2A (CDKN2A), which is a negative modulator for cell proliferation (Pogue et al., 2010). The *miR-17-92* cluster displays enriched expression in cultured oligodendrocytes. Specific deletion of the *miR-17-92* cluster from oligodendrocyte precursor cells (OPCs) results in a decreased number of Olig2-positive oligodendrocytes in the mouse brain (Budde et al., 2010). Overexpression of *miR-17* and *miR-19b* in cultures increases the number of oligodendrocytes. The *miR-17-92* cluster regulates oligodendrocyte development by targeting tumor suppressor *Pten* and activating its downstream Akt signaling pathway.

Moreover, *miR-219* and *miR-338* are identified in the oligodendrocyte lineage in the mouse spinal cord and brain. Overexpression of *miR-219* and *miR-338* in cultured OPCs and in the embryonic chick neural tube promotes differentiation of oligodendrocytes, while knockdown of these two miRNAs in OPC cultures and knockdown of *miR-219* in zebrafish abolish oligodendrocyte maturation (Zhao et al., 2010b). Oligodendrocyte differentiation inhibitors Sox6 and Hes5 are identified as targets of miR-219 and miR-338 during oligodendrocyte development (Zhao et al., 2010b).

Lamin B1 (LMNB1) is reported to be associated with autosomal domination leukodystrophy disease (ADLD), a CNS demyelination disorder (Padiath et al., 2006). Overexpression of Lamin B1 represses expression of oligodendrocyte-specific genes such as myelin basic protein (MBP) and myelin oligodendrocyte glycoprotein (MOG), and leads to impaired oligodendrocyte maturation. A recent study has shown that Lamin B1 is post-transcriptionally regulated by miR-23, a glia-specific miRNA. Overexpression of *miR-23*

results in significantly increased number of oligodendrocytes and rescues the defects of oligodendrocyte differentiation caused by Lamin B1 (Lin and Fu, 2009).

3.5.2 Cell fate determination regulated by lncRNAs

Studies of lncRNA functions on NSC differentiation are emerging. *Dlx-6* is a homeobox containing transcription factor and plays an important role in forebrain neurogenesis (Wang et al., 2010). *Embryonic ventral forebrain-1 (Evf1)* is a 2.7 kb lncRNA transcribed upstream of the mouse *Dlx-6* gene (Kohtz and Fishell, 2004). As an alternatively spliced form of *Evf1*, *Evf2* is transcribed from the intergenic region between the *Dlx-5* and *Dlx-6* loci, and is overlapped with the conserved *Dlx-5/6* intergenic enhancer (Feng et al., 2006; Zerucha et al., 2000) (Fig. 5B). Induced by the Sonic hedgehog (Shh) signaling pathway, *Evf2* has been proven to function as a transcriptional co-activator of *Dlx-2* and activates the *Dlx5/6* enhancer during forebrain development (Feng et al., 2006). Deletion of *Evf2* results in a reduction of GABAergic interneurons and impaired synaptic inhibition in the developing hippocampus (Bond et al., 2009).

Nkx2.2 antisense (Nkx2.2as) is an antisense lncRNA to *Nkx2.2* gene, which is expressed in the developing mammalian forebrain and is required for oligodendrocyte development (Price et al., 1992) (Fig. 5C). Ectopic expression of *Nkx2.2as* in cultured NSCs induces oligodendrocyte differentiation through an upregulation of the *Nkx2.2* mRNA level, suggesting that *Nkx2.2as* regulates NSC differentiation and promotes gliogenesis by modulating protein coding gene *Nkx2.2* expression (Tochitani and Hayashizaki, 2008).

Retinal noncoding RNA 2 (RNCR2), an intergenic lncRNA also known as *Gomafu* and *Miat*, is an abundant polyadenylated RNA in the developing retina (Blackshaw et al., 2004). *RNCR2* is highly expressed in both mitotic and postmitotic retinal progenitors. Knockdown of *RNCR2* leads to an increase of amacrine cells and Müller glial cells in postnatal retina. Mislocalization of *RNCR2* from nuclear to cytoplasm photocopies the effects caused by *RNCR2* knockdown, suggesting that *RNCR2* is required for retinal precursor cell specification (Rapicavoli et al., 2010).

4. Noncoding RNAs as a tool for stem cell-based therapy

Because of the features of self-renewal and the ability to differentiate into many cell types in the CNS, applying NSCs for the treatment of neurological disorders, especially neurodegeneration diseases and injuries in the CNS, has become promising. Directing NSCs into specific cell types and transplanting these cells to replace damaged cells in the CNS have been proven to be successful in some mouse models (Kim and de Vellis, 2009).

Transplantation of NSCs into aged triple transgenic Alzheimer's disease mouse model (3×Tg-AD) rescues the spatial learning and memory defects in these mice (Blurton-Jones et al., 2009). Parkinson's disease (PD) results from a loss of dopaminergic neurons in the substantia nigra. It involves abnormalities in movement variably accompanied by sensory, mood and cognitive changes. Transplantation of undifferentiated human NSCs into PD primate models causes a significant behavioral improvement (Redmond et al., 2007). Directed differentiation of mouse ventral midbrain NSCs in the presence of Shh, FGF8 and Wnt5a produce 10-fold more dopaminergic neurons *in vitro* (Parish et al., 2008).

Transplantation of these pre-differentiated dopaminergic neurons into the brain of PD mouse models results in functional recovery (Parish et al., 2008). Implantation of human NSCs in the rat model of Huntington's disease (HD) is shown improved motor function (McBride et al., 2004). Furthermore, delayed transplantation of adult mouse NSCs surrounding the lesion site of the spinal cord promotes remyelination and functional recovery after spinal cord injuries in rats (Karimi-Abdolrezaee et al., 2006).

Stem cell-based therapeutic applications for neurological disorders also face problems. First, the molecular mechanisms that control NSC proliferation and differentiation into distinct cell types are still unclear. Second, to succeed in clinical applications, transplanting sufficient numbers of NSCs and specific neuronal cell types is critical. Third, to achieve functional recovery from neurological disorders, transplanted cells need to acquire connections with neighbor neurons and restore neural circuitry. Although little studies of using ncRNAs for therapeutic treatment have been done, the emerging reports of ncRNA functions in NSC proliferation and cell fate determination have shown promising future directions. Moreover, due to the technical advances in ncRNA *in vitro* synthesis and delivery, particularly miRNAs, manipulating ncRNA expressions in NSCs will provide a new means for stem cell based therapies for neurological diseases.

5. Acknowledgment

We thank members of the Sun lab for providing thoughtful comments. Owing to space limitations, we apologize for being unable to cite many excellent papers in this field. This work was supported by the Ellison Medical Foundation (T. S.), an award from the Hirschl/Weill-Caulier Trust (T. S.) and an R01-MH083680 grant from the NIH/NIMH (T. S.).

6. References

- Amaral PP, Neyt C, Wilkins SJ, Askarian-Amiri ME, Sunkin SM, Perkins AC, et al. Complex architecture and regulated expression of the Sox2ot locus during vertebrate development. *RNA* 2009; 15: 2013-27.
- Ambros V. The evolution of our thinking about microRNAs. *Nat Med* 2008; 14: 1036-40.
- Andersson T, Rahman S, Sansom SN, Alsio JM, Kaneda M, Smith J, et al. Reversible block of mouse neural stem cell differentiation in the absence of dicer and microRNAs. *PLoS One* 2010; 5: e13453.
- Arvanitis DN, Jungas T, Behar A, Davy A. Ephrin-B1 reverse signaling controls a posttranscriptional feedback mechanism via miR-124. *Mol Cell Biol* 2010; 30: 2508-17.
- Bian S, Sun T. Functions of Noncoding RNAs in Neural Development and Neurological Diseases. *Mol Neurobiol* 2011; 44: 359-73.
- Blackshaw S, Harpavat S, Trimarchi J, Cai L, Huang H, Kuo WP, et al. Genomic analysis of mouse retinal development. *PLoS Biol* 2004; 2: E247.
- Blurton-Jones M, Kitazawa M, Martinez-Coria H, Castello NA, Muller FJ, Loring JF, et al. Neural stem cells improve cognition via BDNF in a transgenic model of Alzheimer disease. *Proc Natl Acad Sci U S A* 2009; 106: 13594-9.
- Bond AM, Vangompel MJ, Sametsky EA, Clark MF, Savage JC, Disterhoft JF, et al. Balanced gene regulation by an embryonic brain ncRNA is critical for adult hippocampal GABA circuitry. *Nat Neurosci* 2009; 12: 1020-7.

- Bonev B, Pisco A, Papalopulu N. MicroRNA-9 reveals regional diversity of neural progenitors along the anterior-posterior axis. *Dev Cell* 2011; 20: 19-32.
- Bonni A, Sun Y, Nadal-Vicens M, Bhatt A, Frank DA, Rozovsky I, et al. Regulation of gliogenesis in the central nervous system by the JAK-STAT signaling pathway. *Science* 1997; 278: 477-83.
- Brennecke J, Malone CD, Aravin AA, Sachidanandam R, Stark A, Hannon GJ. An epigenetic role for maternally inherited piRNAs in transposon silencing. *Science* 2008; 322: 1387-92.
- Brett JO, Renault VM, Rafalski VA, Webb AE, Brunet A. The microRNA cluster miR-106b~25 regulates adult neural stem/progenitor cell proliferation and neuronal differentiation. *Aging (Albany NY)* 2011; 3: 108-24.
- Budde H, Schmitt S, Fitzner D, Opitz L, Salinas-Riester G, Simons M. Control of oligodendroglial cell number by the miR-17-92 cluster. *Development* 2010; 137: 2127-32.
- Carninci P, Kasukawa T, Katayama S, Gough J, Frith MC, Maeda N, et al. The transcriptional landscape of the mammalian genome. *Science* 2005; 309: 1559-63.
- Cheng LC, Pastrana E, Tavazoie M, Doetsch F. miR-124 regulates adult neurogenesis in the subventricular zone stem cell niche. *Nat Neurosci* 2009; 12: 399-408.
- Choi PS, Zakhary L, Choi WY, Caron S, Alvarez-Saavedra E, Miska EA, et al. Members of the miRNA-200 family regulate olfactory neurogenesis. *Neuron* 2008; 57: 41-55.
- Cuellar TL, Davis TH, Nelson PT, Loeb GB, Harfe BD, Ullian E, et al. Dicer loss in striatal neurons produces behavioral and neuroanatomical phenotypes in the absence of neurodegeneration. *Proc Natl Acad Sci U S A* 2008; 105: 5614-9.
- Das PP, Bagijn MP, Goldstein LD, Woolford JR, Lehrbach NJ, Sapetschnig A, et al. Piwi and piRNAs act upstream of an endogenous siRNA pathway to suppress Tc3 transposon mobility in the *Caenorhabditis elegans* germline. *Mol Cell* 2008; 31: 79-90.
- Davis TH, Cuellar TL, Koch SM, Barker AJ, Harfe BD, McManus MT, et al. Conditional loss of Dicer disrupts cellular and tissue morphogenesis in the cortex and hippocampus. *J Neurosci* 2008; 28: 4322-30.
- De Pietri Tonelli D, Pulvers JN, Haffner C, Murchison EP, Hannon GJ, Huttner WB. miRNAs are essential for survival and differentiation of newborn neurons but not for expansion of neural progenitors during early neurogenesis in the mouse embryonic neocortex. *Development* 2008; 135: 3911-21.
- Decembrini S, Bressan D, Vignali R, Pitto L, Mariotti S, Rainaldi G, et al. MicroRNAs couple cell fate and developmental timing in retina. *Proc Natl Acad Sci U S A* 2009; 106: 21179-84.
- Delaloy C, Liu L, Lee JA, Su H, Shen F, Yang GY, et al. MicroRNA-9 coordinates proliferation and migration of human embryonic stem cell-derived neural progenitors. *Cell Stem Cell* 2010; 6: 323-35.
- Denisenko O, Shnyreva M, Suzuki H, Bomsztyk K. Point mutations in the WD40 domain of Eed block its interaction with Ezh2. *Mol Cell Biol* 1998; 18: 5634-42.
- Doe CQ. Neural stem cells: balancing self-renewal with differentiation. *Development* 2008; 135: 1575-87.
- Feng J, Bi C, Clark BS, Mady R, Shah P, Kohtz JD. The Evf-2 noncoding RNA is transcribed from the Dlx-5/6 ultraconserved region and functions as a Dlx-2 transcriptional coactivator. *Genes Dev* 2006; 20: 1470-84.

- Fukagawa T, Nogami M, Yoshikawa M, Ikeno M, Okazaki T, Takami Y, et al. Dicer is essential for formation of the heterochromatin structure in vertebrate cells. *Nat Cell Biol* 2004; 6: 784-91.
- Gage FH. Mammalian neural stem cells. *Science* 2000; 287: 1433-8.
- Gaughwin P, Ciesla M, Yang H, Lim B, Brundin P. Stage-Specific Modulation of Cortical Neuronal Development by Mmu-miR-134. *Cereb Cortex* 2011.
- Giam M, Huang DC, Bouillet P. BH3-only proteins and their roles in programmed cell death. *Oncogene* 2008; 27 Suppl 1: S128-36.
- Gregory RI, Chendrimada TP, Shiekhattar R. MicroRNA biogenesis: isolation and characterization of the microprocessor complex. *Methods Mol Biol* 2006; 342: 33-47.
- Guidi M, Muinos-Gimeno M, Kagerbauer B, Marti E, Estivill X, Espinosa-Parrilla Y. Overexpression of miR-128 specifically inhibits the truncated isoform of NTRK3 and upregulates BCL2 in SH-SY5Y neuroblastoma cells. *BMC Mol Biol* 2010; 11: 95.
- Hebert SS, Papadopoulou AS, Smith P, Galas MC, Planel E, Silahtaroglu AN, et al. Genetic ablation of Dicer in adult forebrain neurons results in abnormal tau hyperphosphorylation and neurodegeneration. *Hum Mol Genet* 2010; 19: 3959-69.
- Houwing S, Kamminga LM, Berezikov E, Cronembold D, Girard A, van den Elst H, et al. A role for Piwi and piRNAs in germ cell maintenance and transposon silencing in Zebrafish. *Cell* 2007; 129: 69-82.
- Huang J, Wu S, Barrera J, Matthews K, Pan D. The Hippo signaling pathway coordinately regulates cell proliferation and apoptosis by inactivating Yorkie, the Drosophila Homolog of YAP. *Cell* 2005; 122: 421-34.
- Kanellopoulou C, Muljo SA, Kung AL, Ganesan S, Drapkin R, Jenuwein T, et al. Dicer-deficient mouse embryonic stem cells are defective in differentiation and centromeric silencing. *Genes Dev* 2005; 19: 489-501.
- Kapranov P, Cheng J, Dike S, Nix DA, Duttagupta R, Willingham AT, et al. RNA maps reveal new RNA classes and a possible function for pervasive transcription. *Science* 2007; 316: 1484-8.
- Karimi-Abdolrezaee S, Eftekharpour E, Wang J, Morshead CM, Fehlings MG. Delayed transplantation of adult neural precursor cells promotes remyelination and functional neurological recovery after spinal cord injury. *J Neurosci* 2006; 26: 3377-89.
- Kawase-Koga Y, Low R, Otaegi G, Pollock A, Deng H, Eisenhaber F, et al. RNAase-III enzyme Dicer maintains signaling pathways for differentiation and survival in mouse cortical neural stem cells. *J Cell Sci* 2010; 123: 586-94.
- Kawase-Koga Y, Otaegi G, Sun T. Different timings of Dicer deletion affect neurogenesis and gliogenesis in the developing mouse central nervous system. *Dev Dyn* 2009; 238: 2800-12.
- Khanna A, Muthusamy S, Liang R, Sarojini H, Wang E. Gain of survival signaling by down-regulation of three key miRNAs in brain of calorie-restricted mice. *Aging (Albany NY)* 2011; 3: 223-36.
- Khraiweh B, Arif MA, Seumel GI, Ossowski S, Weigel D, Reski R, et al. Transcriptional control of gene expression by microRNAs. *Cell* 2010; 140: 111-22.
- Kim SU, de Vellis J. Stem cell-based cell therapy in neurological diseases: a review. *J Neurosci Res* 2009; 87: 2183-200.
- Kohtz JD, Fishell G. Developmental regulation of EVF-1, a novel non-coding RNA transcribed upstream of the mouse *Dlx6* gene. *Gene Expr Patterns* 2004; 4: 407-12.

- Kole AJ, Swahari V, Hammond SM, Deshmukh M. miR-29b is activated during neuronal maturation and targets BH3-only genes to restrict apoptosis. *Genes Dev* 2011; 25: 125-30.
- Kotake Y, Nakagawa T, Kitagawa K, Suzuki S, Liu N, Kitagawa M, et al. Long non-coding RNA ANRIL is required for the PRC2 recruitment to and silencing of p15(INK4B) tumor suppressor gene. *Oncogene* 2010; 30: 1956-62.
- Krichevsky AM, Sonntag KC, Isacson O, Kosik KS. Specific microRNAs modulate embryonic stem cell-derived neurogenesis. *Stem Cells* 2006; 24: 857-64.
- Lagos-Quintana M, Rauhut R, Lendeckel W, Tuschl T. Identification of novel genes coding for small expressed RNAs. *Science* 2001; 294: 853-8.
- Lagos-Quintana M, Rauhut R, Yalcin A, Meyer J, Lendeckel W, Tuschl T. Identification of tissue-specific microRNAs from mouse. *Curr Biol* 2002; 12: 735-9.
- Lander ES, Linton LM, Birren B, Nusbaum C, Zody MC, Baldwin J, et al. Initial sequencing and analysis of the human genome. *Nature* 2001; 409: 860-921.
- Lau NC, Lim LP, Weinstein EG, Bartel DP. An abundant class of tiny RNAs with probable regulatory roles in *Caenorhabditis elegans*. *Science* 2001; 294: 858-62.
- Lau NC, Seto AG, Kim J, Kuramochi-Miyagawa S, Nakano T, Bartel DP, et al. Characterization of the piRNA complex from rat testes. *Science* 2006; 313: 363-7.
- Leaman D, Chen PY, Fak J, Yalcin A, Pearce M, Unnerstall U, et al. Antisense-mediated depletion reveals essential and specific functions of microRNAs in *Drosophila* development. *Cell* 2005; 121: 1097-108.
- Lee RC, Ambros V. An extensive class of small RNAs in *Caenorhabditis elegans*. *Science* 2001; 294: 862-4.
- Leucht C, Stigloher C, Wizenmann A, Klafke R, Folchert A, Bally-Cuif L. MicroRNA-9 directs late organizer activity of the midbrain-hindbrain boundary. *Nat Neurosci* 2008; 11: 641-8.
- Lin ST, Fu YH. miR-23 regulation of lamin B1 is crucial for oligodendrocyte development and myelination. *Dis Model Mech* 2009; 2: 178-88.
- Liu C, Teng ZQ, Santistevan NJ, Szulwach KE, Guo W, Jin P, et al. Epigenetic regulation of miR-184 by MBD1 governs neural stem cell proliferation and differentiation. *Cell Stem Cell* 2010; 6: 433-44.
- Lund E, Dahlberg JE. Substrate selectivity of exportin 5 and Dicer in the biogenesis of microRNAs. *Cold Spring Harb Symp Quant Biol* 2006; 71: 59-66.
- Maiorano NA, Mallamaci A. Promotion of embryonic cortico-cerebral neurogenesis by miR-124. *Neural Dev* 2009; 4: 40.
- Makeyev EV, Zhang J, Carrasco MA, Maniatis T. The MicroRNA miR-124 promotes neuronal differentiation by triggering brain-specific alternative pre-mRNA splicing. *Mol Cell* 2007; 27: 435-48.
- Mancini-Dinardo D, Steele SJ, Levorse JM, Ingram RS, Tilghman SM. Elongation of the Kcnq1ot1 transcript is required for genomic imprinting of neighboring genes. *Genes Dev* 2006; 20: 1268-82.
- Mazo A, Hodgson JW, Petruk S, Sedkov Y, Brock HW. Transcriptional interference: an unexpected layer of complexity in gene regulation. *J Cell Sci* 2007; 120: 2755-61.
- McBride JL, Behrstock SP, Chen EY, Jakel RJ, Siegel I, Svendsen CN, et al. Human neural stem cell transplants improve motor function in a rat model of Huntington's disease. *J Comp Neurol* 2004; 475: 211-9.

- Mercer TR, Dinger ME, Sunkin SM, Mehler MF, Mattick JS. Specific expression of long noncoding RNAs in the mouse brain. *Proc Natl Acad Sci U S A* 2008; 105: 716-21.
- Neilson JR, Sharp PA. Small RNA regulators of gene expression. *Cell* 2008; 134: 899-902.
- Nguyen VT, Kiss T, Michels AA, Bensaude O. 7SK small nuclear RNA binds to and inhibits the activity of CDK9/cyclin T complexes. *Nature* 2001; 414: 322-5.
- Otaegi G, Pollock A, Hong J, Sun T. MicroRNA miR-9 modifies motor neuron columns by a tuning regulation of FoxP1 levels in developing spinal cords. *J Neurosci* 2011; 31: 809-18.
- Padiath QS, Saigoh K, Schiffmann R, Asahara H, Yamada T, Koeppen A, et al. Lamin B1 duplications cause autosomal dominant leukodystrophy. *Nat Genet* 2006; 38: 1114-23.
- Parish CL, Castelo-Branco G, Rawal N, Tonnesen J, Sorensen AT, Salto C, et al. Wnt5a-treated midbrain neural stem cells improve dopamine cell replacement therapy in parkinsonian mice. *J Clin Invest* 2008; 118: 149-60.
- Pasmant E, Laurendeau I, Heron D, Vidaud M, Vidaud D, Bieche I. Characterization of a germ-line deletion, including the entire INK4/ARF locus, in a melanoma-neural system tumor family: identification of ANRIL, an antisense noncoding RNA whose expression coclusters with ARF. *Cancer Res* 2007; 67: 3963-9.
- Pevny L, Placzek M. SOX genes and neural progenitor identity. *Curr Opin Neurobiol* 2005; 15: 7-13.
- Pogue AI, Cui JG, Li YY, Zhao Y, Culicchia F, Lukiw WJ. Micro RNA-125b (miRNA-125b) function in astrogliosis and glial cell proliferation. *Neurosci Lett* 2010; 476: 18-22.
- Pratt AJ, MacRae IJ. The RNA-induced silencing complex: a versatile gene-silencing machine. *J Biol Chem* 2009; 284: 17897-901.
- Price M, Lazzaro D, Pohl T, Mattei MG, Ruther U, Olivo JC, et al. Regional expression of the homeobox gene Nkx-2.2 in the developing mammalian forebrain. *Neuron* 1992; 8: 241-55.
- Rana TM. Illuminating the silence: understanding the structure and function of small RNAs. *Nat Rev Mol Cell Biol* 2007; 8: 23-36.
- Rapicavoli NA, Poth EM, Blackshaw S. The long noncoding RNA RNCR2 directs mouse retinal cell specification. *BMC Dev Biol* 2010; 10: 49.
- Redmond DE, Jr., Bjugstad KB, Teng YD, Ourednik V, Ourednik J, Wakeman DR, et al. Behavioral improvement in a primate Parkinson's model is associated with multiple homeostatic effects of human neural stem cells. *Proc Natl Acad Sci U S A* 2007; 104: 12175-80.
- Reinhart BJ, Slack FJ, Basson M, Pasquinelli AE, Bettinger JC, Rougvie AE, et al. The 21-nucleotide let-7 RNA regulates developmental timing in *Caenorhabditis elegans*. *Nature* 2000; 403: 901-6.
- Renault VM, Rafalski VA, Morgan AA, Salih DA, Brett JO, Webb AE, et al. FoxO3 regulates neural stem cell homeostasis. *Cell Stem Cell* 2009; 5: 527-39.
- Rinn JL, Kertesz M, Wang JK, Squazzo SL, Xu X, Brugmann SA, et al. Functional demarcation of active and silent chromatin domains in human HOX loci by noncoding RNAs. *Cell* 2007; 129: 1311-23.
- Rybak A, Fuchs H, Smirnova L, Brandt C, Pohl EE, Nitsch R, et al. A feedback loop comprising lin-28 and let-7 controls pre-let-7 maturation during neural stem-cell commitment. *Nat Cell Biol* 2008; 10: 987-93.

- Sanchez-Elsner T, Gou D, Kremmer E, Sauer F. Noncoding RNAs of trithorax response elements recruit Drosophila Ash1 to Ultrabithorax. *Science* 2006; 311: 1118-23.
- Sathyan P, Golden HB, Miranda RC. Competing interactions between micro-RNAs determine neural progenitor survival and proliferation after ethanol exposure: evidence from an ex vivo model of the fetal cerebral cortical neuroepithelium. *J Neurosci* 2007; 27: 8546-57.
- Saucedo LJ, Edgar BA. Filling out the Hippo pathway. *Nat Rev Mol Cell Biol* 2007; 8: 613-21.
- Seto AG, Kingston RE, Lau NC. The coming of age for Piwi proteins. *Mol Cell* 2007; 26: 603-9.
- Shamovsky I, Ivannikov M, Kandel ES, Gershon D, Nudler E. RNA-mediated response to heat shock in mammalian cells. *Nature* 2006; 440: 556-60.
- Shi Y, Sun G, Zhao C, Stewart R. Neural stem cell self-renewal. *Crit Rev Oncol Hematol* 2008; 65: 43-53.
- Shibata M, Nakao H, Kiyonari H, Abe T, Aizawa S. MicroRNA-9 regulates neurogenesis in mouse telencephalon by targeting multiple transcription factors. *J Neurosci* 2011; 31: 3407-22.
- Smirnova L, Grafe A, Seiler A, Schumacher S, Nitsch R, Wulczyn FG. Regulation of miRNA expression during neural cell specification. *Eur J Neurosci* 2005; 21: 1469-77.
- Szulwach KE, Li X, Smrt RD, Li Y, Luo Y, Lin L, et al. Cross talk between microRNA and epigenetic regulation in adult neurogenesis. *J Cell Biol* 2010; 189: 127-41.
- Temple S. The development of neural stem cells. *Nature* 2001; 414: 112-7.
- Thompson BJ, Cohen SM. The Hippo pathway regulates the bantam microRNA to control cell proliferation and apoptosis in Drosophila. *Cell* 2006; 126: 767-74.
- Tochitani S, Hayashizaki Y. Nkx2.2 antisense RNA overexpression enhanced oligodendrocytic differentiation. *Biochem Biophys Res Commun* 2008; 372: 691-6.
- Tsai MC, Manor O, Wan Y, Mosammamaparast N, Wang JK, Lan F, et al. Long noncoding RNA as modular scaffold of histone modification complexes. *Science* 2010; 329: 689-93.
- Walker JC, Harland RM. microRNA-24a is required to repress apoptosis in the developing neural retina. *Genes Dev* 2009; 23: 1046-51.
- Wang X, Arai S, Song X, Reichart D, Du K, Pascual G, et al. Induced ncRNAs allosterically modify RNA-binding proteins in cis to inhibit transcription. *Nature* 2008; 454: 126-30.
- Wang XJ, Reyes JL, Chua NH, Gaasterland T. Prediction and identification of Arabidopsis thaliana microRNAs and their mRNA targets. *Genome Biol* 2004; 5: R65.
- Wang Y, Dye CA, Sohal V, Long JE, Estrada RC, Roztocil T, et al. Dlx5 and Dlx6 regulate the development of parvalbumin-expressing cortical interneurons. *J Neurosci* 2010; 30: 5334-45.
- Willingham AT, Orth AP, Batalov S, Peters EC, Wen BG, Aza-Blanc P, et al. A strategy for probing the function of noncoding RNAs finds a repressor of NFAT. *Science* 2005; 309: 1570-3.
- Wutz A, Rasmussen TP, Jaenisch R. Chromosomal silencing and localization are mediated by different domains of Xist RNA. *Nat Genet* 2002; 30: 167-74.
- Yang Z, Zhu Q, Luo K, Zhou Q. The 7SK small nuclear RNA inhibits the CDK9/cyclin T1 kinase to control transcription. *Nature* 2001; 414: 317-22.
- Yu W, Gius D, Onyango P, Muldoon-Jacobs K, Karp J, Feinberg AP, et al. Epigenetic silencing of tumour suppressor gene p15 by its antisense RNA. *Nature* 2008; 451: 202-6.
- Zehir A, Hua LL, Maska EL, Morikawa Y, Cserjesi P. Dicer is required for survival of differentiating neural crest cells. *Dev Biol* 2010; 340: 459-67.

- Zerucha T, Stuhmer T, Hatch G, Park BK, Long Q, Yu G, et al. A highly conserved enhancer in the *Dlx5/Dlx6* intergenic region is the site of cross-regulatory interactions between *Dlx* genes in the embryonic forebrain. *J Neurosci* 2000; 20: 709-21.
- Zhao C, Sun G, Li S, Lang MF, Yang S, Li W, et al. MicroRNA let-7b regulates neural stem cell proliferation and differentiation by targeting nuclear receptor TLX signaling. *Proc Natl Acad Sci U S A* 2010a; 107: 1876-81.
- Zhao C, Sun G, Li S, Shi Y. A feedback regulatory loop involving microRNA-9 and nuclear receptor TLX in neural stem cell fate determination. *Nat Struct Mol Biol* 2009; 16: 365-71.
- Zhao X, He X, Han X, Yu Y, Ye F, Chen Y, et al. MicroRNA-mediated control of oligodendrocyte differentiation. *Neuron* 2010b; 65: 612-26.
- Zheng K, Li H, Zhu Y, Zhu Q, Qiu M. MicroRNAs are essential for the developmental switch from neurogenesis to gliogenesis in the developing spinal cord. *J Neurosci* 2010; 30: 8245-50.

IntechOpen

Neural Stem Cells and Therapy

Edited by Dr. Tao Sun

ISBN 978-953-307-958-5

Hard cover, 440 pages

Publisher InTech

Published online 15, February, 2012

Published in print edition February, 2012

This book is a collective work of international experts in the neural stem cell field. The book incorporates the characterization of embryonic and adult neural stem cells in both invertebrates and vertebrates. It highlights the history and the most advanced discoveries in neural stem cells, and summarizes the mechanisms of neural stem cell development. In particular, this book provides strategies and discusses the challenges of utilizing neural stem cells for therapy of neurological disorders and brain and spinal cord injuries. It is suitable for general readers, students, doctors and researchers who are interested in understanding the principles of and new discoveries in neural stem cells and therapy.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Shan Bian and Tao Sun (2012). Noncoding RNAs in Neural Stem Cell Development, Neural Stem Cells and Therapy, Dr. Tao Sun (Ed.), ISBN: 978-953-307-958-5, InTech, Available from:
<http://www.intechopen.com/books/neural-stem-cells-and-therapy/noncoding-rnas-in-neural-stem-cell-development>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](https://creativecommons.org/licenses/by/3.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen