We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Artificial Insemination and Its Role in Transmission of Swine Viruses

Tanja Opriessnig, Luis G. Giménez-Lirola and Patrick G. Halbur
*Department of Veterinary Diagnostic and Production Animal Medicine,
 College of Veterinary Medicine, Iowa State University, Ames, Iowa,
 USA*

1. Introduction

Artificial insemination (AI) in swine is not a new technique and reports as early as the 1930s (Lush, 1925) describe collecting semen for AI. However, because of farm structure changes, increasing farm sizes and separation of production stages, interest in intensive pig production is growing and AI has become a critical component in modern pig production. In 2001, nearly 60% of North American swine herds utilized AI (Singleton, 2001), a drastic increase from the estimated 5% in the 1990's (Flowers & Esbenshade, 1993). This is still relatively low compared to the 90% or greater use of AI in Western Europe (Madsen, 2005; Maes et al., 2008). The extensive use of AI in pig reproduction in the last decade has facilitated the exchange of desirable genetic characteristics at an international level, allowing producers to make greater use of superior genetics at a lower cost than some natural-service systems (Gerrits et al., 2005). However, the growth in use of AI has increased the risk of quick and widespread transmission of venereally transmissible pathogens (Thacker et al., 1984). It has been reported that the porcine male reproductive tract is highly susceptible to viral infections (Phillips et al., 1972; Spradbrow, 1968). This, coupled with the ability of boars to produce tens to thousands of insemination doses per week and the widespread distribution of the processed semen (both nationally and internationally), further increases the risk of wide transmission of viral pathogens by semen.

Many viruses have been reported to be present in boar semen (Yaeger et al., 1993; Lucas et al., 1974; Madson et al., 2008) and have the potential of being transmitted to susceptible breeding animals during AI, particularly when the boar is viremic or clinically sick. Viral shedding may also continue long after clinical signs have abated. In addition, infertility or reduced reproductive performance have been reported in boars with detectable virus in their semen (Guerin & Pozzi, 2005; Larsen et al., 1980). The potential economic impact and liability associated with transmission of diseases through semen has created great interest and investment in testing boars for viral diseases prior to entry and while at AI centers.

In many countries, foot-and-mouth disease virus (FMDV), porcine reproductive and respiratory syndrome virus (PRRSV), Japanese B encephalitis virus (JBEV), pseudorabies virus (PRV), classical swine fever virus (CSFV) and African swine fever virus (ASFV) are of particular importance. Accurate monitoring of boars in AI stations is essential to reduce the

risk of virus transmission. Monitoring for the presence of other ubiquitous viruses that can be found in semen, such as porcine circovirus type 2 (PCV2), torque teno sus virus (TTSuV), porcine parvovirus (PPV) and porcine adenovirus (PAV) is commonly not a high priority in most countries despite the fact that some of these viruses have been associated with sporadic reproductive failure in breeding animals.

Due to differences in pathogenicity, economic consequences, geographic localization, and epidemiological parameters, different viruses are commonly assigned different levels of importance. The aims of this review are to summarize the information on swine viruses that can be transmitted via AI, their shedding characteristics in boars, effects of these viruses on naïve breeding animals, testing that is currently done to reduce or prevent the risk of transmission, and current best practices being used to reduce or eliminate virus shedding in semen.

2. Impact of viral contamination of semen

Microbial contamination of boar semen not only reduces semen quality but can also reduce conception rate resulting in considerable economic loss. It is not uncommon that a single boar stud serves multiple breeding herds with an average of 200 to 10,000 sows on each site. The risk for breeding herds becoming infected via AI increases with the number of pathogens present in the semen, the dose of the pathogen(s) in the semen, the number of sows inseminated with the contaminated semen, and the level of protective immunity in the breeding herd. From a practical point of view, most economically important viral diseases are associated with clinical signs, so semen collection from sick boars will not always take place, and consequently the risk of pathogen transmission into susceptible breeding herds by this route may be lower than expected. However, when clinical signs are mild or absent the risk of viral transmission via contaminated semen is a concern.

Artificial insemination is now widely recognized as a route for spread of swine diseases (Thacker et al., 1984) and continues to be a major concern to pig breeders and regulatory authorities in countries where AI is practiced. Many viruses known to contaminate semen have been shown to be highly infectious via the uterine route, and wide dissemination of porcine pathogens by semen could occur when semen is shipped globally. Transmission of viruses via the semen route to breeding females has been experimentally proven for CSFV (de Smit et al., 1999), PRRSV (Yaeger et al., 1993), and PPV (Lucas et al., 1974). It is of importance to note that, although virus-contaminated semen indeed constitutes a serious risk for transmission, it does not guarantee that transmission to the sow or gilt by AI will consistently occur (Swenson et al., 1994b; Yaeger et al., 1993). The conditions required for establishment of infection in the breeding female are complex, and lack of transmission might be explained by sow immunity or failure to reach the minimum infectious dose. In this regard, much research has been conducted concerning the risk of transmission of PRRSV by semen, and the minimum dose necessary to establish infection in the sow (Benfield et al., 2000; Prieto et al., 1996b).

Contamination of semen with virus can occur through three possible routes: 1) fecal contamination of semen during collection, 2) systemic viral infection, or 3) local viral infection (testes, accessory gland, etc.). Once virus-contaminated semen is utilized to inseminate a breeding female, the outcome can vary depending on the time of fetal

infection. Early embryonic death may result from direct invasion of the embryo by the pathogen or by uterine epithelial alterations in response to the pathogen (Wrathall & Mengeling, 1979). Until six to seven days after conception, an embryo is surrounded and protected by an impervious barrier called the zona pellucida, which helps the embryo avoid pathogen invasion (Mateusen et al., 2004). However, after entering the blastocyst stage, embryos may become susceptible to viral infections. Fetuses that are infected prior to 70 days of gestation usually die (mummies) and contribute to smaller litter sizes or early termination of pregnancy. Fetuses that are infected around or after 70 days of gestation are able to mount an active immune response against the pathogen and survive intrauterine infection.

As the risk of rapid virus spread via semen is well known, sanitation protocols in AI centers are constantly being reviewed to assure customers that current best practices are in place. A new categorization of viruses has been proposed to further address different risk levels (Guerin & Pozzi, 2005): Category 1 includes viruses for which there is scientific proof of transmission by semen, but without any risks for AI due to official national eradication programs such as FMDV, PRV, CSFV, and ASFV. Category 2 includes viruses for which there is scientific proof of transmission by semen and which can be commonly encountered such as PRRSV. Category 3 includes viruses for which additional scientific proof is needed to better assess the risk of transmission by semen (PCV2, porcine rubulavirus). In addition, some viruses have not been shown to be present in semen and thus are considered non-hazardous to AI.

3. Swine viruses that can be present in boar semen

3.1 Porcine Reproductive and Respiratory Syndrome Virus

Porcine reproductive and respiratory syndrome virus (PRRSV) is a single-stranded, positive-sense, enveloped RNA virus that belongs to the family *Arteriviridae*, genus *Arterivirus* (Cavanagh, 1997). PRRSV was first recognized in the late 1980's (Cavanagh, 1997; Meulenberg et al., 1993) and today is found globally in swine producing countries. As the name of the virus implies, PRRSV infection is associated with reproductive failure in pregnant sows and respiratory disease in pigs of all ages. It is less commonly associated with neonatal diarrhea (Albina et al., 1994; Bierk et al., 2001; Neumann et al., 2005; Rossow, 1998; Wensvoort et al., 1991). Although clinical disease associated with PRRSV in growing pigs can be quite severe, no or mild symptoms are typically seen in boars (Wensvoort et al., 1991). One of the main characteristics of PRRSV is its high transmissibility, making it difficult to maintain pig populations free of PRRSV (Prieto & Castro, 2005). The 50% tissue culture infective dose (TCID₅₀) for exposure via oral, intranasal and AI routes was determined to be 10^{5.3}, 10^{4.0} and 10^{4.5}, respectively (Benfield et al., 2000). Experimental infection in boars has demonstrated seminal shedding of PRRSV (Prieto et al., 1996b; Swenson et al., 1994a), and epidemiological evidence confirms that transmission of PRRSV from fresh semen of acutely infected boars into breeding herds is possible (Robertson, 1992; Yaeger et al., 1993). However, successful transmission is not always achieved (Swenson et al., 1994b; Yaeger et al., 1993) and appears to depend largely on the amount of PRRSV present in the semen (Benfield et al., 2000). In semen of adult boars, PRRSV can persist for variable periods (Christopher-Hennings et al., 1995a; Swenson et al., 1994) suggesting that the virus continues to replicate in the one or more tissues of the reproductive tract. Several

studies have since been carried out to determine the temporal localization of the virus in different organs and tissues (Prieto et al., 2003; Sur et al., 1997) and vascular dissemination and replication of PRRSV in tissues of the reproductive tract was confirmed (Sur et al., 1997). In addition, migration of infected monocytes and macrophages directly from blood and lymph of the reproductive tract into the semen has also been suggested as a mechanism for PRRSV contamination of semen (Christopher-Hennings et al., 1998; Prieto et al., 2003). PRRSV has been detected in semen samples of experimentally infected boars ranging from 4 to 92 days post infection (Christopher-Hennings et al., 1995a, 1998). This marked variability may be due to individual host factors (Christopher-Hennings et al., 1995a, 2001; Swenson et al., 1994a), the breed of the boars (Christopher-Hennings et al., 2001), and the strain of virus used in the experimental inoculation. European and North American PRRSV strains differ genetically and antigenically (Bautista et al., 1993; Meng et al., 1995). A bioassay has been used to determine the presence of infectious PRRSV in semen samples for a period of time lasting from 4 to 42 days post infection (Swenson et al., 1994a). In semen, PRRSV is sometimes associated with drastic changes in quality (reduced motility, abnormal acrosomes, morphological alterations, etc.) and volume of the semen (Teuffert et al., 1998). Conversely, some researchers have found that the quality of semen remains within normal limits after infection with PRRSV (Swenson et al., 1994a; Yaeger et al., 1993). The etiological role of PRRSV in reproductive failure of swine is firmly established (Christianson et al., 1993; Lager & Mengeling, 1995; Mengeling et al., 1994; Prieto et al., 1996a; Terpstra et al., 1991). The effect of PRRSV on reproductive parameters was found to be highly related to strain pathogenicity (Prieto et al., 1997). Sows and gilts bred by infected boars (Gradil et al., 1996; Yaeger et al., 1993) or with experimentally contaminated semen (Prieto et al., 1997; Swenson et al., 1994b) typically show seroconversion to PRRSV, even in the absence of detectable viremia (Christopher-Hennings et al., 1995a, 2001). It has been reported that insemination of seronegative or pre-immunized gilts with boar semen containing PRRSV often has little to no effect on conception rates, but may result in early embryonic infection and death (Prieto et al., 1997). In addition, PRRSV can be a major cause of prenatal death in commercial swine herds and often preweaning mortality is also increased.

3.2 Porcine Parvovirus

Porcine parvovirus (PPV) is a small, single-stranded, non-enveloped DNA virus (Molitor et al., 1984) that belongs to the family *Parvoviridae*, genus *Parvovirus*. PPV is considered ubiquitous in the pig population. The main routes of transmission of PPV are oronasal and transplacental. In general, infection of growing pigs and mature boars with PPV alone is not associated with clinical disease (Allan et al., 2000; Brown, Jr. et al., 1980; Kennedy et al., 2000; Krakowka et al., 2000). There is evidence that PPV-isolates vary in pathogenicity (Mengeling & Cutlip, 1976; Oraveerakul et al., 1993). PPV-isolates have been classified as non-pathogenic (Mengeling & Cutlip, 1976; Paul et al., 1979), pathogenic to non-immunocompetent fetuses leading to death (Mengeling, 1979), pathogenic to immunocompetent fetuses and inducing dermatitis (Choi et al., 1987; Kresse et al., 1985; Lager et al., 1992; Lager & Mengeling, 1994), and enteric PPV-strains (Dea et al., 1985; Duhamel et al., 1991). PPV has previously been isolated from semen of naturally infected boars (Cartwright & Huck, 1967). Boars can shed the virus in semen during the acute phase of infection (Gradil et al., 1990); shedding beyond this phase has not been demonstrated, but the possibility of immunotolerant carriers of PPV as a result of early *in utero* infection has

been suggested (Cartwright et al., 1971). Semen may also become contaminated with PPV from feces or from the male reproductive organs (Biront & Bonte, 1983; Lucas et al., 1974). Changes in sperm output, ejaculate volume, motility, or morphologic defects have not been observed when the semen quality was evaluated after experimental inoculation of boars with PPV (Thacker et al., 1987a). Acute infection of breeding females with PPV is usually subclinical (Johnson et al., 1976). During initial infection, PPV replicates extensively and can be found in many tissues including lymphoid tissues (Cutlip & Mengeling, 1975a). The major and often only clinical response to PPV infection is reproductive failure. Return to estrus, fewer pigs per litter, and increased numbers of mummified fetuses are often observed (Cartwright & Huck, 1967; Joo et al., 1976). The consequence of PPV infection depends on timing of infection of the fetus: Death and reabsorption are usually observed in 10-30 gestation day embryos, death and mummification are seen when 30 to 70 day gestation fetuses are infected, and an active immune response and survival *in utero* is seen in fetuses when infected after 70 days of gestation (Bachmann et al., 1975; Cutlip & Mengeling, 1975b). The role of semen contamination with PPV leading to clinical reproductive problems has not been clearly established (Lucas et al., 1974; Mengeling & Paul, 1986; Thacker et al., 1987b; Wrathall & Mengeling, 1979).

3.3 Pseudorabies Virus

Pseudorabies virus (PRV) is a double-stranded DNA virus that belongs to the subfamily *Alphaherpesvirinae* of the family *Herpesvirus* (Mettenleiter, 2000). PRV is distributed worldwide; however, in recent years eradication efforts have been successful to eliminate PRV from the domestic pig population in parts of Europe, Canada, New Zealand and the United States. PRV replication typically occurs in the nasal and pharyngeal mucosa and primary transmission occurs via the nasal route. Viral replication has also been reported to take place in the genital tract and transmission by copulation is therefore possible (Hall, Jr. et al., 1984a; Vannier & Gueguen, 1979). Differences in pathogenicity and duration of virus shedding exist among PRV strains (Maes et al., 1983). Clinically affected boars are often unable to mount a dummy (Guerin & Pozzi, 2005) and can additionally develop respiratory disease (Hall, Jr. et al., 1984b). PRV can be isolated infrequently from urine, preputial membranes, and semen after either natural (Medveczky & Szabo, 1981) or experimental infection (Vannier & Gueguen, 1979). Very high viral concentrations have been reported in semen ranging from 10^8 to 10^9 TCID₅₀ per ml (Medveczky & Szabo, 1981) to $10^{3.7}$ to $10^{4.9}$ TCID₅₀ per ml (Vannier & Gueguen, 1979). Transient viral excretion for up to 12 days usually occurs during the acute phase of the disease (Ressang, 1973) but viral excretion has been observed for long periods of time after natural infection (Wittmann, 1985). After experimental PRV infection by the intratesticular route, testicular degeneration and transient elevation in sperm abnormalities have been reported (Hall, Jr. et al., 1984a; Larsen et al., 1980). Sows inseminated with contaminated semen may develop vaginitis or endometritis resulting in embryonic death (Maes et al., 2008). Sows infected in the first trimester of pregnancy often reabsorb their fetuses and return to estrus. If sows get infected in the later stages of pregnancy they may abort or have larger numbers of stillborn and weakborn piglets. If susceptible females are infected with PRV close to parturition, there are increased numbers of weakborn pigs with signs of nervous system disease.

3.4 Porcine Rubulavirus

Porcine rubulavirus (PoRV), also known as La-Piedad-Michoacan paramyxovirus (LPMV), is associated with blue eye disease in pigs. This virus was first isolated in Mexico in the early 1980's. PoRV belongs to the genus *Rubulavirus* and family *Paramyxoviridae*. Only one serotype has been recognized to date. Subclinically infected pigs are the primary source for PoRV transmission which occurs primarily via oronasal route. Boars, like other adult animals infected with PoRV, generally do not show clinical signs except for epididymitis and orchitis and in some cases loss of libido (Maes et al., 2008). Transmission of the virus through semen has not been proven experimentally; however, PoRV has been recovered from semen, testes, and other tissues of the reproductive tract for up to 49 days after inoculation (Solis et al., 2007). Based on semen evaluation in herds naturally infected with PoRV, approximately 30% of boars develop temporary or permanent infertility. Semen abnormalities include a decrease in concentration, increased morphologically abnormal sperm, decreased sperm motility and viability, and azoospermia (Maes et al., 2008). Most sows are clinically normal with a few animals developing corneal opacity. Infected breeding herds may experience increased returns to estrus and a reduction in farrowing rates.

3.5 Porcine Enteric Picornaviruses

Porcine enterovirus (PEV) and porcine teschovirus (PTV) are single-stranded positive-sense RNA viruses in the family *Picornaviridae*. PEV and PTV infections are commonly transmitted between pigs by oral exposure to contaminated feces; however, contamination of semen via aerosol during semen collection cannot be excluded (Guerin & Pozzi, 2005). PTV has been isolated from the male genital tract (Phillips et al., 1972); however, insemination of gilts with PTV contaminated semen had no effect on their fertility (De Meurichy & Pensaert, 1977). PEV can cause seminal vesiculitis, sperm abnormalities, and decrease libido (Phillips et al., 1972). Usually, there are no clinical signs in the sows. Although evidence is limited, there are reports that suggest that semen contaminated with PEV and PTV could cause embryonic and neonatal death (Dunne et al., 1969).

3.6 African Swine Fever Virus

African swine fever virus (ASFV) is a double-stranded, linear DNA virus in the genus *Asfivirus* of the family *Asfarviridae* (Tabares et al., 1980). ASFV replicates in monocytes and macrophages of the lymph nodes near the site of viral entry and subsequently spreads through the blood system, the lymphatic system, or both (Maes et al., 2008). ASFV transmission is generally vector-borne via tick bites (Parker et al., 1969); however, infections can also occur through direct interaction of sick and naïve animals (Plowright et al., 1970). Clinical signs range from subclinical to severe acute systemic disease that can resemble other hemorrhagic diseases in pigs. In naïve pigs, acute disease is characterized by high fever, loss of appetite, and hemorrhages in the skin (Moulton & Coggins, 1968). ASFV has been experimentally isolated from semen (Schlafer & Mebus, 1987; Thacker et al., 1984). Effect on semen quality and volume has to the authors knowledge not been reported. Under experimental conditions, ASFV has been shown to induce abortion when sows were infected between 38 and 92 days of gestation (Schlafer & Mebus, 1987). It was determined that abortion resulted from the effect of the virus on the dam (clinical illness) rather than direct effect on the fetuses (Schlafer & Mebus, 1987).

3.7 Classical Swine Fever Virus

Classical swine fever virus (CSFV) is a small, enveloped, positive-sense, single-stranded RNA virus in the genus *Pestivirus* of the family *Flaviviridae* (Becher et al., 1999). CSFV has been eradicated from many countries including Australia, New Zealand, North America, and Western Europe; however, it has been periodically reintroduced into domestic pigs that are in contact with wild boars. During the CSFV epizootic of 1997-1998 in the Netherlands (de Smit et al., 1999), two AI centers became infected and more than 100,000 sows from approximately 1,700 farms were affected (Hennecken et al., 2000) highlighting the economic importance of disease spread through AI centers. The main transmission route of CSFV is oronasally with primary virus replication in tonsils. Data from natural infection (Hennecken et al., 2000) and experimental inoculations (Choi & Chae, 2003; Floegel et al., 2000) have demonstrated that CSFV can be excreted in semen from infected boars. Boars experimentally infected with CSFV shed the virus in semen for up to 53 days post-infection (Choi & Chae, 2003). The virus does not affect the semen quality and motility and concentration are within normal range. Sows that were inseminated with contaminated semen seroconverted; however, the virus was shown to cross the placental barrier and infected the fetuses causing embryonic mortality (de Smit et al., 1999). CSFV continues to be important, especially in areas where CSFV is endemic, because the virus is highly contagious and infection of pregnant sows, in contrast to acute infections in piglets, may not be apparent (Hare et al., 1985; Ressang, 1973). If clinically normal CSFV immunotolerant piglets are born, they can spread virus for months without showing signs of disease or developing an antibody response (de Smit et al., 1999).

3.8 Foot and Mouth Disease Virus

Foot and mouth disease virus (FMDV) is a small single-stranded positive-sense RNA virus which belongs to the genus *Aphthovirus* in the family *Picornaviridae* (Carrillo et al., 1990). FMDV is endemic in Asia, some areas in South America and in Africa. FMDV can be isolated from all excretions and secretions including urine, feces, milk and semen (Lubroth & Brown, 1995). Infected pigs develop fever, lethargy and lameness which is associated with vesicular lesions in the hoof area (Mebus, 1978). Infection with FMDV leads to viremia, with subsequent systemic dissemination of the virus, including the genital tract and the skin around the preputial orifice (Alexandersen et al., 2001). FMDV has been detected in boar semen before and during manifestation of clinical signs of the disease. The viral concentration in semen has been found to be low (Guerin & Pozzi, 2005). After natural infection, FMDV has been isolated from infected boar semen for up to 9 days, but AI with contaminated semen failed to transmit the disease to sows (McVicar et al., 1977). To the authors' knowledge, an effect on semen quality has not been described. Abortion storms can be observed and are mainly due to the high fever and clinical illness of the dams.

3.9 Japanese B Encephalitis Virus

Japanese B encephalitis virus (JBEV) is a member of the *Flavivirus* genus of the family *Flaviviridae* (Solomon et al., 2003). JBEV is a mosquito-borne pathogen affecting humans and animals. This virus represents an economically important reproductive pathogen of breeding pigs, especially in Asia and Northern Australia, and is a common cause of

infertility in Japanese pigs (Habu et al., 1977). Infection of susceptible boars resulted in edematous, congested testes and semen with numerous abnormal spermatozoa and significantly decreased total and motile sperm counts (Habu et al., 1977). These changes are usually temporary and most boars recover completely. JBEV has been isolated from the testicles of boars with orchitis and also can be shed in the semen for 5 weeks (Habu et al., 1977). JBEV infection can easily be transmitted if gilts are inseminated with infected semen (Guerin & Pozzi, 2005; Habu et al., 1977).

3.10 Porcine Circovirus type 2

Porcine circovirus type 2 (PCV2) is a small, single-stranded, ambi-sense DNA virus that belongs to the family *Circoviridae*, genus *Circovirus* (Tischer et al., 1982). PCV2 is a ubiquitous virus and most herds worldwide are seropositive for PCV2 (Dulac & Afshar, 1989; Edwards & Sands, 1994; Segalés et al., 2008; Tischer et al., 1986). When first described in 1998, PCV2 was linked to disease mainly characterized by wasting and generalized lymphadenopathy in growing pigs (Allan et al., 1998; Morozov et al., 1998). Since that time, PCV2 has been associated with several disease manifestations in pigs commonly referred to as PCV-associated disease (PCVAD) which includes systemic disease, respiratory disease, enteric disease, porcine dermatitis and nephropathy syndrome (PDNS) and reproductive failure in dams (late term abortions and stillbirths) (Choi & Chae, 2001; Harms et al., 2002; Kim et al., 2004a; Kim et al., 2004b). PCV2-associated reproductive disease under field conditions is rare. The main route of transmission has been postulated to be the fecal-oral route (Segalés et al., 2005); however, due to the rapid spread and the extensive use of AI, semen transmission has been suggested as a potentially significant route of dissemination of PCV2 (Horlen et al., 2007; Lawton et al., 2004; Rose et al., 2003; Sibila et al., 2004; West et al., 1999). Mature boars infected with PCV2 generally lack clinical signs and lesions (Larochelle et al., 2000; Madson et al., 2008). The virus has been detected in semen of naturally and experimentally infected boars, even after the appearance of antibodies in the serum (Larochelle et al., 2000). In the acute phase of infection, PCV2 DNA was detected in serum and semen at two and five days post inoculation, respectively (Larochelle et al., 2000; Madson et al., 2008). Detection of PCV2 viremia commonly precedes the detection of semen-associated virus (Madson et al., 2008), but semen shedding has been reported in the absence of viremia (Larochelle et al., 2000). Anti-PCV2 antibodies typically develop by two weeks post inoculation (Larochelle et al., 2000; Madson et al., 2008). Following intranasal inoculation, intermittent semen shedding of PCV2 DNA was observed during a 47 day observation period (Larochelle et al., 2000). Intermittent semen shedding was confirmed by a different study (Grasland et al., 2008) after testing semen samples for 56 days in four inoculated boars. In contrast, continuous semen shedding was observed following intranasal and intramuscular inoculation following a 90 day observational period (Madson et al., 2008). In addition, naturally infected boars were found to sporadically shed PCV2 DNA in semen for up to 27.3 weeks in a PCV2 positive boar stud (McIntosh et al., 2006). Peak PCV2 shedding in semen occurs between nine and 20 days post inoculation (Grasland et al., 2008; Madson et al., 2008). Changes in morphology, motility, or quantity are not commonly associated with natural or experimental PCV2 infection (Madson et al., 2008; McIntosh et al., 2006). Clinical signs of PCV2 infection in the dam are typically absent or unapparent; however, a low percentage of females may abort due to systemic illness.

Pyrexia and anorexia are frequently observed in aborting dams (Park et al., 2005). The virus is also able to replicate in the zona pellucida-free embryos, leading to embryonic death (Mateusen et al., 2004, 2007). Delayed farrowing (>118 days of gestation) (Ladekjær-Mikkelsen et al., 2001) or pseudopregnancy (Josephson & Charbonneau, 2001) is less frequently observed with PCV2-associated reproductive failure. PCV2 affected litters are commonly composed of increased numbers of non-viable fetuses (mummified and stillborn) at parturition (Madson & Opriessnig, 2011).

3.11 Other viruses that are of minor importance

3.11.1 Swine Vesicular Disease Virus

Swine vesicular disease virus (SVDV) is a small, non-enveloped single-stranded positive-sense RNA virus, in the family *Picornaviridae*, genus *Enterovirus* (Nardelli et al., 1968). Documented outbreaks of this disease have been limited to selected countries in Asia, Europe and Central America. Clinical signs include appearance of vesicles around the coronary bands, snout, tongue and lip (Kanno et al., 1995) making this disease an important differential for FMDV. After natural infection, SVDV has been isolated from infected boar semen for up to four days, but AI with contaminated semen failed to transmit the disease to sows (McVicar et al., 1977).

3.11.2 Non-Classical Swine Fever Virus pestiviruses

Pigs are also susceptible to non-CSFV pestiviruses, including bovine viral diarrhea virus (BVDV) and border disease virus (BDV), which are associated with disease in cattle and sheep, respectively. Pigs congenitally infected with these viruses may shed large amounts of virus. Previously, BVDV has been isolated from oropharyngeal fluid, urine and semen of a congenitally infected, infertile boar (Terpstra & Wensvoort, 1997).

3.11.3 Porcine retroviruses

Retroviruses are RNA viruses that exist in two main groups: endogenous or exogenous retroviruses. Endogenous retroviruses are thought to be present in all vertebrates accounting for approximately 8% of their genomes (Tucker et al., 2006). All pigs carry PERV in their genome (Wilson, 2008) and three porcine endogenous retrovirus (PERV) subtypes have been identified based on their envelope sequence and tropism in cell culture (Le et al., 1997; Wilson et al., 2000): PERV-A, PERV-B and PERV-C (Martin et al., 2000a, 2000b; Patience et al., 1997; Specke et al., 2001; Takeuchi et al., 1998). More recently, the possible existence of exogenous porcine retrovirus in pigs was proposed (Scobie et al., 2004; Wood et al., 2004). It was demonstrated that a human-tropic recombination between PERV-A and PERV-C (designated as PERV-A/C) was not a product of *in vitro* recombination; instead PERV-A/C appeared to exist *in vivo* as an exogenous virus (Wood et al., 2004). Furthermore, a PERV-A/C recombinant was isolated from porcine peripheral blood mononuclear cells and was not present in a proviral form in the miniature swine genome (Scobie et al., 2004a; Wood et al., 2004). The PERV A/C virus is a recombination within the *env* region and is thought to arise from exogenous recombination of mRNA (Martin et al., 2006). Shedding via the semen route occurs, since PERVs are embedded in the genome.

3.11.4 Swine Influenza Virus and transmissible gastroenteritis virus

Swine influenza virus (SIV), a negative-sense, single-stranded RNA virus in the genus *influenzavirus A* of the family *Orthomyxoviridae* (Lamb & Krug, 2001; Wright & Webster, 2001), and transmissible gastroenteritis virus (TGEV), a positive-sense, single-stranded RNA virus in the *Coronaviridae* family (Lai & Cavanagh, 1997) have not been isolated from semen. Contamination of semen and transmission of SIV and TGEV have not been demonstrated to date; however, these viruses are highly contagious and could be easily transmitted via aerosol at the time of collection and preparation of semen. Boars in the acute stages of SIV or TGEV infection are often clinically ill on examination (Guerin & Pozzi, 2005).

3.11.5 Porcine Cytomegalovirus

Porcine cytomegalovirus (PCVM), a double-stranded DNA virus, is classified in the genus *Cytomegalovirus* in the subfamily *Betaherpesvirinae* of the family *Herpesviridae* (Roizman, 1982). Infection with PCMV is usually subclinical in adults including boars. Following infection in boars, the virus was detected in the testis and the epididymis (Shirai et al., 1985); however, semen shedding of the virus has not been determined.

3.11.6 Porcine Adenovirus

Porcine adenovirus (PAV) is a double-stranded, linear DNA virus which belongs to the family *Adenoviridae*. PAV is distributed worldwide and associated with gastroenteric disease. As such, its importance in AI is due to the potential of fecal contamination of semen.

4. Diagnostic approaches to prevent virus transmission by the semen route

The concern of spreading diseases on a global scale via semen has placed restrictions on international movement of semen. From the early 1990s onwards, major improvements in detection methods particularly in terms of sensitivity have better enabled diagnostic laboratories to detect viruses in boar semen. Today, various methods are available for demonstration of viruses in boars and include monitoring for presence of clinical signs, demonstration of virus (virus isolation, bioassay), nucleic acids, and demonstration of antibodies against the virus. However, it needs to be considered that the results of any of these methods may vary substantially, depending on individual sample type and the sensitivity and specificity of a chosen diagnostic method. Accurate diagnosis of semen associated viruses (correctly identified positive or negative status of any given sample) is extremely important as false positive as well as false negative test results can cause substantial economic damage. For example, with PRRSV, any false positive sample out of a boar stud has the following consequences: (1) Immediate hold on all semen samples from the stud to breeding herds resulting in missed cycles and associated production losses in hundreds to thousands of sows supplied by the boar stud. (2) Retesting of boars. (3) Culling of suspect positive boars.

4.1 Importance of sample type

Samples that are commonly collected in boar studs for routine surveillance include semen (raw or extended), blood swabs, serum and more recently oral fluids. Blood swabs have

become a popular sample type in recent years. For blood swab collection, the ear is pricked with a needle during semen collection and a cotton swab is used to catch the resulting blood drop (Reicks et al., 2006). Pathogens that cause systemic disease are typically identified in serum or blood first before they enter the semen and this should be taken into consideration when collecting appropriate samples. A negative result on semen only means that the tested sample does not contain virus, and that the particular semen sample is likely to be virus-free. It does not provide certainty that there will be no risk of contamination in the future. On the other hand, a negative result on serum does not guarantee that the semen is free of the virus as the viremic stage may have been very short.

4.2 Pooling of samples

Economic aspects are important to consider. For interpretation of diagnostic results at the boar stud level, in addition to the detection limit of the diagnostic method used, it is extremely important to test a representative number of semen samples, and to include evaluation of more than one parameter, such as presence of antibodies, viremia, or clinical symptoms. Because of the potential high cost of this approach, producers commonly rely on testing pooled samples. This strategy allows testing a larger number of animals while running the same number of tests, thus decreasing the cost per boar stud. Pooling can be successfully applied if the assay utilized has a high analytical sensitivity; however, because of a dilution effect, the sensitivity of the test when run on pooled samples is lower than its sensitivity when run on individual samples (Munoz-Zanzi et al., 2006).

4.3 Detection methods

4.3.1 Clinical evaluation of the boar

As a very basic and broad approach, in order to avoid spread of disease via contaminated semen, the health status of the animals should be checked daily. In case of any abnormality or clinical disease signs, semen collection should be halted until the animal has recovered. However, as indicated before, clinical examination alone is insufficient for most viral infections, since clinically normal boars can shed pathogens in their semen for extended periods of time.

4.3.2 Detection of viable virus

For demonstration of viable virus in semen two options exist including virus isolation and conducting bioassays. Virus isolation is often difficult as bacterial contamination of semen can be substantial. Virus isolation can be further complicated by the existence of cytotoxic factors in semen that destroy cell culture systems, and antiviral factors that nonspecifically neutralize virus (Christopher-Hennings et al., 1995b, 1997; Prieto et al., 1996b, 2003). In many cases swine bioassays have been found to be more sensitive than cell culture systems for demonstrating viable viruses in semen. However, animal inoculations cannot be justified to be used routinely for large numbers of samples and turnaround time for test results is poor.

4.3.3 Direct detection of RNA/DNA

Much progress has been made in recent years to improve the quality of molecular diagnostic assays used on semen. Numerous PCR techniques are now available for detecting

pathogens in boar semen. Many of these are highly sensitive, specific and rapid (Christopher-Hennings et al., 1995b). In the case of PRRSV, reverse transcriptase-PCR is today considered to be the most sensitive diagnostic technique. It allows for the detection of as little as 10^0 TCID₅₀ per seminal dose (Christopher-Hennings et al., 1995b; Gradil et al., 1996), 20 times less virus than what has been shown experimentally to result in transmission of PRRSV in gilts (Shin et al., 1997).

4.3.4 Detection of antibodies

Serology is a simple and relatively inexpensive way to survey for pathogens that should not be present in the boar stud. The main disadvantage of serology is the time period between pathogen exposure and detectable levels of antibodies in a boar which may range from a few days to weeks during which time the boar is not identified as being positive and may shed the virus via semen.

5. Prevention of introduction of viruses into boar studs

The best way to prevent disease transmission via semen is to assure pathogens are not introduced to the boar studs by maintaining very strict biosecurity measures (Madec et al., 1999) and routine surveillance. Due to the high risk of dissemination of disease via AI, the most important goal is to provide pathogen-free semen, which is feasible with adequate control measures.

5.1 Biosecurity

Strict biosecurity measures should be maintained, involving all of the following: (1) *Location of the boar stud*: To avoid introduction of pathogens via aerosol routes, boar studs are often located in remote areas away from pig dense populations. (2) *Protection from aerosol contamination*: High traffic areas and location next to heavily travelled roads have shown to be a risk factor for exposure to airborne viruses (Otake et al., 2002). Boar stud owners often attempt to protect the boars by implementing filter systems for incoming air in AI centers to safeguard boars against the entry of airborne pathogens (Dee et al., 2005, 2006). (3) *Incoming replacement boars*: Replacement boars are typically isolated for 30 to 60 days and housed in a separate off-site facility. During the isolation time, the health status of the boars is evaluated. Animals that test positive for any disease of concern are typically not allowed to enter. During the isolation time, incoming boars are often vaccinated against diseases present in the resident boars. The annual boar replacement rate in AI centers is typically 60% (Singleton, 2001). (4) *Employees and visitors*: Strict regulations are typically in place for people entering AI centers. Often, there is 24 to 72 hour down-time required during which time people are not allowed to have contact with other live pigs. In addition, there is often a shower in/shower out requirement for all visitors and employees. (5) *Deliveries*: Any deliveries (clothes, mail, supplies, feed, tools, and equipment) are brought to a special entrance with no direct contact with the boar stud and are fumigated prior to entry into the boar studs. (6) *Rodent and insect control*: Rodents and insects have been shown to be capable of carrying several pig viruses (Lorincz et al., 2010; Otake et al., 2003a, 2003b; Plowright et al., 1970) and appropriate prevention measures need to be in place. (7) *Implementation of a regular and appropriate cleaning and disinfectant procedure*.

5.2 Routine surveillance testing

Commercial AI centers need be regularly checked for conformity with specific criteria, assuring that their products are free of certain pathogens and contain a minimal or acceptable number of microorganisms (Guerin & Pozzi, 2005; Prieto et al., 2004). Many veterinary diagnostic laboratories took this need into account and started to offer daily service for boar studs.

5.3 General semen handling and storage

To reduce the unavoidable presence of bacteria in the ejaculate and to prolong *in vitro* longevity of sperm, use of antimicrobials in semen is a common part of most semen extenders. Apart from a possible dilution effect of pathogens, semen processing and addition of antimicrobials does not eliminate viruses. The use of effective antiviral agents to render semen virus-free has so far not been adopted in the swine AI industry. However, several best practices for handling, storage conditions, and use of fresh semen have been described to reduce the potential for transmission of viral pathogens (Guerin & Pozzi, 2005). For boars, the immediate use of fresh semen increases the risk of pathogen transmission (Prieto et al., 2004).

5.4 Vaccination

The use of modified-live and inactivated-virus vaccines in boars can be highly effective in eliminating or decreasing shedding of viruses and with that decreasing the risk of virus transmission by AI. For example, vaccination against PPV may help to reduce shedding of the virus following infection. This has also been shown for PCV2 under experimental conditions (Opriessnig et al., 2011). In many countries, vaccination is done as part of a PRV eradication program (Siegel & Weigel, 1999). In the case of PRRSV, the use of a modified live virus vaccine shortened or eliminated virus shedding in boars challenged with wild-type virus for 50 days after vaccination (Christopher-Hennings et al., 1997; Nielsen et al., 1997). However, based on other studies it appears that PRRSV vaccination provides only partial protection (Nielsen et al., 1997). Moreover, semen shedding has been demonstrated after vaccination with a modified live vaccine virus (Christopher-Hennings et al., 1997). In addition, the modified live PRRSV vaccine virus has been shown to be shed in the semen in low levels (Christopher-Hennings et al., 1995b). In contrast, an inactivated vaccine, did not clearly reduce subsequent shedding of wild-type virus in semen (Christopher-Hennings et al., 1997; Stevenson et al., 1994). The presence of PRRSV in semen was demonstrated by PCR in most of the vaccinated boars during an interval of 7–21 days post-vaccination, although some boars sporadically shed the virus for longer periods of time (Christopher-Hennings et al., 1997). Similarly, when a swine bioassay was used, the presence of infectious virus in semen samples of vaccinated boars was confirmed 14 days post-vaccination (Nielsen et al., 1997).

6. Summary

Virus contamination of boar semen poses a great risk for breeding herds worldwide due the possibility of fast introduction of viruses into large naïve and susceptible populations.

Producers in some countries have led the way in establishing and maintaining specific pathogen-free AI centers in which incoming and resident boars are systematically screened for specific viruses to prevent introduction of viruses and if viruses are introduced to prevent spread of the virus within the boar studs and to breeding herds that receive semen from the studs. Serious attention is being given to appropriate location of boar studs and major investments are being made in air filtration equipment and building design and strict adherence is given to biosecurity protocols to further decrease the risks of introduction of diseases, particularly PRRSV, to boar studs in North America.

7. References

- Albina, E., Madec, F., Cariolet, R., & Torrison, J. (1994). Immune response and persistence of the porcine reproductive and respiratory syndrome virus in infected pigs and farm units. *Veterinary Record*, Vol. 134, No. 22, pp. 567-573.
- Alexandersen, S., Oleksiewicz, M.B., & Donaldson, A.I. (2001). The early pathogenesis of foot-and-mouth disease in pigs infected by contact: a quantitative time-course study using TaqMan RT-PCR. *Journal of General Virology*, Vol. 82, No. 4, pp. 747-755.
- Allan, G.M., McNeilly, F., Ellis, J., Krakowka, S., Meehan, B., McNair, I., Walker, I., & Kennedy, S. (2000). Experimental infection of colostrum deprived piglets with porcine circovirus 2 (PCV2) and porcine reproductive and respiratory syndrome virus (PRRSV) potentiates PCV2 replication. *Archives of Virology*, Vol. 145, No. 11, pp. 2421-2429.
- Allan, G.M., McNeilly, F., Kennedy, S., Daft, B., Clarke, E.G., Ellis, J.A., Haines, D.M., Meehan, B.M., & Adair, B.M. (1998). Isolation of porcine circovirus-like viruses from pigs with a wasting disease in the USA and Europe. *Journal of Veterinary Diagnostic Investigation*, Vol. 10, No. 1, pp. 3-10.
- Bachmann, P.A., Sheffy, B.E., & Vauhan, J.T. (1975). Experimental in utero infection of fetal pigs with a porcine parvovirus. *Infection and Immunity*, Vol. 12, No. 3, pp. 455-460.
- Bautista, E.M., Goyal, S.M., & Collins, J.E. (1993). Serologic survey for Lelystad and VR-2332 strains of porcine respiratory and reproductive syndrome (PRRS) virus in US swine herds. *Journal of Veterinary Diagnostic Investigation*, Vol. 5, No. 4, pp. 612-614.
- Becher, P., Orlich, M., Kosmidou, A., Konig, M., Baroth, M., & Thiel, H.J. (1999). Genetic diversity of pestiviruses: identification of novel groups and implications for classification. *Virology*, Vol. 262, No. 1, pp. 64-71.
- Benfield, D., Nelson, C., Steffen, M., & Rowland, R. (2000). Transmission of PRRSV by artificial insemination using extended semen seeded with different concentrations of PRRSV. *Proceedings of the American Association of Swine Practitioners*, Vol. 31, pp. 405-408.
- Bierk, M.D., Dee, S.A., Rossow, K.D., Otake, S., Collins, J.E., & Molitor, T.W. (2001). Transmission of porcine reproductive and respiratory syndrome virus from persistently infected sows to contact controls. *Canadian Journal of Veterinary Research*, Vol. 65, No. 4, pp. 261-266.
- Biront, P. & Bonte, P. (1983). Porcine parvovirus (P.P.V.): infection in boars. I. Possibility of a genital localization in the boar after oronasal infection. *Zentralblatt für Veterinärmedizin Reihe B*, Vol. 30, No. 7, pp. 541-545.

- Brown, T.T., Jr., Paul, P.S., & Mengeling, W.L. (1980). Response of conventionally raised weanling pigs to experimental infection with a virulent strain of porcine parvovirus. *American Journal of Veterinary Research*, Vol. 41, No. 8, pp. 1221-1224.
- Carrillo, C., Plana, J., Mascarella, R., Bergada, J., & Sobrino, F. (1990). Genetic and phenotypic variability during replication of foot-and-mouth disease virus in swine. *Virology*, Vol. 179, No. 2, pp. 890-892.
- Cartwright, S.F. & Huck, R.A. (1967). Viruses isolated in association with herd infertility, abortions and stillbirths in pigs. *Veterinary Record*, No. 81, pp. 196-197.
- Cartwright, S.F., Lucas, M., & Huck, R.A. (1971). A small haemagglutinating porcine DNA virus. II. Biological and serological studies. *Journal of Comparative Pathology*, Vol. 81, No. 1, pp. 145-155.
- Cavanagh, D. (1997). Nidovirales: a new order comprising Coronaviridae and Arteriviridae. *Archives of Virology*, Vol. 142, No. 3, pp. 629-633.
- Choi, C. & Chae, C. (2001). Colocalization of porcine reproductive and respiratory syndrome virus and porcine circovirus 2 in porcine dermatitis and nephrology syndrome by double-labeling technique. *Veterinary Pathology*, Vol. 38, No. 4, pp. 436-441.
- Choi, C. & Chae, C. (2003). Detection of classical swine fever virus in boar semen by reverse transcription-polymerase chain reaction. *Journal of Veterinary Diagnostic Investigation*, Vol. 15, No. 1, pp. 35-41.
- Choi, C.S., Molitor, T.W., Joo, H.S., & Gunther, R. (1987). Pathogenicity of a skin isolate of porcine parvovirus in swine fetuses. *Veterinary Microbiology*, Vol. 15, No. 1-2, pp. 19-29.
- Christianson, W.T., Choi, C.S., Collins, J.E., Molitor, T.W., Morrison, R.B., & Joo, H.S. (1993). Pathogenesis of porcine reproductive and respiratory syndrome virus infection in mid-gestation sows and fetuses. *Canadian Journal of Veterinary Research*, Vol. 57, No. 4, pp. 262-268.
- Christopher-Hennings, J., Holler, L.D., Benfield, D.A., & Nelson, E.A. (2001). Detection and duration of porcine reproductive and respiratory syndrome virus in semen, serum, peripheral blood mononuclear cells, and tissues from Yorkshire, Hampshire, and Landrace boars. *Journal of Veterinary Diagnostic Investigation*, Vol. 13, No. 2, pp. 133-142.
- Christopher-Hennings, J., Nelson, E.A., Hines, R.J., Nelson, J.K., Swenson, S.L., Zimmerman, J.J., Chase, C.L., Yaeger, M.J., & Benfield, D.A. (1995a). Persistence of porcine reproductive and respiratory syndrome virus in serum and semen of adult boars. *Journal of Veterinary Diagnostic Investigation*, Vol. 7, No. 4, pp. 456-464.
- Christopher-Hennings, J., Nelson, E.A., Nelson, J.K., & Benfield, D.A. (1997). Effects of a modified-live virus vaccine against porcine reproductive and respiratory syndrome in boars. *American Journal of Veterinary Research*, Vol. 58, No. 1, pp. 40-45.
- Christopher-Hennings, J., Nelson, E.A., Nelson, J.K., Hines, R.J., Swenson, S.L., Hill, H.T., Zimmerman, J.J., Katz, J.B., Yaeger, M.J., Chase, C.C., & Benfield, D.A. (1995b). Detection of porcine reproductive and respiratory syndrome virus in boar semen by PCR. *Journal of Clinical Microbiology*, Vol. 33, No. 7, pp. 1730-1734.
- Christopher-Hennings, J., Nelson, E.A., Nelson, J.K., Rossow, K.D., Shivers, J.L., Yaeger, M.J., Chase, C.C., Garduno, R.A., Collins, J.E., & Benfield, D.A. (1998).

- Identification of porcine reproductive and respiratory syndrome virus in semen and tissues from vasectomized and nonvasectomized boars. *Veterinary Pathology*, Vol. 35, No. 4, pp. 260-267.
- Cutlip, R.C. & Mengeling, W.L. (1975a). Experimentally induced infection of neonatal swine with porcine parvovirus. *American Journal of Veterinary Research*, Vol. 36, No. 8, pp. 1179-1182.
- Cutlip, R.C. & Mengeling, W.L. (1975b). Pathogenesis of in utero infection: experimental infection of eight- and ten-week-old porcine fetuses with porcine parvovirus. *American Journal of Veterinary Research*, Vol. 36, No. 12, pp. 1751-1754.
- De Meurichy, W. & Pensaert, M. (1977). Effect of an enterovirus in gilts inseminated with a semen-virus mixture. *Zentralblatt für Veterinärmedizin Reihe B*, Vol. 24, No. 2, pp. 97-103.
- De Smit, A.J., Bouma, A., Terpstra, C., & Van Oirschot, J.T. (1999). Transmission of classical swine fever virus by artificial insemination. *Veterinary Microbiology*, Vol. 67, No. 4, pp. 239-249.
- Dea, S., Elazhary, M.A., Martineau, G.P., & Vaillancourt, J. (July 1985). Parvovirus-like particles associated with diarrhea in unweaned piglets. *Canadian Journal of Comparative Medicine*, Vol. 49, No. 3, pp. 343-345.
- Dee, S., Batista, L., Deen, J., & Pijoan, C. (2005). Evaluation of an air-filtration system for preventing aerosol transmission of porcine reproductive and respiratory syndrome virus. *Canadian Journal of Veterinary Research*, Vol. 69, No. 4, pp. 293-298.
- Dee, S.A., Deen, J., Cano, J.P., Batista, L., & Pijoan, C. (July 2006). Further evaluation of alternative air-filtration systems for reducing the transmission of Porcine reproductive and respiratory syndrome virus by aerosol. *Canadian Journal of Veterinary Research*, Vol. 70, No. 3, pp. 168-175.
- Duhamel, G.E., Bargar, T.W., Schmitt, B.J., Molitor, T.W., & Lu, W. (1991). Identification of parvovirus-like virus particles in intestinal crypt epithelial cells of pigs with diarrhea. *Journal of Veterinary Diagnostic Investigation*, Vol. 3, No. 1, pp. 96-98.
- Dulac, G.C. & Afshar, A. (1989). Porcine circovirus antigens in PK-15 cell line (ATCC CCL-33) and evidence of antibodies to circovirus in Canadian pigs. *Canadian Journal of Veterinary Research*, Vol. 53, No. 4, pp. 431-433.
- Dunne, H.W., Wang, J.T., Clark, C.D., Hokanson, J.F., Morimoto, T., & Bubash, G.R. (1969). The effects of in utero viral infection on embryonic, fetal, and neonatal survival: a comparison of SMEDI (porcine picorna) viruses with hog cholera vaccinal virus. *Canadian Journal of Comparative Medicine*, Vol. 33, No. 4, pp. 244-252.
- Edwards, S. & Sands, J.J. (1994). Evidence of circovirus infection in British pigs. *Veterinary Record*, Vol. 134, No. 26, pp. 680-681.
- Floegel, G., Wehrend, A., Depner, K.R., Fritzemeier, J., Waberski, D., & Moennig, V. (2000). Detection of classical swine fever virus in semen of infected boars. *Veterinary Microbiology*, Vol. 77, No. 1-2, pp. 109-116.
- Flowers, W.L. & Esbenshade, K.L. (1993). Optimizing management of natural and artificial matings in swine. *Journal of Reproduction and Fertility Supplements*, Vol. 48, pp. 217-228.

- Gerrits, R.J., Lunney, J.K., Johnson, L.A., Pursel, V.G., Kraeling, R.R., Rohrer, G.A., & Dobrinsky, J.R. (2005). Perspectives for artificial insemination and genomics to improve global swine populations. *Theriogenology*, Vol. 63, No. 2, pp. 283-299.
- Gradil, C., Dubuc, C., & Eaglesome, M.D. (1996). Porcine reproductive and respiratory syndrome virus: seminal transmission. *Veterinary Record*, Vol. 138, No. 21, pp. 521-522.
- Gradil, C., Molitor, T., Harding, M., & Crabo, B. (1990). Excretion of porcine parvovirus through the genital tract of boars. *American Journal of Veterinary Research*, Vol. 51, No. 3, pp. 359-362.
- Grasland, B., Blanchard, P., Jan, B., Oger, A., Rose, N., Madec, F., Jestin, A., & Cariolet, R. (2008). Transmission of porcine circovirus of type 2 (PCV2) with semen. *Proceedings of the 20th International Pig Veterinary Society Congress*, Vol. 2, p. 56.
- Guerin, B. & Pozzi, N. (2005). Viruses in boar semen: detection and clinical as well as epidemiological consequences regarding disease transmission by artificial insemination. *Theriogenology*, Vol. 63, No. 2, pp. 556-572.
- Habu, A., Murakami, Y., Ogasa, A., & Fujisaki, Y. (1977). [Disorder of spermatogenesis and viral discharge into semen in boars infected with Japanese encephalitis virus (author's transl)]. *Uirusu*, Vol. 27, No. 1, pp. 21-26.
- Hall, L.B., Jr., Kluge, J.P., Evans, L.E., Clark, T.L., & Hill, H.T. (1984a). Testicular changes observed in boars following experimental inoculation with pseudorabies (Aujeszky's) virus. *Canadian Journal of Comparative Medicine*, Vol. 48, No. 3, pp. 303-307.
- Hall, L.B., Jr., Kluge, J.P., Evans, L.E., & Hill, H.T. (1984b). The effect of pseudorabies (Aujeszky's) virus infection on young mature boars and boar fertility. *Canadian Journal of Comparative Medicine*, Vol. 48, No. 2, pp. 192-197.
- Hare, W.C., Mitchell, D., Singh, E.L., Bouillant, A.M., Eaglesome, M.D., Ruckerbauer, G.M., Bielanski, A., & Randall, G.C. (1985). Embryo transfer in relation to bovine leukemia virus control and eradication. *Canadian Veterinary Journal*, Vol. 26, No. 8, pp. 231-234.
- Harms, P.A., Halbur, P.G., & Sorden, S.D. (2002). Three cases of porcine respiratory disease complex associated with porcine circovirus type 2 infection. *Journal of Swine Health and Production*, Vol. 10, pp. 27-30.
- Henneken, M., Stegeman, J.A., Elbers, A.R., van, N.A., Smak, J.A., & Verheijden, J.H. (2000). Transmission of classical swine fever virus by artificial insemination during the 1997-1998 epidemic in The Netherlands: a descriptive epidemiological study. *Veterinary Quarterly*, Vol. 22, No. 4, pp. 228-233.
- Horlen, K.P., Schneider, P., Anderson, J., Nietfeld, J.C., Henry, S.C., Tokach, L.M., & Rowland, R.R. (2007). A cluster of farms experiencing severe porcine circovirus associated disease: Clinical features and association with the PCV2b genotype. *Journal of Swine Health and Production*, Vol. 15, No. 5, pp. 270-278.
- Johnson, R.H., Donaldson-Wood, C., & Allender, U. (Feb. 1976). Observations on the epidemiology of porcine parvovirus. *Australian Veterinary Journal*, Vol. 52, No. 2, pp. 80-84.
- Joo, H.S., Donaldson-Wood, C.R., & Johnson, R.H. (1976). Observations on the pathogenesis of porcine parvovirus infection. *Archives of Virology*, Vol. 51, No. 1-2, pp. 123-129.

- Josephson, G. & Charbonneau, G. (2001). Case report of reproductive problem in a new startup operation. *Swine Health and Production*, Vol. 9, No. 5, pp. 258-259.
- Kanno, T., Inoue, T., Wang, Y., Sarai, A., & Yamaguchi, S. (1995). Identification of the location of antigenic sites of swine vesicular disease virus with neutralization-resistant mutants. *Journal of General Virology*, Vol. 76, pp. 3099-3106
- Kennedy, S., Moffett, D., McNeilly, F., Meehan, B., Ellis, J., Krakowka, S., & Allan, G.M. (2000). Reproduction of lesions of postweaning multisystemic wasting syndrome by infection of conventional pigs with porcine circovirus type 2 alone or in combination with porcine parvovirus. *Journal of Comparative Pathology*, Vol. 22, No. 1, pp. 9-24.
- Kim, J., Ha, Y., Jung, K., Choi, C., & Chae, C. (2004a). Enteritis associated with porcine circovirus 2 in pigs. *Canadian Journal of Veterinary Research*, Vol. 68, No. , pp. 218-221.
- Kim, J., Jung, K., & Chae, C. (2004b). Prevalence of porcine circovirus type 2 in aborted fetuses and stillborn piglets. *Veterinary Record*, Vol. 155, No. 16, pp. 489-492.
- Krakowka, S., Ellis, J.A., Meehan, B., Kennedy, S., McNeilly, F., & Allan, G. (2000). Viral wasting syndrome of swine: experimental reproduction of postweaning multisystemic wasting syndrome in gnotobiotic swine by coinfection with porcine circovirus 2 and porcine parvovirus. *Veterinary Pathology*, Vol. 37, No. 3, pp. 254-263.
- Kresse, J.I., Taylor, W.D., Stewart, W.W., & Eernisse, K.A. (1985). Parvovirus infection in pigs with necrotic and vesicle-like lesions. *Veterinary Microbiology*, Vol. 10, No. 6, pp. 525-531.
- Ladekjær-Mikkelsen, A.S., Nielsen, J., Storgaard, T., Bøtner, A., Allan, G., & McNeilly, F. (2001). Transplacental infection with PCV-2 associated with reproductive failure in a gilt. *Veterinary Record*, Vol. 148, No. 24, pp. 759-760.
- Lager, K.M. & Mengeling, W.L. (1994). Porcine parvovirus associated with cutaneous lesions in piglets. *Journal of Veterinary Diagnostic and Investigation*, Vol. 6, No. 3, pp. 357-359.
- Lager, K.M. & Mengeling, W.L. (1995). Pathogenesis of in utero infection in porcine fetuses with porcine reproductive and respiratory syndrome virus. *Canadian Journal of Veterinary Research*, Vol. 59, No. 3, pp. 187-192.
- Lager, K.M., Mengeling, W.L., & Liu, W. (1992). Comparison of the virulence of two isolates of porcine parvovirus in 72-day-old porcine fetuses. *Journal of Veterinary Diagnostic and Investigation*, Vol. 4, No. 3, pp. 245-248.
- Lai, M.M. & Cavanagh, D. (1997). The molecular biology of coronaviruses. *Advances in Virus Research*, Vol. 48, pp. 1-100.
- Lamb, R.A. & Krug, R.M. (2001). Orthomyxoviridae: The viruses and their replication, In: *Field's Virology*, Knipe, D.M., Howley, P.M., Griffin, D.E., Lamb, R.A., Martin, M.A., Roizman, B., & Straus, S.E., pp. 1487-1531, Lippincott-Raven Publishers, ISBN Philadelphia
- Larochelle, R., Bielanski, A., Müller, P., & Magar, R. (2000). PCR detection and evidence of shedding of porcine circovirus type 2 in boar semen. *Journal of Clinical Microbiology*, Vol. 38, No. 12, pp. 4629-4632.

- Larsen, R.E., Shope, R.E., Jr., Leman, A.D., & Kurtz, H.J. (1980). Semen changes in boars after experimental infection with pseudorabies virus. *American Journal of Veterinary Research*, Vol. 41, No. 5, pp. 733-739.
- Lawton, D.E., Morris, R.S., & King, C.M. (2004). PMWS in New Zealand part 2: Epidemiological evidence for a novel agent. *18th International Pig Veterinary Society Congress*, Vol. 1, p. 128.
- Le, T.P., Stoye, J.P., Takeuchi, Y., Patience, C., & Weiss, R.A. (1997). Two sets of human-tropic pig retrovirus. *Nature*, Vol. 389, No. 6652, pp. 681-682.
- Lorincz, M., Csagola, A., Biksi, I., Szeredi, L., Dan, A., & Tuboly, T. (2010). Detection of porcine circovirus in rodents - short communication. *Acta Veterinaria Hungarica*, Vol. 58, No. 2, pp. 265-268.
- Lubroth, J. & Brown, F. (1995). Identification of native foot-and-mouth disease virus non-structural protein 2C as a serological indicator to differentiate infected from vaccinated livestock. *Research in Veterinary Science*, Vol. 59, No. 1, pp. 70-78.
- Lucas, M.H., Cartwright, S.F., & Wrathall, A.E. (1974). Genital infection of pigs with porcine parvovirus. *Journal of Comparative Pathology*, Vol. 84, No. 3, pp. 347-350.
- Lush, J.L. (1925). The possibility of sex control by artificial insemination with centrifuged spermatozoa. *Journal of Agricultural Research*, Vol. 30, No. 10, pp. 893-913.
- Madec, F., Eveno, E., Morvan, P., Hamon, L., Morvan, H., Albina, E., Truong, C., Hutet, E., Cariolet, R., Arnould, C., & Jestin, A. (1999). La maladie de l'amaigrissement du porcelet (MAP) en France 1 - Aspects descriptifs, impact en élevage. [Porcine wasting disease (PWD) in France: 1 - Description of the disease and impact in affected herds]. *Journées Recherche en Porcine France*, Vol. 31, No. 347-354.
- Madsen, K.S. (2005). Management of disease control and epidemics in AI in Denmark. *Theriogenology*, Vol. 63, No. 2, pp. 585-594.
- Madson, D.M. & Opriessnig, T. (2011). Effect of porcine circovirus type 2 (PCV2) infection on reproduction: disease, vertical transmission, diagnostics and vaccination. *Animal Health Research Reviews*, Vol. 12, No. 1, pp. 47-65.
- Madson, D.M., Ramamoorthy, S., Kuster, C., Pal, N., Meng, X.J., Halbur, P.G., & Opriessnig, T. (2008). Characterization of shedding patterns of porcine circovirus types 2a and 2b in experimentally inoculated mature boars. *Journal of Veterinary Diagnostic Investigation*, Vol. 20, No. 6, pp. 725-734.
- Madson, D.M., Ramamoorthy, S., Kuster, C., Pal, N., Meng, X.J., Halbur, P.G., & Opriessnig, T. (2009). Infectivity of porcine circovirus type 2 DNA in semen from experimentally-infected boars. *Veterinary Research*, Vol. 40, No. 1, p. 10.
- Maes, D., Nauwynck, H., Rijsselaere, T., Mateusen, B., Vyt, P., de, K.A., & Van, S.A. (2008). Diseases in swine transmitted by artificial insemination: an overview. *Theriogenology*, Vol. 70, No. 8, pp. 1337-1345.
- Maes, R.K., Kanitz, C.L., & Gustafson, D.P. (1983). Shedding patterns in swine of virulent and attenuated pseudorabies virus. *American Journal of Veterinary Research*, Vol. 44, No. 11, pp. 2083-2086.
- Martin, S.I., Wilkinson, R., & Fishman, J.A. (2006). Genomic presence of recombinant porcine endogenous retrovirus in transmitting miniature swine. *Virology Journal*, Vol. 3, p. 91.

- Martin, U., Winkler, M.E., Id, M., Radecke, H., Arseniev, L., Groteluschen, R., Simon, A.R., & Steinhoff, G. (2000a). Transmission of pig endogenous retrovirus to primary human cells. *Transplantation Proceedings*, Vol. 32, No. 5, p. 1157.
- Martin, U., Winkler, M.E., Id, M., Radecke, H., Arseniev, L., Takeuchi, Y., Simon, A.R., Patience, C., Haverich, A., & Steinhoff, G. (2000b). Productive infection of primary human endothelial cells by pig endogenous retrovirus (PERV). *Xenotransplantation*, Vol. 7, No. 2, pp. 138-142.
- Mateusen, B., Maes, D.G., Van, S.A., Lefebvre, D., & Nauwynck, H.J. (2007). Effect of a porcine circovirus type 2 infection on embryos during early pregnancy. *Theriogenology*, Vol. 68, No. 6, pp. 896-901.
- Mateusen, B., Sanchez, R.E., Van, S.A., Meerts, P., Maes, D.G., & Nauwynck, H.J. (2004). Susceptibility of pig embryos to porcine circovirus type 2 infection. *Theriogenology*, Vol. 61, No. 1, pp. 91-101.
- McIntosh, K.A., Harding, J.C., Parker, S., Ellis, J.A., & Appleyard, G.D. (2006). Nested polymerase chain reaction detection and duration of porcine circovirus type 2 in semen with sperm morphological analysis from naturally infected boars. *Journal of Veterinary Diagnostic Investigation*, Vol. 18, No. 4, pp. 380-384.
- McVicar, J.W., Eisner, R.J., Johnson, L.A., & Pursel, V.G. (1977). Foot-and-mouth disease and swine vesicular disease viruses in boar semen. *Proceedings of the Annual Meeting of the United States Animal Health Association*, Vol. 81, pp. 221-230.
- Mebus, C.A. (1978). Ulcerative diseases of animals with an infectious etiology. *Journal of Oral Pathology*, Vol. 7, No. 6, pp. 365-371.
- Medveczky, I. & Szabo, I. (1981). Isolation of Aujeszky's disease virus from boar semen. *Acta Veterinaria Hungarica*, Vol. 29, No. 1, pp. 29-35.
- Meng, X.J., Paul, P.S., Halbur, P.G., & Lum, M.A. (1995). Phylogenetic analyses of the putative M (ORF 6) and N (ORF 7) genes of porcine reproductive and respiratory syndrome virus (PRRSV): implication for the existence of two genotypes of PRRSV in the USA and Europe. *Archives of Virology*, Vol. 140, No. 4, pp. 745-755.
- Mengeling, W.L. (1979). Prenatal infection following maternal exposure to porcine parvovirus on either the seventh or fourteenth day of gestation. *Canadian Journal of Comparative Medicine*, Vol. 43, No. 1, pp. 106-109.
- Mengeling, W.L. & Cutlip, R.C. (1976). Reproductive disease experimentally induced by exposing pregnant gilts to porcine parvovirus. *American Journal of Veterinary Research*, Vol. 37, No. 12, pp. 1393-1400.
- Mengeling, W.L., Lager, K.M., & Vorwald, A.C. (1994). Temporal characterization of transplacental infection of porcine fetuses with porcine reproductive and respiratory syndrome virus. *American Journal of Veterinary Research*, Vol. 55, No. 10, pp. 1391-1398.
- Mengeling, W.L. & Paul, P.S. (1986). Interepizootic survival of porcine parvovirus. *Journal of the American Veterinary Medical Association*, Vol. 188, No. 11, pp. 1293-1295.
- Mettenleiter, T.C. (2000). Aujeszky's disease (pseudorabies) virus: the virus and molecular pathogenesis--state of the art, June 1999. *Veterinary Research*, Vol. 31, No. 1, pp. 99-115.

- Meulenbergh, J.J., Hulst, M.M., de Meijer, E.J., Moonen, P.L., den, B.A., de Kluyver, E.P., Wensvoort, G., & Moormann, R.J. (1993). Lelystad virus, the causative agent of porcine epidemic abortion and respiratory syndrome (PEARS), is related to LDV and EAV. *Virology*, Vol. 192, No. 1, pp. 62-72.
- Molitor, T.W., Joo, H.S., & Collett, M.S. (1984). Porcine parvovirus DNA: characterization of the genomic and replicative form DNA of two virus isolates. *Virology*, Vol. 137, No. 2, pp. 241-254.
- Morozov, I., Sirinarumitr, T., Sorden, S.D., Halbur, P.G., Morgan, M.K., Yoon, K.J., & Paul, P.S. (1998). Detection of a novel strain of porcine circovirus in pigs with postweaning multisystemic wasting syndrome. *Journal of Clinical Microbiology*, Vol. 36, No. 9, pp. 2535-2541.
- Moulton, J. & Coggins, L. (1968). Comparison of lesions in acute and chronic African swine fever. *Cornell Vet*, Vol. 58, No. 3, pp. 364-388.
- Munoz-Zanzi, C., Thurmond, M., Hietala, S., & Johnson, W. (2006). Factors affecting sensitivity and specificity of pooled-sample testing for diagnosis of low prevalence infections. *Preventive Veterinary Medicine*, Vol. 74, No. 4, pp. 309-322.
- Nardelli, L., Lodetti, E., Gualandi, G.L., Burrows, R., Goodridge, D., Brown, F., & Cartwright, B. (1968). A foot and mouth disease syndrome in pigs caused by an enterovirus. *Nature*, Vol. 219, No. 5160, pp. 1275-1276.
- Neumann, E.J., Kliebenstein, J.B., Johnson, C.D., Mabry, J.W., Bush, E.J., Seitzinger, A.H., Green, A.L., & Zimmerman, J.J. (2005). Assessment of the economic impact of porcine reproductive and respiratory syndrome on swine production in the United States. *Journal of the American Medical Association*, Vol. 227, No. 3, pp. 385-392.
- Nielsen, T.L., Nielsen, J., Have, P., Baekbo, P., Hoff-Jorgensen, R., & Botner, A. (1997). Examination of virus shedding in semen from vaccinated and from previously infected boars after experimental challenge with porcine reproductive and respiratory syndrome virus. *Veterinary Microbiology*, Vol. 54, No. 2, pp. 101-112.
- Opriessnig, T., Madson, D.M., Schalk, S., Brockmeier, S., Shen, H.G., Beach, N.M., Meng, X.J., Baker, R.B., Zanella, E.L. & Halbur, P.G. (2011). Porcine circovirus type 2 (PCV2) vaccination is effective in reducing disease and PCV2 shedding in semen of boars concurrently infected with PCV2 and *Mycoplasma hyopneumoniae*. *Theriogenology*, Vol. 76, No. 2, pp. 351-360.
- Oraveerakul, K., Choi, C.S., & Molitor, T.W. (1993). Tissue tropisms of porcine parvovirus in swine. *Archives of Virology*, Vol. 130, No. 3-4, pp. 377-389.
- Otake, S., Dee, S.A., Jacobson, L., Torremorell, M., & Pijoan, C. (2002). Evaluation of aerosol transmission of porcine reproductive and respiratory syndrome virus under controlled field conditions. *Veterinary Record*, Vol. 150, No. 26, pp. 804-808.
- Otake, S., Dee, S.A., Moon, R.D., Rossow, K.D., Trincado, C., Farnham, M., & Pijoan, C. (2003a). Survival of porcine reproductive and respiratory syndrome virus in houseflies. *Canadian Journal of Veterinary Research*, Vol. 67, No. 3, pp. 198-203.
- Otake, S., Dee, S.A., Moon, R.D., Rossow, K.D., Trincado, C., & Pijoan, C. (2003b). Evaluation of mosquitoes, *Aedes vexans*, as biological vectors of porcine reproductive and respiratory syndrome virus. *Canadian Journal of Veterinary Research*, Vol. 67, No. 4, pp. 265-270.

- Park, J.-S., Kim, J., Ha, Y., Jung, K., Choi, C., Lim, J.-K., Kim, S.-H., & Chae, C. (2005). Birth abnormalities in pregnant sows infected intranasally with porcine circovirus 2. *Journal of Comparative Pathology*, Vol. 132, No. 2-3, pp. 139-144.
- Parker, J., Plowright, W., & Pierce, M.A. (1969). The epizootiology of African swine fever in Africa. *Veterinary Record*, Vol. 85, No. 24, pp. 668-674.
- Patience, C., Takeuchi, Y., & Weiss, R.A. (1997). Infection of human cells by an endogenous retrovirus of pigs. *Nature Medicine*, Vol. 3, No. 3, pp. 282-286.
- Paul, P.S., Mengeling, W.L., & Brown, T.T., Jr. (1979). Replication of porcine parvovirus in peripheral blood lymphocytes, monocytes, and peritoneal macrophages. *Infection and Immunity*, Vol. 25, No. 3, pp. 1003-1007.
- Phillips, R.M., Foley, C.W., & Lukert, P.D. (1972). Isolation and characterization of viruses from semen and the reproductive tract of male swine. *Journal of the American Veterinary Medical Association*, Vol. 161, No. 11, pp. 1306-1316.
- Plowright, W., Perry, C.T., Peirce, M.A., & Parker, J. (1970). Experimental infection of the argasid tick, *Ornithodoros moubata* porcinus, with African swine fever virus. *Archiv für die Gesamte Virusforschung*, Vol. 31, No. 1, pp. 33-50.
- Prieto, C. & Castro, J.M. (2005). Porcine reproductive and respiratory syndrome virus infection in the boar: a review. *Theriogenology*, Vol. 63, No. 1, pp. 1-16.
- Prieto, C., Garcia, C., Simarro, I., & Castro, J.M. (2003). Temporal localization of porcine reproductive and respiratory syndrome virus in reproductive tissues of experimentally infected boars. *Theriogenology*, Vol. 60, No. 8, pp. 1505-1514.
- Prieto, C., Garcia, C., Simarro, I., & Castro, J.M. (2004). Temporal shedding and persistence of porcine reproductive and respiratory syndrome virus in boars. *Veterinary Record*, Vol. 154, No. 26, pp. 824-827.
- Prieto, C., Sanchez, R., Martin-Rillo, S., Suarez, P., Simarro, I., Solana, A., & Castro, J.M. (1996a). Exposure of gilts in early gestation to porcine reproductive and respiratory syndrome virus. *Veterinary Record*, Vol. 138, No. 22, pp. 536-539.
- Prieto, C., Suarez, P., Bautista, J.M., Sanchez, R., Rillo, S.M., Simarro, I., Solana, A., & Castro, J.M. (1996b). Semen changes in boars after experimental infection with porcine reproductive and respiratory syndrome (PRRS) virus. *Theriogenology*, Vol. 45, No. 2, pp. 383-395.
- Prieto, C., Suarez, P., Simarro, I., Garcia, C., Martin-Rillo, S., & Castro, J.M. (1997). Insemination of susceptible and preimmunized gilts with boar semen containing porcine reproductive and respiratory syndrome virus. *Theriogenology*, Vol. 47, No. 3, pp. 647-654.
- Reicks, D.L., Munoz-Zanzi, C., & Rossow, K. (2006). Sampling of adult boars during early infection with porcine reproductive and respiratory syndrome virus for testing by polymerase chain reaction using new blood collection technique (blood swab method). *Journal of Swine Health and Production*, Vol. 14, pp. 258-264.
- Ressang, A.A. (1973). Studies on the pathogenesis of hog cholera. I. Demonstration of hog cholera virus subsequent to oral exposure. *Zentralblatt für Veterinärmedizin Reihe B*, Vol. 20, No. 4, pp. 256-271.
- Robertson, I.B. (1992). Transmission of blue-eared pig disease. *Veterinary Record*, Vol. 130, No. 21, p. 478.

- Roizman, B. (1982). The family Herpesviridae: General description, taxonomy, and classification, In: *The Herpesviruses*, Vol. 1, pp. 1-23, Plenum Press, ISBN New York
- Rose, N., Larour, G., Le, D.G., Eveno, E., Jolly, J.P., Blanchard, P., Oger, A., Le, D.M., Jestin, A., & Madec, F. (2003). Risk factors for porcine post-weaning multisystemic wasting syndrome (PMWS) in 149 French farrow-to-finish herds. *Preventive Veterinary Medicine*, Vol. 61, No. 3, pp. 209-225.
- Rossow, K.D. (1998). Porcine reproductive and respiratory syndrome. *Veterinary Pathology*, Vol. 35, No. 1, pp. 1-20.
- Schlafer, D.H. & Mebus, C.A. (1987). Abortion in sows experimentally infected with African swine fever virus: pathogenesis studies. *American Journal of Veterinary Research*, Vol. 48, No. 2, pp. 246-254.
- Scobie, L., Taylor, S., Wood, J.C., Suling, K.M., Quinn, G., Meikle, S., Patience, C., Schuurman, H.J., & Onions, D.E. (2004). Absence of replication-competent human-tropic porcine endogenous retroviruses in the germ line DNA of inbred miniature Swine. *Journal of Virology*, Vol. 78, No. 5, pp. 2502-2509.
- Segalés, J., Allan, G.M., & Domingo, M. (2005). Porcine circovirus diseases. *Animal Health Research Reviews*, Vol. 6, No. 2, pp. 119-142.
- Segalés, J., Olvera, A., Grau-Roma, L., Charreyre, C., Nauwynck, H., Larsen, L., Dupont, K., McCullough, K., Ellis, J., Krakowka, S., Mankertz, A., Fredholm, M., Fossum, C., Timmusk, S., Stockhofe-Zurwieden, N., Beattie, V., Armstrong, D., Grassland, B., Baekbo, P., & Allan, G. (2008). PCV-2 genotype definition and nomenclature. *Veterinary Record*, Vol. 162, No. 26, pp. 867-868.
- Shin, J., Torrison, J., Choi, C.S., Gonzalez, S.M., Crabo, B.G., & Molitor, T.W. (1997). Monitoring of porcine reproductive and respiratory syndrome virus infection in boars. *Veterinary Microbiology*, Vol. 55, No. 1-4, pp. 337-346.
- Shirai, J., Narita, M., Iijima, Y., & Kawamura, H. (1985). A cytomegalovirus isolated from swine testicle cell culture. *Nippon Juigaku Zasshi*, Vol. 47, No. 5, pp. 697-703.
- Sibila, M., Calsamiglia, M., Segalés, J., Blanchard, P., Badiella, L., Le, D.M., Jestin, A., & Domingo, M. (2004). Use of a polymerase chain reaction assay and an ELISA to monitor porcine circovirus type 2 infection in pigs from farms with and without postweaning multisystemic wasting syndrome. *American Journal of Veterinary Research*, Vol. 65, No. 1, pp. 88-92.
- Siegel, A.M. & Weigel, R.M. (1999). Herd factors affecting the selection and success of intervention strategies in the program for eradication of pseudorabies (Aujeszky's disease) virus from Illinois swine farms. *Preventive Veterinary Medicine*, Vol. 40, No. 3-4, pp. 243-259.
- Singleton, W.L. (2001). State of the art in artificial insemination of pigs in the United States. *Theriogenology*, Vol. 56, No. 8, pp. 1305-1310.
- Solis, M., Ramirez-Mendoza, H., Mercado, C., Espinosa, S., Vallejo, V., Reyes-Leyva, J., & Hernandez, J. (2007). Semen alterations in porcine rubulavirus-infected boars are related to viral excretion and have implications for artificial insemination. *Research in Veterinary Science*, Vol. 83, No. 3, pp. 403-409.

- Solomon, T., Ni, H., Beasley, D.W., Ekkelenkamp, M., Cardoso, M.J., & Barrett, A.D. (2003). Origin and evolution of Japanese encephalitis virus in southeast Asia. *Journal of Virology*, Vol. 77, No. 5, pp. 3091-3098.
- Specke, V., Tacke, S.J., Boller, K., Schwendemann, J., & Denner, J. (2001). Porcine endogenous retroviruses: in vitro host range and attempts to establish small animal models. *Journal of General Virology*, Vol. 82, pp. 837-844.
- Spradbrow, P.B. (Sept. 1968). The isolation of infectious bovine rhinotracheitis virus from bovine semen. *Australian Veterinary Journal*, Vol. 44, No. 9, pp. 410-412.
- Stevenson, G.W., Van Alstine, W.G., & Kanitz, C.L. (1994). Characterization of infection with endemic porcine reproductive and respiratory syndrome virus in a swine herd. *Journal of the American Medical Association*, Vol. 204, No. 12, pp. 1938-1942.
- Sur, J.H., Doster, A.R., Christian, J.S., Galeota, J.A., Wills, R.W., Zimmerman, J.J., & Osorio, F.A. (1997). Porcine reproductive and respiratory syndrome virus replicates in testicular germ cells, alters spermatogenesis, and induces germ cell death by apoptosis. *Journal of Virology*, Vol. 71, No. 12, pp. 9170-9179.
- Swenson, S.L., Hill, H.T., Zimmerman, J.J., Evans, L.E., Landgraf, J.G., Wills, R.W., Sanderson, T.P., McGinley, M.J., Brevik, A.K., & Ciszewski, D.K. (1994a). Excretion of porcine reproductive and respiratory syndrome virus in semen after experimentally induced infection in boars. *Journal of the American Medical Association*, Vol. 204, No. 12, pp. 1943-1948.
- Swenson, S.L., Hill, H.T., Zimmerman, J.J., Evans, L.E., Wills, R.W., Yoon, K.J., Schwartz, K.J., Althouse, G.C., McGinley, M.J., & Brevik, A.K. (1994b). Artificial insemination of gilts with porcine reproductive and respiratory syndrome (PRRS) virus-contaminated semen. *Swine Health and Production*, Vol. 2, No. 6, pp. 19-23.
- Tabares, E., Marcotegui, M.A., Fernandez, M., & Sanchez-Botija, C. (1980). Proteins specified by African swine fever virus. I. Analysis of viral structural proteins and antigenic properties. *Archives of Virology*, Vol. 66, No. 2, pp. 107-117.
- Takeuchi, Y., Patience, C., Magre, S., Weiss, R.A., Banerjee, P.T., Le, T.P., & Stoye, J.P. (1998). Host range and interference studies of three classes of pig endogenous retrovirus. *Journal of Virology*, Vol. 72, No. 12, pp. 9986-9991.
- Terpstra, C. & Wensvoort, G. (1997). A congenital persistent infection of bovine virus diarrhoea virus in pigs: clinical, virological and immunological observations. *Veterinary Quarterly*, Vol. 19, No. 3, pp. 97-101.
- Terpstra, C., Wensvoort, G., & Pol, J.M. (1991). Experimental reproduction of porcine epidemic abortion and respiratory syndrome (mystery swine disease) by infection with Lelystad virus: Koch's postulates fulfilled. *Veterinary Quarterly*, Vol. 13, No. 3, pp. 131-136.
- Teuffert, J., Schluter, H., & Muller, T. (1998). [Boar semen--a possible risk factor in infection occurrence of porcine reproductive and respiratory syndrome]. *Deutsche Tierärztliche Wochenschrift*, Vol. 105, No. 9, pp. 340-345.
- Thacker, B.J., Joo, H.S., Winkelman, N.L., Leman, A.D., & Barnes, D.M. (1987a). Clinical, virologic, and histopathologic observations of induced porcine parvovirus infection in boars. *American Journal of Veterinary Research*, Vol. 48, No. 5, pp. 763-767.

- Thacker, B.J., Joo, H.S., Winkelman, N.L., Leman, A.D., & Barnes, D.M. (May 1987b). Clinical, virologic, and histopathologic observations of induced porcine parvovirus infection in boars. *American Journal of Veterinary Research*, Vol. 48, No. 5, pp. 763-767.
- Thacker, B.J., Larsen, R.E., Joo, H.S., & Leman, A.D. (Sept. 1984). Swine diseases transmissible with artificial insemination. *J Am.Vet Med Assoc.*, Vol.185, No.5, pp.511-516.
- Tischer, I., Gelderblom, H., Vettermann, W., & Koch, M.A. (1982). A very small porcine virus with circular single-stranded DNA. *Nature*, Vol. 295, No. 5844, pp. 64-66.
- Tischer, I., Miels, W., Wolff, D., Vagt, M., & Griem, W. (1986). Studies on epidemiology and pathogenicity of porcine circovirus. *Archives of Virology*, Vol. 91, No. 3-4, pp. 271-276.
- Tucker, A.W., Mellencamp, M.M., Donadeu, M., & Scobie, L. (2006). Retroviremia in commercial pigs and its preliminary association with poor health. *Journal of Clinical Microbiology*, Vol. 44, No. 10, pp. 3846-3847.
- Vannier, P. & Gueguen, B. (1979). Excrétion du virus de la maladie d'Aujeszky par les voies génitales mâles du porc. *Journee Recherche Porcine*, pp. 4001-406
- Wensvoort, G., Terpstra, C., Pol, J.M., ter Laak, E.A., Bloemraad, M., de Kluyver, E.P., Kragten, C., van, B.L., den, B.A., Wagenaar, F., & . (1991). Mystery swine disease in The Netherlands: the isolation of Lelystad virus. *Veterinary Quarterly*, Vol. 13, No. 3, pp. 121-130.
- West, K.H., Bystrom, J.M., Wojnarowicz, C., Shantz, N., Jacobson, M., Allan, G.M., Haines, D.M., Clark, E.G., Krakowka, S., McNeilly, F., Konoby, C., Martin, K., & Ellis, J.A. (1999). Myocarditis and abortion associated with intrauterine infection of sows with porcine circovirus 2. *Journal of Veterinary Diagnostic Investigation*, Vol. 11, No. 6, pp. 530-532.
- Wilson, C.A. (2008). Porcine endogenous retroviruses and xenotransplantation. *Cellular and Molecular Life Sciences*, Vol. 65, No. 21, pp. 3399-3412.
- Wilson, C.A., Wong, S., VanBrocklin, M., & Federspiel, M.J. (2000). Extended analysis of the in vitro tropism of porcine endogenous retrovirus. *Journal of Virology*, Vol. 74, No. 1, pp. 49-56.
- Wittmann, G. (1985). Aujeszky's disease: factors important for epizootiology and control. *Review of Science Technology OIE*, Vol. 4, pp. 5-44.
- Wood, J.C., Quinn, G., Suling, K.M., Oldmixon, B.A., Van Tine, B.A., Cina, R., Arn, S., Huang, C.A., Scobie, L., Onions, D.E., Sachs, D.H., Schuurman, H.J., Fishman, J.A., & Patience, C. (2004). Identification of exogenous forms of human-tropic porcine endogenous retrovirus in miniature Swine. *Journal of Virology*, Vol. 78, No. 5, pp. 2494-2501.
- Wrathall, A.E. & Mengeling, W.L. (1979). Effect of inseminating seropositive gilts with semen containing porcine parvovirus. *British Veterinary Journal*, Vol. 135, No. 5, pp. 420-425.
- Wright, P.F. & Webster, R.G. (2001). Orthomyxoviruses, In: *Field's Virology*, Knipe, D.M., Howley, P.M., Griffin, D.E., Lamb, R.A., Martin, M.A., Roizman, B., & Straus, S.E., pp. 1533-1579, Lippincott-Raven Publishers, ISBN Philadelphia.

Yaeger, M.J., Prieve, T., Collins, J., Christopher-Hennings, J., Nelson, E., & Benfield, D. (1993). Evidence for the transmission of porcine reproductive and respiratory syndrome (PRRS) virus in boar semen. *Swine Health and Production*, Vol. 1, No. 5, pp. 7-9.

IntechOpen

IntechOpen

A Bird's-Eye View of Veterinary Medicine

Edited by Dr. Carlos C. Perez-Marin

ISBN 978-953-51-0031-7

Hard cover, 626 pages

Publisher InTech

Published online 22, February, 2012

Published in print edition February, 2012

Veterinary medicine is advancing at a very rapid pace, particularly given the breadth of the discipline. This book examines new developments covering a wide range of issues from health and welfare in livestock, pets, and wild animals to public health supervision and biomedical research. As well as containing reviews offering fresh insight into specific issues, this book includes a selection of scientific articles which help to chart the advance of this science. The book is divided into several sections. The opening chapters cover the veterinary profession and veterinary science in general, while later chapters look at specific aspects of applied veterinary medicine in pets and in livestock. Finally, research papers are grouped by specialisms with a view to exploring progress in areas such as organ transplantation, therapeutic use of natural substances, and the use of new diagnostic techniques for disease control. This book was produced during World Veterinary Year 2011, which marked the 250th anniversary of the veterinary profession. It provides a fittingly concise and enjoyable overview of the whole science of veterinary medicine.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Tanja Opriessnig, Luis G. Giménez-Lirola and Patrick G. Halbur (2012). 14_Artificial Insemination and Its Role in Transmission of Swine Viruses, A Bird's-Eye View of Veterinary Medicine, Dr. Carlos C. Perez-Marin (Ed.), ISBN: 978-953-51-0031-7, InTech, Available from: <http://www.intechopen.com/books/a-bird-s-eye-view-of-veterinary-medicine/artificial-insemination-and-its-role-in-transmission-of-swine-viruses>

INTech
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](https://creativecommons.org/licenses/by/3.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen