We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

***In Vitro* Selection of Salt Tolerant Calli Lines and Regeneration of Salt Tolerant Plantlets in Mung Bean (*Vigna radiata* L. Wilczek)**

Srinath Rao* and Prabhavathi Patil

Department of Botany, Gulbarga University, Gulbarga Karnataka, India

1. Introduction

Salinity is one of the important abiotic factors in limiting plant productivity (Munns, 2002). 19.5% of the irrigated agricultural land is considered saline (Flowers & Yeo, 1995). Although minerals are essential for plants, their excess quantity in the soil is injurious to plants. Plants exposed to saline environment suffer from ion excess or water deficit and oxidative stress linked to the production of reactive oxygen species (ROS), which cause damage to lipids, proteins and nucleic acids (Hernandez, *et al.*, 2000). Oxidative stress is considered to be one of the major damaging factors in plant cells exposed to salinity (Gossette, *et al.*, 1994; Hernandez, *et al.*, 1995; Khan & Panda, 2002; Queiros *et al.*, 2007). The process of salt response and tolerance has been studied at the whole plant level (Hasegawa *et al.*, 2000; Jogeshwar, *et al.*, 2006). However the structural complexity of the whole plant makes it difficult to separate systemic from cellular salinity tolerance mechanism (Hawkins & Lips, 1997). The importance of plant tissue culture in the improvement of salt tolerance in plants has been pointed long back (Dix, 1993; Hasegawa, *et al.*, 1994; Nabors *et al.*, 1980; Tal 1994). In recent years tissue culture techniques are being used as a useful tool to elucidate the mechanism involved in salt tolerance by using *in vitro* selected salt tolerant cell lines (Davenport, *et al.*, 2003; Gu, *et al.* 2004; Lutts *et al.*, 2004 ; Naik & Harinath, 1998; Rao & Patil. 1999; Venkataiah. *et al.*, 2004). Besides, these lines have been used to regenerate salt tolerant plants (Chen *et al.*, 200; Jaiwal & Singh, 2001; Miki *et al.* 2001; Ochatt *et al.*, 1999; Rao & Krupanidhi, 1996; Shankhdhar, *et al.*, 2000). The selection of crop varieties for greater tolerance to saline environment will allow greater productivity from large saline lands.

2. Materials and methods

2.1 Callus induction and culture

Seeds of mung bean cultivar (cv) PS-16 were obtained from Agriculture Research Station Gulbarga, India. Seeds were washed thoroughly and imbibed in sterile water for six hours and then soaked in 70% (v/v) ethanol for 4 min and then surface sterilized for 1 min by 0.1% HgCl₂ (m/v). The seeds were germinated on sterile filter paper soaked in sterile water in

culture tubes and allowed to germinate for 5 days at 16/8h (light/dark) photoperiod at $42 \mu \text{mol m}^{-2} \text{s}^{-1}$ irradiance provided by cool fluorescent tubes and incubated at $26 \pm 1^\circ \text{C}$. Cotyledons from 5 days old seedlings were used as source of explants for callus induction. The cotyledons were cut along the four side (i.e. leaf apex, petiole region and leaf margin of both the sides) with a sterile scalpel and cultured on Murashige and Skkog's (MS) medium supplemented with $9.09 \mu \text{M}$ 2, 4-Dichlorophenoxy acetic acid (2,4-D) and $5.76 \mu \text{M}$ kinetin(Kn) solidified with 8% (m/v) agar, before solidifying the pH of the media was adjusted to 5.6, the cultures were incubated under 16/8h (light/dark) photoperiod and allowed to grow for four weeks.

2.2 Selection of salt tolerant lines

Callus tissue grown on MS medium as described above was directly exposed to different concentrations of NaCl (50, 100, 150 and 200 mM) and the concentration at which growth was completely inhibited was determined. Cell survival was nonexistent at 200 mM. Consequently a concentration of 150 mM was used for selection and a portion of calli developed at this concentration was thereafter sub- cultivated on fresh MS medium containing same concentration (150mM) of NaCl and allowed to grow for 8 weeks. After this, calli were then transferred on MS medium without NaCl for 4 weeks. To test the stability of salt tolerant trait. Callus line growing healthily, on medium without NaCl were again transferred to MS medium containing 150 mM NaCl for 4 weeks, surviving callus lines showing good growth at this stage was considered to be NaCl tolerant. The salt tolerant calli was picked up and were further grown for a period of 4 weeks and used for enzyme, protein and proline estimation.

2.3 Assessment of growth

Callus tissue (~250 mg) of both selected and non-selected were grown MS medium supplemented with $9.09 \mu \text{M}$ 2, 4-D and $5.76 \mu \text{M}$ Kin containing 0-150 mM NaCl. The increase in fresh weight of the callus was determined after four weeks. Three replicates each consisting of 10 culture tubes were maintained per treatment, the relative growth rate of callus was calculated as the $(\text{FM}_f - \text{FM}_i) / \text{FM}_i$, where FM_f and FM_i were the final and initial fresh weight respectively. Samples of callus tissues were stored in -80°C for further analysis.

2.4 Lipid peroxidation

Lipid peroxidation was determined in terms of Malondialdehyde (MDA) content using the Thiobarbitric acid reaction (TBARS) according to Heath and Packer (1968). Frozen callus tissue was ground to a fine powder in liquid nitrogen with a mortar and pestle. Subsequently about 100 mg of powder was homogenized in 5 cm^3 of TCA (0.1% w/v) and centrifuged at 10,000g for 10min at 5°C . 1 cm^3 of supernatant was mixed with 4 cm^3 of 0.5% TBA reagent in 20% TCA. The mixture was then heated at 95°C for 30 min, cooled over ice and centrifuged at 10,000g for 10 min. The absorbance of the supernatant was recorded at 532 nm and corrected for non specific absorbance at 600 nm. MDA content was calculated using an extinction coefficient (ϵ) of $155 \text{ mM}^{-1} \text{ cm}^{-1}$ and expressed as $\text{nmol g}^{-1} \text{FW}$.

2.5 Determination of ascorbate

Ascorbic acid (As A) was extracted in 5% (m/v) methophosphoric acid with sand at 4 °C. The homogenate was then centrifuged at 3000 g for 20 min at 4 °C. Ascorbate content was quantified in the supernatant as described by Shukla, *et al.*, (1979). An aliquot of 1 cm³ of the supernatant was mixed with 2.5 cm³ of 1% (v/v) freshly diluted Folin - Ciocalteu reagent. The reaction mixture was allowed to stand for 40 min at room temperature, then the absorbance was recorded at 730 nm, using ascorbic acid as standard.

2.6 Superoxide dismutase activity (SOD) (EC 1.15.11)

The activity of SOD was determined based on its ability to inhibit the auto oxidation of pyrogallol. The measurement was based on the modified method of Markland and Markland, (1947). The reaction mixture consisted of 0.25 M pyrogallol in 0.1 M sodium phosphate buffer (pH 7.4) and 0.1 cm³ of enzyme extract. The reaction was initiated by light illumination and the rate of oxidation was measured spectrophotometrically at 429 nm as per the procedure described by Nakono & Asada, (1981). SOD activity is expressed in units/mg protein. One unit is defined as the amount of the enzyme which caused 50% inhibition of pyrogallol oxidation.

2.7 Ascorbate peroxidase activity (APX)

The activity of APX was determined by measuring the decrease in absorbance at 290 nm and the amount of Ascorbate oxidized to dehydroascorbate was calculated from the extinction coefficient 2.8 (Nakano and Asada, 1981). The reaction mixture (2 cm³) consisted of 25 mM phosphate buffer (pH 7.0), 0.1 mM EDTA, 0.25 mM ascorbic acid, 1.0 mM H₂O₂ and 0.2 cm³ enzyme extract.

2.8 Catalase activity (CAT; EC 1.11.1.6)

Catalase activity was measured according to method of Chandlee and Scandalios (1984). One gram of frozen callus was homogenized in a pre chilled pestle & mortar with 5 ml of ice cold 50 mM phosphate buffer. The extract was centrifuged at 4°C for 20 Min. at 12,500 X g. The supernatant was used for enzyme assay. The assay mixture contained 2.6 ml of 50 mM potassium phosphate buffer (pH 7.0) 0.4 cm² 15 mM H₂O₂ and 0.1 cm² of enzyme extract. The decomposition of H₂O₂ was followed by decline in absorbance at 240 nm.

2.9 Peroxidase activity (POX; EC 1.11.1.7)

Peroxidase activity was assayed by the method of Kumar and Khanna (1982). Assay mixture contained 2 ml of 0.1 M Phosphate buffer (pH 6.8) 1ml of 0.01M pyrogallol, 1 ml of 0.005 M H₂O₂ and 0.5 ml of enzyme extract. The solution was incubated for 5 min at 25 °C after which the reaction was terminated by adding 1 ml of 2.5N H₂SO₄. The purpurogallin formed was determined by measuring the absorbance at 420 nm against a blank prepared by adding the extract after the addition of 2.5N H₂SO₄ at zero times. The activity was expressed in unit mg⁻¹ protein. One unit (U) is defined as the change in the observance by 0.1 min⁻¹ mg⁻¹ protein.

2.10 Protein estimation

Samples of frozen callus tissue were ground in Tris-HCl (60 mM, pH 6.8) at 4°C, using prechilled mortar and pestle. The extract was centrifuged (4000g for 15 min at 4°C) and the supernatant was used for protein estimation according to Bradford (1976).

2.11 Proline estimation

Proline was extracted and determined calorimetrically by the method of Bates, *et al.*, (1973). Callus tissue (~200 mg) was homogenized with 4 cm³ of 50 mM phosphate buffer (pH 7.8) containing 1% (w/v) polyvinyl pyrrolidone (PVP) 0.01% (w/v) Triton X-100, and centrifuged at 800 rpm for 15 min. Proline was determined in the supernatant by measuring the absorbance of proline-ninhydrin product formed, at 520 nm in a spectrophotometer using toluene as a solvent.

2.12 Regeneration of plantlets from NaCl tolerant calli

Plantlets were regenerated from salt tolerant and non-tolerant (control callus) callus. For regeneration of plantlets, callus was transferred in fragments of about 300 mg directly to MS medium supplemented with 0.5-2 mg/l BAP, alone or in combination with 0.25 and 0.5 mg/l NAA. After the shoots reached a height of 3-4 cm individual shoots were excised and transferred to rooting media which consisted of MS medium supplemented with 0.5 and 1 mg/l IBA.

2.13 NaCl tolerance test of regenerated plantlets

After the regenerated shoots produced roots, they were transferred to Erlenmeyer flasks containing the same rooting medium supplemented with 150 mM NaCl (Chen, *et al.*, 2001). The survival rate of plantlets regenerated from non selected callus and NaCl selected callus was determined by using the formula

$$\frac{\text{The number of surviving plantlets} \times 100}{\text{Total number of tested plantlets}}$$

2.14 Statistical analysis

The results presented are an average over three independent experiments. Data were processed with analysis of variance (ANOVA) and the means were compared using students test at a significance level of = 0.05.

3. Results and discussion

3.1 Establishment of salt tolerant lines

The NaCl tolerant mungbean cell lines obtained in this study were selected from callus cultures initiated from cotyledon explants. The direct recurrent selection procedures were successful in selection of mungbean cell lines exhibiting tolerance to 150 mM NaCl.

Callus tissue cultured on medium containing 50 mM NaCl showed good cell proliferation and appeared morphologically similar to control, where as callus tissue cultured on medium

containing 100 mM NaCl showed moderate cell proliferation and small portions showing brownish color indicating early stages of necrosis. When callus tissue was cultured on medium supplemented with 150 mM NaCl, a great portion of the cells did not show any sign of growth and turned brown within 2 weeks of culture. However, small clusters of cells survived on 150 mM NaCl containing medium. Subsequently the cell clumps that proliferated in the presence of the salt were picked up and sub cultivated on NaCl free medium for 4 weeks. The salt tolerant cell lines were again transferred on to a fresh medium supplemented with 150 mM NaCl. On this medium cell clumps proliferated in to a mass of callus exhibiting friable nature. The selection processes resulted in establishment of NaCl-tolerant callus lines showing good proliferation of callus.

Cell lines exhibiting tolerance to NaCl have been previously selected from a range of legumes Gosal & Bajaj,1984) pasture legumes, (Smith & McComb, 1981) chick pea, (Pandey & Ganapathy,1984,1985) Alfalfa (Wincicov, 1991) red gram (Rao & Krupanidhi,1996) pea (Ochatt, *et al.*, 1999) soybean (Liu & Van Staden,2000) ground nut (Jain, *et al.*, 2001). And in several other plant species such as, canola (Tyagi & Rangaswamy, 1997, Jain, *et al.*, 1991a, b) finger millet, (Pius, *et al.* 1993) Cotton (Gosset, *et al.*, 1994) Potato (Olmos & Hellin, 1996; Queiros, *et al.*, 2007) rice (Dang & Nguyen, 2003; Shankdhar *et al.*, 2000) Sunflower (Davenport, *et al.*, 2003) chilli (Venkataiah, *et al.*, 2004) sugarcane (Gandonou, *et al.*, 2006).

3.2 Growth of callus

Growth rate of adapted callus line in NaCl supplemented media was steady and sustainable but it was lower than the growth of the control callus grown on stress-free medium.

Reduction in the growth is a common phenomenon in cultured cells grown on medium supplemented with NaCl (Cushman *et al.*, 1990; Greenway & Munns, 1980; Jain *et al.*, 1991 a, b; Luts *et al.*, 2004; Rus *et al.* 1999; Shankdhar *et al.*, 2000; Thomas *et al.*, 1992; Venkataiah *et al.*, 2004) and it has been interpreted that a certain amount of the total energy available for tissue metabolism is channeled to resist the stress (Cushman, *et al.*, 1990). However 200 mM NaCl tolerant calli obtained either by direct or indirect selection process did not sustained a regular growth on salt supplemented media, these calli suffered with time a decrease in growth and all callus tissues died when the culture period was prolonged beyond one month. A similar response was reported in salt tolerant potato calli lines (Queiros, *et al.*, 2007) and in *Chrysanthemum* sp. (Hossain, *et al.*, 2007).

However, at a given concentration of NaCl, selected calli maintained higher growth rate than non-selected calli when grown on NaCl supplemented medium (Table-1). Similar to our findings, Kumar and Sharma (1989), Patil & Rao (1999) & Gulati & Jaiwal (2010) reported that NaCl tolerant lines maintained higher growth rate when compared to non selected calli in this species. Most of the previously published reports in legumes indicated that selected cell lines maintained higher rate of growth on saline media when compared to their wild type for example, Pea (Hassan & Wilkins, 1988; Olms, *et al.*, 1995) red gram (Krupanidhi & Rao,1996) and in other plants like tobacco (Hasegawa *et al.*, 1980) colt cherry (Ochatt & Power, 1989) rice (Basu, *et al.*, 2002; Kishor & Reddy, 1986) finger millet (Pius, *et al.*, 1993) sugar cane (Gandonou, *et al.*, 2006).

Serial No	NaCl (µm)	non- selected callus (mg)	selected callus (mg)	MDA (nmol g ⁻¹ freshweight)	
				Non-selected callus (mg)	Selected callus (mg)
1	00 (Control)	778.0 ± 0.67 ^b	780.0 ± 0.45 ^b		
2	50	806.6 ± 0.34 ^a	865.3 ±0.83 ^a	84.0±0.75 ^c	82.6±0.13 ^c
3	100	647.6± 0.97 ^c	740.0 ±0.56 ^c	90.0 ± 0.86 ^c	82.3±0.21 ^c
4	150	449.3 ± 0.53 ^d	614.6±0.36 ^d	140.6 ±0.54 ^b	88.0±0.43 ^b

Data represents average of three replicates; each replicate consists of 10 cultures. Mean ± Standard error. Mean followed by the same superscript in a column is not significantly different at P = 0.05 levels.

Table 1. Effect of NaCl on fresh weight and MDA content of selected and non-selected callus of *Vigna radiata* L.

3.3 Analysis of proline and protein contents

Proline and protein content in the selected and non-selected callus line is presented in table-2. Both protein and proline contents were significantly ($p < 0.05$) affected with an increase in the concentration of NaCl. NaCl tolerant callus line maintained increased protein levels than control callus when exposed to 150 mM NaCl. Increased protein content in NaCl tolerant callus has been reported earlier (Chen, *et al.*, 2001; Queiros, *et al.*, 2007). Extra protein bands in *In Vitro* raised NaCl tolerant *V. radiata* plants have been reported by (Hassan, *et al.*, 2004). Similarly a gradual increase in the Proline content of tolerant callus line was noticed with an increase in the concentration of NaCl from 100 to 150 mM, maximum being at 150 mM NaCl level. (Kumar & Sharma, 1989; Patil & Rao, 1999) reported higher proline in selected calli when compared to non selected calli in this species. Accumulation of proline has been widely advocated for use as parameter of selection of cell lines for salt tolerance and it has been shown that proline over producing lines are more tolerant than their parent cultivars (Kavi Kishor *et al.*, 1995). Increased proline levels in NaCl tolerant callus has been earlier reported in many legumes like, soybean (Liu, & Van Staden, 2000) ground nut (Jain, *et al.*, 2001) and in other plants like tobacco (Wattad, *et al.*, 1983) canola (Jain, *et al.*, 1991 a,b; SashiMadan, *et al.*, 1995) tobacco (Gangopadhyay, *et al.*, 1997b) barley (Chaudhuri, *et al.*, 1997) *Citrus* (Piqueras, *et al.*, 1996) alfalfa (Petrusa, *et al.*, 1997) basmati rice (Basu, *et al.*, 2002) bamboo (Singh, *et al.*, 2003) chili (Venkataiah, *et al.*, 2004) sugarcane (Gandonou, *et al.*, 2006). On the contrary some workers did not observe any appreciable increase in free proline content (Dix & Pearce, 1981, Errabii, *et al.*, 2006; Jain, *et al.*, 1987). Higher proline levels in salt tolerant callus may be due to an increased rate of synthesis or a decreased rate of oxidation of this compound as suggested by Wyn-Jones & Gorham (1984). Patnaik & Dabata (1997) suggested that salinity induced proline accumulation could be due to putrescence oxidation. There are divergent reports concerning the role of proline in salt tolerance. High levels of proline in NaCl selected cells have been suggested as a factor conferring salt tolerance (Kumar & Sharma, 1989; Pandey & Ganapathy, 1985; Wattad *et al.*, 1983). In contrast proline accumulation has also been excluded as the mechanism of tolerance in some cases (Dix & Pearce, 1981; Errabii *et al.*, 2007; Tal *et al.* 1979). The significant levels of proline observed in our study supports the former. It is well known fact that proline stabilizes the structure and function of various macromolecules (Kavi Kishor, *et al.*, 2005; Rhodes, 1987; Smirnoff & Cumbes, 1989).

Serial No.		Protein mg g ⁻¹ FW		Proline mg g ⁻¹ FW		Ascorbic acid mg g ⁻¹ FW	
		Non-selected callus	Selected callus (mg)	Non-selected callus	Selected callus (mg)	Non-selected callus	Selected callus (mg)
1	00 (Control)	22.6 ±0.43 ^c	27.2 ± 0.76 ^c	7.6±0.53 ^c	8.8 ± 0.54 ^d	8.4 ± 0.44 ^a	22.9 ± 0.45 ^c
2	50	28.2 ±0.87 ^b	42,0±0.56 ^b	10.5 ±0.19 ^b	18.2 ±0.76 ^b	8.6±0.35 ^a	26.5 ± 0.45 ^b
3	100	32.2 ± 0.56 ^a	47.0 ± 0.23 ^a	16.2±0.65 ^a	28.2±0.55 ^a	6.4±0.87 ^b	42.7±0.50 ^a
4	150	16.4±0.32 ^d	19.2 ±0.65 ^d	5.9 ±0.25 ^d	12.2±0.22 ^c	5.5 ± 0.43 ^c	27.1±0.25 ^b

Data represents average of three experiments; Mean± standard error
Mean followed by the same super script in a column is not significantly different at
P= 0.05 level.

Table 2. Effect of NaCl on protein, proline, and Ascorbic acid content in non-selected and selected callus on *Vigna radiate* L.

3.4 Lipid peroxidation

Oxidative damage to lipids was determined as lipid peroxidation in terms of amount of malondialdehyde (MDA). MDA content of tolerant callus was less when compared to that of sensitive callus line (Table-1). Sumithra, *et al.*,2006) reported low levels of lipid peroxidation in salt tolerant *V. radiata* cv Pusa bold than in salt sensitive cv CO 4. Jain, *et al.*, (2001) reported 4 fold increases in MDA content of sensitive callus of ground nut when grown on saline medium where as in the tolerant callus line the increase was only 1.1 fold. The results indicate that NaCl tolerant callus maintained membrane integrity when growing in saline environment however sensitive calls line were unable to maintain membrane integrity under salinity stress resulting in decreased growth and metabolic imbalance. Changes in the cell wall have been shown to be important for salt adaptation (Binzel, *et al.*, 1985; Curz, *et al.*, 1992). Salt stress is known to result in extensive lipid per oxidation (Davenport, *et al.*, 2003; Hernandez, *et al.*, 2000; Khan & Panda, 200;; Kholova *et al.*, 2009; Queiros *et al.*, 2007). In the present investigations it was noticed that the extent of lipid peroxidation was higher in NaCl sensitive cell lines than NaCl adopted cell lines, such observations were also made in ground nut (Jain *et al.*, 2001) wheat (Sarin, *et al.*, 2005) eggplant (Yaser, *et al.*, 2006). It seems that due to low peroxidation in NaCl tolerant callus line, membrane integrity might have been maintained, thus preventing protein denaturation which is supported by the fact that NaCl tolerant callus lines had higher protein content. Lipid membranes are vulnerable targets for stress induced cellular damage and the extent of damage is commonly used as a measure of stress (Gadallah, 1999; Zhou, *et al.*, 1992).

3.5 Ascorbic acid content

Ascorbic acid content increased gradually in NaCl tolerant calli line than in the control line. An increase of about 18% was found in callus tissue grown at 50 mM NaCl, while in callus grown on medium supplemented with 100 and 150 mM NaCl ascorbic acid content

significantly increased to 30% and 59% respectively (Table-2). It was previously reported that the response of many plants cells to salinity is the increased synthesis of ascorbic acid. A positive correlation between ascorbic acid content and salinity was also reported in NaCl tolerant cell lines in several species (Gosset, *et al.*, 1996; Sarin, *et al.*, 2005; Qlmos & Hellin, 1996; Queiros, *et al.*, 2007). This antioxidant compound is one of the most effective free radical scavengers implicated in the adaptations of plants to stress (Shigeoka, *et al.*, 2002; Smirnoff, *et al.*, 2000; Queiros, *et al.*, 2007).It is reported that, high .levels of endogenous ascorbate is essential to effectively maintain the anti-oxidant system that protects plants from oxidative damage due to abiotic stress (Shigeoka, *et al.*, 2002).

3.6 Antioxidant enzymes

NaCl stress affect plant processes that lead to the formation of reactive oxygen species (ROS) super oxide radical (O₂) Hydrogen peroxide (H₂O₂) and hydroxide radicals (OH). The ROS cause oxidative damage to membrane lipids, proteins and nucleic acid. Accelerated detoxification is fundamental in development of tolerance to NaCl. To control the levels of ROS and protect the cells from injury under stress conditions, it is important that ROS should be scavenged.

Plants protect cells and sub cellular systems from the effects of reactive oxygen species (ROS) by enzymes such as Super oxide dismutase (SOD) Ascorbate peroxidase (APX) and Catalase (CAT). In the present investigation it was noticed that NaCl significantly altered the activities of SOD, APX & CAT with an increase in the levels of NaCl from 50 to 150 mM NaCl level. Maximum activity of these enzymes (Table-3) was noticed in callus tissue tolerant to 150 mM NaCl, where as in sensitive callus the activity of these enzymes was highly inhibited.

Serial No.	NaCl (µm)	SOD (EU mg ⁻¹ protein)		APX (EU mg ⁻¹ protein)		CAT (EU mg ⁻¹ protein)	
		Non-selected callus (mg)	Selected callus (mg)	Non-selected callus (mg)	Selected callus (mg)	Non-selected callus (mg)	Selected callus (mg)
1	00 (Control)	40.0 ± 0.76 ^b	40.6±0.34 ^c	0.95 ±0.23 ^b	0.88 ± 0.67 ^d	22.3 ± 0.56 ^a	22.9±0.41.6 ^c
2	50	42.8 ± 0.45 ^a	50.5 ± 0.56 ^b	1.40±0.12 ^a	2.10 ±0.56 ^b	20.9 ± 0.75 ^b	26.5 ± 0.45 ^b
3	100	38.2 ± 0.36 ^c	75.8 ± 0.65 ^a	0.71 ±0.34 ^c	2.94 ± 0.67 ^a	16.4 ± 0.67 ^c	42.7± 0.50 ^a
4	150	32.6±0.65 ^d	50. 6 ± 0.48 ^b	0.58 ± 0.56 ^b	1.80 ± 0.56	12.6 ± 0.45 ^d	27.1± 0.25 ^b

Data represents average of three experiments. Mean ± standard error.
Mean followed by the same superscript in a column is not significantly different at P= 0.05 levels.

Table 3. Effect of NaCl on antioxidant enzymes in non-selected and selected callus in *Vigna radiata* L.

Serial No.	Growth regulators (mg/l)	Frequency of shoot induction (%)	Number of shoot buds per explants.	Average length of plant let (cm)
1	BAP 0.5	0.0	00	00
2	BAP 1.0	21.66 ± 1.6 a	2.90 ± 0.16 a	3.8 ± 0.25 a
3	BAP 2.0	26.00 ± 1.4 b	8.20 ± 0.16 b	5.8 ± 0.34 b
4	BAP 2+NAA0.25	30.10 ± 1.8 c	12.80 ± 0.34 c	7.2 ± 0.55 c
5	BAP 2 +NAA 0.5	40.13 ± 1.8 d	14.80± 0.64 d	8.6 ± 0.68 c

Data represents average of three experiments. Mean ± standard error. Mean followed by the same superscript in a column is not significantly different at P= 0.05 levels.

Table 4. Effect of growth regulators on Frequency, Number of shoots, and mean shoot length from cotyledon derived callus in *Vigna radiata*.

Fig. 1. Photographs showing effect of NaCl on callus cultures of *Vigna radiata*

SOD, APX and CAT are considered as useful enzymes to help plants defend salt stress (Koca, *et al.* 2006, Shen, *et al.*, 2002). Sumithra *et al.*, (2006) have reported increased SOD and CAT activity in NaCl tolerant cultivars in this species. There are previous reports indicating increased activity of SOD in pea (Hernandez, *et al.*, 1993) wheat (Sairam, *et al.*, 2003) cotton (Gossett, *et al.*, 2004) potato (Queiros, *et al.*, 2007) sweet potato (He, *et al.*, 2008). Apart from SOD activity considerable increase in the activity of the enzymes APX and CAT was noticed in tolerant callus line than in sensitive callus when grown on different concentrations of NaCl. increase in the activities of SOD, CAT, APX in callus tissue raised from NaCl tolerant callus in cotton than in plants obtained from sensitive callus was reported by Gossett, *et al.*, (2004). Elkahoui, *et al.*, (2004) reported increased APX activities in NaCl adopted cell lines in periwinkle and Mittova, *et al.*, (2002) in salt tolerant wild tomatoes.

- A) Initiation of multiple leafy shoot buds on MS + 2 mg/l BAP + 0.5 mg/l NAA
- B) Further elongation of shoot after 25 days of culture
- C) Rooting of the shoot on MS + 1 mg/l IBA.
- D) A potted plant in a polycup containing 3 : 1 garden soil and sand.

Fig. 2. Photographs showing organogenesis from NaCl tolerant callus

Catalase is a common enzyme found in all living organisms which are exposed to oxygen, where it functions to catalyze the decomposition of H_2O_2 to water and oxygen (Chelikani, *et*

al., 2004). H_2O_2 is a harmful by product formed as a result of stress conditions to prevent damage it must be quickly converted into less dangerous substance like gaseous oxygen and water molecules. Hence the role of catalase is important in reducing stress related damages. Willekens, et al., (1997) suggested that the function of catalase in the cell is to remove the bulk of H_2O_2 where as peroxidase would be involved mainly in scavenging H_2O_2 that is not taken by catalase. In the present investigations it was noticed that both catalase and peroxidase activity increased in NaCl tolerant callus. From this observation it can be inferred that catalase/peroxidase might have acted co-operatively to remove H_2O_2 at a minimum expense of reducing power.

3.7 Regeneration of plantlets from NaCl selected and non selected callus

For regeneration of plantlets calli pieces approximately 250 ± 10 mg were transferred to MS medium supplemented with 0.5-2 mg/l BAP alone or in combination with 0.25 and 0.5 mg/l NAA. On MS medium without growth regulators and 0.5 mg/l shoot bud initiation was not observed; however on MS medium supplemented with 1 & 2 mg/l BAP shoots were initiated and the frequency of shoot formation was 21.66 and 26.00% respectively. Supplementing NAA at 0.25-0.5 mg/l further enhanced the frequency of shoot formation increased to 30.10 & 40.13% respectively.

Regeneration from various seedling explants viz shoot tip, cotyledons, cotyledonary node of mung on BAP supplemented medium has been reported earlier (Mathews, 1987; Gulati & Jaiwal, 1992; Kaviraj & Rao, 2009; Mendoza, et al., 1999). There is only one report of regeneration from callus cultures in mungbean (Rao, et al., 2005), similarly plantlets have been regenerated from NaCl tolerant callus in many species like *Citrus sinensis* (Ben-Hyyim & Goffer, 1989) *Brassica juncea* (Jain, et al., 1991; Kerthi, et al., 1989; Sashimadan, et al., 1995) *Triticum durum* (Zair, et al., 2003) *Ipomea batatas* (He, et al., 2009). In the present study however, it was noticed that the formation of shoots from NaCl tolerant callus was lower (14.00% than that from NaCl-non selected calli (40.13%) and the number of multiple shoots per callus piece was reduced to 3.33 from 14.80, a phenomenon reported earlier by few workers (Basu, et al., 1997; Chen et al., 2001; Ochatt, et al., 1999; Queiros, et al., 2007; Zair et al., 2003).

4. Conclusions

In conclusion NaCl stress induces severe oxidative stress in mungbean callus where the antioxidant defense system seemingly fails to combat with the stress induced oxidative damage. However it is possible to select callus line tolerant to elevated levels of NaCl stress by sudden exposure to high concentration of NaCl, accordingly a NaCl tolerant cell line was selected from cotyledon derived callus of mung bean which proved to be a true cell line variant. It can also be concluded that, the salt tolerant cell line could overcome the adverse effect of NaCl and maintained better growth on NaCl supplemented medium when compared to salt sensitive cell lines (controls). This conclusion is based on the following observations (a) cells which have been removed from the selection pressure for at least four passages retained tolerance to NaCl after transferring to NaCl (150 mM) medium (b) High accumulation of proline was noticed in salt tolerant cell lines compared to salt sensitive cell lines. (c) The activity of anti-oxidant enzymes was high. The results presented in this paper suggest that the mechanism of enhanced tolerance in mungbean callus is via improved

synergistic and protective effects of antioxidant enzyme like SOD, APX and CAT. It is also concluded that the callus lines were able to grow in elevated concentration (150mM NaCl) by maintaining membrane integrity, high levels of protein and proline, which supports the hypothesis that proline plays a significant role in protecting the cells from oxidative stress as reported in other grain legumes like chickpea and pigeonpea.

5. References

- Al-Naggar, AMM. Saker, MM. Shabana, R. Ghanem, SA. Reda, AH. & Eid, S. 2008 *In vitro* selection and molecular characterization of salt tolerant canola plantlets *Arab. J Biotech* 11(2): 207-218,
- Basu, S., Gangopadhyay, G., Gupta, S. & Mukherjee, BB. 1996. Screening for cross tolerance against related osmotic stress in adapted calli of salt sensitive and tolerance varieties of rice. *Phytomorphology*. 46(4): 357-364
- Basu, S., gangopadhyay, G., Mukherjee, BB. & Gupta, S. Plant regeneration of salt adapted callus of indica rice (var. Basmati370) in saline conditions. *Plant Tissue & Organ Culture* 50: 153-159.
- Ben-Hayyim, G. & Kochba, J. 1983 Aspects of salts Tolerance in a NaCl-Selected Stable Cell Line of *Citrus sinensis*. *Plant Physiol*. 72: 685-690
- Bhaskaran, S., Smith, RH. & Schertz, K. 1983 Sodium chloride tolerant callus of *Sorghum bicolor* (L.). *Z Pflanzenphysiol*. 112: 459-463
- Binzel, ML. Hasegawa, PM. Rhodes, RH. Handa, D. Handa, AK. & Raym, AB. 1987 Solute Accumulation in Tobacco Cells adapted to NaCl. *Plant Physiol*. 84: 1408-1415
- Binzel, MC. Hasegawa, PM. Handa, AK. & Bressan, RA. 1985 Adaptation of tobacco cells to NaCl. *Plant Physiol*. 79: 118-125
- Borochoy, NH. & Borochoy, A. 1991. Response of melon plants to salt 1: Growth. Morphology and root membrane properties", *J. Plant Physiol*. Vol. 139: Pp. 100-105
- Bradford, M.M.: A rapid and sensitive method for quantification of microgram quantities of protein utilizing the principle of protein dye binding. *Anal Biochem* 72: 248-254 1976
- Chaudhuri P Sengupta RK Ghosh P 1997 *In Vitro* assessment of salinity tolerance in *Hordeum vulgare*. *Indian J Plant Physiol* 2 (2): 123-126
- Chen, R., Gyokusen, K. & Saito, A. 2001. Selection, regeneration and protein profile characteristics of NaCl-Tolerant callus of *Robinia pseudoacacia* L. *J For Res* 6: 43-48
- Croughan, TP., Stavarek, SJ., Rains, DW. 1978. Selection of a NaCl tolerant line of cultured alfalfa cells. *Crop Sci* 18: 959-963
- Dix, PJ. 1993. The role of mutant cell lines in studies on environmental stress tolerance an assessment. *Plant J* 3: 309-313
- Dix, PJ. & Pearce, RS. 1981 Proline accumulation in NaCl-resistant and sensitive lines of *Nicotiana glauca*. *Zeitschrift fur Pflanzen-schutzs Physiologie* 102: 243-248
- Datta, KS., Kumar, A., Varma, SK. & Angrish, R. 1996 Effects of salinity on water relations and ion uptake in three tropical forage crops. *Indian J Plant Physiol* 1: 102-108
- Errabii, T., Gandonou, CB., Eassalmani, H. Abrini, J. Idaomar, M. & Skali-Senhaji, N. 2007. Effect of NaCl and mannitol induced stress on sugarcane (*Saccharum Sp.*) callus cultures. *Acta Physiol Plant* 29: 95-102

- Gadallah MAA 1999 Effect of proline and glycine-betaine on *Vicia faba* responses to salt stress. *Biol Plant* 42: 247-249,
- Gandonou CB Errabii T Abrani J Idamora M Skali-senhaji 2006 N Selection of callus cultures of sugarcane (*Saccharum* Sp.) tolerant to NaCl and their response to salt stress. *Plant Cell Tissue Organ Cult.* 87: 9-16
- Gangopadhyay G Basu S Mukherjee BB and Gupta S 1997 Effects of Salt and osmotic Shocks on unadapted and adapted callus lines of tobacco. *Plant Cell, Tissue Organ Cult.* 49: 45-52
- Gosal SS and Bajaj YPS 1984. Isolation of sodium chloride resistant cell lines in some grain legumes. *Indian Journal of Experimental Biology* 22: 209-214
- Gosset DR Millhollon EP Cranlucs M Banks SW and Marney M 1994 The effects of NaCl on antioxidant enzyme activities in callus tissue of salt tolerant and salt sensitive cotton cultivars. *Plant Cell Rep.* 13: 498-503
- Gossett DR Millhollon EP Lucas MC 1994 Antioxidant response to NaCl stress in salt-tolerant and salt-sensitive cultivars of cotton. *Crop Sci* 34: 706-714
- Greenway H Munns R 1980 Mechanism of salt tolerance in non- halophytes. *Ibid.* 31: 149-190
- Gu, R., Liu, Q., Pie, D. & Jiang, X. 2004 Understanding saline and osmotic tolerance of *Populus euphratica* suspended cells. *Plant Cell Tissue & Organ Culture* 78: 261-265
- Gulati, A. & Jaiwal, P.K. 1992 *In Vitro* induction of multiple shoots and regeneration from shoot tips of mung bean (*Vigna radiata* L. Wilczke). *Plant Cell Tissue & Organ Culture* 29: 199-205
- Hasegawa, P.M., Bressan, R.A., Nelson, D.E., Samaras, Y & Rhodes D. 1994. Tissue culture in the improvement of salt tolerance in plants. In: Monographs on Theoretical and Applied Genetics: Breeding plants with resistance to problem soils
- Hasegawa, P.M., Bressan, R.A. & Handa, A.K. 1980 Growth characteristics of NaCl selected and non-selected cells of *Nicotiana tabacum* L. *Plant and Cell Physiol.* 21 (8): 1347-1355
- Hassan NS and Willkins DA 1988 *In vitro* selection for salt tolerant lines in *Lycopersicon peruvianum*. *Plant Cell Reports.* 7: 463-466
- Hassan NM Serag MS El-Feky FM 2004 Changes in nitrogen content and protein profiles following *In vitro* selection of NaCl resistant mung bean and tomato. *Acta Physiologiae Plantarum.* 26(2): 165-175
- Hawkins HJ Lips SH 1997 Cell suspension cultures of *Solanum tuberosum* L. as a model system for N and salinity response. Effect of salinity on NO₃⁻ uptake and PM-ATPase activity. *J Plant Physiol* 150: 103-109
- Heath RL Packer L 1968 Photo peroxidation in isolated chloroplasts 1. Kinetics and stoichiometry of fatty acid peroxidation. *Arch Biochem Bio Phys* 125: 189-198
- Hernandez, J.A., Olmos, E., Corpas, F.J., Sevilla, F., Del Rio, L.A Salt induced oxidative stress in chloroplasts of pea plants *Plant Sci.* 105: 151-167, 1995.
- Hernandez JA Jimenez A Mullineaux P Sevilla F 2000 Tolerance of pea (*Pisum sativum* L.) to long term salt stress is associated with induction of antioxidant defenses. *Plant Cell Environ.* 23: 853-862
- He, S., Han, Y. Zhai, H. & Liu, Q. 2007 *In vitro* selection and identification of sweetpotato (*Ipomoea batatas* (L.) Lam.) Plants tolerant to NaCl. *Plant Cell Tissue Organ Cult.* 96:69-74

- Hossain, Z., Azad, AK., Datta, M SK. & Biswas, AK. 2009 Development of NaCl-tolerant line in *Chrysanthemum morifolium* Ramat. Through shoot organogenesis of selected callus line. *Biotechnology*. 129(4): 658-667
- Jain, RK., Dhawan, RS., Sharma, DR. & Chowdhury, JB. 1987 salt tolerance and proline accumulation: a comparative study in salt tolerant and wild type cultured cells of egg plant. *Plant Cell Reports* 6: 382-384
- Jain, M., Mathur, G., Koul, S. & Sarin, NB. 2001 Ameliorative effects of proline on salt stress-induced lipid peroxidation in cell lines of groundnut (*Arachis hypogaea* L.) *Plant Cell Rep* 20: 463-468
- Jaiswal R Singh NP 2001 Plant Regeneration from NaCl Tolerant Callus/Cell Lines of Chickpea. *ICPN* 8: 21-23
- Karadimova M Djambova G 1993 Increased NaCl - tolerance in wheat (*Triticum aestivum* L. and *T. durum* desf.) through *in vitro* selection. *In vitro Cell Dev Biol* 29: 180-182.
- Kavi Kishor, PB & Reddy, GM. 1985. Resistance of rice Callus Tissues to Sodium Chloride and Polyethylene Glycol. 59:1129-1131
- Kavi Kishor, PB. ; Sangam. S.; Amrutha, RN.; Sri Laxmi, P. ; Naidu, KR.; Rao, KRSS., Rao, S. Reddy, KJ., Theriappan & Sreenivasulu, N. 2005 Regulation of proline biosynthesis, degradation, uptake and transport in higher plants: Its implications in plant growth and abiotic stress tolerance. 88(3): 424-43.
- Kaviraj, CP. & Rao, S. 2009. High frequency of plant regeneration from cotyledonary node explants of green gram (*Vigna radiata* L.)-A recalcitrant grain legume. *The Bioscan* 4(2): 267-271.
- Kumar V Sharma DR Isolation and characterization of sodium chloride resistant callus cultures of *Vigna radiata* (L.) Wilczek. var. radiata. *J Expt Botany* 40: 143-147
- Kumar KB Khan PA Peroxidase and Polyphenol oxidase in excised ragi (*Elaeusine coracana* cv. 2020 leaves during senescence *Indian J Expt Botany*. 20: 412-416 1982
- Liu, T & Vanstadn, J. 2000 Selection and characterization of sodium chloride-tolerant callus of *Glycine max* (L.) Merr cv. Acme. *Plant Growth Regulation* 31: 195-207
- Lutts, S., Majerus, V. & Kinet, JM. 1999. NaCl effects on proline metabolism in rice (*Oryza sativa*) seedlings. *Physiol. Plant*. 105: 450-458.
- Miki Y Hashiba M Hisajima S 2001 Establishment of salt stress tolerant rice plants through step-up NaCl treatment *in vitro*. *Biol. Plant*. 44: 391-395
- Mathews, H., 1987 Morphogenetic responses from *In Vitro* cultured seedling explants of mungbean (*Vigna radiata* L. Wilczek) plant cell Tissue & organ Culture. 11, 233-240.
- Murashige, T. & Skoog, F. 1962 A revised medium for rapid growth and bio assays with tobacco tissue cultures. *Plant Physiol*. 15: 473-497,
- Nabors, MW., Gibbs, SE., Bernstein, CS. & Meis ME. 1980. NaCl tolerant plants from cultured cells. *Z. Pflanzenphysiol*. 97: 13-17.
- Naik, GR. & Harinath Babu, K. 1987 *In vitro* response of sugarcane to salinity tolerance. 37th convention D.S.T.A. 187-194.
- Nakona, Y., Asada, K.: Hydrogen peroxide is scavenged by ascorbic specific peroxidase in spinach chloroplasts. *Plant Cell Physiol*. 22: 867-880, 1981.
- Ochatt, SJ ; Marconi, P.L., Radice, S., Arnozis, P.A., Caso, O.H.: *In vitro* recurrent selection of potato: production and characterization of salt tolerant cell lines and plants. *Plant Cell, Tissue and Organ Culture*. 55: 1-8, 1999.

- Pandey, R., Ganapathy, P.S.: Effect of Sodium Chloride stress on callus cultures of *Cicer arietinum* L. Cv. BG. 203: Growth and Ion accumulation. *Journal. Exp. Botany*. 157 (35): 1194-1197, 1984
- Patil, P. & Rao S. 1999 Selection and characterization of NaCl tolerant callus cultures of *Vigna radiata* (L.) Wilczek. *Plant Tissue Culture and Biotechnology Emerging Trends*.
- Petrusa, LM. & Winicov, I. 1997 Proline status in salt tolerant and salt sensitive alfalfa cell line and adapted plants in response to NaCl. *Plant Physiol Biochem* 35: 303-310.
- Pius, J., Eapen, S., George, L. & Rao, P.S. Isolation of sodium chloride tolerant cell lines and plants in finger millet. *Biologia Plantarum*. 35(2): 267-271 1993.
- Queiros, F., Fidalgo, F., Santos, I., Salema, R. *In vitro* selection of salt tolerant cell lines in *Solanum tuberosum* L. *Biologia Plantarum*. 51(4): 728-734, 2007.
- Rao Srinath, Patil, P. 1998 Effect of sodium chloride on callus culture of *Cicer arietinum* L. *Bulletin of pure and Applied Sciences*. 17B (2): 81-87.
- Rao, Srinath, Patil, P. & Kaviraj, CP. 2005 Callus induction and organogenesis from various explants in *Vigna radiata* (L.) Wilczek. *Indian Journal of Biotechnology*. 4: 556-560.
- Rout, G.R., Senapatil, S.K., Panda, J.J.: Selection of Salt tolerant Plants of *Nicotiana tabacum* L. Through *In vitro* and its biochemical characterization. *Acta Biologica Hungarica*. 59(1): 77-92, 2008.
- Rhodes, D., Verslues, P.E., Sharp, R.E.: Role of amino acids in abiotic stress resistance. In: Singh, B.K. (Ed.) *Plant amino acids: Biochemistry and Biotechnology*. Arcel Dekker, N.Y. pp 319-356.
- Rus, A.M., Panoff, M., Perez-Alfocea, F., Bolarin, M.C.: NaCl response in tomato calli and whole plants. *J. Plant Physiol*. 155: 727-733, 1999.
- Sairam, R.K., Srivastava, G.C., Agarawal, S., Meena, R.C. Differences in antioxidant activity in response to salinity stress in tolerant and susceptible wheat genotypes. *Biologia Plantarum* 49 (1): 85-91, 2005.
- Shigeoka, S., Ishikawa, T., Tamoi, M., Miagawa, Y., Takeda, T., Yabuta, Y., Yoshimora, K.: Regulation and function of ascorbate peroxidase isoenzymes. *J. Exp. Bot.* 53: 1305-1319, 2000.
- Shashi Madan, Nainawatee, HS., Jain, RK. & Chowdhury, JB. 1995. Proline and Proline metabolizing enzymes *in-vitro* Selected NaCl-tolerant *Brassica juncea* L. under Salt Stress. *Annals of Botany*. 76:51-57
- Singh, M., Jaiswal, U., Jaiswal, V.S.: *In Vitro* selection of NaCl-Tolerant callus lines and regeneration of plantlets in a bamboo (*Dendrocalamus strictus* Ness.) *Biol-Plant*. 39: 229-233, 2003.
- Subhashini, K. & Reddy, GM. *In vitro* selection for salinity and regeneration of plants in rice. *Current Science*. 58: 584-586.
- Sumithra, K., Jutur, K., Dalton-Caramel, B. & Reddy, A.R. 2006. Salinity induced changes in two cultivars of *Vigna radiata*: response of antioxidative and proline metabolism. *Plant Growth Regul.* 50: 11-22.
- Tal, M., Katz, A., Heikin, M. & Dechan, K. 1979 Salt tolerance in the wild relatives of the cultivated tomato: Proline accumulation in *Lycopersicon esculentum* Mill. *L.peruvianum* Mill., and *Solanum pennelli* Cor. Treated with NaCl and polyethylene glycol. *New Phytol.* 82: 349-355
- Tal, M. 1994. *In vitro* selection for salt tolerance in crop plants: Theoretical and practical considerations. *In vitro Cell Dev. Biol.* 30: 175-180.

- Thomas, JC., Armond, RL. and Bonnert, HJ. 1992. Influence of NaCl on growth, proline and phosphoenol pyruvate carboxylase levels in *Mesembryanthemum crystallinum* suspension cultures. *Plant Physiol* 98: 626-631.
- Tyagi, RK. & Rangaswamy, NS. 1997. *In vitro* induction and selection for salt tolerance in oilseed brassicas and regeneration of the salt tolerant soma clones. *Phytomorphology*. 47(2): 209-220
- Venkataiah, P., Christopher, T. & Subhash K. 2004. Selection and characterization of sodium chloride and mannitol tolerant callus lines of red pepper (*Capsicum annuum* L.) *Plant Physiol*. 9(2): 158-163
- Watad, AA., Reinhold, L., Lerner, HR. 1983. Comparison between a Stable NaCl-Selected *Nicotiana* Cell Line and the Wild Type. *Plant Physiol* 73: 624-629
- Winicov, I. 1991. Characterization of salt tolerant alfalfa (*Medicago sativa* L.) plants regenerated from salt tolerant cell lines. *Plant Cell Reports* 10: 561-564
- Winicov, I. & Bastola, DR. 1997. Salt tolerance in crop plants: new approaches through tissue culture and gene regulation. *Acta Physiol Plant* 19(4): 435-449
- Yasar, Ellialyoglu, S. and Kusvuran, S. 2006. Ion and lipid peroxide content in sensitive and tolerant eggplant callus cultured under salt stress. *Europ J Hort Sci* 71(4): 169-172
- Zair, I., Chlyah, A., Sabounji, K., Tittahsen, M. & Chlyah, H. 2003. Salt tolerance improvement in some wheat cultivars after application of *in vitro* selection pressure. *Plant Cell Tissue & Organ Culture* 73: 237-244
- Zhao, Y., Aspinall, D. & Palge, LG. 1992. Protection of membrane integrity in *Medicago sativa* (L.) by glycinebetaine against the effects of freezing. *J Plant Physiol* 140: 541-543

IntechOpen

Biotechnology - Molecular Studies and Novel Applications for Improved Quality of Human Life

Edited by Prof. Reda Sammour

ISBN 978-953-51-0151-2

Hard cover, 250 pages

Publisher InTech

Published online 14, March, 2012

Published in print edition March, 2012

This book deals with the importance of application of molecular biology as an approach of biotechnology for improvement of the quality of human life. One of the interesting topics in this field, is the identification of the organisms that produce bioactive secondary metabolites. It also discusses how to structure a plan for use and preservation of those species that represent a potential source for new drug development, especially those obtained from bacteria. The book also introduces some novel applications of biotechnology, such as therapeutic applications of electroporation, improving quality and microbial safety of fresh-cut vegetables, producing synthetic PEG hydro gels to be used as an extra cellular matrix mimics for tissue engineering applications, and other interesting applications.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Srinath Rao and Prabhavathi Patil (2012). In Vitro Selection of Salt Tolerant Calli Lines and Regeneration of Salt Tolerant Plantlets in Mung Bean (*Vigna radiata* L. Wilczek), *Biotechnology - Molecular Studies and Novel Applications for Improved Quality of Human Life*, Prof. Reda Sammour (Ed.), ISBN: 978-953-51-0151-2, InTech, Available from: <http://www.intechopen.com/books/biotechnology-molecular-studies-and-novel-applications-for-improved-quality-of-human-life/in-vitro-selection-of-salt-tolerant-calli-lines-and-regeneration-of-salt-tolerant-plantlets-in-mung-bean>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](https://creativecommons.org/licenses/by/3.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen