We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Influence of Acidification on Dough Rheological Properties

Daliborka Koceva Komlenić, Vedran Slaćanac and Marko Jukić
*Faculty of Food Technology, Josip Juraj Strossmayer University of Osijek,
 Croatia*

1. Introduction

Due to increasing consumer demands for more natural, tasty and healthy food, the traditional process of sourdough bread production has been enjoyed renewed success in recent years, (Brümmer & Lorenz, 1991; Thiele *et al.*, 2002; Lopez *et al.*, 2003). Sourdough was traditionally used as leavening agents until it was replaced by baker's yeast in the 19th century (Corsetti & Setanni, 2007). Today, sourdough is employed in the manufacture of a number of products, such as breads, cakes and crackers (De Vuyst & Gänzle, 2005). Some of these products have strictly regional and artisanal character; well some of these are widely distributed on the world market (De Vuyst & Neysens, 2005). Many wheat breads and cakes are characteristic for the Mediterranean countries, the San Francisco bay and Southern America, whereas a number of bakery preparations made with rye, wheat, barley, or mixtures of these flours are typical for Central Europe, Eastern Europe and Scandinavia (Stephan & Neumann, 1999).

In general, sourdough is mixture of flour and water that is than fermented with lactic acid bacteria (LAB) (Hammes & Gänzle, 1998). Ordinarily, these LAB are heterofermentative strains which produce lactic and acetic acid in the mixture, and resulting in a sour taste of the dough (Vogel *et. al.*, 1999). The acidification process affected by the application of sourdoughs is mainly used to improve quality, taste and flavour of wheat breads (Brümmer & Lorenz, 1991; Katina *et al.*, 2006a; Arendt *et al.*, 2007), and the slow staling (Katina *et al.*, 2006b; Plessas *et al.*, 2007).

Sourdough can be freshly produced in bakeries, or can be obtained from the commercial suppliers in the some of the following applying forms: living, liquid sourdough or dried, non-fermenting sourdough (Böcker *et. al.*, 1990). Considerably, doughs could be acidified biologically, chemically and naturally preferment.

The addition of sourdough during production of wheat bread causes major changes in the dough characteristics (Clarke *et al.*, 2002; Clarke *et al.*, 2004; Ketabi *et al.*, 2008), especially in the flavour and structure (Clarke *et. al.*, 2002). The effect of the fermentation process of wheat doughs containing lactic acid bacteria are complex and depending on variations between sourdoughs regard to the type of starter culture, dough yield and fermentation regime used (Wehrle *et. al.*, 1997). This propounds requirements for the detail microbiological investigations on sourdough microflora. Study of sourdough from

microbiological point of view barely started a hundred years ago (Salovaara, 1998). Today, microbial population of different types of sourdough is rather known. Many inherent properties of sourdough rely on the metabolic activities of its resident LAB: lactic fermentation, acetic fermentation, proteolysis, synthesis of volatile compounds, anti-mould activity and antiropiness are among the most important activities during sourdough fermentation (Hammess and Gänzle, 1998). The most of the effects of sourdough have been considered by pH value decrease which has been caused with organic acid production.

Between many of important effects, the drop in pH value caused by the organic acids produced influences the viscoelastic behaviour of doughs, respectively to sourdoughs rheological properties (Wehrle & Arendt, 1998). A correct description of the changes in dough behaviour is necessary to maintain handling and machinability in industrial production (Wehrle et. al., 1997). Furthermore, following the identification and classification of LAB from cereal fermentation, basic and applied today face the challenge of identifying functional characters of these bacteria to completely exploit their microbial metabolic potential from the production of baked goods (Vogel et. al., 2002). In European countries, production of sourdoughs from wheat flour has been predominant. Most fundamental studies on wheat sourdough have been conducted in Western Europe, and the results have been published in German-language journals and books (Brümmer & Lorenz, 1991).

Rheological properties, acidification and flavour development are the most important parameters in fermentation process control (Hames et. al., 2005). Rheological properties of dough have been determined by a number of methods, such as dynamic rheological measurements, extensigraphs, alveographs, lubricated uniaxial compression, oscillatory probe rheometers etc. (Hoseney, 1994). The rheological characteristics of dough have been considerably changed with fermentation. Types of microorganisms, metabolic activity and time-dependent development pH value effect on rheological properties (Wehrle & Arendt, 1998). Acids strongly influence on the mixing properties of dough. Doughs with lower pH requires a slightly shorter mixing time and have less stability than dough with normal pH level (Hoseney, 1994). Changes in pH values caused by production of lactic acid also influence on the rheological properties of dough (Wehrle *et al.*, 1997). Dough with containing acid has been characterized by increased phase angle and reduced complex modulus indicative of overmixing (Wehrle et al 1997). Small physical and chemical changes in the gluten network can result in significant changes in rheological properties. Clarke et al. (2002) were concluded that addition of sourdough prepared either from a single strain starter culture or a mixed strain starter culture had significant impact on the rheological properties of wheat flour dough.

Koceva Komlenić et al (2010) investigated the influence of chemical and biological acidification on dough rheological properties. According to their experimental results, rheological properties strongly depend on acidification type. Dough with lower pH value showed less stability during mixing, decreased extensibility and gelatinization maximum. In general, the rheological properties of dough greatly improved when the sourdough was added.

Regard to all facts mentioned above, fermentation of dough with LAB greatly effects on the properties of many bakery products. Structure properties are one of the most important. From that reason, rheological measurements play important role in definition of quality of products from sourdough. In this paper, influence of sourdough fermentation on the structure of sourdough was observed.

2. Microbiology of sourdough

It is well known that the type of bacterial flora developed in each fermented food depends on water activity, pH (acidity), minerals concentration, gas concentration, incubation temperature and composition of food matrix (Font de Valdez et. al. 2010). The microflora of raw cereals is composed of bacteria, yeast and fungi ($10^4 - 10^7$ CFU/g), while flour usually contains $2 \times 10^4 - 6 \times 10^6$ CFU/g (Stolz, 1999). In sourdough fermentation major role play heterofermentative species of LAB (Salovaara, 1998; Corsetti & Settani, 2007), especially when sourdoughs are prepared in a traditional manner (Corsetti et. al., 2003).

Type Ia	Type Ib	Type Ic	Type II	Type III
Obligate heterofermentative	Obligate heterofermentative	Obligate heterofermentative	Obligate heterofermentative	Obligate heterofermentative
<i>Lb. sanfranciscensis</i>	<i>Lb. brevis</i> <i>Lb. brevis</i> <i>Lb. buchneri</i> <i>Lb. fermentum</i> <i>Lb. fructivorans</i> <i>Lb. pontis</i> <i>Lb. reuteri</i> <i>Lb. sanfranciscensis</i> <i>W. cibara</i>	<i>Lb. pontis</i> <i>Lb. fermentum</i> <i>Lb. reuteri</i>	<i>Lb. brevis</i> <i>Lb. fermentum</i> <i>Lb. frumenti</i> <i>Lb. pontis</i> <i>Lb. panis</i> <i>Lb. reuteri</i> <i>Lb. sanfranciscensis</i> <i>W. confusa</i>	<i>Lb. brevis</i>
	Facultative heterofermentative <i>Lb. alimentarius</i> <i>Lb. casei</i> <i>Lb. paralimentarius</i> <i>Lb. plantarum</i>			Facultative heterofermentative <i>Lb. plantarum</i> <i>P. pentosaceus</i>
	Obligate homofermentative <i>Lb. acidophilus</i> <i>Lb. delbrueckii</i> <i>Lb. farciminis</i> <i>Lb. mindensis</i>	Obligate homofermentative <i>Lb. amylovorus</i>	Obligate homofermentative <i>Lb. acidophilus</i> <i>Lb. delbrueckii</i> <i>Lb. amylovorus</i> (rye) <i>Lb. farciminis</i> <i>Lb. johnsonii</i>	

Table 1. Classification of sourdoughs and the corresponding chracteristic microflora (De Vuyst & Neysens, 2005)

Sourdough LAB generally belongs to the genus *Lactobacillus*, *Leuconostoc*, *Pediococcus* or *Weisella* (Ehrmann & Vogel, 2005; De Vuyst & Neysens, 2005). A few less than 50 different species of LAB from sourdough have been reported by De Vuyst & Neysens (2005). Although species belonging *Leuconostoc*, *Pediococcus*, *Weisella*, *Enterococcus* and *Streptococcus* genera have been isolated from sourdough, *Lactobacillus* strains are the most frequently observed bacteria in sourdoughs (Corsetti & Settanni, 2007). Characteristic microflora of sourdough is presented in Table 1. Undesirable bacteria such as *Staphylococcus aureus* and *Bacillus cereus*, as well as other bacteria, may be present in minor concentration (De Vuyst and Neysens, 2005). Corsetti & Settani (2007) reported that the recently described species *Lactobacillus frumenti*, *Lactobacillus mindensis*, *Lactobacillus paralimentarius*, *Lactobacillus*

spicheri, *Lactobacillus rossiae*, *Lactobacillus acidifarinae*, *Lactobacilus zymae*, *Lactobacillus hammesii*, *Lactobacillus nantensis* and *Lactobacillus siliginis* were first isolated from the sourdough matrices. 15 lactobacilli species known to occur in sourdough are also known to live in human and animal intestines. *Lactobacillus reuteri*, *Lactobacillus acidophilus* and *Lactobacillus plantarum* are some of these intestinal lactobacilli (Hammes & Gänzle, 1998).

Kumar et. al. (2004) reported a phylogram based on 16S rRNA gene sequences of sourdough lactobacilli. A total of 41 species are included in phylogram tree, some of them, such as *Lactobacillus rhamnosus*, have been rarely or only once reported to be found in sourdough (Spicher & Löner, 1985). Hammes et. al. (2005) reported that about 30 species are considered to be typical of sourdough environments. *Lactobacillus sanfranciscensis*, *Lactobacillus brevis* and *Lactobacillus plantarum* are the most frequently lactobacilli isolated from sourdough (Gobbetti, 1998; Corsetti et. al. 2001; Valmorri et. al., 2006; Corsetti & Settanni, 2007). *Lb. sanfranciscensis* was first reported in the San Francisco sourdough French bread process, to be responsible for acid production (Corsetti et. al., 2001). *Lb. sanfranciscensis* has been predominant bacteria in traditional production by various stages of continuous production and production by commercial starter cultures (Gobbetti & Corsetti, 1997). Gobbetti et. al. (1998) reported on the *Lb. sanfranciscensis* – *Lb. plantarum* association in Italian wheat sourdough. *Lb. alimentarius*, which probably belong to *Lb. paralimentarius*, were first isolated from Japanese sourdough (Cai et. al., 1999). *Lb. brevis* and *Lb. plantarum* have been associated with *Lb. fermentum* in Russian sourdoughs (Kazayanska et. al., 1983). *Lb. fermentum* dominates Swedish sourdoughs and German type II sourdoughs (Meroth et al., 2004.) Gobbetti et. al. (1994b) reported that *Lb. acidophilus* is found in Umbrian (Italian region), while Corsetti et. al. (2003) described a new sourdough associated species, *Lb. rossae*, in sourdoughs of central and southern Italy (Settanni et. al., 2005a). African sorghum sourdoughs, which are produced at higher temperature (> 35 °C) contain obligate heterofermentative *Lb. fermentum*, *Lb. pontis* and *Lb. reuteri* species, as well as the obligate homofermentative *Lb. amylovorus* (De Vuyst & Neysens, 2005).

According to all facts mentioned above, it is so clear that large biodiversity in lactoballi composition of the sourdough exists, regarding to type of sourdough (Table 2), as well as to some cultural, geographical and traditional reasons.

Country	Product/method of isolation and identification	Lactic acid bacteria
Belgium	Wheat/rye sourdoughs polyphasic approach	<i>Lb. brevis</i> , <i>Lb. plantarum</i> , <i>Lb. sanfranciscensis</i> , <i>Lb. paralimentarius</i> , <i>P. pentosaceus</i> , <i>Lb. helveticus</i>
Finland	Rye sourdough phenotypical	<i>Lb. acidophilus</i> , <i>Lb. plantarum</i> , <i>Lb. casei</i>
Denmark	Sour rye dough phenotypical	<i>Lb. reuteri</i> , <i>Lb. panis</i> , <i>Lb. amylovorus</i>
France	Wheat bread phenotypical	<i>Lb. plantarum</i> , <i>Lb. casei</i> , <i>Lb. delbrueckii</i> subsp. <i>delbrueckii</i> , <i>Lb. acidophilus</i> , <i>Lb. brevis</i> , <i>Leuc. mesenteroides</i> subsp. <i>mesenteroides</i> , <i>Leuc. mesenteroides</i> subsp. <i>dextranicum</i> , <i>P. pentosaceus</i> , <i>Lb. Curvatus</i>
Germany	Wheat sourdough phenotypical	<i>Lb. delbrueckii</i> , <i>Lb. planlarum</i> , <i>Lb. casei</i> , <i>Lb. fermentum</i> , <i>Lb. buchneri</i> , <i>Lb. brevis</i>
	Rye bread phenotypical	<i>Lb. acidophilus</i> , <i>Lb. farciminis</i> , <i>Lb. alimentarius</i> , <i>Lb. casei</i> , <i>Lb. plantarum</i> , <i>Lb. brevis</i> , <i>Lb. sanfranciscensis</i> , <i>Lb. fructivorans</i> , <i>Lb. fermentum</i> , <i>Lb. buchneri</i>

Country	Product/method of isolation and identification	Lactic acid bacteria
	Rye sourdough phenotypical	<i>Lb. acidophilus</i> , <i>Lb. casei</i> , <i>Lb. plantarum</i> , <i>Lb. farciminis</i> , <i>Lb. alimentarius</i> , <i>Lb. brevis</i> , <i>Lb. buchneri</i> , <i>Lb. fermentum</i> , <i>Lb. fructivorans</i> , <i>Lb. sanfranciscensis</i> , <i>Pediococcus</i> spp.
	Wheat sourdoughs (Panettone, wheat bread) phenotypical	<i>Lb. plantarum</i> , <i>Lb. casei</i> , <i>Lb. farciminis</i> , <i>Lb. homohiochii</i> , <i>Lb. brevis</i> , <i>Lb. hilgardii</i> (spontaneous); <i>Lb. sanfranciscensis</i> , <i>Lb. brevis</i> , <i>Lb. hilgardii</i> , <i>W. viridescens</i> (masa madre)
	Rye sourdough RAPD-PCR	<i>Lb. amylovorus</i> , <i>Lb. pontis</i> , <i>Lb. frumenti</i> , <i>Lb. reuteri</i>
	Rye bran PCR-DGGE	<i>Lb. sanfranciscensis</i> , <i>Lb. mindensis</i> (type I rye sourdough); <i>Lb. crispatus</i> , <i>Lb. pontis</i> , <i>Lb. panis</i> , <i>Lb. fermentum</i> , <i>Lb. frumenti</i> (type II rye sourdough); <i>Lb. johnsonii</i> , <i>Lb. reuteri</i> (type II rye bran sourdough)
Greece	Wheat sourdoughs polyphasic approach	<i>Lb. sanfranciscensis</i> , <i>Lb. brevis</i> , <i>Lactobacillus</i> spp. ^a , <i>Lb. paralimentarius</i> , <i>W. cibaria</i>
Italy	Panettone phenotypical	<i>Lb. brevis</i> , <i>Lb. plantarum</i>
	Panettone, Brioche phenotypical	<i>Lb. sanfranciscensis</i> , <i>Lb. fermentum</i> , <i>Lb. plantarum</i> , <i>Leuc. mesenteroides</i> , <i>Pediococcus</i> spp.
	Umbrian wheat sourdoughs phenotypical	<i>Lb. sanfranciscensis</i> , <i>Lb. plantarum</i> , <i>Lb. farciminis</i>
	Pizza (Naples) phenotypical	<i>Lb. sakei</i> , <i>Lb. plantarum</i> , <i>Leuc. gelidum</i> , <i>Leuc. mesenteroides</i>
	Verona sourdoughs RAPD-PCR	<i>Lb. sanfranciscensis</i>
	Lombardian mother sponges	<i>Lb. sanfranciscensis</i>
	speciesspecific PCR	
	Apulian wheat sourdoughs 16S rDNA sequencing 16S/23S rRNA spacer region PCR	<i>Lb. sanfranciscensis</i> , <i>Lb. alimentarius</i> , <i>Lb. brevis</i> , <i>Leuc. citreum</i> , <i>Lb. plantarum</i> , <i>Lb. lactis</i> subsp. <i>lactis</i> , <i>Lb. fennentum</i> , <i>Lb. acidophilus</i> , <i>W. confusa</i> , <i>Lb. delbmeckii</i> subsp. <i>bulgaricus</i>
Iran	Sangak phenotypical	<i>Leuc. mesenteroides</i> , <i>Lb. plantarum</i> , <i>Lb. brevis</i> . <i>P. cerevisiae</i>
Mexico	Pozol (maize) 16S rDNA sequencing	<i>Lb. lactis</i> , <i>S. suis</i> . <i>Lb. plantarum</i> , <i>Lb. casei</i> . <i>Lb. alimentarius</i> . <i>Lb. delbrueckii</i>
Morocco	Sourdough ferments traditional starter sponges phenotypical	<i>Lb. plantarum</i> , <i>Lb. brevis</i> , <i>Lb. buchneri</i> , <i>Lb. casei</i> . <i>Leuc. mesenteroides</i> , <i>Pediococcus</i> sp.
	Soft wheat flour phenotypical	<i>Lb. plantarum</i> , <i>Lb. delbrueckii</i> , <i>Lb. buchneri</i> , <i>Lb. casei</i> , <i>Lb. sanfranciscensis</i> , <i>Leuc. mesenteroides</i> , <i>P. pentosaceus</i>
Portugal	Broa phenotypical	<i>Leuconostoc</i> spp., <i>Lb. brevis</i> , <i>Lb. curvatus</i> , <i>Lb. delbrueckii</i> , <i>Lb. lactis</i> subsp. <i>lactis</i> , <i>E. casseliflavus</i> , <i>E. durans</i> , <i>E. faecium</i> , <i>S. constellatus</i> , <i>S. equinus</i>
Russia	Rye sourdough phenotypical	<i>Lb. plantarum</i> , <i>Lb. brevis</i> , <i>Lb. fermentum</i>
Spain	Wheat sourdough phenotypical	<i>Lb. brevis</i> , <i>Lb. plantarum</i>
	Wheat sourdough phenotypical	<i>Lb. brevis</i> , <i>Lb. plantarum</i> , <i>Lb. cellobiosus</i> . <i>Leuc. mesenteroides</i>
Sudan	Kisra (sorghum sourdough)	<i>Lb. fermentum</i> , <i>Lb. reuteri</i> , <i>Lb. amylovorus</i>
	Kisra RAPD	<i>E. faecalis</i> , <i>Lb. laclis</i> , <i>Lb. fermentum</i> , <i>Lb. reuteri</i> , <i>Lb. vaginalis</i> , <i>Lb. helveticus</i>
Sweden	Rye/wheat phenotypical	<i>Lb. fermentum</i> . <i>Lb. delbrueckii</i> , <i>Lb. acidophillus</i> , <i>Lb. plantarum</i> , <i>Lb. rhamnosus</i> , <i>Lb. farciminis</i> . <i>Lb. fermentum</i> , <i>Lb. sanfranciscensis</i> , <i>Lb. brevis</i> . <i>W. viridescens</i>

Country	Product/method of isolation and identification	Lactic acid bacteria
USA	Rye sourdough phenotypical	<i>Lactobacillus</i> sp., <i>P. pentosaceus</i>
	San Francisco sourdough French bread phenotypical	<i>Lb. sanfranciscensis</i>

Table 2. Biodiversity of sourdough lactic acid bacteria in sourdough of different origin (De Vuyst & Neysens, 2005)

The following yeasts have been detected in cereals (9×10^4 CFU/g) and flour (2×10^3 CFU/g): *Candida*, *Cryptococcus*, *Pichia*, *Rodothorula*, *Torulaspora*, *Trychoporon*, *Saccharomyces* and *Sporobolomyces*. *Saccharomyces cerevisiae* is not found in the raw materials. Its occurrence in sourdough has been explained by the application of baker’s yeast in most daily bakery practice (Corsetti et. al., 2001). Among fungi (ca. 3×10^4), *Alternaria*, *Cladosporium*, *Drechslera*, *Fusarium*; *Helminthosporium*, *Ulocladium*; *Aspergillus* and *Penicillium* were found in raw cereals and flour. Possibility to dough be contaminated with these species exist, but in the most of cases it could be avoided (De Vuyst & Neysens, 2005).

3. Role of LAB in sourdough fermentation

The metabolic activities of lactobacilli during sourdough fermentation improve dough properties, bread texture and flavour; retard the stalling process of bread; and prevent bread from mould and bacterial spoilage (Gerez et. al., 2009; de Valdez et. al., 2010). Additionally, LAB could contribute to nutritive value and healthiness of bread (Gobbetti et. a., 2007). The sourdough microflora is usually composed of stable associations of yeasts and lactobacilli.

3.1 Carbohydrate metabolism

In dough fermentation carbonhydrate metabolism varies depending on involved LAB strains, type of sugars, presence of yeasts and processing conditions (Gobetti et. al., 1994a). The importance of antagonistic and synergistic interactions between lactobacilli and yeasts are based on the metabolism of carbon hydrates and amino acids and the production of carbon dioxide (Gobetti & Corsetti 1997). Lactic and acetic acid are predominant products of sourdough fermentation (Figure 1).

Diversity in metabolic pathways between different sourdoughs is large. Recipe and geographical origin make the difference in the aroma of breads consumed in the different areas (Hansen & Schieberle, 2005). All these parameters influence on characteristics and give originality to different breads from sourdoughs. It is the basic reason why bread from biologically fermented dough posses a superior sensory quality than bread produced from chemically acidified dough (Kirchhoff & Schieberle, 2002). The ratio between lactic and acetic acid is an important factor that might affect the aroma profile and structure of final product. Acetic acid, produced by heterofermentative LAB, is responsible for a shorter and harder gluten, while lactic acid can gradually account for a more elastic gluten structure (Lorenz, 1983; Corsetti & Settani, 2007). *Lb. sanfranciencis*, isolated from traditional Italian sourdoughs ferment only glucose and maltose. However, *Lb. sanfranciencis* strains isolated from some other sourdoughs use other sugars, such as sucrose, raffinose, galactose, melobiose, ribose and fructose (Tieking et al., 2003). Ginés et. al. (1997) presented metabolic potential of *Lb. reuteri* CRL1100, a strain isolated from homemade sourdough. This

microorganism metabolizes glucose and galactose, but not fructose and cellobiose. The most of lactobacilli strain in sourdough are unable to ferment sucrose, a disaccharide whose metabolism by heterofermentative lactobacilli in sourdough appears to be attributable to glycosyl-transferases rather than to invertase activities (Font de Valdez et. al., 2010). Recently, it has been reported that certain lactic acid bacteria are able to produce exopolysaccharides (EPS), which might have a positive affect on bread volume and shelf life (Font de Valdez et. al., 2010). Following EPS have been produced during fermentation of sourdough: fructans (levan, inulin) and glucans (reuteran, dextran, xanthan). EPS influence significantly to the texture improvement of sourdough and bread (Tieking et. al., 2003). The production of EPS in sufficient amounts during sourdough fermentation would create the possibility to replace hydrocolloids in baking (Tieking et. al., 2003). Hydrocolloids have been reported to improve bread quality through stabilization of water-flour association in sourdoughs matrix (Tieking et. al., 2003; Font de Valdez et. al., 2010).

Fig. 1. Fate of potential electron acceptors upon carbohydrates fermentation by *Lb. sanfranciscensis* (Corsetti et al, 2007)

3.2 Metabolism of proteins

During sourdough fermentations protein modification and protein degradation have been occurred. Protein degradation that occurs during sourdough fermentations is among the key metabolic route that directly effects on dough texture and flavour of sourdough. During dough fermentation, LAB releases small peptides and free amino acids important for microbial growth and acidification of dough (Rollán & Font de Valdez, 2001). Also, small peptides and free amino acids are important as precursors for flavour development of the leavened baked products (Thiele et. al., 2002). According to the results of studies performed by Gerez et. al. (2006) 13 nine lactobacilli and four pediococci were able to use gluten as a nitrogen source. Gerez et. al. (2006) also reported an increase in essential amino acids (treonine, valine, lysine and phenylalanine) in a gluten based medium fermented by LAB strains. Thiele et al. (2004) reported that the degree of protein degradation in wheat

sourdough is usually similar to that observed in chemically acidified dough. However, bacterial proteolysis during sourdough fermentation was shown to contribute much more to the development of typical sourdough flavours of baked breads compared to breads produced from acidified or yeasted dough (Hansen et. al., 1989a, Hansen et. al., 1989b). Gluten proteins determine, to a great extent, the rheological properties of wheat dough. Substantial hydrolysis of gliadin and glutenin proteins occurs during sourdough fermentation. Proteolytic activity in sourdough originates not only from LAB enzymes, but also derives also from the cereal materials present in sourdough (Thiele, 2002; Thiele, 2004). Except activity of own enzymes, LAB contribute to overall proteolysis during sourdough fermentation by creating optimum (acidic) conditions for activity of cereal proteinases (Vermeulen et al. 2006). The partial hydrolysis of glutenins during sourdough fermentation results in depolymerisation and solubilisation of the gluten macro peptide (GMP). After 24 hours of fermentation with defined lactobacilli strains, all gluten proteins were SDS-soluble (Thiele et. al., 2003). Glutathione (GSH) is the most relevant reducing agent in wheat doughs (Grosh & Wieser, 1999). Heterofermentative lactobacilli express glutathione reductase during growth in dough and reduce extracellular oxidized glutathione (GSSG) (Jänsch et. al., 2007). The continuous transformation of GSSG to GSH by LAB metabolism maintains high SH levels in wheat doughs, and increase the amount of SH-groups in gluten proteins (Vermeulen et. al., 2006)

Increased proteolysis during sourdough fermentation leads to the liberation of amino acids in wheat dough. Importance of amino acid conversion to typical flavour volatile compounds has been reported by a number of researchers (Juillard et. al., 1998; Tamman et. al., 2000; Zotta et al. 2006). Sourdough fermentation with LAB results in an increase of amino acids concentration during fermentation, whereas dough fermentation with yeast reduces the concentration of free amino acids (Thiele et. al., 2002). Furthermore Gassenmayer & Schieberle (1995) reported that addition of amino acids (e.g. ornithine, leucine and phenylalanine) to dough resulted in an enhanced conversion to flavour compounds. The level of individual amino acids in wheat dough depends on the pH level of dough, fermentation time and the consumption of amino acids by the fermentative microflora (Thiele et. al., 2002). In wheat sourdoughs, *Lb. brevis linderi*, *Lb. safransciensis*, *Lb. brevis* and *Lb. plantarum* have been reported to increase the levels of aliphatic, dicarboxylic and hydroxyl amino acids (Gobbetti et. al., 1994a, Gobbetti et. al., 1994b). The yeasts, *S. cerevisiae* and *S. exiguus* decrease the total level of amino acids.

3.3 Development of aroma compounds

The key degradation reaction of amino acids during fermentation of sourdough is the Erlich pathway, leading to aldehydes or the corresponding alcohol, while during baking takes place the Strecker reaction which also lead to aldehydes, but also to the corresponding acids (Hoffmann & Schieberle, 2000). Differences other than acetic acid production, in the overall aroma profile of final bread depend to the type of dominating lactobacilli, as well as to the type of flour. E. g., firm wheat fermented with heterofermentative strains had higher content of ethyl acetate and hexyl acetate compared to cases when the homofermentative strains were used (Lund et. al., 1989). On the other hand, higher content of aldehydes was higher in rye sourdough fermented with homofermentative LAB (Lund et. al., 1989).

The arginine metabolism by *Lb. pontis* and *Lb. sanfransciensis* has been demonstrated to have an impact on bread flavour (Thiele et. al., 2002; De Angelis et. al., 2002). Phenylalanine metabolism was studied in *Lb. plantarum* and *Lb. sanfransciensis*, besides producing phenyllactic acid and 4-hydroxy compounds, which have documented antifungal activity (Vermeulen et. al., 2006). Gerez et. al. (2006) demonstrated that gluten-breakdown with lactobacilli and pediococci, beside reducing gluten-allergen compounds. Furthermore increased the basic amino acid concentration in broth cultures, mainly due to an increase in the ornithine amount, which is considered to be key flavour precursor in wheat bread, generating 2-acetyl-pyrolin (Gassenmeyer & Schieberle, 1995; Thiele et. a., 2002). Czerny & Schieberle (2002) reported that during fermentation of dough, LAB did not generate new aroma compounds than those present in raw materials. However, they demonstrated that many compounds as acetic acid and 3-methylbutanal were increased, whereas aldehydes were decreased.

From all referred above, LAB play very important role in overall sourdough fermentation process. Together with yeasts is responsible for unique quality of the baking bread. More research, however, is needed to clarify all metabolic pathways during fermentation of sourdough.

4. Classification of sourdoughs

Sourdoughs have been classified on the base of procedures during their production (Böcker et. al., 1995). Sourdoughs have been classified into three types:

1. type I sourdoughs or traditional sourdoughs
2. type II sourdoughs or accelerated sourdoughs
3. type III sourdoughs or dried sourdoughs

Each type of sourdough is characterised by a specific sourdough LAB microflora (Table 1).

4.1 Type I sourdough

Type I sourdoughs are produced with traditional techniques, and are characterized by continuous (daily) refreshments to keep the microorganisms in an active state. Type I sourdough is indicated by high metabolic activity, above all regard to leavening, i. e. gas production. The process is conducted at room temperature (20 – 30 °C) and the pH is approximately 4.0. Examples of baked breads with type I sourdough are Francisco sourdough French bread, Panettone and other brioches, Toscanon and Altamura bread, Pugliese, and three-stage sourdough rye bread. Traditional, type I sourdough encompass pure culture, pasty sourdough starter preparations from different origin (type Ia), spontaneously developed, mixed culture sourdoughs made from wheat and rye or mixture thereof and prepared through multiple stage fermentation processes (type Ib), and sourdough made in tropical regions fermented at high temperatures (Stolz, 1999). Pure culture sourdoughs (type Ia) are derived from natural sourdough fermentations.

4.2 Type II sourdough

Type II of sourdough has been prepared in semi-fluid silo condition. Those bakery pre-products serve mainly as dough acidifiers. Several modified, accelerated processes with

continuous propagation and long-term one-step fermentations are common now. They guarantee more production reliability and flexibility. A recent trend of industrial bakeries exists in the instalment of continuous sourdough fermentation plants (Stolz & Böcker, 1996).

Typical type II process last for 2-5 days and are often carried out at increased fermentation temperature (usually $> 30^{\circ}\text{C}$) to speed up the process. Those sourdoughs exhibit a high acid content at a pH of < 3.5 after 24 hours of fermentation. The microorganisms are commonly in the late stationary phase and therefore exhibit restricted metabolic activity. The high dough yields of these preparations permit pumping of the dough. They are frequently used in local bakeries. Those sourdoughs are stored fresh until use (up to one week), they can be produced in large quantities. In industry, they are applied for the production of dried sourdough products as well (De Vuyst & Neysens, 2005).

4.3 Type III sourdough

Type III sourdoughs are dried doughs in powder form, which are initiated by defined starter cultures. They are used as acidifier supplements and aroma carries during bread-making. They mostly contain LAB that are resistant to drying and are able to survive in that form, e.g. heterofermentative *Lb. brevis*, facultative heterofermentative *P. pentosaceus* and *Lb. plantarum* strains. The drying process (spray-drying or drum-drying) also leads to an increased shelf-life of the sourdough and turns it into a stock product until further use. Dried sourdoughs are convenient, simple in use, and result in standardized end products. They can be distinguished in colour, aroma and acid content (De Vuyst & Neysens, 2005).

5. Rheology of the sourdough: Influence of LAB action

5.1 Effects of LAB to dough structure

Cereal grains contain starch and non-starch polysaccharides, the latter composed of glucose (β -glucans), fructose (polyfructan), xylose and arabinose (arabinoxylan) (Belitz & Grosh, 1999). Starch is partially digestible, while some other polysaccharides are not, and would represent dietary fibres. Wheat flours contain, in addition to polyfructans, also nystose, kestose and other fructooligosaccharides of the inulin type (Campbell et al., 1999). Their prebiotic potential has been well examined and documented (Van Loo et al., 1999; Corsetti & Settani, 2007). The structural effects of sourdough in wheat-based system may first be due to the direct influence of low pH on structure-forming dough components, such as gluten, starch, arabinoxylan etc. (Angioloni et. al., 2006). Dough is very sensitive to changes in ionic strength and pH and such changes could have direct impact on the constituents of dough (Clarke et al., 2002). The drop in pH value caused by the produced organic acids influences the viscoelastic behaviour of dough. A correct description of the changes in dough behaviour is necessary to maintain handling and machinability in industrialized production (Wehrle et. al., 1997). A number of earlier studies have examined influence of acids and different pH values on the dough properties. All of these confirmed that changes in the absolute pH value of sourdough significantly influence sourdough components. It has been well documented (Tsen, 1966; Wehrle et. al., 1997; Takeda et. al., 2001; Tieking et. al., 2003) and supported by the findings of the many rheological studies which indicated differences between the nonacidified and chemically or biologically acidified doughs (Clarke et. al., 2002). Osborne (1907) reported a century ago that the presence of acids increased the

solubility of the glutenin fraction extracted from the wheat flour. Barber et. al. (1992) reported that there could be mild acid hydrolysis on starch in sourdough system.

The pH profile may affect the time frame during which the acid influences the constituent ingredients of the dough. The changing pH values during sourdough fermentation period may also afford passage through a range of pH values close to the optimum for various enzymes present in the dough system. It is so-called secondary (indirect) effect of sourdough acidification (Clarke et al., 2004). The activity of proteolytic and amylolytic enzyme present may be influenced to a greater degree by the pH profile of the biological acidification fermentation period in contrast to the rather instantaneous nature of the chemically acidified regime. Optimum activity of these enzymes, which play significant role in changes of dough constituents, achieve optimum activity at pH 4-5 for the proteolytic and pH 3.6 – 6.2 for the amylolytic enzymes (Belitz & Grosh, 1992). Other enzymes that might affect the structural components of the dough the activity of which is pH dependent include peroxidases, catalases, lipoxigenases and polyphenol oxydases (Belitz & Grosh, 1992; Clarke et. al., 2002). Results obtained by the the fundamental rheological tests, baking tests, and farinograms show that activity of some enzymes in the biologically acidified dough led to structural changes in the dough (Corsetti et. al., 2000; Clarke et. al., 2002; Clarke et. al., 2004). Corsetti et. al. (2000) also reported that even limited photolytic degradation of wheat proteins affects the physical properties of gluten, which in turn can have a major effect on bread firmness and staling.

Gas production during fermentation of sourdough has been marked as a one of the most important parameters to affect on sourdough structure and rheology. Hammes and Gänzle (1998) have noted that the contribution of yeasts and LAB to the overall gas volume differs with the type of starters and the dough technology applied. These authors have also reported that gas formation by the sourdough microflora is only of minor importance if baker's yeast added. Clarke et. al. (2002) proved that the amount of gas produced by the sourdough microorganisms does not contribute remarkably to the increase in loaf specific volume. However, Hammes & Vogel (1995) indicated certain differences connected to the type of used LAB starter. Thus, obligately heterofermentative *Lb. pontis* significantly higher influenced to the gas formation in sourdough than facultatively hetrofermentative *Lb. plantarum*.

Exopolysaccharides (EPS) in sourdough are products of LAB metabolism. EPS have been recognized as the one of the important structure stability factors (Tieking et. al., 2003). Two classes of EPS from LAB can be distinguished, extracellular synthesized homopolysaccharides (HoPS), composed of only one type of monosaccharide and are synthesised by extracellular glucan and fructosyltransferases (glycosyltransferases) using sucrose as the glycosyl donor, and heteropolysaccharides (HePS) with irregular repeating units (Corsetti & Settani, 2007). HoPS are today generally applied to improve the texture of baked goods, while HePS usually have been used only in fermented dairy industry (Laws & Marshall, 2001). Sourdough lactobacilli have not been found to produce HePS.

One of the key sourdough LAB starters, *Lb. sanfranciscensis*, has been well known and characterized for its contribution to the enhancement of polysaccharide content due to the production of EPS (Korakli et. al., 2001; Korakli et. al., 2003; Tieking et. al., 2003). Formation of EPS is a well accepted characteristic of sourdough LAB, since this feature influencing on the viscosity of sourdough (Vogel et. al., 2002).

Such as presented, through the many microbiological and biochemical reaction starter microorganisms in sourdough affect on its important structural changes. From that reason, choose of adequate starters and control of structural changes during process of fermentation and baking is essential.

5.2 Dough structure

The main constituents are those derived from the flour, most importantly the proteins, both soluble and insoluble. The carbohydrates, which include starch, sugars, soluble and insoluble polysaccharides, are the most abundant. The lipids form a small, but significant part of the flour. The molecules of some minor constituents are composed of the lipid part (glycolipids) or are protein part (lipoproteins). These molecules are interesting because of their possible role in the interaction between more abundant constituents. Finally, flour enzymes may affect the dough properties (Chavan & Chavan, 2011). Water, one of the basic constituents, plays a key role in the formation of dough. During dough mixing, some air is occluded. The air forms the nuclei of the gas cells, which expand during fermentation and oven rise (Gan et. al., 1995). The remaining constituents include all other ingredients that are added to dough, including yeasts, salt, malt, enzymes, flour improvements, sugar, fats, emulsifiers, milk or soy solids, mould inhibitors, or, in the case of sourdough LAB starters in the one of the mentioned forms.

Dough is composed of a continuous phase in which gas cells are dispersed. This continuous phase is called "dough phase" (Bloksma & Bushuk 1988). The following constituents can be distinguished:

1. Starch granules, occupying about 60% of the volume of the dough phase. Large elliptical granules occur side by side with small spherical granules. Size of this granule is significantly differing between normal wheat dough and sourdough (Brummer & Lorenz, 1991);
2. Swollen protein
3. Yeast cells, with the diameter of about 2 μm
4. LAB cells (in the case of sourdough)
5. Lipids
6. Irregularly shaped remnants of cell organelles and wheat grain tissues (Pomeranz, 1988). Immediately after mixing, the gas cells form spherical holes with diameters between 10 and 100 μm . Their number at that stage is estimated to be between 10^{11} and 10^{13} m^{-3} . Clarke et al. (2002) reported that LAB starters do not significantly influencing on gas transformation in dough, but certain differences between wheat dough and sourdough has been revealed by Hammes & Vogel (1995).

Wheat flour dough, same like sourdough, is viscoelastic, exhibits both flow and elastic recovery. When a piece of dough is placed on a flat surface, if the humidity is high enough so it will flow. The amount of flow depends on the balance of viscous and elastic properties (Faridi & Faubion, 1986). In fact, dough is not truly elastic such as rubber or similar artificial materials. If a piece of dough be stretched rapidly and the force be released immediately, it will only partially recover its original shape (Hoseney, 1994). In dough, the cross-links between molecules are secondary (noncovalent) bonds that are constantly breaking and reforming other units (Rao, 1984). This appears that starch is a not an inert ingredient in

flour-water systems. Instead starch appears to act as filler in the gluten polymer. Filled polymers are known to have a larger modulus than their unfilled counterparts.

In general, sourdough structure could be described similar to the model of the only yeast-fermented dough. However, according to all facts mentioned in previous text, it is so obvious that sourdough have different structure, due to added LAB starters (Koceva Komlenić et. al., 2010).

5.3 Fundamentals in dough rheology

Rheology is the study of how materials deform, flow, or fail when force is applied. If a material viscosity is constant regardless of stirring or flowing through pipe, these materials are called Newtonian. However, in many food systems, including flour-water systems, the viscosity changes (decreases) as the shear rate is increased (Bagley, 1992). These systems show more complicated non-Newtonian behaviour, and we cannot define the system by a single viscosity value, but must give viscosity at each shear rate. Additionally, viscosity can also be affected by the time involved in making the measurement (Hoseney, 1994). Many different kind of rheological moduli exist in applied rheology. In general, modulus refer to the stiffness of the material and is proportionally constant relating stress to strain. It tells how much force is required to produce a specific deformation of the material under test (Rao, 1986).

In cereal technology, farinograph, mixograph, extensograph and amylograph have been referred to as rheological measurements (Hoseney, 1994). These instruments measure how doughs deform and flow. The problem with the use of these instruments for rheological studies is that cannot define the stress at any moment of time during the test. This is not to say that above mentioned instruments are not useful. They have stood the test of time and can give much useful information. They are particularly used when used to characterize flour. It has been important to know whether the mixing properties of the flour used today are similar to or different from those of the flour used yesterday. The mixograph or farinograph can easily and rapidly detect this (Faridi & Faubion, 1986). Unlike, from fundamental measurements we can learn much more about rheological behaviour of dough. We can find effect of various interactions and how the properties of the dough change as a function of time or temperature. Additionally, the measurements can often be made on the complete dough system so the results are easy to interpret. The second reason is, that it has now become much easier to obtained good rheological data with the advent of minicomputers incorporated to the measurements and their related equipments (Hoseney, 1994).

Rheological measurements recently used for the characterization of dough rheology are described below.

5.4 Rheological tests on dough

The physical characteristics of doughs are important in relation to the uses of flours. In the case of sourdough, LAB fermentation is another factor. Pseudo-rheological characteristics are investigated mainly with the following methods:

1. The Brabender Farinograph measures and records the resistance of dough to mixing as it is formed from flour and water, developed and broken down. This resistance is called consistency. The maximum consistency of the dough is adjusted to a fixed value by

altering the quantity of water added. This quantity, the water absorption may be used to determine a complete mixing curve, the various features of which are guide to the strength of the flour (Figure 2)

Fig. 2. Representative farinogram showing some commonly measured indexes. A consistency of 500 farinograph units (FU) corresponds to a power of 68 and 81 W per kilogram of dough in mixtures for 300 and 50 g of flour, respectively (Bloksma & Bushuk, 1988).

- 2. The Brabender Extensograph records the resistance of dough to stretching and the distance the dough stretches before breaking. A flour-salt-water dough is prepared under standard conditions in the Brabender Farinograph and moulded on the Extensograph into a standard shape. After a fixed period the dough is stretched and a curve drawn, recording the extensibility of the dough and its resistance to stretching (Fig. 3). The dough is removed and subjected to a further two stretches. The Extensograph has replaced the Extensometer in the Brabender instrument range.

Fig. 3. Extensigram, showing extensibility (E), resistance at a constant extension of 5 cm (R_5) and maximum resistance (R_m). (Hoseney, 1994).

- 3. The Chopin Alveograph uses air pressure to inflate a bubble of dough until it bursts. The instrument continuously records the air pressure and the time that elapses before the dough breaks (Figure 4).
- 4. The Brabender Amylograph continuously measures the resistance to stirring of a 10% suspension of flour in water while the temperature of the suspension is raised at a constant rate of 1.5 °C/min from 20° - 95 °C. It is of use in testing flour for soups (etc.),

for which purpose the viscosity of the product after gelatinization is an important characteristic, as well as for adjusting the malt addition to flours for breadmaking

Fig. 4. A representative alveogram. P = overpressure (mm), L = abscissa at rupture (mm), G = swelling index (ml), V = volume of air (ml), W = deformation energy (10^{-4} J), h = maximum height (mm) and S = area under the curve (cm^2). (Hoseney, 1994).

5. The Rapid Visco Analyzer (RVA), may be regarded as a derivative of the Amylograph. Measurements of viscosity are made using small samples, containing 3-4 g of starch, in periods which may be as short as 2 min. Use of disposable containers and mixer paddles eliminates the needs for careful washing of the parts between tests. As with the Amylograph, the characteristics of starch pastes and the effects of enzyme of them can be recorded on charts. They can also be transferred in digital form direct to a data-handling computer.
6. True dynamic rheological instruments. In recent years frustration with instrument-dependent units obtained with some of the above methods, together with the poor reproducibility, from one instrument to another of the same type, has led cereal chemists to turn to true rheological measurements. Suitable instruments for use with doughs, slurries and gels, derived from flours, include Bohlin VOR (viscometric, oscillation and relaxation), the Carri Med CSL Rheometer and the Rheometrics RDA2. In addition to providing excellent reproducibility, these instruments, which are also used in many non-foods, and many non-cereal food materials, allow comparisons to be made across a wide range of substances. Although they are expensive, they will undoubtedly enable the development of tests that can be performed on simpler, dedicated instruments (Faridi & Faubion, 1990; Kent & Evers, 1994).

5.5 Documented rheological results on the sourdough: Recapitulation

In a number of previous studies differences between simple yeast-fermented doughs, chemically acidified (using lactic or acetic acid) doughs and biologically acidified (using a starter culture) dough were investigated by the use of empirical and dynamic rheology measurements (Zeng et. al., 1997; Wehrle et. al., 1997; Wehrle & Arendt, 1998; Lee et. al.,

2001; Clarke et. al. 2002; Clarke et. al., 2004; Angioloni et. al., 2006). Hoseney (1994) reported that acid developed during fermentation with LAB strongly influence the mixing behaviour of doughs, whereby doughs with lower pH values require a slightly shorter mixing time and have less stability than normal doughs. The water absorption of flour is an important factor influencing the handling properties and machinability of dough in large mechanized bakeries and is related to the quality of the finished baked product (Caterall, 1998). Incorporation of sourdough changed the mixing behaviour, resulting in a significant decrease in water absorption relative to control (Maher Galal et. al., 1978; Wehrle et. al., 1997; Clarke et. al., 2002; Koceva Komlenić et. al., 2010). The addition of sourdough prepared with starter culture significantly reduced the stability of the dough relative to both the control (yeast fermented) and chemically acidified dough (Wehrle et. al., 1997; Clarke et. al., 2002; Clarke et. al., 2004). A large difference was found for the degree of softening. Addition of sourdough or acid significantly increased the degree of softening (Maher Galal et. al., 1978; Clarke et. al., 2002). Clark et. al. (2002) reported that biologically acidified doughs (sourdoughs) showed a significantly greater degree of softening than the chemically acidified (with lactic or acetic acid) doughs (Table 3. and Figure 5.).

Sample**	Water absorption (g/100g)			Dough development time (min)			Stability (min)			Degree of softening (BU)		
Control I	62.15	±	0.06 g	2.25	±	0.29 e	2.03	±	0.05 a	61.25	±	2.50 e
LAB I	60.23	±	0.05 h	3.25	±	0.29 c	0.90	±	0.23 e	82.50	±	2.89 b
DS I	64.28	±	0.05 e	2.80	±	0.23 d	1.70	±	0.12 b	66.25	±	2.50 cd
LA I	62.35	±	0.06 f	2.15	±	0.17 e	1.40	±	0.01 c	62.50	±	2.89 de
Control II	67.75	±	0.06 c	3.98	±	0.05 b	1.28	±	0.05 cd	65.00	±	5.77 cde
LAB II	65.58	±	0.05 d	4.63	±	0.10 a	0.65	±	0.06 f	122.50	±	2.89 a
DS II	69.23	±	0.05 a	4.20	±	0.14 b	1.05	±	0.29 e	81.25	±	2.50 b
LA II	68.10	±	0.12 b	4.03	±	0.05 b	0.90	±	0.23 e	67.50	±	2.89 c

*Values are mean ± SD of four independent determinations; Mean values followed by common letter within the same column are not significantly different (*p* < 0.05)
**Control, control dough; I and II, flour types: T-550 and T-110, respectively; LAB, biologically acidified dough by addition of sourdough prepared with *Lactobacillus brevis* L-75; DS, biologically acidified dough by addition of dry sourdough; LA, chemically acidified dough by addition of lactic acid

Table 3. Farinogram results* for different wheat doughs (control, biologically and chemically acidified doughs). (Koceva Komlenić et al. 2010).

The effect of sourdough addition on extensibility of dough were also investigated (Tsen, 1966; Corsetti et. al., 2000; Clarke et. al., 2002; Katina et. al. 2006a; Katina et. al. 2006b, Arendt et. al., 2007; Koceva Komlenić et. al., 2010). The Extensograph gives information about about a dough extensibility and resistance to extension (Walker & Haazelton, 1996). Generally, the addition of sourdough or acid significantly reduced extensibility of the dough. Clarke et. al. (2002) demonstrated the higher reduction of extensibility when the mixed-strain starter culture was used compared to single strain used (Figure 6). It is in agreement with the results obtained by Tsen (1966).

Fig. 5. Farinograms of doughs: Control I (prepared with flour type I) and LAB I (prepared with flour type I and *L. brevis* preferment). (Koceva Komlenić et al. 2010).

Fig. 6. Extensograph analysis of doughs with sourdough and lactic acid: I and II (different flour type), LAB (biologically acidified dough by addition of sourdough prepared using a starter culture of *L. brevis* L-75), DS (biologically acidified dough by addition of dry sourdough), LA (chemically acidified dough by addition of lactic acid). (Koceva Komlenić et al. 2010)

Dynamic oscillatory measurements examined the effects of sourdough or acid addition on the viscoelastic properties of the dough. The rheological characteristics of fermented dough are determined by many factors. At the beginning of the mixing process, physical actions, such as hydration, take place. The gluten network is formed by proteins, and starch granules absorb water. Enzyme activity of amylases, proteases and hemicellulases causes breakdown of several flour components. Changes of pH level also alter the rheological behaviour of the dough (Wehrle et al. 1997). Even small chemical and physical changes in gluten network can lead to significant changes in rheological characteristics. Formation of gas during fermentation of sourdough leads to an increased volume and decreased density.

According to the most of the authors, main variables of viscoelastic properties of the dough are phase angle (δ) and complex modulus G^* (Figure 7 and 8.). The results of fundamental oscillatory measurements gave the key attributes of rheological behaviour of the dough. The addition of sourdough prepared by LAB starters did increase the phase angle values relative to the control or the chemically acidified dough significantly (Clarke et al., 2002). It means that addition of sourdough reduced the elasticity of the dough in contrast to the chemical acidification, which did not have the same effect (Hoseney, 1994). Some authors observed that there were no significant differences between G^* values for the control and chemically acidified doughs over all frequencies, while the values for the control were significantly higher than those obtained from sourdough over all frequencies. The same was true for the chemically acidified doughs which yielded greater G^* values than the sourdoughs. An increase in phase angle and decrease in G^* due to addition of sourdough indicated that the dough was less elastic and became simultaneously less firm at low rate of strain applied (0.1%). These results also clearly suggest that chemical acidification was not directly comparable with biological acidification. Thiele et al. (2002) reported that, with regard to the levels of amino acids in wheat dough, may influence on sourdough rheology. Clarke et al. (2004) also reported a dramatic loss of firmness and elasticity in preferments during 24-hours fermentation of sourdough. These results suggest that cereal proteases with acidic optima play a central role in the rheological changes taking place during sourdough fermentation.

Wehrle & Arendt (1998) analyzed rheological characteristics of controlled and spontaneous fermented sourdough. During the first 5 hours of fermentation, phase angles did not change significantly. After 5 hours phase angles of sourdough increased sharply. Maximum phase angles were achieved after fermentation time of 20 hours. A high phase angle indicates viscous behaviour of material. Fermented sourdough after 20 hours of fermentation is an entirely viscous manner, with phase angles close to 90° , which indicates an ideal fluid. Phase angle was closely related to CO_2 production by heterofermentative microorganisms in the dough. Results of Wehrle & Arendt indicate the beginning of gas production in the sourdough after 5th hour of fermentation (Table 4). Clarke et al. (2004) also observed a dramatic loss of firmness and elasticity during 24-hr period irrespective of the presence of acid or lactic acid bacteria. The presence of acid, however, enhanced the effects. Wehrle et al. (1997) reported that doughs with acid initially were firmer and more viscous than doughs without acids, but showed less stability during mixing. Addition of acids intensified decrease in G^* and increase in δ with longer mixing times when measured immediately after mixing.

Fig. 7. Phase angle (δ) as a function of frequency for wheat dough formulations: control (nonacidified), LA (chemically acidified with lactic acid), SS1 (added sourdough prepared using a single strain starter culture of *Lactobacillus brevis* L-62), SS2 (added sourdough prepared using a single strain starter culture of *L. plantarum* L2-1), MS (added sourdough prepared using a mixed strain starter culture, Böcker Reinzucht-Sauerteig Weizen). Mean value $\pm$ standard deviation of three replicates (Clarke et al. 2002).

Fig. 8. Absolute value of the complex dynamic modulus ($|G^*|$) as a function of frequency for wheat dough formulations: control (nonacidified), LA (chemically acidified with lactic acid), SS1 (added sourdough prepared using a single strain starter culture of *Lactobacillus brevis* L-62), SS2 (added sourdough prepared using a single strain starter culture of *L. plantarum* L2-1), MS (added sourdough prepared using a mixed strain starter culture, Böcker Reinzucht-Sauerteig Weizen). Mean value $\pm$ standard deviation of three replicates. (Clarke et al. 2002).

Rheological properties, as well as acidification and flavour development, are important parameters in controlling fermentation process. Types of microorganisms, metabolic activity, and time-dependent development of pH levels have an effect on the final rheological properties.

	Control	LA ^b	SS1 ^c	SS2 ^d	MS ^e
Dough development curve parameters					
Hm ^f (mm)	87.0 ± 1.0b	79.0 ± 2.6ab	75.2 ± 1.2a	76.0 ± 3.3a	87.8 ± 6.3b
h ^g (mm)	86.3 ± 0.4b	78.3 ± 3.8ab	71.8 ± 2.3a	72.7 ± 3.6a	87.3±6.9b
(Hm – h)/Hm ^h (%)	0.7±0.8ab	0.9 ± 1.6abc	4.5 ± 2.1c	4.3 ± 0.8bc	0.6 ± 0.7a
T1 ⁱ (min)	177 ± 3b	179 ± 2b	140 ± 22a	132 ± 6a	179 ± 1b
Gaseous release curve parameters					
T ^j (min)	114.0 ± 9.1b	117.0 ± 4.0b	76.5 ± 10.4a	96.0 ± 21.3ab	67.0 ± 5.7a
Vt ^k (mL)	1,942 ± 50ab	2,019 ± 56b	1,986 ± 52ab	2,021 ± 16b	1,870 ± 75a

^a Mean value ± standard deviation of two replicates. Mean values followed by a common letter within the same row are not significantly different (P < 0.05).

^b Chemically acidified by the addition of lactic acid.

^c Added sourdough prepared using a single strain starter culture of *Lactobacillus brevis* L-62.

^d Added sourdough prepared using a single strain starter culture of *L. plantarum* L2-1.

^e Added sourdough prepared using a mixed strain starter culture, Böcker Reinzucht-Sauerteig Weizen.

^f Maximum height of dough development curve.

^g Dough development height at the end of 3-hr test period.

^h % Reduction in dough development height at the end of test period relative to T1.

ⁱ Time of maximum height of dough development curve.

^j Time of maximum height of gaseous release curve.

^k Total volume of carbon dioxide released by dough.

Table 4. Development and Gaseous Release Characteristics of Doughs from Rheofermentometer Testinga (Clarke et al. 2002)^a

6. Perspectives

The composition and processing of cereals grains, substrate formulation, growth capability and productivity of starter culture, stability of probiotic strains during storage, organoleptic properties and nutritional value of the final product are key parameters to be considered (Charalampopoulos, 2002). Probiotics bacteria from the sourdough starters must meet the criteria not only for good survival during fermentation process but also for fermentation and symbiosis with other starter cultures used (Kedia et. al., 2007). Nowadays, the use of probiotic bacteria as starter cultures in baked goods is at its early stage and merits further investigation (Font de Valdes et. al., 2010). Since sourdough systems are so complex, and the determination of rheological properties of dough should be comprehensive to reveal influence of sourdough composition and all components of flour (gluten, starch, lipids, water soluble proteins and pentosans). Regardless of whether the empirical or fundamental rheological testing is used and despite the decades of bread dough rheology testing there is still no universal equipment or method that can provide the data that are 100% accurate in evaluating performance during dough processing. Over time, numerous models of dough rheological behavior have been developed to simplify describing the rheological changes in bread dough system and they are still emerging.

Recently, Tanner *et al.* (2008) introduced the use of a simple Lodge-type model (Lodge, 1964), including a power law memory function, and a damage function (assumed to be a function of strain) to represent the breakdown of the molecular structure within the dough, for a description of the rheological behavior of wheat dough. The model produced a

satisfactory reconstruction of stress data in shear start up and elongation flow at low deformation rates. The model has relatively few parameters, all of which can easily be found from simple experiments. The sequence of step strains, with both reversing and steadily increasing steps, is reasonably well described by the model. The model can therefore describe small-strain motions, steady shearing, steady uniaxial and biaxial elongations, recoil, stress relaxation and step strain motions. Since then, several modifications of this model have been developed. Hicks *et al.* (2011) find that the damage term is explicitly a function of strain, a concept that may carry over to elongation flow and that rapid decrease of the damage function post material fracture is a representation of the transition of dough from solid- to liquid-like rheological behavior. Tanner *et al.* (2011) described an improved damage function model for bread dough rheology that enables describing uniaxial and biaxial stretching with the same damage function derived from shear data.

Further investigations of dough rheological behavior is required to obtain quality results and to develop models that accurately simulate processing situations and parameters and provides a platform for comprehensive process control.

7. References

- Angioloni, A., Romani, S. Pinnavaia, G. G., Dalla Rosa, M. (2006). Characteristics of bread making doughs: influence of sourdough fermentation on the fundamental rheological properties. *European Food Research and Technologie*, 222, pp. 54-57
- Arendt, E.K., Ryan, L.A. M. & Dal Bello, F. (2007). Impact of sourdough on the texture of bread. *Food Microbiology*, 24, pp. 165-174.
- Bagley, E. B. (1992). Mechanistic basis of rheological behaviour of foods. In: *Physical Chemistry of Foods*. Schwartzberg H. G. & Hartel R. W. (eds.), Marcel Dekker, New York.
- Barber, B., Ortolá, C., Barber, S., & Fernández, F. (1992). Storage of packaged white bread. III. Effects of sour dough and addition of acids on bread characteristics. *Zeitschrift für Lebensmittel Untersuchung und Forschung*, 194, pp. 442-449.
- Belitz, H. D., & Grosh, W. (1999). *Food chemistry*. Springer-Verlag: Berlin, Germany
- Belitz, H.-D., & Grosch, W. (1992). *Lehrbuch der Lebensmittelchemie*. 4th ed. Springer-Verlag: Berlin, Germany.
- Bloksma, A. H., & Bushuk, W. (1988). Rheology and chemistry of dough. In: *Wheat chemistry and technology*, Vol. 2, Pomeranz, Y. (Edt). American Association of Cereal Chemists, St. Paul, Minnesota. pp. 131-217.
- Böcker, G., Stolz, P., & Hammes, W. P. (1995). Neue Erkenntnisse zum Ökosystem Sauerteig und zur Physiologie des Sauerteig-Typischen Stämme *Lactobacillus sanfrancisco* und *Lactobacillus pontis*. *Getreide Mehl und Brot*, 49, pp. 370-374.
- Böcker, G., Vogel, R. F., & Hammes, W. P. (1990). *Lactobacillus sanfrancisco* als stabiles Element in einem Reinzucht-Sauerteig- Präparat. *Getreide Mehl und Brot*, 44, pp. 269-274.
- Brümmer, J.-M. & Lorenz, K. (1991). European developments in wheat sourdoughs. *Cereal Foods World*, 36, pp. 310-314.

- Cai, Y., Okada, H., Mori, H., Benno, Y., & Nakase, T. (1999). *Lactobacillus paralimentarius* sp. nov., isolated from sourdough. *International Journal of Systematic and Evolutionary Microbiology*, 49, pp. 1451-1455.
- Campbell, J. M., Bauer, L. L., Fahey, G. C., Hogarth, A. J. C. L., Wolf, B. W., & Hunter, D. E. (1997). Selected fructooligosaccharide (1-kestose, nystose, and 1F-b-fructofuranosylnystose) composition of foods and feeds. *Journal of Agricultural and Food Chemistry*, 45, pp. 3076-3082.
- Catterall, P. (1998). Flour milling. In: *Technology of Breadmaking*. S. P. Cauvain and L. S. Young, eds. Blackie Academic and Professional: London. pp. 296-329.
- Charalampopoulos, D., Pandiella, S.S., and Webb, C. (2002). Growth studies of potentially probiotic lactic acid bacteria in cereal-based substrates. *Journal of Applied Microbiology*, 92, pp. 851 - 859.
- Chavan, R. S. & Chavan, S. R. (2011). Sourdough technology - A traditional way for wholesome foods: A review. *Comprehensive Reviews in Food Science and Food Safety*. 11, pp. 170-183
- Clarke, C. I., Schober, T. J., Dockery, P., O'Sullivan, K. & Arendt, E. K. (2004). Wheat sourdough fermentation: effects of time and acidification on fundamental rheological properties. *Cereal Chemistry*, 81, pp. 409-417.
- Clarke, C.I., Schober, T.J. & Arendt, E.K. (2002). Effect of single strain and traditional mixed strain starter cultures on rheological properties of wheat dough and on bread quality. *Cereal Chemistry*, 79, pp. 640-647.
- Corsetti, A. & Settanni, L. (2007). *Lactobacilli* in sourdough fermentation. *Food Research International* 40, pp. 539-558
- Corsetti, A., De Angelis, M., Dellaglio, F., Paparella, A., Fox, P. F., Settanni, L., & Gobbetti, M. (2003). Characterization of sourdough lactic acid bacteria based on genotypic and cell-wall protein analyses. *Journal of Applied Microbiology*, 94, pp. 641-654.
- Corsetti, A., Gobbetti, M., De Marco, B., Balestrieri, F., Paoletti, F., Russi, L., and Rossi, J. (2000). Combined effect of sourdough lactic acid bacteria and additives on bread firmness and staling. *Journal of Agricultural and Food Chemistry*, 48, pp. 3044-3051.
- Corsetti, A., Lavermicocca, P., Morea, M., Baruzzi, F., Tosti, N., & Gobbetti, M. (2001). Phenotypic and molecular identification and clustering of lactic acid bacteria and yeasts from wheat (species *Triticum durum* and *Triticum aestivum*) sourdoughs of southern Italy. *International Journal of Food Microbiology*, 64, pp. 95-104.
- Czerny, M., & Schieberle, P. (2002). Important aroma compounds in freshly ground wholemeal and white wheat flour-identification and quantitative changes during fermentation. *Journal of Agricultural and Food Chemistry*, 50, pp. 6835-6840.
- De Angelis, M., Mariotti, L., Rossi, J., Servili, M., Fox, P. F., Rollán, G. C., et al. (2002). Arginine catabolism by sourdough lactic acid bacteria: purification and characterization of the arginine deiminase pathway enzymes from *Lactobacillus sanfranciscensis* CB1. *Applied and Environmental Microbiology*, 68, pp. 6193-6201.
- De Vuyst, L., & Gänzle, M. (2005). Second international symposium on sourdough: from fundamentals to applications. *Trends in Food Science and Technology*, 16, pp. 2-3.
- De Vuyst, L., & Neysen, P. (2005). The sourdough microflora: biodiversity and metabolic interactions. *Trends in Food Science and Technology*, 16, pp. 43-56.

- Ehrmann, M.A. , and Vogel , R.F. (2005) Molecular taxonomy and genetics of sourdough lactic acid bacteria. *Trends in Food Science and Technology* 16 , pp. 31 - 42 .
- Faridhi, H. & Faubion, J. M. (1990) *Dough Rheology and Baked Product Texture*. Van Norstrand Rheinhold. New York.
- Faridi H. & Faubion J. M. (1986). *Fundamentals of dough Rheology*. America Association of Cereal Chemists, St. Paul, Minnesota
- Font de Valdez, G., Gerez, C. L., Torino, M. I. & Rollán, G. (2010). New Trends in Cereal - based Products Using Lactic Acid Bacteria. In F. Mozzi, R. R. Raya, G. M. Vignolo (Eds.), *Biotechnology of lactic acid bacteria: novel applications*. Wiley-Blackwell, Iowa, pp. 273-287
- Gan Z., Ellis P. R., Schofield J. D. (1995). Mini review: Gas cell stabilisation and gas retention in wheat bread dough. *Journal Cereal Science*, 21, pp. 215-230
- Gassenmeier, K., & Schieberle, P. (1995). Potent aromatic compounds in the crumb of wheat bread (French-type)-influence of pre-ferments and studies on the formation of key odorants during dough processing. *Zeitschrift fur Lebensmittel Untersuchung und Forschung*, 201, pp. 241-248.
- Gerez , C.L., Torino, M.I., Rollán , G. & Font de Valdez , G. (2009). Prevention of bread mould spoilage by using lactic acid bacteria with antifungal properties. *Food Control* 20, pp. 144 - 148 .
- Gerez, C.L., Rollán, G. & Font de Valdez, G. (2006). Gluten breakdown by lactobacilli and pediococci strains isolated from sourdough. *Letters in Applied Microbiology*, 42, pp. 459 - 464.
- Ginés , S. , Bárcena , J.M., Ragout , A., & Font de Valdez, G. (1997) Effect of the carbon source on the fermentation balance of *Lactobacillus reuteri*. *Microbiologie, Aliments, Nutrition*, 15 , pp. 23 - 27 .
- Gobbetti, M. (1998). The sourdough microflora: interactions of lactic acid bacteria and yeasts. *Trends in Food Science and Technology*, 9, pp. 267-274.
- Gobbetti, M., & Corsetti, A. (1997). *Lactobacillus sanfrancisco* a key sourdough lactic acid bacterium: a review. *Food Microbiology*, 14, pp. 175-187.
- Gobbetti, M., Corsetti, A., Rossi, J., La Rosa, F., & De Vincenzi, S. (1994a). Identification and clustering of lactic acid bacteria and yeasts from wheat sourdoughs of central Italy. *Italian Journal of Food Science*, 6, pp. 85-94.
- Gobbetti, M., Corsetti, A., & Rossi, J. (1994b). The sourdough microflora. Interactions between lactic acid bacteria and yeasts: metabolism of carbohydrates. *Applied Microbiology and Biotechnology*, 41, pp. 456-460.
- Gobbetti, M., Rizzello, J.C., Di Cagno, R. & De Angelis, M. (2007) Sourdough lactobacilli and celiac disease. *Food Microbiology*, 24, pp. 187-196.
- Grosh, W., and Wieser, H. (1999). Redox reactions in wheat dough as affected by ascorbic acid. *Journal of Cereal Science*, 29, pp. 1-16.
- Hammes, W. P. & Vogel, R. F. (1995). The genus *Lactobacillus*. Pp. 19-54 in: *The Lactic Acid Bacteria*. Vol. 2. B. J. B. Wood and W. H. Holzapfel, eds. Chapman and Hall: London.

- Hammes, W. P., Brandt, M. J., Francis, K. L., Rosenheim, M., Seitter, F. H., & Vogelmann, S. (2005). Microbial ecology of cereal fermentations. *Trends in Food Science and Technology*, 16, pp. 4-11.
- Hammes, W.P. & Gänzle, M.G. (1998). Sourdough breads and related products. In: *Microbiology of fermented foods*. Vol 1, 2nd Ed, (edited by B.J.B. Wood). Pp. 199-216. London, UK: Blackie Academic & Professional
- Hansen, A., & Schieberle, P. (2005). Generation of aroma compounds during sourdough fermentation: applied and fundamental aspects. *Trends in Food Science and Technology*, 16, pp. 85-94.
- Hansen, A., Lund, B., & Lewis, M. J. (1989a). Flavour production and acidification of sour doughs in relation to starter culture and fermentation temperature. *Lebensmittel Wissenschaft und Technologie*, 22, pp. 145-149.
- Hansen, A., Lund, B., & Lewis, M. J. (1989b). Flavour of sourdough rye bread crumb. *Lebensmittel Wissenschaft und Technologie*, 22, pp. 141-144.
- Hicks, C.I., See, H., Ekwebelam, C. (2011). The shear rheology of bread dough: modeling. *Rheologica Acta*. 50, pp. 701-710.
- Hoffmann, T., & Schieberle, P. (2000). Formation of aroma-active Strecker aldehydes by a direct oxidation of Amadori compounds. *Journal of Agricultural and Food Chemistry*, 48, pp. 4301-4305.
- Hoseney, R.C. (1994). *Principles of cereal science and technology*. American Association of Cereal Chemists, Inc. St. Paul Minnesota. USA .
- Jänsch, A., Korakli, M., Vogel, R. F. & Gänzle, M. G. (2007). Glutathione Reductase from *Lactobacillus sanfranciscensis* DSM20451: Contribution to Oxygen Tolerance and Thiol Exchange Reactions in Wheat Sourdoughs. *Applied And Environmental Microbiology*, 73, pp. 4469-4476
- Juillard, V., Guillot, A., Le Bars, D., & Gripon, J. C. (1998). Specificity of milk peptide utilization by *Lactococcus lactis*. *Applied and Environmental Microbiology*, 64, pp. 1230-1236.
- Katina, K., Heiniö, R.L., Autio, K. & Poutanen, K. (2006a). Optimization of sourdough process for improved sensory profile and texture of wheat bread. *LWT - Food Science and Technology*, 39, pp. 1189-1202.
- Katina, K., Salmenkallio-Marttila, M., Partanen, R., Forssell, P. & Autio, K. (2006b). Effects of sourdough and enzymes on staling of high-fibre wheat bread. *LWT - Food Science and Technology*, 39, pp. 479-491.
- Kazanskaya, L. N., Afanasyeva, O. V., & Patt, V. A. (1983). Microflora of rye sours and some specific features of its accumulation in bread baking plants of the USSR. In J. Holas & F. Kratochvil (Eds.), *Developments in food science. Progress in cereal chemistry and technology* (pp. 759-763). London: Elsevier.
- Kedia, G., Wang, R., Patel, H., and Pandiella, S.S. (2007) Use of mixed cultures for the fermentation of cereal - based substrates with potential probiotic properties. *Process Biochemistry*, 42, pp. 65 - 70 .
- Kent, N. L. & Evers, A. D. (1994). *Technology of Cereals*. Elsevier Science, Oxford.

- Ketabi, A., Soleimanian-Zad, S., Kadivar, M., Sheikh-Zeinoddin, M. (2008). Production of microbial exopolysaccharides in the sourdough and its effects on the rheological properties of dough. *Food Research International*, 41, 10, pp. 948-951
- Kirchhoff, E., & Schieberle, P. (2002). Quantitation of odor-active compounds in rye flour and rye sourdough using a stable isotope dilution assay. *Journal of Agricultural and Food Chemistry*, 50, pp. 5311-5378.
- Koceva Komlenić, D., Ugarčić-Hardi, Ž., Jukić, M., Planinić, M., Bucić-Kojić, A., Strelec, I. (2010). Wheat dough rheology and bread quality effected by *Lactobacillus brevis* preferment, dry sourdough and lactic acid addition. *International journal of food science & technology*. 45, pp. 1417-1425
- Korakli, M., Pavlovic, M., Gänzle, M. G., & Vogel, R. F. (2003). Exopolysaccharide and kestose production by *Lactobacillus sanfranciscensis* LTH2590. *Applied and Environmental Microbiology*, 69, pp. 2073-2079.
- Korakli, M., Rossman, A., Gänzle, M. G., & Vogel, R. F. (2001). Sucrose metabolism and exopolysaccharide production in wheat and rye sourdough by *Lactobacillus sanfranciscensis*. *Journal of Agricultural and Food Chemistry*, 49, pp. 5194-5200.
- Kumar, S., Tamura, K., & Nei, M. (2004). MEGA3: Integrated software for molecular evolutionary genetics analysis and sequence alignment. *Briefings in Bioinformatics*, 5, pp. 150-163.
- Laws, A., & Marshall, V. M. (2001). The relevance of exopolysaccharides to the rheological properties in milk fermented with ropy strains of lactic acid bacteria. *International Dairy Journal*, 11, pp. 709-721.
- Lee, L., Ng, P. K. W., Whallon, J. H. & Steffe, J. F. (2001). Relationship between rheological properties and microstructural characteristics of nondeveloped, partially developed, and developed doughs. *Cereal Chemistry*. 78, pp. 447-452
- Lodge AS (1964) Elastic liquids. Academic, London
- Lopez, H.W., Duclos, V., Coudray, C., Krespine, V., Feillet-Coudray, C., Messenger, A., Demigné, C. & Rémésy, C. (2003). Making bread with sourdough improves mineral bioavailability from reconstituted whole wheat flour in rats. *Nutrition*, 19, pp. 524-530.
- Lorenz, K. (1983). Sourdough processes. Methodology and biochemistry. *Baker's Digest*, 55, pp. 85-91.
- Lund, B., Hansen, A., & Lewis, M. J. (1989). The influence of dough yield on acidification and production of volatiles in sour doughs. *Lebensmittel Wissenschaft und Technologie*, 22, pp. 150-153.
- Maher Galal, A., Varriano-Marston, E. & Johnson, J. A. (1978). Rheological dough properties as affected by organic acids and salt. *Cereal Chemistry*. 55, pp. 683-691.
- Meroth, C. B., Hammes, W. P., & Hertel, C. (2004). Characterisation of the microbiota of rice sourdoughs and description of *Lactobacillus spicheri* sp. nov. *Systematic and Applied Microbiology*, 27, pp. 151-159.
- Osborne, T. B. (1907). The proteins of the wheat kernel. Carnegie Institute of Washington publication 84. Judd and Detweiler: Washington, DC.

- Plessas, S., Trantallidi, M., Bekatorou, A., Kanellaki, M., Nigam, P. & Koutinas, A.A. (2007). Immobilization of kefir and *Lactobacillus casei* on brewery spent grains for use in sourdough wheat bread making. *Food Chemistry*, 105, pp. 187-194.
- Rao, V. N. M. (1984). Dynamic force deformation properties of foods. *Food Technology* 38, pp. 103-109
- Rollán, G. & Font de Valdez, G. (2001). The peptide hydrolase system of *Lactobacillus reuteri*. *International Journal of Food Microbiology*, 70, pp. 303 - 307.
- Salovaara, H. (1998). Lactic acid bacteria in cereal-based products. In S. Salminen & A. von Wright (Eds.), *Lactic acid bacteria microbiology and functional aspects* (pp. 115-138). New York: Marcel Dekker.
- Settanni, L., Van Sinderen, D., Rossi, J., & Corsetti, A. (2005a). Rapid differentiation and in situ detection of 16 sourdough *Lactobacillus* species by multiplex PCR. *Applied and Environmental Microbiology*, 71, pp. 3049-3059.
- Spicher, G., & Lönner, C. (1985). Die mikroflora des sauerteiges. XXI. Mitteilung: die in sauerteigen schwedischer bäckereien vorkommenden lactobacillen. *Zeitschrift für Lebensmittel Untersuchung und Forschung*, 181, pp. 9-13.
- Stephan, H., & Neumann, H. (1999a). Technik der Roggen-Sauerteigführung. In G. Spicher, & H. Stephan (Eds.), *Handbuch Sauerteig: Biologie, Biochemie, Technologie* (5th ed.) (pp. 161-245). Hamburg: Behr's Verlag.
- Stephan, H., & Neumann, H. (1999b). Technik der Weizenvorteig und Weizensauerteigführung. In G. Spicher, & H. Stephan (Eds.), *Handbuch Sauerteig: Biologie, Biochemie, Technologie* (5th ed.) (pp. 247-275). Hamburg: Behr's Verlag.
- Stolz, P. (1999). Mikrobiologie des Sauerteiges. In G. Spicher & H. Stephan (Eds.), *Handbuch Sauerteig: Biologie, Biochemie, Technologie* (pp. 35-60). Hamburg: Behr's Verlag.
- Stolz, P. (1999). Mikrobiologie des Sauerteiges. In G. Spicher, & H. Stephan (Eds.), *Handbuch Sauerteig: Biologie, Biochemie, Technologie* 5th ed. (pp. 35-60). Hamburg: Behr's Verlag.
- Stolz, P., Hammes, W. P., & Vogel, R. F. (1996). Maltosephosphorylase and hexokinase activity in lactobacilli from traditionally prepared sourdoughs. *Advances in Food Science*, 18, pp. 1-6.
- Takeda, K., Matsumura, Y. & Shimizu, M. (2001). Emulsifying and surface properties of wheat gluten under acidic conditions. *Journal of Food Science*, 66, pp. 393-399.
- Tamman, J. D., Williams, A. G., Novle, J., & Lloyd, D. (2000). Amino acid fermentation in non-starter *Lactobacillus* spp. isolated from cheddar cheese. *Letters in Applied Microbiology*, 2000, pp. 370-374.
- Tanner, R.I., Qi, F., Dai, S. (2011). Bread dough rheology: an improved damage function model. *Rheologica Acta*. 50, pp. 75-86.
- Tanner, R.I., Qi, F., Dai, S. (2008). Bread dough rheology and recoil: I. Rheology. *Journal of Non-Newtonian Fluid Mechanics*. 148, pp. 33-40.
- Thiele, C., Gänzle, M. G., & Vogel, R. F. (2003). Fluorescence labeling of wheat proteins for determination of gluten hydrolysis and depolymerisation during dough processing and sourdough fermentation. *Journal of Agricultural and Food Chemistry*, 51, pp. 2745-2752.

- Thiele, C., Gänzle, M.G., & Vogel, R.F. (2002). Contribution of sourdough lactobacilli, yeast and cereal enzymes to the generation amino acids in dough relevant for bread flavour. *Cereal Chemistry*, 79, pp. 45-51.
- Thiele, C., Grassl, S. & Gänzle, M. G. (2004). Gluten hydrolysis and depolymerization during sourdough fermentation. *Journal of Agricultural and Food Chemistry*, 52, pp. 1307-1314.
- Tieking, M., Korakli, M., Ehrmann, M. A., Gänzle, M. G., & Vogel, R. F. (2003). In situ production of exopolysaccharides during sourdough fermentation by cereal and intestinal isolates of lactic acid bacteria. *Applied and Environmental Microbiology*, 69, pp. 945-952.
- Tieking, M., Korakli, M., Ehrmann, M. A., Gänzle, M. G., & Vogel, R. F. (2003). In situ production of exopolysaccharides during sourdough fermentation by cereal and intestinal isolates of lactic acid bacteria. *Applied and Environmental Microbiology*, 69, pp. 945-952.
- Tsen, C. C. (1966). A note on effects of pH on sulfhydryl groups and rheological properties of dough and its implication with the sulfhydryl-disulfide interchange. *Cereal Chemistry*, 43, pp. 456-460.
- Valmorri, S., Settanni, L., Suzzi, G., Gardini, F., Vernocchi, P., & Corsetti, A. (2006). Application of a novel polyphasic approach to study the lactobacilli composition of sourdoughs from the Abruzzo region (central Italy). *Letters in Applied Microbiology*, 43, pp. 343-349
- Van Loo, J., Cummings, J., Delzenne, N., Englyst, H., Franck, A., Hopkins, M., et al. (1999). Functional Food Properties of Non-digestible Oligosaccharides: a Consensus Report from theENDOProject (DGXII AIRII-CT-1095). *British Journal of Nutrition*, 81, pp. 121-132.
- Vermeulen, N., Gänzle, M. G., & Vogel, R. F. (2006). Influence of peptide supply and cosubstrates on phenylalanine metabolism of *Lactobacillus sanfranciscensis* DSM20451T and *Lactobacillus plantarum* TMW1.468. *Journal of Agricultural and Food Chemistry*, 54, pp. 3832-3839.
- Vogel, R. F., Ehrmann, M. A., & Gänzle, M. G. (2002). Development and potential of starter lactobacilli resulting from exploration of the sourdough ecosystem. *Antonie van Leeuwenhoek*, 81, pp. 631-638.
- Vogel, R. F., Knorr, R., Müller, M. R. A., Steudel, U., Gänzle, M. G., & Ehrmann, M. A. (1999). Non-dairy lactic fermentations: the cereal world. *Antonie van Leeuwenhoek*, 76, pp. 403-411.
- Walker, C. E., & Hazelton, J. L. (1996). Dough rheological testing. *Cereal Foods World*, 41, pp. 23-28.
- Wehrle, K. & Arendt, E.K. (1998). Rheological changes in wheat sourdough during controlled and spontaneous fermentation. *Cereal Chemistry*, 75, 882-886.
- Wehrle, K., Grau, H., & Arendt, E.K. (1997). Effects of lactic acid, acetic acid and table salt on fundamental rheological properties of wheat dough. *Cereal Chemistry*, 74, pp. 739-744.

- Zeng, M., Morris, C. F., Batey, I. L., & Wrigley, C. W. (1997). Sources of variation for starch gelatinization, pasting, and gelation properties on wheat. *Cereal Chemistry*, 74, pp. 63-71.
- Zotta, T., Piraino, P., Ricciardi, A., McSweeney, P. L., & Parente, E. (2006). Proteolysis in model sourdough fermentations. *Journal of Agricultural and Food Chemistry*, 54, pp. 2567-2574.

IntechOpen

IntechOpen

Rheology

Edited by Dr. Juan De Vicente

ISBN 978-953-51-0187-1

Hard cover, 350 pages

Publisher InTech

Published online 07, March, 2012

Published in print edition March, 2012

This book contains a wealth of useful information on current rheology research. By covering a broad variety of rheology-related topics, this e-book is addressed to a wide spectrum of academic and applied researchers and scientists but it could also prove useful to industry specialists. The subject areas include, polymer gels, food rheology, drilling fluids and liquid crystals among others.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Daliborka Koceva Komlenić, Vedran Slaćanac and Marko Jukić (2012). Influence of Acidification on Dough Rheological Properties, Rheology, Dr. Juan De Vicente (Ed.), ISBN: 978-953-51-0187-1, InTech, Available from: <http://www.intechopen.com/books/rheology/influence-of-acidification-on-dough-rheological-properties->

INTech
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen