We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Characterization of Repolarization Alternans in the Coronary Artery Disease

Laura Burattini and Roberto Burattini
*Department of Information Engineering,
 Polytechnic University of Marche, Ancona,
 Italy*

1. Introduction

Repolarization alternans (RA), an electrophysiologic phenomenon consisting on every-other-beat changes of the repolarization morphology (amplitude, shape, and, sometimes, polarity) unaccompanied by gross changes in the heart-cycle length, is generally recognized as a promising electrocardiographic (ECG) predictor of sudden cardiac arrest (SCA; Bloomfield et al., 2006, Chow et al. 2006, Ikeda et al., 2006, Klingenheben et al., 2000, Leino et al., 2009, Maeda et al., 2009, Narayan, 2006, Rosenbaum et al., 1994, Sakaki et al., 2009, Salerno-Uriarte et al., 2007, Stein et al., 2008), that is one of the leading causes of death in developed countries (over 300,000 lives per year in the U.S.; Montagnana et al., 2008, Zheng et al., 2001). On the ECG tracing, RA consists of an alternation of either the ST-segment (Nearing et al., 1991, Rosenbaum et al., 1994), or the T-wave, or even the U-wave (Habbab & el-Sherif, 1992). Macroscopic and, thus, visible RA (like that reported in Fig. 1) is quite rare. After being first reported by Hering in the early 1900s (Hering, 1909), RA has later on been occasionally observed in patients affected by various diseases, among which ischemia (Kleinfeld & Rozanski, 1977), long QT syndrome (Schwartz & Malliani, 1975, Zareba et al., 1994) and ventricular arrhythmias (Verrier et al., 2003). Recent investigations on RA rely on automatic detection of microvolt RA from the digital ECG signal by means of specifically designed techniques, among which the spectral method (implemented in the commercial ECG machine CH2000 and Heartwave, Cambridge Heart Inc., Bedford, MA; Smith et al., 1988, Rosenbaum et al., 1994, Rosenbaum et al., 1996), the complex demodulation method (Nearing et al., 1991, Nearing & Verrier, 1993), the modified moving average method (implemented in the commercial ECG machine CASE-8000, GE Medical Systems, Milwaukee, WI; Nearing & Verrier, 2002), the Laplacian likelihood ratio method (Martínez & Olmos, 2005, Martínez et al., 2006), our adaptive match filter method (Burattini et al., 2008, Burattini et al., 2009b) and others (Burattini et al., 1999, Martínez & Olmos, 2005). Microvolt RA has been found to be much more common than visible RA (Adachi et al., 1999, Chow et al., 2007, Ikeda et al., 2006, Klingenheben et al., 2000, Narayan et al., 2006, Narayan et al., 2007, Reddy et al., 1984) and linked to inducible (Narayan and Smith, 2000, Rosenbaum et al., 1994, Smith et al., 1988) as well as spontaneous (Bloomfield et al., 2006, Klingenheben et al., 2000, Maeda et al., 2009, Narayan, 2006, Verrier et al., 2003) ventricular arrhythmias.

The pathophysiology underlying microvolt RA in humans has not been delineated yet. Several experimental settings have shown that RA may reflect spatial (Banville & Gray, 2002, Chinushi et al., 1998, Narayan, 2006) or temporal (Narayan, 2006, Pastore et al., 1999) dispersion of repolarization, both preceding ventricular fibrillation (Weiss et al., 1999, Smith & Cohen, 1984). Indeed, spatial variations in repolarization (action potential duration, APD) or conduction velocity may prevent depolarization in myocytes that are still repolarizing from their last cycle, causing the typical 2:1 behavior (Banville & Gray, 2002, Chinushi et al., 1998). RA may also result from an alternation of the APD (temporal dispersion of repolarization), which can occur when the relationship of the APD to its preceding diastolic interval (i.e. the APD restitution curve) has a slope greater than one. Under these conditions, mainly occurring in the presence of fast heart rates, small changes of the diastolic interval may cause large APD fluctuations that facilitate alternans (Pastore et al., 1999).

Fig. 1. ECG tracing affected by macroscopic RA.

Alternans derived from the cardiac spatial and temporal dispersion of repolarization projects onto body-surface ECG RA, which shows (Selvaraj et al., 2007) an heterogeneous temporal location along the JT segment, constituted by the ST-segment and T-wave complex. Characterization of this heterogeneity is a matter of major interest, since controversial results have been reported after attempting to identify an association of RA to different diseases. According to Narayan & Smith (Narayan & Smith, 1999) RA has been found to be more specific for inducible ventricular tachycardia when distributed late, rather

than elsewhere, throughout the JT segment. In other studies (Martínez et al., 2006, Nearing et al., 1994) RA has been found to be located within the ST segment and the first half of the T wave in patients undergoing left anterior descending artery occlusion (LAD) and left circumflex artery occlusion (LCX), and to occur a little further on along the JT segment in patients undergoing right coronary artery occlusion (RCA). Eventually, RA heterogeneity along the entire JT segment has been observed in patients with cardiomyopathy (Selvaraj et al., 2007). To the best of our knowledge, quantitative investigations on RA location along the repolarization segment in the coronary artery disease have not been reported yet. Such an investigation was the aim of the present study.

Considering that our adaptive match filter (AMF) based method has been successfully used in previous clinical (Burattini et al., 2008, Burattini et al., 2009a, Burattini et al., 2010) and simulation (Burattini et al., 2006, Burattini et al., 2009b, Burattini et al., 2011) studies for characterization of T-wave alternans (which can be considered a special case of RA limited to the T-wave), this method was applied in the present study to Holter ECGs recordings from coronary artery disease (CAD) patients and control healthy (CH) subjects in order to: A) automatically identify and quantitatively characterize RA in terms of both amplitude and location with respect to the T-wave apex; and B) define a physiological RA region which allows discrimination of abnormal (RA+) from normal RA cases.

2. Methods

2.1 Study populations and clinical data

Our clinical data belong to the Intercity Digital Electrocardiology Alliance (IDEAL) Study databases, available at the Telemetric and Holter ECG Warehouse database (<http://thew-project.org>). The IDEAL protocol was approved by the Research Subject Review Board of the University of Rochester (Rochester, NY) and the study was conducted following required rules for human subjects' research principles, according to the Declaration of Helsinki, as well as to Title 45, U.S. Code of Federal Regulations, Part 46, Protection of Human Subjects, Revised November 13, 2001, effective December 13, 2001.

Two populations were considered, which consist of 201 coronary artery disease (CAD; 166 men) patients and 167 control-healthy (CH; 86 men) subjects, respectively. Namely, a patient was classified as belonging to the CAD-group if:

1. having positive angiogram (at least one vessel with critical narrowing >75%) with either exercise induced ischemia on ECG or evidence of previous myocardial infarction;
2. being in stable phase of the ischemic heart disease with digital ECG recordings performed on an outpatients basis. Stable post-MI patients should be enrolled at least 2 month after the index event;
3. not having evidence of congenital heart failure. Patients with clinical or echocardiographic evidence for congestive heart failure (LVD >60 mm, EF <40%) were excluded;
4. being in sinus rhythm (no atrial flutter/fibrillation, no pacemaker rhythm, no atrioventricular block, no sick sinus syndrome);
5. not having dilated cardiomyopathy (left ventricular diameter >60 mm and ejection fraction <40%);
6. not having coronary instability (unstable angina requiring hospitalization with evolving ischemic changes in standard ECG);

7. not having congestive heart failure (left ventricular diameter >60 mm and ejection fraction <40%);
8. not having had coronary artery bypass surgery (CABG) in the past. Patients with a history of non-CABG coronary revascularization (percutaneous transluminal coronary angioplasty, stent, atherectomy) were eligible to be enrolled;
9. not having major comorbidity such as malignancy, severe hepatic, renal or cerebral disease.

Instead, a subject was classified as belonging to the CH-group when fulfilling the following criteria:

1. not having overt cardiovascular disease or history of cardiovascular disorders (including stroke, transient ischemic attack, and peripheral vascular disease);
2. not having history of high blood pressure (>150/90 mmHg);
3. not taking medication;
4. not being affected by other chronic illness (e.g. diabetes, asthma, chronic obstructive pulmonary disease, etc.);
5. being diagnosed as being healthy if evaluated by a physician for cardiovascular-related syndrome (chest pain, palpitation, syncope);
6. having normal physical examination;
7. having sinus rhythm in 12-lead ECG without any suspicious abnormalities (e.g. signs of ventricular hypertrophy, inverted T-wave, intraventricular conduction disturbances);
8. having normal echo and normal ECG exercise testing in presence of suspicious ECG changes;
9. not being pregnant.

Clinical parameters such as age, body mass index (BMI), systolic (SAP) and diastolic (DAP) arterial pressure were measured in each subject, whereas the ventricular ejection fraction (LVEF, in %) was determined only for the CAD patients. Therapeutic treatments were also reported. In addition, a 20-minute, 3-lead (pseudo-orthogonal configuration, with bipolar leads X, Y, Z corresponding to limb lead I, augmented limb lead aVF, and precordial lead V3, respectively) digital Holter ECG recording was obtained from each individual in supine and resting conditions, making use of Burdick recorders (Burdick Inc., Milton, WI; sampling frequency: 200 Hz, resolution: 10 μ V). ECG tracings were used to determine the heart rate (mRR; computed as the mean time interval RR between two consecutive sinus beats in a 5-min window) and heart-rate variability (sdRR; determined as the standard deviation of the RR intervals in a 5-min window) as well as RA parameters (see below).

2.2 RA identification and characterization

Repolarization alternans (RA) was identified in the first 5 minutes of each ECG recording. More specifically, ECG segments consisting of 16 consecutive heart beats were recursively (every 2 s) submitted to a preprocessing stage performing noise removal, R-peak detection, baseline removal and identification, and replacement of ectopic or noisy beats (Burattini et al., 2006, Martínez. & Olmos, 2005), before being processed by our heart-rate adaptive match filter for RA identification (see below). Only 16-beat ECG strings characterized by a stable heart rate (RR-interval standard deviation less than 10% mean RR interval) and by the presence of a low level of noise or ectopic beats (no more than 1 replaced beats in each ECG string) were considered eligible for the subsequent RA analysis.

In the presence of a fixed heart rate (and, thus, of a constant time-interval RR between two consecutive sinus beats) RA is, by definition, characterized by a specific frequency of half heart rate: $f_{RA}=0.5$ cycles per beat, or $f_{RA}=1/(2 \times RR)$ Hz. In clinical cases in which heart rate may be considered stable but presenting some physiological variation, the RA phenomenon is assumed to be characterized by a narrow frequency band, $2 \times df_{RA}$ wide, centered around $f_{RA}=1/(2 \times \text{meanRR})$ (Fig. 2; Burattini et al., 2006, Burattini et al., 2008, Burattini et al., 2009a, Burattini et al., 2010). On this basis, our AMF is designed as a pass-band filter with its passing band centered in f_{RA} (Burattini et al., 2006, Burattini et al., 2008).

Technically, the AMF is implemented as a 6th order bidirectional Butterworth band-pass filter having the passing band 0.12 Hz wide (i.e. $df_{RA}=0.06$ Hz) and centered at a frequency that adapts to mean RR interval. More specifically, our AMF is a cascade of a low-pass filter (LPF) with cut-off frequency $f_{LPF}=f_{RA}+df_{RA}$, and a high-pass filter (HPF) with a cut-off frequency $f_{HPF}=f_{RA}-df_{RA}$. The squared module of the AMF transfer function is expressed by the following equation:

$$|H_{AMF}(f)|^2 = |H_{LPF}(f)|^2 \cdot |H_{HPF}(f)|^2 = \frac{1}{1 + \left(\frac{f}{f_{LPF}}\right)^6} \cdot \frac{\left(\frac{f}{f_{HPF}}\right)^6}{1 + \left(\frac{f}{f_{HPF}}\right)^6} \quad (1)$$

Being the AMF applied in a bidirectional fashion, no group delay occurs. Thus, the AMF is expected to detect RA by filtering out noise as well as every other ECG component but the RA typical one.

The input signal of the AMF is the 16-beat ECG tracing over which f_{RA} has been computed. In the absence of RA, the output of the AMF, called RA signal (Fig. 2), is a zero constant signal. Instead, in the presence of RA, the RA signal is a sinusoidal signal, possibly amplitude-modulated, characterized by the same length of the ECG and by a frequency equal to f_{RA} .

The time occurrences of the sinusoid maxima or minima, which are expected to fall inside the JT intervals when pertaining to RA, provide the center of mass of the alternations in each beat, and, thus, a localization of the alternans inside the repolarization (Fig. 3). After having identified a reference point (T_{ref}), inside the repolarization segment, as the T-wave apex for monophasic T waves, or as the amplitude-weighted mean point between T apexes for biphasic T waves, an RA delay (RAD, in ms) parameter was computed, for each beat, as the difference between the instant at which T_{ref} occurs and the instant at which the sinusoidal RA signal maximum or minimum occurs. Thus, a negative RAD value indicates the presence of early repolarization, which mostly involves the ST segment or the first half of the T wave. Instead, an RAD value close to zero indicates the presence of central RA, which mostly occurs over the T-wave apex. Eventually, a positive RAD value indicates the presence of late RA, characterized by the alternation of the final portion of the T wave. Within the same beat, the amplitude value of the sinusoidal RA signal provides an estimate of the RA amplitude (RAA, in μV).

Fig. 2. Block diagram describing the AMF-based procedure for characterization of RA in terms of RAA and RAD parameters.

In the present study, RA analysis was initially performed in each ECG lead as follows. RAD and RAA values were first averaged over the 16 beats of each single-lead ECG segment (M16b_RAD_X and M16b_RAA_X, M16b_RAD_Y and M16b_RAA_Y, and M16b_RAD_Z and M16b_RAA_Z for leads X, Y, and Z, respectively); the resulting values were then averaged over the entire 5 minutes of each single-lead recording (M5m_RAD_X and

M5m_RAA_X, M5m_RAD_Y and M5m_RAA_Y, and M5m_RAD_Z and M5m_RAA_Z for leads X, Y, and Z, respectively). Eventually, the latter values were averaged over the three leads (M5m_RAD and M5m_RAA) for a comprehensive RA characterization relative to a single patient.

Fig. 3. Simulated ECG tracing (solid lines) and relative RA signal (dotted lines) at the output of the heart-rate adaptive match filter (AMF) in the presence of early (panel A), central (panel B), and late (panel C) repolarization alternans.

2.3 Definition of an RA normality region

The M5m_RAD and M5m_RAA distributions over the CH population were used to identify an RA normality region delimited by three thresholds. Two of them (THR_RADmin and THR_RADmax) were defined for the M5m_RAD parameter as the 0.5th and the 99.5th percentiles of the M5m_RAD distribution, respectively. The third one (THR_RAA) was defined for the non-negative M5m_RAA parameter as the 99.5th percentile of the M5m_RAA distribution. A subject, independently of the belonging population, was classified as characterized by normal RA levels if the following conditions were simultaneously satisfied:

$$\begin{cases} M5m_RAD \geq THR_RADmin \\ M5m_RAD \leq THR_RADmax \\ M5m_RAA \leq THR_RAA \end{cases} \quad (2)$$

These conditions define a physiologic RA region in the M5m_RAD vs. M5m_RAA plane (Fig. 4). Thus, no satisfaction of at least one condition denotes an abnormal RA case (RA+).

Fig. 4. Definition of a physiologic RA region in the M5m_RAD vs. M5m_RAA plane (shaded area in the upper panel). RA cases falling outside the physiologic RA region are considered abnormal RA cases (RA+). The ECG beat in the lower panel shows RA location along the JT interval.

2.4 Statistics

To be independent of normal distributions, non-parametric tests were used to perform comparisons among quantities. In the specific, comparisons between the distributions of the clinical and ECG parameters over the two populations (CAD vs. CH) were performed using the Wilcoxon rank sum test for equal medians. The Kruskal-Wallis test was used to perform the one-way ANOVA test to evaluate if, in a population, the RA parameters distributions over the three ECG leads were characterized by the same median value. Information about which pairs of leads had different median values was obtained using the multiple comparison procedure. The χ^2 test was used to compare number of treated subjects between the two populations. Eventually, the existence of a possible relationship of the RA parameters with the clinical or the other ECG parameters was evaluated computing the correlation coefficient ρ . Statistical significance level was set at 5%.

3. Results

A summary of the clinical and ECG parameter values characterizing the CAD and the CH populations are reported in Table 1, where pharmaceutical treatments are also specified. Compared to the CH subjects, the CAD patients were significantly older and characterized by significantly higher BMI, SAP, and DAP, and by significantly lower heart rate (longer mRR) and heart-rate variability (lower sdRR).

	CAD patients (201)	CH subjects (167)	P value
Clinical parameters:			
Age (years)	58±11	38±15	<10 ⁻³²
BMI (kg/m ²)	27±4	24±5	<10 ⁻¹³
SAP (mmHg)	129±18	118±12	<10 ⁻¹¹
DAP (mmHg)	79±10	75±8	<10 ⁻³
LVEF	58±11	NA	-
Treatments:			
Beta blocker	150	0	<0.005
Digoxin	0	0	NS
Diuretic	11	0	<0.01
ACE inhibitor	57	0	<0.005
Antiarrhythmic	6	0	NS
ECG parameters:			
mRR (ms)	960±155	904±140	<10 ⁻³
sdRR (ms)	37±40	43±24	<10 ⁻⁶
M5m_RAD_X (ms)	-29±52	-25±29	NS
M5m_RAA_X (μV)	17±10	15±5	NS
M5m_RAD_Y (ms)	-34±51	-24±30	NS
M5m_RAA_Y (μV)	21±12	20±8	NS
M5m_RAD_Z (ms)	-38±44	-32±31	NS
M5m_RAA_Z (μV)	18±9	15±5	<10 ⁻³
M5m_RAD(ms)	-33±37	-27±23	NS
M5m_RAA (μV)	19±9	17±15	<0.05

Table 1. Clinical parameters, therapeutic treatments and ECG parameters relative to the CAD patients and the CH subjects. NA: not available; - : not applicable; NS: not significant.

Some levels of RA were detected in all 201 CAD patients and 167 CH subjects. In both populations, RA amplitude values were significantly greater in lead Y (CAD: M5m_RAA_Y=21±12 μV; CH: M5m_RAA_Y=20±8 μV) than in leads X and Z (CAD: M5m_RAA_X=17±10 μV, M5m_RAA_Z=18±9 μV, P<10⁻⁵; CH: M5m_RAA_X=15±5 μV, M5m_RAA_Z=15±5 μV, P<10⁻⁹). Instead, comparable values of RA delay were measured over the leads in the CAD patients (M5m_RAD_X=-29±52 ms, M5m_RAD_Y=-34±51 ms, M5m_RAD_Z=-38±44 ms), while the CH subjects were characterized by a significantly lower RA delay value in lead Z (M5m_RAD_Z=-32±31 ms) compared to both lead X and Y (M5m_RAD_X=-25±29 ms, M5m_RAD_Y=-24±30 ms, P<0.01). Although RA amplitude appears systematically higher for the CAD patients than for the CH subjects in all leads, statistically significant differences were observed only in correspondence of lead Z (M5m_RAA_Z). No significant differences were found between the two populations in terms of RAD parameters, independently of the lead.

A comprehensive description of the RA phenomenon in the two populations was provided by M5m_RAA and M5m_RAD parameters. In the specific, the CAD population showed RA episodes characterized by higher amplitude (CAD M5m_RAA: $19\pm9\text{ }\mu\text{V}$, CH M5m_RAA: $17\pm5\text{ }\mu\text{V}$, $P<0.05$; Table 1) and comparable delay (CAD M5m_RAD: $-33\pm37\text{ ms}$, CH M5m_RAD: $-27\pm23\text{ ms}$; Table 1). Still, M5m_RAD and M5m_RAA standard deviation values (Table 1) as well as the their distribution histograms (Fig. 5) indicate a greater M5m_RAD and M5m_RAA variability among the CAD patients than the CH subjects. According to these results, RA is localized over the first half of the T wave in most cases of the CH population, and only occasionally overcomes the T-wave apex. Instead, among the CAD patients, RA is distributed along the entire repolarization segment (ST/T wave), with several cases occurring over the ST segment or over the T-wave right-hand side.

Fig. 5. Histograms of M5m_RAD and M5m_RAA distributions for the CAD (panels A and B) and CH (panels C and D) populations showing a greater M5m_RAD and M5m_RAA variability among the CAD patients than the CH subjects.

The definition of an RA normality region (Fig. 6) delimited by $\text{THR_RADmin}=-82\text{ ms}$, $\text{THR_RADmax}=28\text{ ms}$, and $\text{THR_RAA}=35\text{ }\mu\text{V}$ (as defined in Methods) allowed the identification of 29 (14.4%) RA+ CAD patients with abnormal M5m_RAD. Specifically, 22 (10.9%) had abnormally low M5m_RAD ($\text{M5m_RAD}<\text{THR_RADmin}$) while the remaining 7 (3.5%) had abnormally high M5m_RAD ($\text{M5m_RAD}>\text{THR_RADmax}$). RA+ CAD patients characterized by abnormally high M5m_RAA ($\text{M5m_RAA}>\text{THR_RAA}$) were 11 (5.5%). Among these, 4 were also characterized by abnormally low M5m_RAD. Thus, altogether, 36 (17.9%) RA+ cases were identified among the CAD patients. By contrast, only 3 (1.8%) cases at the verge of abnormal condition were identified among the CH subjects (Fig. 6).

Correlative analysis showed that, in both CAD and CH populations, RA parameters distributions were not significantly linked to the other clinical and ECG ($|\rho| \leq 0.44$; Table 2) parameters distributions. Moreover, CAD patients under pharmaceutical treatments showed RA levels comparable to those characterizing not-treated patients (Table 3).

	CAD patients (201)		CH subjects (167)	
	M5m_RAD (ms)	M5m_RAA (μ V)	M5m_RAD (ms)	M5m_RAA (μ V)
Clinical parameters:				
Age (years)	0.11	-0.01	0.20*	-0.28*
BMI (kg/m ²)	0.19*	-0.10	-0.09	-0.30*
SAP (mmHg)	0.11	-0.08	-0.06	-0.11
DAP (mmHg)	0.16*	-0.07	-0.11	-0.16*
LVEF	-0.17*	0.01	-	-
ECG parameters:				
mRR (ms)	-0.44*	0.29*	-0.42*	0.20*
sdRR (ms)	0.00	0.10	-0.35*	0.39*

Table 2. Correlation coefficient values linking RA parameters with the other clinical and ECG parameters. - : not applicable; *: statistically significant ($P < 0.05$).

Fig. 6. Representation of the RA cases for CH subjects (o) and the CAD patients (*) in relation to the RA normality region (internal solid-line square) in the M5m_RAD vs. M5m_RAA plane.

	Beta blocker		Diuretic		ACE inhibitor		Anti-arrhythmic	
	Treated (N=150)	Not Treated (N=51)	Treated (N=11)	Not Treated (N=190)	Treated (N=57)	Not Treated (N=144)	Treated (N=6)	Not Treated (N=195)
Clinical parameters:								
Age (years)	58±10	59±12	65±11	58±11	59±10	58±11	54±13	58±11
BMI (kg/m ²)	27±4	27±4	28±5	27±3	27±3	27±4	29±3	27±4
SAP (mmHg)	130±18	127±16	130±14	129±18	134±21	127±16	133±14	129±18
DAP (mmHg)	80±10	76±10	76±9	79±10	81±12	78±9	82±12	79±10
LVEF	58±11	58±10	49±12*	59±10	58±12	58±10	51±11	58±10
ECG parameters:								
mRR (ms)	979±156*	903±137	924±158	962±155	943±143	967±159	865±122	963±155
sdRR (ms)	37±41	35±35	71±72	35±36	42±39	35±40	30±31	37±40
M5m_RAD (ms)	-34±39	-33±32	-19±41	-34±37	-30±31	-35±39	-13±23	-34±37
M5m_RAA (µV)	19±7	19±11	20±7	19±9	19±9	19±8	21±8	19±9

Table 3. Comparison of clinical and ECG parameters between CAD patients under vs. not under pharmaceutical treatment. *: statistical significance (P<0.05) when comparing treated vs. not treated CAD patients groups.

4. Discussion

Our study represents the first attempt to quantify RA heterogeneity in the coronary artery disease (CAD) in terms of the traditional RA amplitude as it combines with a temporal parameter that allows RA localization along the JT segment as a delay from the T-wave apex. This characterization was accomplished by analyzing RA in 5-minute resting ECG recordings by means of our AMF-based technique, previously tested for T-wave alternans detection and quantification in both simulated and clinical settings (Burattini et al., 2006, Burattini et al., 2008, Burattini et al., 2009a, Burattini et al., 2009b, Burattini et al., 2010, Burattini et al., 2011).

Previous studies reported by others identified temporal RA localization in cases other than CAD, either by computing the time-interval from the R peak to the RA instant of occurrence (Martínez et al., 2006, Nearing et al., 1994, Selvaraj et al., 2007) or by identifying three windows of half JT duration (localized at the beginning, at the middle and at the end of repolarization, respectively) and, subsequently, determining which of these was more affected by RA (Narayan & Smith, 1999). The former approach has the major limitation of not taking into account the physiological variability of the RR interval and the JT duration, so that the same time interval localizing RA can be associated to different alternating portions of the JT segment. The latter approach tries to overcome this limitation by use of three windows half-overlapped. However, because of the large variability characterizing the ST segment in clinical cases, the middle window does not necessarily include the T-wave apex, as it is expected to do. Our AMF-based approach resolves these limitations by measuring the “delay” (RAD) of the alternans with respect to the T-wave apex, so that large negative values of RAD are necessarily due to RA occurring along the ST segment and the T-wave left-hand side. Instead, RAD values close to zero indicate the presence of RA mainly localized over the T wave, while positive values of RAD indicate alternation of the T-wave right-hand side. Since each portion of the JT segment pertains to a specific phase of the ventricular repolarization, our AMF-based method is expected to provide a more accurate identification of the ventricular repolarization phase involved in the alternation.

Besides RA heterogeneity characterization, our AMF-based method has the further peculiarity of relying on the RA continuity hypothesis. Other techniques, such as the spectral method, the RA identification technique most commonly used in clinics (Smith et al., 1988, Rosenbaum et al., 1994, Rosenbaum et al., 1996), and the Laplacian likelihood ratio method (Martínez & Olmos, 2005, Martínez et al., 2006) ascribe to noise the low-TWA levels, not associated with cardiac instability, which are even detected under physiological conditions. This methodological approach aroused from the early hypothesis that RA could be an on-off phenomenon, usually not present in health (Bloomfield et al., 2002). After the recent experimental study by Pruvot et al. (Pruvot et al., 2004) who demonstrated the possibility of inducing various levels of RA, some of which not necessarily associated to cardiac instability, the hypothesis that RA is a phenomenon characterized by a continuously changing amplitude and/or location from physiological to pathological condition has gained increasing consideration. This assumption requires the set up of methods, like our AMF-based one, that allow identification of an RA normality region to improve reliability in the discrimination of non-physiological or abnormal RA (RA+) levels potentially at risk of cardiac instability. In the present study the RA normality region was delimited by three thresholds (THR_RADmin, THR_RADmax, THR_RAA, respectively), computed from the

RA parameters distribution over the CH population (Burattini et al., 2008, Burattini et al., 2009a). More specifically, THR_RADmin and THR_RADmax were defined as the 0.5th and 99.5th percentiles of the M5m_RAD distribution, whereas one other threshold was defined as the 99.5th percentile over the THR_RAA distribution, being it a non-negative parameter. Use of percentiles, rather than mean and SD, makes the procedure independent of the assumption of normal distribution for parameter estimates. These threshold definitions strongly optimize specificity rather than sensitivity. The rationale for our choice is that, as mentioned above, RA was initially supposed not be present in healthy conditions (Bloomfield, 2002), so that the number of positive detections among healthy subjects was forced to be negligible (Burattini et al., 2008). Due to our recent studies highlighting the hypothesis that RA is a phenomenon characterized by an amplitude continuously changing from physiological to abnormal conditions (Burattini et al., 2009a, Burattini et al., 2010), identification of thresholds levels at 0.5th and 99.5th percentile may be too restrictive and, thus, not appropriate for an optimal identification of abnormal RA cases. Optimization of thresholds for RA+ identification in relation to risk stratification, however, is beyond the scope of this work, which is mainly focused on RA characterization in the coronary artery disease.

With this aim, our AMF-based technique was applied to ECG tracings from CAD patients, who, compared to CH subjects, are known to show increased levels of T-wave alternans (Bigger & Bloomfield, 2007, Bloomfield et al., 2006, Burattini et al., 2009a, Burattini et al., 2010, Ikeda et al., 2006, Zacks et al., 2007) and to be prone to experience major cardiac events related to it (Narayan et al., 2006). Our results indicate that, in agreement with what previously found (Burattini et al., 2008, Burattini et al., 2009a, Burattini et al., 2010), RA affects all subjects, either diseased or healthy, with different characteristics though. First of all, the CAD patients were, on average, characterized by higher RA amplitude than the CH subjects (CAD: 19 ± 9 μ V, CH: 17 ± 15 μ V; $P < 0.05$). The most interesting results, however, concern the RA localization, which was distributed all over the entire JT segment (ST segment, or T-wave left-hand side, or T-wave right-hand side) in the CAD patients, whereas it occurred mainly over the T-wave left-hand side, and occasionally in correspondence of the T-wave apex, among the CH subjects. Similarly to our CAD patients, RA was found to be distributed all along the JT segment also in the cardiomyopathy (Selvaraj et al., 2007). In spite of the fact that our choice of the threshold values delimiting the RA normality region tends to minimize the number of RA+ cases, the greater RA heterogeneity observed in our CAD patients compared with CH subjects allowed discrimination of a relevant percentage (17.9%) of RA+ cases in the coronary artery disease (Fig. 6). In these abnormal cases RA was characterized by high amplitude (5.5%), as previously reported in other studies on the CAD (Narayan et al., 2006), or occurred either early (11.0%), as in other diseased states (Martínez et al., 2006, Nearing et al., 1994), or late (3.5%) along the JT segment. According to Narayan & Smith (1999), RA occurring late is mainly associated to ventricular tachycardia. Thus, the characterization of RA in terms of both RA amplitude (generally used to discriminate RA+ cases in most studies using other techniques) and RA delay (a parameters which instead is a peculiarity of our AMF-based method) allowed a significant increment in the identification of RA+ cases among the CAD patients, which raised from 11 (identified when using only the RA amplitude parameter) to 36.

According to our results, increased RA heterogeneity observed in the CAD population was not associated to other clinical factors (age, BMI, SAP, DAP and LVEF) or pharmaceutical treatments (beta blocker, digoxin, diuretic, and ACE inhibitor), with the only exception of heart rate. Indeed, a significant, though low ($-0.44 \leq \rho \leq -0.42$; $P < 0.05$), correlation was found between M5m_RAD and RR in both populations, indicating that early RA tends to characterize patients with prolonged RR. Instead, a negligible correlation ($0.20 \leq \rho \leq 0.29$, $P < 0.05$) was found between M5m_RAA and RR distributions. This result is complementary and not in contradiction with the findings from several other studies, which have previously related RA amplitude to heart rate (Bloomfield et al., 2002, Narayan & Smith, 1999, Narayan & Smith, 2000, Smith et al., 1988). Such studies have indeed demonstrated that, within a subject, RA increases when heart rate is forced to increase (by pacing, exercise or drug) over 90 beats/min (Bloomfield et al., 2002). Identification of RA at rest, as in the present study, was often neglected because it requires sophisticated RA identification methods, such as our AMF-based one (Burattini et al., 2011), suitable to detect low levels of RA even in the presence of physiological levels of heart-rate variability (heart-rate variability is indeed strongly reduced at high heart rates).

5. Conclusion

Our quantitative investigation of RA heterogeneity in CAD patients yields the conclusion that, compared to the CH subjects, the former population shows higher RA amplitude and greater variability of RA localization along the JT segment. Especially, in the coronary artery disease, the RA occurs not only over the left-hand side of the T wave, as it is generally observed in health, but also along the ST segment (early repolarization) or along the T-wave right-hand side (late repolarization). Identification of patients characterized by RA occurring in early or late repolarization is relevant since in the literature these kinds of RA phenomena are associated to an increased risk of ventricular arrhythmias. Thus, our AMF-based method appears as a useful tool to identify RA characteristics that help enhancing pathophysiological interpretation and, with it, the ability to discriminate patients at increased risk of sudden cardiac death, who might be appropriately treated before they experience a major cardiac event.

6. References

- Adachi, K., Ohnishi, Y., Shima, T., Yamashiro, K., Takei, A., Tamura, N. & Yokoyama, M. (1999). Determinant of microvolt-level T-wave alternans in patients with dilated cardiomyopathy. *Journal of the American College of Cardiology*, Vol.34, No.2, (August 1999), pp. 374-380, ISSN 0735-1097.
- Banville, I. & Gray, R.A. (2002). Effect of action potential duration and conduction velocity restitution and their spatial dispersion on alternans and the stability of arrhythmias. *Journal of Cardiovascular Electrophysiology*, Vol. 13, No.11, (November 2002), pp. 1141-1149, ISSN 1045-3873.
- Bigger, J.T. & Bloomfield D.M. (2007). Microvolt T-wave alternans: An effective approach to risk stratification in ischemic cardiomyopathy? *Nature Clinical Practice Cardiovascular Medicine*, Vol. 4, No.6, (June 2007), pp. 300-301, ISSN 1743-4297.

- Bloomfield, D.M., Hohnloser, S.H. & Cohen, R.J. (2002). Interpretation and classification of microvolt T wave alternans tests. *Journal of Cardiovascular Electrophysiology*, Vol.13, No.5 (May 2002), pp.502-512, ISSN 1045-3873.
- Bloomfield, D.M., Bigger, J.T., Steinman, R.C., Namerow, P.B., Parides, M.K., Curtis, A.B., Kaufman, E.S., Davidenko, J.M., Shinn, T.S. & Fontaine, J.M. (2006). Microvolt T-wave alternans and the risk of death or sustained ventricular arrhythmias in patients with left ventricular dysfunction. *Journal of the American College of Cardiology*, Vol.47, No.2, (January 2006), pp. 456-463, ISSN 0735-1097.
- Burattini, L., Zareba, W. & Moss, A.J. (1999). Correlation method for detection of transient T-wave alternans in digital Holter ECG recordings. *Annals of Noninvasive Electrocardiology*, Vol. 4, No.4, (October, 1999), pp. 416-424, ISSN 1082-720X.
- Burattini, L., Zareba, W. & Burattini, R. (2006). Automatic detection of microvolt T-wave alternans in Holter recordings: Effect of baseline wandering. *Biomedical Signal Processing and Control*, Vol.1, (2006), pp.162-168, ISSN 1746-8094.
- Burattini, L., Zareba, W. & Burattini, R. (2008). Adaptive match filter based method for time vs. amplitude characterization of microvolt ECG T-wave alternans. *Annals of Biomedical Engineering*, Vol.36, No.9, (September 2008), pp. 1558-1564, ISSN 0090-6964.
- Burattini, L., Zareba, W. & Burattini, R. (2009a). Assessment of physiological amplitude, duration and magnitude of ECG T-wave alternans. *Annals of Noninvasive Electrocardiology*, Vol.14, No.4, (October 2009), pp. 366-374, ISSN 1082-720X.
- Burattini, L., Bini, S. & Burattini, R. (2009b). Comparative analysis of methods for automatic detection and quantification of microvolt T-wave alternans. *Medical Engineering & Physics*, Vol. 31, No.10, (December 2009), pp. 1290-1298, ISSN 1350-4533.
- Burattini, L., Zareba, W. & Burattini, R. (2010). Identification of gender-related normality regions for T-wave alternans. *Annals of Noninvasive Electrocardiology*, Vol.15, No.4, (October 2010), pp. 328-336, ISSN 1082-720X.
- Burattini, L., Bini, S. & Burattini, R. (2011). Automatic microvolt T-wave alternans identification in relation to ECG interferences surviving preprocessing. *Medical Engineering & Physics*, Vol.33, No.1, (January 2011), pp. 17-30, ISSN 1350-4533.
- Chinushi, M., Restivo, M., Caref, E.B. & El-Sherif, N. (1998). Electrophysiological basis of arrhythmogenicity of QT/T alternans in the long-QT syndrome: tridimensional analysis of the kinetics of cardiac repolarization. *Circulation Research*, Vol.83, No.6, (September 1998), pp. 614-628, ISSN 0009-7330.
- Chow, T., Kereiakes, D.J., Bartone, C., Booth, T., Schloss, E.J., Waller, T., Chung, E.S., Menon, S., Nallamothu, B.K. & Chan, P.S. (2006). Prognostic utility of microvolt T-wave alternans in risk stratification of patients with ischemic cardiomyopathy. *Journal of the American College of Cardiology*, Vol.47, No.9, (May 2006), pp. 1820-1827, ISSN 0735-1097.
- Chow, T., Kereiakes, D.J., Bartone, C., Booth, T., Schloss, E.J., Waller, T., Chung, E., Menon, S., Nallamothu, B.K. & Chan, P.S. (2007). Microvolt T-wave alternans identifies patients with ischemic cardiomyopathy who benefit from implantable cardioverter-defibrillator therapy. *Journal of the American College of Cardiology*, Vol.49, No.1, (January 2007), pp.50-58, ISSN 0735-1097.

- Habbab, M.A. & El-Sherif N. (1992). TU alternans, long QTU, and torsade de pointes: clinical and experimental observations. *Pacing and Clinical Electrophysiology*, Vol.15, No.6, (June 1992), pp. 916-931, ISSN 0147-8389.
- Hering H. (1909). Experimentelle studien an saugentieren uber das electrocardiogram. *Zchr Exper Path Therapie*, Vol. 7. p. 363.
- Ikeda, T., Yoshino, H., Sugi, K., Tanno, K., Shimizu, H., Watanabe, J., Kasamaki, Y., Yoshida, A. & Kato, T. (2006). Predictive value of microvolt T-wave alternans for sudden cardiac death in patients with preserved cardiac function after acute myocardial infarction: results of a collaborative cohort study. *Journal of the American College of Cardiology*, Vol.48, No.11, (December 2006), pp. 2268-2274, ISSN 0735-1097.
- Kleinfeld, M.J. & Rozanski J.J. (1977). Alternans of the ST segment in Prinzmetal's angina. *Circulation*, Vol.55, No.4, (April 1977), pp. 574-577, ISSN 0009-7322.
- Klingenhoben, T., Zabel, M., D'Agostino, R.B., Cohen, R.J. & Hohnloser, S.H. (2000). Predictive value of T-wave alternans for arrhythmic events in patients with congestive heart failure. *Lancet*, Vol.356, No. 9230, (August 2000), pp. 651-652, ISSN 0140-6736.
- Leino, J., Minkkinen, M., Nieminen, T., Lehtimäki, T., Viik, J., Lehtinen, R., Nikus, K., Kööbi, T., Turjanmaa, V., Verrier, R.L. & Kähönen, M. (2009). Combined assessment of heart rate recovery and T-wave alternans during routine exercise testing improves prediction of total and cardiovascular mortality: the Finnish Cardiovascular Study. *Heart Rhythm*, Vol.6, No.12, (December 2009), pp.1765-1771, ISSN 1547-5271.
- Maeda, S., Nishizaki, M., Yamawake, N., Ashikaga, T., Shimada, H., Asano, M., Ihara, K., Murai, T., Suzuki, H., Fujii, H., Sakurada, H., Hiraoka, M. & Isobe, M. (2009). Ambulatory ECG-based T-wave alternans and heart rate turbulence predict high risk of arrhythmic events in patients with old myocardial infarction. *Circulation Journal*, Vol.73, No.3, (December 2009), pp. 2223-2228, ISSN 1346-9843.
- Martínez, J.P. & Olmos, S. (2005). Methodological principles of T wave alternans analysis: a unified framework. *IEEE Transactions on Bio-Medical Engineering*, Vol.52, No.4, (April 2005), pp.599-613, ISSN 0018-9294.
- Martínez, J.P., Olmos, S., Wagner, G. & Laguna, P. (2006). Characterization of repolarization alternans during ischemia: time-course and spatial analysis. *IEEE Transactions on Bio-Medical Engineering*, Vol.53, No.4, (April 2006), pp. 701-711, ISSN 0018-9294.
- Montagnana, M., Lippi, G., Franchini, M., Targher, G. & Cesare Guidi, G. (2008). Sudden cardiac death: prevalence, pathogenesis, and prevention. *Annals of Medicine*, Vol.40, No.5, (2008), pp. 360-375, ISSN 0785-3890.
- Narayan, S.M. & Smith, J.M. (1999). Differing rate dependence and temporal distribution of repolarization alternans in patients with and without ventricular tachycardia. *Journal of Cardiovascular Electrophysiology*, Vol.10, No.1, (January 1999), pp.61-71, ISSN 1045-3873.
- Narayan, S.M. & Smith, J.M. (2000). Exploiting rate-related hysteresis in repolarization alternans to improve risk stratification for ventricular tachycardia. *Journal of the American College of Cardiology*, Vol.35, No.6, (May 2000), pp. 1485-1492, ISSN 0735-1097.

- Narayan, S.M. (2006). T-wave alternans and the susceptibility to ventricular arrhythmias. *Journal of the American College of Cardiology*, Vol.47, No.2, (2006), pp. 269-281, ISSN 0735-1097.
- Narayan, S.M., Smith, J.M., Lindsay, B.D., Cain, M.E. & Dávila-Román, V.G. (2006). Relation of T-wave alternans to regional left ventricular dysfunction and eccentric hypertrophy secondary to coronary heart disease. *The American Journal of Cardiology*, Vol.97, No.6, (March 2006), pp. 775-780, ISSN 0002-9149.
- Narayan, S.M., Franz, M.R., Lalani, G., Kim, J. & Sastry, A. (2007). T-wave alternans, restitution of human action potential duration, and outcome. *Journal of the American College of Cardiology*, Vol.50, No.25, (December 2007), pp. 2385-2392, ISSN 0735-1097.
- Nearing, B.D., Huang, A.H. & Verrier, R.L. (1991). Dynamic tracking of cardiac vulnerability by complex demodulation of the T wave. *Science*, Vol.252, No.5004, (April 1991), pp.437-440, ISSN 0036-8075.
- Nearing, B.D. & Verrier, R.L. (1993). Personal computer system for tracking cardiac vulnerability by complex demodulation of the T wave. *Journal of Applied Physiology*, Vol.74, No.5, (May,1993), pp. 2606-2612, ISSN 8750-7587.
- Nearing, B.D., Oesterle, S.N. & Verrier, R.L. (1994). Quantification of ischaemia induced vulnerability by precordial T wave alternans analysis in dog and human. *Cardiovascular Research*, Vol.28, No.9, (September 1994), pp. 1440-1449, ISSN 0008-6363.
- Nearing, B.D. & Verrier, R.L. (2002). Modified moving average analysis of T-wave alternans to predict ventricular fibrillation with high accuracy. *Journal of Applied Physiology*, Vol.92, No.2, (February 2002), pp.541-549, ISSN 8750-7587.
- Pastore, J.M., Girouard, S.D., Laurita, K.R., Akar, F.G. & Rosenbaum, D.S. (1999). Mechanism linking T-wave alternans to the genesis of cardiac fibrillation. *Circulation*, Vol.99, No.10, (March 1999), pp.1385-94, ISSN 0009-7322.
- Pruvot, E.J., Katra, R.P., Rosenbaum, D.S. & Laurita, K.R. (2004). Role of calcium cycling versus restitution in the mechanism of repolarization alternans. *Circulation Research*, Vol.94, No. 8, (April 2004), pp. 1083-1090, ISSN 0009-7330.
- Reddy, C.V., Kiok, J.P., Khan, R.G. & El-Sherif, N. (1984). Repolarization alternans associated with alcoholism and hypomagnesemia. *The American Journal of Cardiology*, Vol.53, No.2, (January 1984), pp. 390-391, ISSN 0002-9149.
- Rosenbaum, D.S., Jackson, L.E., Smith, J.M., Garan, H., Ruskin, J.N. & Cohen, R.J. (1994). Electrical alternans and vulnerability to ventricular arrhythmias. *The New England Journal of Medicine*, Vol.330, No. 4, (January 1994), pp. 235-241, ISSN 0028-4793.
- Rosenbaum, D.S., Albrecht, P. & Cohen, R.J. (1996). Predicting sudden cardiac death from T wave alternans of the surface electrocardiogram: promises and pitfalls, *Journal of Cardiovascular Electrophysiology*, Vol.7, No.11, (November 1996), pp. 1095-1111, ISSN 1045-3873.
- Sakaki, K., Ikeda, T., Miwa, Y., Miyakoshi, M., Abe, A., Tsukada, T., Ishiguro, H., Mera, H., Yusu, S. & Yoshino, H. (2009). Time-domain T-wave alternans measured from Holter electrocardiograms predicts cardiac mortality in patients with left ventricular dysfunction: a prospective study. *Heart Rhythm*, Vol.6, No.3, (March 2009), pp. 332-337, ISSN 1547-5271.

- Salerno-Uriarte, J.A., De Ferrari, G.M., Klersy, C., Pedretti, R.F., Tritto, M., Sallusti, L., Libero, L., Pettinati, G., Molon, G., Curnis, A., Occhetta, E., Morandi, F., Ferrero, P., Accardi, F. & ALPHA Study Group Investigators (2007). Prognostic value of T-wave alternans in patients with heart failure due to nonischemic cardiomyopathy: results of the ALPHA Study. *Journal of the American College of Cardiology*, Vol.50, No. 19, (November 2007), pp. 1896-1904, ISSN 0735-1097.
- Schwartz, P.J. & Malliani, A. (1975). Electrical alternation of the T-wave: clinical and experimental evidence of its relationship with the sympathetic nervous system and with the long Q-T syndrome. *American Heart Journal*, Vol.89, No.1, (January 1975), pp.45-50, ISSN 0002-8703.
- Selvaraj, R.J., Picton, P., Nanthakumar, K., Mak, S. & Chauhan, V.S. (2007). Endocardial and epicardial repolarization alternans in human cardiomyopathy: evidence for spatiotemporal heterogeneity and correlation with body surface T-wave alternans. *Journal of the American College of Cardiology*, Vol.43, No.3 (January 2007), pp. 338-346, ISSN 0735-1097.
- Smith, J.M. & Cohen, R.J. (1984). Simple finite-element model accounts for wide range of cardiac dysrhythmias. *Proceedings of the National Academy of Sciences of the United States of America*, Vol. 81, No.1, (January 1984), pp. 233-237.
- Smith, J.M., Clancy, E.A., Valeri, C.R., Ruskin, J.N. & Cohen, R.J. (1988). Electrical alternans and cardiac electrical instability. *Circulation*. Vol.77, No.1,(January 1988), pp. 110-121, ISSN 0009-7322.
- Stein, P.K., Sanghavi, D., Domitrovich, P.P., Mackey, R.A. & Deedwania, P. (2008). Ambulatory ECG-based T-wave alternans predicts sudden cardiac death in high-risk post-MI patients with left ventricular dysfunction in the EPHEsus study. *Journal of Cardiovascular Electrophysiology*, Vol.19, No.10, (October 2008), pp. 1037-1042, ISSN 1045-3873.
- Verrier, R.L., Nearing, B.D., La Rovere, M.T., Pinna, G.D., Mittleman, M.A., Bigger, J.T. Jr, Schwartz, P.J. & ATRAMI Investigators. (2003). Ambulatory electrocardiogram-based tracking of T wave alternans in postmyocardial infarction patients to assess risk of cardiac arrest or arrhythmic death. *Journal of Cardiovascular Electrophysiology*, Vol.14, No.7, (July 2003), pp. 705-711, ISSN 1045-3873.
- Weiss, J.N., Garfinkel, A., Karagueuzian, H.S., Qu, Z. & Chen, P.S. (1999). Chaos and the transition to ventricular fibrillation: a new approach to antiarrhythmic drug evaluation. *Circulation*, Vol.99, No.21, (June 1999), pp. 2819-2826, ISSN 0009-7322.
- Zacks, E.S., Morin, D.P., Ageno, S. Janik, M., Mauer, A.C., Markowitz, S.M., Mittal, S., Iwai, S., Shah, B.K., Lerman, B.B. & Stein, K.M. (2007). Effect of oral betablocker therapy on microvolt T-wave alternans and electrophysiology testing in patients with ischemic cardiomyopathy. *American Heart Journal*, Vol.153, No.3, (March 2007), pp. 392-397, ISSN 0002-8703.
- Zareba, W., Moss, A.J., le Cessie, S. & Hall, W.J. (1994). T wave alternans in idiopathic long QT syndrome. *Journal of the American College of Cardiology*, Vol.23, No.7,(June 1994), pp. 1541-1546, ISSN 0735-1097.

Zheng, Z.J., Croft, J.B., Giles, W.H. & Mensah, G.A. (2001). Sudden cardiac death in the United States, 1989 to 1998. *Circulation*, Vol.104, No.18, (October 2001), pp. 2158-2163, ISSN 0009-7322.

IntechOpen

IntechOpen

Coronary Artery Diseases

Edited by Dr. Illya Chaikovsky

ISBN 978-953-51-0238-0

Hard cover, 332 pages

Publisher InTech

Published online 07, March, 2012

Published in print edition March, 2012

This book has "wide geography" both literally and figuratively. First of all, this book brings together contributions from around the world, both from post-industrial countries and developing world. This is natural, because coronary artery disease is becoming pandemic worldwide. CAD is the single most frequent cause of death in developed countries, causes about 1 in every 5 deaths. Mortality from cardiovascular disease is predicted to reach 23.4 million in 2030. Moreover, in the developing world, cardiovascular disease tends to affect people at a younger age and thus could negatively affect the workforce and economic productivity. The morbidity, mortality, and socioeconomic importance of CAD make its diagnosis and management fundamental for all practicing physicians. On another hand, the book widely represents "geography" of CAD itself, i.e. many various aspects of its pathophysiology, epidemiology, diagnosis, treatment are touched in this book. This book does not pretend on complete and integral description of the Coronary artery disease. Rather, it contains selected issues on this complex multifactorial disease. Nevertheless, we hope that readers will find Coronary Artery Disease useful for clinical practice and further research.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Laura Burattini and Roberto Burattini (2012). Characterization of Repolarization Alternans in the Coronary Artery Disease, *Coronary Artery Diseases*, Dr. Illya Chaikovsky (Ed.), ISBN: 978-953-51-0238-0, InTech, Available from: <http://www.intechopen.com/books/coronary-artery-diseases/identification-of-abnormal-repolarization-alternans-in-coronary-artery-disease>

INTech
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](https://creativecommons.org/licenses/by/3.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen