We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Chemical Composition, Antioxidant and Antimicrobial Activities of Essential Oil of *Warionia saharae* from Oases of Morocco

K. Sellam¹, M. Ramchoun², C. Alem², Farid Khallouki²
B. El Moualij² and L. El Rhaffari^{1*}

¹Environment and Health Research Team (EHRT),
Faculty of the Sciences and Technology, Er-Rachidia Morocco,

²Research Team in Natural Product Biochemistry,
Faculty of the Sciences and Technology, Er-Rachidia Morocco,
Morocco

1. Introduction

The *Warionia saharae* which belongs to the important composite's family is an endemic species of North Africa, characterized by a discerning odour (Bonnet and Maurry, 1889). *Warionia saharae* was reported for the first time in the Oranais Sahara (Beni ounif in Algeria) by Dr. Warrion as a shrub of 1 to 3 m of height. The thick trunk, is covered of a gray peel, structural of very wavy terminal leaf bouquets, and of capitulate of yellow flowers (Photos 1 and 2). The picking of stems leafed of this bush, clear a very heady and spicy odour; the latex that flows out of injuries of the peel, glue to hands in a very tenacious way (Lebrun, 1979).

In Morocco, *Warionia saharae* is growing wild in various regions (Benabid and Fennane, 1994). The habitat is between schistose rocks (Watillon et al., 1988). *Warionia saharae* is known in Morocco by the Berber vernacular names of "âfessas" and "Tazart n-îfiss". In the Moroccan traditional medicine, the leaves of the plant are used to treat inflammatory diseases, such as rheumatoid arthritis, and for gastrointestinal disorders (Bellakhdar, 1997), inflammation of the womb, colds, Jaundice and cardiac pains (El Rhaffari et Zaid, 2002).

This work aims to study and characterise a new bioactive natural products from medicinal plants of Moroccan oases. Information concerning in vitro antioxidant, antimicrobial activities of the essential oil from the *Warionia saharae* has not been reported earlier.

2. Materials and methods

2.1 Plant material

The aerial part of *Warionia saharae* was collected in Er-rachidia (Morocco), during the flowering period (April/June, 2009). A duplicate specimen is held at the FST Er-rachidia. The dried plant material is stored in the laboratory at room temperature (25°C) and in the shade before the extraction.

Photo 1. *Warionia saharae* (Plant) (Taken y El Rhaffari L.)

Photo 2. *Warionia saharae* (Flower) (Taken y El Rhaffari L.)

2.2 Steam distillation apparatus and procedure

The extraction of essential oil of the aerial part of *Warionia saharae* was conducted by steam distillation in a Clevenger apparatus. The essential oil obtained was dried under anhydrous sodium sulfate and stored at 4°C in the dark before analysis.

2.3 Essential oil analysis

Analyses of volatile compounds were run on a Trace GC Ultra gas chromatography equipped with a VB-5 capillary column (Methylpolysiloxane with 5% phenyl; 25 m x 0.2 mm; film thickness 0.2µm) was directly coupled to the mass spectrometer (Polaris Q. MS). Helium (1 mL/min) was used as carrier gas. The program was 2 min isothermal at 40 °C, then the temperature increased by 4 °C/min to 180 °C and 2 min isothermal at 180 °C. The injection port temperature was 250 °C and that of the detector was 280 °C. Ionization of sample components was performed in the EI mode (70 eV).

Kovat's retention indices were calculated using cochromatographed standards hydrocarbons. The individual compounds were identified by MS and their identity was confirmed by comparing their retention indices relatives to C8–C32 n-alkanes and by comparing their mass spectra and retention times with those of authentic samples or with data already available in the NIST library and literature (Adams, 1989 et 2001).

2.4 Antimicrobial activity

The antimicrobial activity was evaluated by paper disc diffusion and dilution methods against three selected Gram-positive and Gram-negative species: *Staphylococcus aureus* ATCC 29213, *Escherichia coli* ATCC 35218, *Pseudomonas aeruginosa* ATCC 27853, and against the phytopathogenic fungi *Penicillium* sp. and *Candida albicans*.

The qualitative antimicrobial essay of the volatile fraction of *W. saharae* was carried out by the disc diffusion method (Anonymous, 1995).

The minimal inhibitory concentration (MIC) and minimal bactericidal concentration (MBC) of tested volatile fractions were determined using the Mueller Hinton broth (MHB) dilution method (Anonymous, 1995).

All samples were tested in triplicate. The MIC was defined as the lowest concentration preventing visible growth (May et al., 2000; Burt, 2004).

2.5 Antioxidant activity

The antioxidant activity was assessed by 1,1-diphenyl-2-picrylhydrazyl (DPPH), ferric reducing ability (FRAP) and β -carotene bleaching method systems. Data collected for each assay was an average of three experiments.

The method is based on the radical scavenging activity of the antioxidant (Mensor et al., 2001; Tepe et al., 2004).

The FRAP assay was performed as described by Benzie and Nilsson (Benzie and Strain, 1996; Nilsson et al., 2005).

The β -carotene method was carried out according to Shahidi technique (Shahidi et al., 2001).

2.6 Statistical analysis

For antioxidant assays the results were expressed as value±error using statistical analysis formulas referring to the value of $p < 0.05$ of confidence interval for the Student’s t-test law.

3. Results and discussion

3.1 Chemical composition of the essential oil

The essential oil was extracted by the hydrodistillation of the dried leaves of *Warionia saharae* from Errachidia region (Morocco), were analyzed by GC-MS.

The essential oil yields, calculated on a dry weight basis, was about 1.1%(w/w). The identified combinations in essential oil, retention time (RT) and quantitative percentage of the compounds are presented in table 1. A total of 32 compounds, amounting 92.5% of the oils, were identified.

Peak number	RT ^b	Compound ^a	Composition (%)
1	6.63	α-Thujene	0.9
2	6.75	α-Pinene	0.6
3	10.19	Camphene	0.99
4	10.68	Sabinene	2.22
5	11.79	α-Phellandrene	1.50
6	12.24	α-Carene	0.60
7	12.54	p-Cymene	3.77
8	12.77	1,8-Cineole	6.12
9	13.80	3-Carene	0.77
10	14.88	α-Terpinene	0.41
11	15.49	Linalool	16.79
12	18.12	Terpinen-4-ol	3.40
13	18.60	α-Fenchol	2.59
14	18.86	Ethyl 8-fluorooctanoate	0.55
15	20.84	Geraniol	1.14
16	22.56	Carvacrol	1.34
17	23.92	Ethyl 3-phenylpropionate	0.85
18	23.99	Ocimenyl acetate	0.81
19	28.51	Tridecan-1-ol	0.55
20	28.73	Longipinane	2.05
21	29.45	Allo-Aromadendrene	0.44
22	30.20	α-Farnesene	0.51
23	30.75	Trans nerolidol	17.95
24	31.19	Caryophyllene oxide	1.91
25	33.24	β-Eudesmol	23.74
Total			92.5

^a Compounds listed in order of elution from a DB-5 column.

^b Retention time (as minutes).

Table 1. Composition of essential oil of *Warionia saharae* (%).

The major components of *Warionia saharae* essential oils were β -Eudesmol (23.74%), Trans-Nerolidol (17.95%), Linalool (16.79%), 1,8 cineole (6.12%).

The high content of oxygenated identified compounds might explain the characteristic and fragrant odor of the oil (Figure 1 and Table 1).

Fig. 1. Chromatogram *Warionia saharae* volatile fractions run on VB-5 capillary column.

3.2 Antimicrobial activity

Antimicrobial activities of *Warionia saharae* essential oil were evaluated by a paper disc diffusion method against bacterial strains including Gram+ and Gram-, and fungal strains (Table 2).

microorganism	source	Inhibition (mm) ^a					MIC (μ g/ml)		
		Essential oil (EO)			STA ^b (μ g/disk)		EO	Amp	Nys
		5 μ g /disc	10 μ g /disc	15 μ g /disc	A10	N10			
E. coli	ATCC25922	12	13	16	16	Nd	20	25	Nd
P. aeruginosa	ATCC27853	16	20	22	18	Nd	10	5	Nd
S. aureus	ATCC25923	10	14	15	14	Nd	5	10	Nd
Candida albicans	BTCE	18	18	20	Nd	22	2.5	Nd	5
Penicillium sp.	BTCE	14	16	20	Nd	18	5	Nd	10

^a Diameter of the zone of inhibition (mm) including disk diameter of 6mm – average of three experiments.

^b standards antibiotic; A10 : Ampicillin (Amp), N10 : Nystatin (Nys).

MIC, minimum inhibitory concentration; values given as mg/ml for the essential oils and as μ g/ml for antibiotics.

BTCE, Blood Transfusion Center Errachidia (Morocco).

Nd, not determined.

Table 2. Antimicrobial activity of *Warionia saharae* essential oil.

The comparison between in vitro bacteriostatic activity of essential oil of *Warionia saharae*, and the inhibition zone formed by standard antibiotic disc (Ampicillin (A10) and Nystatin (N10)) showed that the essential oils were more active against the microorganisms tested. These results suggest that the volatile oil of *Warionia saharae* would probably be a good therapeutic agent against these bacteria and fungus.

The bacteriostatic properties of the oil are suspected to be associated with the high Eudesmol and Nerolidol content, which has been tested previously and was found to have a significant antibiotic activity (Brehm-Stecher and Johnson, 2003; Lattaoui and Tantaoui-Elaraki, 1994).

3.3 Antioxidant activity

The study of the antioxidant activities showed that the essential oil have a higher antioxidant capacity relative to the antioxidant of reference BHT and BHA ($94.2 \pm 0.4\%$).

In contrast to β -carotene/linoleic acid system, essential oil showed a moderate activity in this system ($65.2 \pm 0.4\%$).

The ferric reducing ability of plasma (FRAP) assay was used for assessing “antioxidant power” of *Warionia saharae* essential oil. The antioxidant power of the leaves oil samples were compared with BHA and BHT as reference antioxidants.

The results reported here can be considered as the first information on the antimicrobial and antioxidant properties of *Warionia saharae*. Further studies are needed to evaluate the in vivo potential of these oils in animal models.

4. Conclusion

This work interested *Warionia saharae* from Er-rachidia region (south eastern Morocco). It showed that the essential oil yield is about 1.1% (w/w) in dried leaves.

The essential oils is analysed by GC-MS. Thirty two compounds were identified, they are amounting 92.5% of the oils. The major components of essential oils were β -Eudesmol (23.74%), Trans-Nerolidol (17.95%), Linalool (16.79%), 1,8 cineole (6.12%), p-Cymene (3.77%) and terpinen 4-ol (3.40%).

Warionia oil exhibited a significant antimicrobial activity against *Staphylococcus aureus*, *Candida albicans* and *Bacillus cereus*. In each case the activity of essential oil was higher than those of the standard antibiotic.

The essential oil showed a relatively high radical scavenging ability and antioxidant activity determined by 1,1-diphenyl-1-picrylhydrazyl (DPPH) assay, ferric reducing (FRAP) assay and β -carotene bleaching test.

The GC and GC-MS analysis was useful in order to identify the components partially involved in antimicrobial and antioxidant activities of the essential oils obtained from *Warionia saharae* leaves.

Therefore, *Warionia saharae* essential oil could be a source of pharmaceutical materials required for the preparation of new therapeutic and antimicrobial agents. This is the object of our future investigations.

5. References

- Adams, R.P., 1989. Identification of Essential Oil by Ion Trap Mass Spectroscopy. Academic Press, San Diego.
- Adams, R.P., 2001. Identification of Essential Oil Components by Gas Chromatography/QUADrupole Mass Spectroscopy. Allured, Illinois.
- Anonymous: National Committee for Clinical Laboratory, 1995. Standards methods for dilution antimicrobial susceptibility tests for bacteria that grow aerobically. Wayne Pa., vol. 15, p. 15 (approved standards no. M7-A3).
- Bellakhdar, J. (1997). La pharmacopée marocaine traditionnelle, Médecine arabe ancienne et savoirs populaires, Ibis press, p. 208.
- Benabid, A. and Fennane, M. (1994). Connaissance sur la végétation du Maroc: phytogéographie, phytosociologie et séries de végétations. Lazaroa 14: 21-97.
- Benzie, I.F., Strain, J.J., 1996. The ferric reducing ability of plasma (FRAP) as a measure of "Antioxidant Power": The FRAP Assay. Anal. Biochem. 239, 70-76.
- Bonnet and Maurry, P. (1889). Etude sur le *Warionia saharae*. Benth & coss. Assoc. Fr. Avanc. Sc. Congrès de Paris.
- Brehm-Stecher B F and Johnson E A. (2003). Sensitization of *Staphylococcus aureus* and *Escherichia coli* to antibiotics by the sesquiterpenoids nerolidol, farnesol, bisabolol and apritone. Antimicrobial Agents Chemotherapy 47(10): 3357-3360.
- Burt S. (2004). Essential oils: their antibacterial properties and potential applications in foods-a review. Int J Food Microbiol. 94:223-53.
- El Rhaffari L. et Zaid A. 2002 Pratique de la phytothérapie dans le sud-est du Maroc (Tafilalet). Un savoir empirique pour une pharmacopée rénovée. Actes du 4^{ème} congrès Européen d'Ethnopharmacologie: origine des pharmacopées traditionnelles et élaboration des pharmacopées savantes, Metz, 11-13 mai 2000. Publiés par le CRD Montpellier, 2002, pp 295-304.
- Hicheri F, Ben Jannet H, Cheriaa J, Jegham S, & Mighri Z. (2003) Antibacterial activities of a few prepared derivatives of oleanolic acid and of other natural triterpenic compounds. C R Chim .6:473-83.
- Lattaoui N, Tantaoui-Elaraki A. (1994). Individual and combined antibacterial activity of the main components of three thyme essential oils. Revista Italiana Eppos 13: 13-19.
- Lebrun, J.P. (1979). Eléments pour un Atlas des plantes vasculaires de l'Afrique sèche, 2: 11-12. J.E.M.V.P.T., Bot 6.
- May J, Chan CH, King A, Williams L, & French GL. (200). Time-kill studies of tea tree oils on clinical isolates. J Antimicrob Chemother. 45:639-43.
- Mensor, L.L., Menezes, F.S., Leitão, G.G., Reis, A.S., Dos Santos, T.C., Coube, C.S., & Leitao, S.G., 2001. Screening of Brazilian plant extracts for antioxidant activity by the use of DPPH free radical method. Phytotherapy Res. 15, 127-130.
- Nilsson, J., Pillai, D., Önning, G., Persson, C., Nilsson, A., & Akesson, B., 2005. Comparison of the 2,2'-azinobis-3-ethylbenzothiazoline-6-sulphonic acid (ABTS) and ferric reducing antioxidant power (FRAP) methods to assess the total antioxidant capacity in extracts of fruit and vegetables. Mol. Nutrit. Food Res. 49, 239-246.

- Shahidi F, Chavan UD, Naczki M & Amarowicz R (2001). Nutrient distribution and phenolic antioxidants in air-classified fractions of beach pea (*Lathyrus maritimus* L.). J. Agric. Food Chem. 49: 926-933.
- Tepe, B., Daferera, D., Sokmen, M., Polissiou, M., & Sokmen, A., 2004. In vitro antimicrobial and antioxidant activities of the essential oils and various extracts of *Thymus eigi* Zohary. J. Agric. Food Chem. 52, 1132-1137.
- Watillon, C., Gaspar, T., Hofinger, M. & Ramaut, J.L. (1988). La micropropagation de *Warionia saharae*. Benth & Coss. Al Biruniya. 4 (1): 35-38.

Gas Chromatography - Biochemicals, Narcotics and Essential Oils

Edited by Dr. Bekir Salih

ISBN 978-953-51-0295-3

Hard cover, 236 pages

Publisher InTech

Published online 09, March, 2012

Published in print edition March, 2012

Gas Chromatography involves the study of various vaporizable molecules in chemistry and the other related research fields. This analytical method has a number of features and advantages that make it an extremely valuable tool for the identification, quantification and structural elucidation of organic molecules. This book provides detailed gas chromatography information to applications of biochemicals, narcotics and essential oils. The details of the applications were briefly handled by the authors to increase their comprehensibility and feasibility. This guide should be certainly valuable to the novice, as well as to the experienced gas chromatography user who may not have the enough experience about the specific applications covered in this book. We believe this book will prove useful in most laboratories where modern gas chromatography is practiced.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

K. Sellam, M. Ramchoun, C. Alem, Farid Khallouki, B. El Moualij and L. El Rhaffari (2012). Chemical Composition, Antioxidant and Antimicrobial Activities of Essential Oil of Warionia saharae from Oases of Morocco, Gas Chromatography - Biochemicals, Narcotics and Essential Oils, Dr. Bekir Salih (Ed.), ISBN: 978-953-51-0295-3, InTech, Available from: <http://www.intechopen.com/books/gas-chromatography-biochemicals-narcotics-and-essential-oils/chemical-composition-antioxidant-and-antimicrobial-activities-of-essential-oil-of-warionia-saharae-f>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](https://creativecommons.org/licenses/by/3.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen