We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Lipoprotein (a) and Cardiovascular Risk

José Antonio Díaz Peromingo

*Short Stay Medical Unit, Department of Internal Medicine,
Hospital Clínico Universitario, Santiago de Compostela,
Spain*

1. Introduction

First epidemiological studies of Lp(a) and CHD were reported at the end of the last century (1-3) but the investigation of this lipoprotein as a potential cardiovascular risk factor has been hampered by the lack of consistent approaches to its measurement for decades. Lp(a) laboratory standardization emerged in 2000 (4) and was accepted by the World Health Organization in 2004 (5). Another challenge associated to its measurement is the fact that population differences can also contribute to variation in Lp(a) serum concentration (6). Since Lp(a) characterization, evidences favoring its association with cardiovascular risk have been reported. At the same time, studies against this association have also been published leading to some confusion regarding to the possible role of Lp(a) in cardiovascular disease. The last years have clarified somewhat this issue and evidences of Lp(a) as an independent cardiovascular risk factor have been proposed (7-13). Several key points such as its homology with plasminogen, differences among the apo(a) isoforms, genetic considerations as well as special circumstances such as the relationship of Lp(a) and atrial fibrillation, dialysis, alcohol consumption and blood coagulation have been investigated. In this chapter, Lp(a) metabolism, epidemiological and genetic considerations, association with coronary heart disease and stroke, special situations as well as controversies and current treatment options are related.

2. Lipoprotein (a) metabolism

Lipoprotein (a), Lp (a), is a low density lipoprotein (LDL)-like particle synthesized in the liver by hepatocytes and then secreted into plasma. It was first described by Berg in 1963 (14). It consists of an apolipoprotein B100 (apoB100) molecule that is linked covalently by a disulfide bond to a large glycoprotein known as apolipoprotein (a), [apo(a)] (15). Lp(a) metabolic route is shown in figure 1. Its molecular weight ranges from 200 kDa to more than 800 kDa (16). The apo(a) gene (*LPA*) is a major determinant of the plasma concentration of Lp(a), including variations in the kringle region-coding repeats, with accounts for the size polymorphism of apo(a) leading to different apo(a) sizes (17). This fact is very important because small size isoforms seem to be associated to worse cardiovascular profile. Apo(a) chain contains 5 cysteine-rich domains known as kringles, and especially Kringle IV (KIV) is very similar to plasminogen (18,19). This particle is not only located in the plasma but also has been shown to enter the arterial intima of humans and has an increased affinity by the

extracellular matrix (20). This issue confers a greater opportunity to Lp(a) oxidation (21) and interaction of Lp(a) with macrophages (22,23). Recently, it has been suggested that Lp(a) could be a preferential carrier of oxidized phospholipids in human plasma (24). These oxidized Lp(a) have a greater atherosclerotic effect as compared to native Lp(a) and this action may be increased by hyperglucemia (25). Different Lp(a) subtypes have been proposed regarding to apo(a) isoforms and these apo(a) isoforms predict the risk for CHD independently of the ethnic group (26). These isoforms are classified in order to their different size (16). Table 1 shows classification of these isoforms and its relation with KIV repeats.

Fig. 1. Metabolic route of Lp(a).

3. Epidemiological aspects

Plasma levels of Lp(a) show great diversity regarding to different ethnical groups but a plasmatic concentration greater than 30 mg/dl is currently considered an independent cardiovascular risk factor (27). In this sense, African-Americans have higher Lp(a) concentrations than Caucasians. These levels may also be very different even in individuals carrying apo(a) of the same size polymorphism. It has been suggested the possibility of the presence of additional factors affecting this ethnical differences or the existence of high risk-Lp(a) or low risk-Lp(a) (28,29). By the other hand, not all ethnic groups show the same relation with Lp(a). In American-Indians, Lp(a) level has been reported to be low and non independently associated with cardiovascular disease (30).

<i>Repeats (No.)</i>	<i>Molecular weight (kDa)</i>
5-12	<400
13-20	400-500
20-25	500-650
>25	>700

Table 1. Relation between KIV₂ repeats and apo(a) isoforms size

Respecting to apo(a) isoforms, it has been suggested a most important pathogenic role of Lp(a) particles with smaller apo(a) isoforms (18,31). This is probably due to several factors. First, an increased capacity to bind oxidized phospholipids, second, the ability to localize in blood vessel walls, and eventually related to its thrombogenic effect by increasing inhibition of plasmin activity. Apo(a) size heterogeneity is related to a copy number variation in the protein domain kringle IV type 2 (KIV₂) (32) (Table 1). This copy number variation (5-50 identically repeated copies) confers heterogeneity in the molecular mass of apo(a) ranging between 200 and 800 kDa. Ethnic differences in the frequency distribution of apo(a) KIV repeated alleles have been reported (33,34). In all ethnic groups, Caucasians, Asians and African-Americans, higher levels of circulating Lp(a) concentrations tend to be associated with smaller apo(a) isoforms (35,36). This finding could explain partially the association of higher Lp(a) levels and cardiovascular disease. People with smaller apo(a) isoforms have an approximately 2-fold higher risk of coronary artery disease and ischemic stroke than those with larger apo(a) isoforms. Furthermore, isoforms with less KIV repetitions (isoforms F, B, S1 and S2) have the greater analogy with plasminogen being associated with higher coronary risk (37,38).

4. Genetic considerations

Apo(a) gen (6q2.6-q2.7) (39,40) have different kringle domains that show a high degree of homology to the kringle domains IV and V of plasminogen (41).

Genetic variants associated with Lp(a) level have been associated with coronary disease (42). More specifically, the apo(a) gen is the major determinant of variation in some populations like African-Americans modulating the plasmatic concentration of Lp(a) (43). It has been reported that apo(a) gene accounts for greater than 90% of the variation of plasmatic Lp(a) concentrations (28). Apo(a) gen polymorphisms as well certain gene cluster associated to *LPA* have been shown to modulate Lp(a) concentrations leading to an increase in the risk for coronary artery disease (44). The genetic basis for apo(a) isoform variation is a segment existing in multiple repeats (KIV₂ polymorphism) located in the *LPA* gene (41). Variations in nucleotide polymorphisms in *LPA* may be an important contributor to the observed Lp(a) between-population variance and increase Lp(a) level in some populations (45-47). Once again, ethnic differences have been reported in people of European continental ancestry where apo(a) isoform polymorphism contributes between 40% and 70% of the variation of Lp(a) concentration showing fewer number of KIV₂ repeats (41,46), (Table 1).

5. Evidences favoring association with cardiovascular disease

- **Coronary heart disease:** Circulating Lp(a) concentration is associated with risk of coronary heart disease (CHD) independently from other conventional risk factors including total cholesterol concentration. Lp(a) excess has been independently associated to myocardial infarction and unstable angina (48), restenosis after coronary angioplasty (49), and coronary bypass grafting (50) respectively. Prospective epidemiological studies have reported positive association of baseline Lp(a) concentration with CHD risk. Based on this epidemiological data, a relative risk of 1.5 has been reported involving those patients with mean Lp(a) values of 50 mg/dL, especially in patients with premature coronary disease (51). Continuous associations of Lp(a) with the risk of coronary artery disease have been reported and this association is similar regarding to coronary death and non-fatal myocardial infarction (52-54). This association is not significantly affected by sex, non-HDL or HDL cholesterol, triglycerides, blood pressure, diabetes, or body mass index. These results are consistent mainly in Caucasians but studies in non-Caucasians are needed to corroborate also this issue in other populations (33). The association of Lp(a) concentrations with CHD is only slightly reduced after adjustment for long-term average levels of lipids and other established risk factors. This situation increases the likelihood that Lp(a) is an independent risk factor for CHD (53). The strength of Lp(a) as coronary risk factor is relatively modest as compared with non-HDL cholesterol. This is somewhat different when the level of Lp(a) is very high leading to a proportionally most important role for Lp(a) as CHD risk factor (52). Trying to associate fibrinolysis and myocardial ischemic disease, it has been suggested that Lp(a) may inhibit fibrinolysis of coronary artery thrombus (55). This is because higher levels of Lp(a) have been reported in survivors of myocardial infarction in whom recanalization of infarct artery failed as compared with patients with a patent artery (56). Other prospective studies have not shown relationship between high levels of Lp(a) or apo(a) isoforms and cardiovascular risk (57-59) contributing to some degree of controversy.
- **Stroke:** Serum Lp(a) concentration is also associated independently with risk of ischemic stroke (60,61). Current data in relation to Lp(a) concentration and stroke are sparse but seem to be similar than those for CHD. Serum Lp(a) level was demonstrated to predict stroke in elderly people in a large longitudinal (62) and in a case-control study (63). It has been shown that high levels of Lp(a) are associated with ischemic stroke in patients with atrial fibrillation especially when left atrial thrombus is present (64). Unhealthy dietary fat intake and a high serum Lp(a) level have been shown to predict fatal and nonfatal stroke or transient ischemic attack independently of established risk factors in a study of a community-based sample of middle-age men (65). Lp(a) has also been detected in intraparenchymal cerebral vessels suggesting a potential inflammatory role in acute stroke for Lp(a) (66). Other studies have not found statistical relationship between higher level of Lp(a) and thrombotic stroke (67).

6. Special situations

There are some common medical conditions that may be influenced by the level of Lp(a). Conversely, serum Lp(a) levels can be modified by the existence of some medical disorders. These medical conditions are summarized as follows:

- **Lp(a) and dialysis:** It is well known that atherosclerosis is more prevalent among patients with end-stage renal disease (68). Hemodialysis procedure “per se” has been shown to modify serum levels of Lp(a) increasing them after hemodialysis procedure (69). It has been proposed that inflammation, a very important condition in hemodialysis patients, could play an important role in this Lp(a) increase (70-72). Basal serum levels of Lp(a) are increased in dialysis patients and the level is elevated in almost 70% of patients (73). Even more, in patients with continuous ambulatory peritoneal dialysis, Lp(a) level is significantly higher as compared with patients on hemodialysis (74) pointing to a possible modulating effect of Lp(a) concentration by the different dialysis procedures. Particularly, high serum Lp(a) levels and the low molecular weight apo(a) phenotype have been associated with adverse clinical outcomes in dialysis patients (75).
- **Lp(a) and atrial fibrillation:** Higher serum Lp(a) level in ischemic stroke patients associated with atrial fibrillation and left atrial thrombus formation or in acute myocardial infarction has been reported (76,77). Lp(a) elevation and reduced left atrial appendage flow velocities have been shown to be independently risk factors for thromboembolism in chronic nonvalvular atrial fibrillation (55). Probably, the association of Lp(a) is stronger in the presence of atrial thrombus instead of atrial fibrillation itself, because of the plasminogen inhibitory action of Lp(a) (64). In this sense, other studies have not found association between higher levels of Lp(a) and non-valvular atrial fibrillation (78).
- **Lp(a) and blood coagulation:** the genetic homology in the cDNA sequence of human apo(a) with plasminogen, the zymogen for the major fibrinolytic serine protease plasmin (79), has been related with the cardiovascular pathogenicity of Lp(a) (80). There is a major difference in the kringle structure between plasminogen and Lp(a) that is a single aminoacid exchange (R560S) that prevents apo(a) from enzymatic cleavage such as the action of tissue-type plasminogen activator (t-PA) or urokinase plasminogen activator (u-PA). This molecular mimicry between plasminogen and Lp(a) contribute to the role of Lp(a) in atherogenesis binding Lp(a) to the tissue factor pathway inhibitor (TFPI), docking to diverse lipoprotein receptors (especially those affecting LDL or very low density lipoprotein (VLDL) and by the entrapment of Lp(a) into extracellular proteins (81). This situation leads to a retention of Lp(a) and recruitment of monocytes, upregulating the expression of the plasminogen activator inhibitor 2 in these monocytes (82). It has also been reported that Lp(a) modulates endothelial cell surface fibrinolysis contributing to the increase in atherosclerotic risk (83).
- **Lp(a) and alcohol intake:** Many epidemiological and clinical studies have shown that light-to-moderate alcohol consumption is associated with reduced risk of CHD and total mortality in the middle-age and elderly of both genders (84,85). Lipid levels are modified by alcohol in different forms but it is not completely clear the way they are. In alcohol abuse patients, levels of Lp(a) have been reported to decrease and this has been related to the time of abstinence (86). In other study an increased level among table wine drinkers has been described (87). A special situation is the association of alcohol intake, Lp(a) level and vascular disease. In this sense, high serum Lp(a) concentration and heavy drinking were found independently associated with larger infrarenal aortic diameters (88) and abdominal aortic aneurysms (89), probably due to the capability of Lp(a) to inhibit elastolysis in the vessels wall (90).

7. Treatment

Treatment possibilities are scarce at present when the aim is to reduce Lp(a) plasma concentration. Only niacin, in a dose dependent fashion, and certain inhibitors of cholesteryl ester transfer protein have shown limited effect ranging between 20%-40% lowering from baseline levels (91,92). Other drugs such as acetylsalicylic acid and L-carnitine can decrease mildly elevated Lp(a) concentrations (91,93,94). Contradictory findings have been reported with statins (95-98). Promising molecules like mipomersen, an antisense oligonucleotide directed to human apoB₁₀₀ have been shown to reduce Lp(a) concentrations by 70% in transgenic mice (99). Similar molecules such as eprotirone, tibolone and proprotein convertase subtilisin/kexin type 9 (PCSK-9) inhibitors can also decrease Lp(a) concentrations being currently under development (91,100-102). Nevertheless, the most dramatic change in Lp(a) concentrations can be achieved with regular lipid apheresis (103,104). Table 2 shows the efficacy of different treatment options in reducing Lp(a) plasmatic level.

<i>Treatment</i>	<i>Change in Lp(a) concentration (%)</i>
Diet and exercise	0
Resins	0
Fibrates	5-10
Statins	5
Nicotinic acid	35
Neomicine	25
Estrogen substitutive therapy	15-40
Apheresis	40-60

Table 2. Effect of different pharmacological therapies on Lp(a) serum concentration.

8. Controversies

The risk associated to Lp(a) concentration is only about one-quarter of that seen with LDL cholesterol so any clinical implication of this moderate association currently appeared limited. The role of specific Lp(a) subtypes could help to clarify the vascular risk. Particularly, smaller apo(a) isoforms could act associated with other factors such as small-dense LDL and oxidized LDL particles in the vessel wall increasing inflammation and accelerating atherosclerotic disease. This fact needs for more investigation.

Studies reporting association of apo(a) isoforms size variations with the risk of vascular disease have reported divergent relative risks, involve wide confidence intervals and the number of individuals included has been small. If smaller apo(a) isoforms are relevant to

vascular disease independent from Lp(a) concentration is not completely clear at present. Moreover, many studies have used different cut-offs to define smaller apo(a) size.

The effect of the change in Lp(a) level and its relation with inflammation as well as its influence on endothelial function are unknown at present.

It has been suggested that Lp(a) is associated with CHD only at very high concentrations but this affirmation remains somewhat controversial making very important to identify possible ethnical differences as well as an adequate cut-off level we can rely on.

9. Conclusions

Lp(a) results from the association of apo(a) and LDL particles. Since first studies linking Lp(a) and cardiovascular disease, an important amount of clinical and laboratory evidences have supported the fact that Lp(a) is an independent cardiovascular risk factor, especially in younger people with premature cardiovascular disease.

Many ethnical differences and variations in apo(a) size have been reported. Moreover, small apo(a) size isoforms have been related with an increased cardiovascular risk. Its relation with the number of KIV repeats determines genetically variation in apo(a) size. Several studies including methanalysis have related higher levels of Lp(a) with CHD and stroke.

It seems also that Lp(a) is elevated in patients under dialysis, and possibly in those with atrial fibrillation increasing the cardiovascular risk of these patients, normally already high.

An interesting link between laboratory and clinical effects of Lp(a) is its action modulating the fibrinolytic system because of the great homology between Lp(a) and plasminogen.

The association between higher levels of Lp(a) and alcohol intake remains more controversial at present.

Current treatment options are not very useful except for niacin and plasma apheresis but both therapies are not easy to use because of toxicity, tolerability and availability.

Finally, large prospective studies are needed focusing on Lp(a)-associated small apo(a) isoforms and cardiovascular disease, and also in order to ensure treatment approaches.

10. References

- [1] Schriewer H, Assmann G, Sandkamp M, Schulte H. The relationship of lipoprotein (a) (Lp(a)) to risk factors of coronary heart disease: initial results of the prospective epidemiological study on company employees in Westfalia. *J Clin Chem Clin Biochem.* 1984;22:591-596.
- [2] Schernthaner G, Kostner GM, Dieplinger H, Prager R, Mühlhauser I. Apolipoproteins (A-I, A-II, B), Lp(a) lipoprotein and lecithin: cholesterol acyltransferase activity in diabetes mellitus. *Atherosclerosis.* 1983;49:277-293.
- [3] Murai A, Miyahara T, Fujimoto N, Matsuda M, Kameyama M. Lp(a) lipoprotein as a risk factor for coronary heart disease and cerebral infarction. *Atherosclerosis.* 1986;59:199-204.

- [4] Marcovina SM, Albers JJ, Scanu AM, Kennedy H, Giaculli F, Berg K, Couderc R, Dati F, Rifai N, Sakurabayashi I, Tate JR, Steinmetz A. Use of a reference material proposed by the International Federation of Clinical Chemistry and Laboratory Medicine to evaluate analytical methods for the determination of plasma lipoprotein(a). *Clin Chem*. 2000;46:1956-1967.
- [5] Dati F, Tate JR, Marcovina SM, Steinmetz A; International Federation of Clinical Chemistry and Laboratory Medicine; IFCC Working Group for Lipoprotein(a) Assay Standardization. First WHO/IFCC International Reference Reagent for Lipoprotein(a) for Immunoassay--Lp(a) SRM 2B. *Clin Chem Lab Med*. 2004;42:670-676.
- [6] Scanu AM, Lawn RM, Berg K. Lipoprotein(a) and atherosclerosis. *Ann Intern Med*. 1991;115:209-218.
- [7] Orsó E, Ahrens N, Kilalić D, Schmitz G. Familial hypercholesterolemia and lipoprotein(a) hyperlipidemia as independent and combined cardiovascular risk factors. *Atheroscler Suppl*. 2009;10:74-78.
- [8] Ariyo AA, Thach C, Tracy R; Cardiovascular Health Study Investigators. Lp(a) lipoprotein, vascular disease, and mortality in the elderly. *N Engl J Med*. 2003;349:2108-2115.
- [9] Nordestgaard BG, Chapman MJ, Ray K, Borén J, Andreotti F, Watts GF, Ginsberg H, Amarencio P, Catapano A, Descamps OS, Fisher E, Kovanen PT, Kuivenhoven JA, Lesnik P, Masana L, Reiner Z, Taskinen MR, Tokgözoğlu L, Tybjaerg-Hansen A; European Atherosclerosis Society Consensus Panel. Lipoprotein(a) as a cardiovascular risk factor: current status. *Eur Heart J*. 2010;31:2844-2853.
- [10] Marcovina SM, Koschinsky ML, Albers JJ, Skarlato S. Report of the National Heart, Lung, and Blood Institute Workshop on Lipoprotein(a) and Cardiovascular Disease: recent advances and future directions. *Clin Chem*. 2003;49:1785-1796.
- [11] Boffa MB, Marcovina SM, Koschinsky ML. Lipoprotein(a) as a risk factor for atherosclerosis and thrombosis: mechanistic insights from animal models. *Clin Biochem*. 2004;37:333-343.
- [12] Emerging Risk Factors Collaboration, Erqou S, Kaptoge S, Perry PL, Di Angelantonio E, Thompson A, White IR, Marcovina SM, Collins R, Thompson SG, Danesh J. Lipoprotein(a) concentration and the risk of coronary heart disease, stroke, and nonvascular mortality. *JAMA*. 2009;302:412-423.
- [13] Anuurad E, Boffa MB, Koschinsky ML, Berglund L. Lipoprotein(a): a unique risk factor for cardiovascular disease. *Clin Lab Med*. 2006;26:751-772.
- [14] Berg K. A new serum type system in man. The LP system. *Acta Pathol Microbiol Scand*. 1963;59:369-382.
- [15] Utermann G. The mysteries of lipoprotein(a). *Science*. 1989;246:904-910.
- [16] Erqou S, Thompson A, Di Angelantonio E, Saleheen D, Kaptoge S, Marcovina S, Danesh J. Apolipoprotein(a) isoforms and the risk of vascular disease: systematic review of 40 studies involving 58,000 participants. *J Am Coll Cardiol*. 2010;55:2160-2167.
- [17] Rubin J, Kim HJ, Pearson TA, Holleran S, Berglund L, Ramakrishnan R. The apolipoprotein(a) gene: linkage disequilibria at three loci differs in African Americans and Caucasians. *Atherosclerosis*. 2008;201:138-147.
- [18] Simó JM, Joven J, Vilella E, Ribas M, Pujana MA, Sundaram IM, Hammel JP, Hoover-Plow JL. Impact of apolipoprotein(a) isoform size heterogeneity on the lysine

- binding function of lipoprotein(a) in early onset coronary artery disease. *Thromb Haemost.* 2001;85:412-417.
- [19] Kraft HG, Lingenhel A, Pang RW, Delport R, Trommsdorff M, Vermaak H, Janus ED, Utermann G. Frequency distributions of apolipoprotein(a) kringle IV repeat alleles and their effects on lipoprotein(a) levels in Caucasian, Asian, and African populations: the distribution of null alleles is non-random. *Eur J Hum Genet.* 1996;4:74-87.
- [20] Williams KJ, Flees GM, Petrie KA, Snyder ML, Brocia RW, Swenson TL. Mechanisms by which lipoprotein lipase alters cellular metabolism of lipoprotein(a), low density lipoprotein, and nascent lipoproteins. Roles for low density lipoprotein receptors and heparin sulfate proteoglycans. *J Biol Chem.* 1992;267:13284-13292.
- [21] Naruscewicz M, Selinger E, Davignon J. Oxidative modification of lipoprotein(a) and the effect of beta-carotene. *Metabolism.* 1992;41:1215-1224.
- [22] Zioncheck TF, Powell LM, Rice GC, Eaton DL, Lawn RM. Interaction of recombinant apolipoprotein(a) and lipoprotein(a) with macrophages. *J Clin Invest.* 1991;87:767-771.
- [23] Tsimikas S, Witztum JL. The role of oxidized phospholipids in mediating lipoprotein(a) atherogenicity. *Curr Opin Lipidol.* 2008;19:369-377.
- [24] Bergmark C, Dewan A, Orsoni A, Merki E, Miller ER, Shin MJ, Binder CJ, Hörkkö S, Krauss RM, Chapman MJ, Witztum JL, Tsimikas S. A novel function of lipoprotein [a] as a preferential carrier of oxidized phospholipids in human plasma. *J Lipid Res.* 2008;49:2230-2239.
- [25] Kotani K, Yamada S, Uurtuya S, Yamada T, Taniguchi N, Sakurabayashi I. The association between blood glucose and oxidized lipoprotein(a) in healthy young women. *Lipids Health Dis.* 2010;9:103.
- [26] Sandholzer C, Saha N, Kark JD, Rees A, Jaross W, Dieplinger H, Hoppichler F, Boerwinkle E, Utermann G. Apo(a) isoforms predict risk for coronary heart disease. A study in six populations. *Arterioscler Thromb.* 1992;12:1214-1226.
- [27] Wittrup HH, Tybjaerd-Hansen A, Nordestgaard BG. Lipoprotein lipase mutations, plasma lipids and lipoproteins, and risk of ischaemic heart disease. A meta-analysis. *Circulation.* 1999;99:2901-2907.
- [28] Boerwinkle E, Leffert CC, Lin J, Lackner C, Chiesa G, Hobbs HH. Apolipoprotein(a) gene accounts for greater than 90% of the variation in plasma lipoprotein(a) concentrations. *J Clin Invest.* 1992;90:52-60.
- [29] Mooser V, Scheer D, Marcovina SM, Wang J, Guerra R, Cohen J, Hobbs HH. The Apo(a) gene is the major determinant of variation in plasma Lp(a) levels in African Americans. *Am J Hum Genet.* 1997;61:402-417.
- [30] Wang W, Hu D, Lee ET, Fabsitz RR, Welty TK, Robbins DC, J L Yeh, Howard BV. Lipoprotein(a) in American Indians is low and not independently associated with cardiovascular disease. The Strong Heart Study. *Ann Epidemiol.* 2002;12:107-114.
- [31] Zeljkovic A, Bogavac-Stanojevic N, Jelic-Ivanovic Z, Spasojevic-Kalimanovska V, Vekic J, Spasic S. Combined effects of small apolipoprotein (a) isoforms and small, dense LDL on coronary artery disease risk. *Arch Med Res.* 2009;40:29-35.
- [32] Kraft HG, Lingenhel A, Pang RW, Delport R, Trommsdorff M, Vermaak H, Janus ED, Utermann G. Frequency distributions of apolipoprotein(a) kringle IV repeat alleles and their effects on lipoprotein(a) levels in Caucasian, Asian, and African

- populations: the distribution of null alleles is non-random. *Eur J Hum Genet.* 1996;4:74-87.
- [33] Guyton JR, Dahlen GH, Patsch W, Kautz JA, Gotto AM Jr. Relationship of plasma lipoprotein Lp(a) levels to race and to apolipoprotein B. *Arteriosclerosis.* 1985;5:265-272.
- [34] Chiu L, Hamman RF, Kamboh MI. Apolipoprotein A polymorphisms and plasma lipoprotein(a) concentrations in non-Hispanic Whites and Hispanics. *Hum Biol.* 2000;72:821-835.
- [35] Boomsma DI, Knijff P, Kaptein A, Labeur C, Martin NG, Havekes LM, Princen HM. The effect of apolipoprotein(a)-, apolipoprotein E-, and apolipoprotein A4-polymorphisms on quantitative lipoprotein(a) concentrations. *Twin Res.* 2000;3:152-158.
- [36] Paultre F, Pearson TA, Weil HF, Tuck CH, Myerson M, Rubin J, Francis CK, Marx HF, Philbin EF, Reed RG, Berglund L. High levels of Lp(a) with a small apo(a) isoform are associated with coronary artery disease in African American and white men. *Arterioscler Thromb Vasc Biol.* 2000;20:2619-2624.
- [37] Seman LJ, McNamara JR, Schaefer EJ. Lipoprotein(a), homocysteine, and remnantlike particles: emerging risk factors. *Curr Opin Cardiol.* 1999;14:186-191.
- [38] Lamon-Fava S, Marcovina SM, Albers JJ, Kennedy H, Deluca C, White CC, Cupples LA, McNamara JR, Seman LJ, Bongard V, Schaefer EJ. Lipoprotein(a) levels, apo(a) isoform size, and coronary heart disease risk in the Framingham Offspring Study. *J Lipid Res.* 2011;52:1181-1187.
- [39] Kraft HG, Köchl S, Menzel HJ, Sandholzer C, Utermann G. The apolipoprotein (a) gene: a transcribed hypervariable locus controlling plasma lipoprotein (a) concentration. *Hum Genet.* 1992;90:220-30.
- [40] Ober C, Nord AS, Thompson EE, Pan L, Tan Z, Cusanovich D, Sun Y, Nicolae R, Edelstein C, Schneider DH, Billstrand C, Pfaffinger D, Phillips N, Anderson RL, Philips B, Rajagopalan R, Hatsukami TS, Rieder MJ, Heagerty PJ, Nickerson DA, Abney M, Marcovina S, Jarvik GP, Scanu AM, Nicolae DL. Genome-wide association study of plasma lipoprotein(a) levels identifies multiple genes on chromosome 6q. *J Lipid Res.* 2009;50:798-806.
- [41] Crawford DC, Peng Z, Cheng JF, Boffelli D, Ahearn M, Nguyen D, Shaffer T, Yi Q, Livingston RJ, Rieder MJ, Nickerson DA. LPA and PLG sequence variation and kringle IV-2 copy number in two populations. *Hum Hered.* 2008;66:199-209.
- [42] Clarke R, Peden JF, Hopewell JC, Kyriakou T, Goel A, Heath SC, Parish S, Barlera S, Franzosi MG, Rust S, Bennett D, Silveira A, Malarstig A, Green FR, Lathrop M, Gigante B, Leander K, de Faire U, Seedorf U, Hamsten A, Collins R, Watkins H, Farrall M; PROCARDIS Consortium. Genetic variants associated with Lp(a) lipoprotein level and coronary disease. *N Engl J Med.* 2009;361:2518-2528.
- [43] Mooser V, Scheer D, Marcovina SM, Wang J, Guerra R, Cohen J, Hobbs HH. The Apo(a) gene is the major determinant of variation in plasma Lp(a) levels in African Americans. *Am J Hum Genet.* 1997;61:402-417.
- [44] Marcovina SM, Albers JJ, Wijsman E, Zhang Z, Chapman NH, Kennedy H. Differences in Lp[a] concentrations and apo[a] polymorphs between black and white Americans. *J Lipid Res.* 1996;37:2569-2585.

- [45] Chretien JP, Coresh J, Berthier-Schaad Y, Kao WH, Fink NE, Klag MJ, Marcovina SM, Giaculli F, Smith MW. Three single-nucleotide polymorphisms in LPA account for most of the increase in lipoprotein(a) level elevation in African Americans compared with European Americans. *J Med Genet*. 2006;43:917-923.
- [46] Dumitrescu L, Glenn K, Brown-Gentry K, Shephard C, Wong M, Rieder MJ, Smith JD, Nickerson DA, Crawford DC. Variation in LPA is associated with Lp(a) levels in three populations from the Third National Health and Nutrition Examination Survey. *PLoS One*. 2011;6:e16604.
- [47] Kraft HG, Haibach C, Lingenhel A, Brunner C, Trommsdorff M, Kronenberg F, Müller HJ, Utermann G. Sequence polymorphism in kringle IV 37 in linkage disequilibrium with the apolipoprotein (a) size polymorphism. *Hum Genet*. 1995;95:275-282.
- [48] Dangas G, Ambrose JA, D'Agate DJ, Shao JH, Chockalingham S, Levine D, Smith DA. Correlation of serum lipoprotein(a) with the angiographic and clinical presentation of coronary artery disease. *Am J Cardiol*. 1999;83:583-585.
- [49] Desmarais RL, Sarembock IJ, Ayers CR, Vernon SM, Powers ER, Gimple LW. Elevated serum lipoprotein(a) is a risk factor for clinical recurrence after coronary balloon angioplasty. *Circulation*. 1995;91:1403-1409.
- [50] Hoff HF, Beck GJ, Skibinski CI, Jürgens G, O'Neil J, Kramer J, Lytle B. Serum Lp(a) level as a predictor of vein graft stenosis after coronary artery bypass surgery in patients. *Circulation*. 1988;77:1238-1244.
- [51] Bostom AG, Cupples LA, Jenner JL, Ordovas JM, Seman LJ, Wilson PW, Schaefer EJ, Castelli WP. Elevated plasma lipoprotein(a) and coronary heart disease in men aged 55 years and younger. A prospective study. *JAMA*. 1996;276:544-548.
- [52] Bennet A, Di Angelantonio E, Erqou S, Eiriksdottir G, Sigurdsson G, Woodward M, Rumley A, Lowe GD, Danesh J, Gudnason V. Lipoprotein(a) levels and risk of future coronary heart disease: large-scale prospective data. *Arch Intern Med*. 2008;168:598-608.
- [53] Danesh J, Collins R, Peto R. Lipoprotein(a) and coronary heart disease. Meta-analysis of prospective studies. *Circulation*. 2000;102:1082-1085.
- [54] Marcovina SM, Koschinsky ML. A critical evaluation of the role of Lp(a) in cardiovascular disease: can Lp(a) be useful in risk assessment? *Semin Vasc Med*. 2002;2:335-344.
- [55] Igarashi Y, Kasai H, Yamashita F, Sato T, Inuzuka H, Ojima K, Aizawa Y. Lipoprotein(a), left atrial appendage function and thromboembolic risk in patients with chronic nonvalvular atrial fibrillation. *Jpn Circ J*. 2000;64:93-98.
- [56] Moliterno DJ, Lange RA, Meidell RS, Willard JE, Leffert CC, Gerard RD, Boerwinkle E, Hobbs HH, Hillis LD. Relation of plasma lipoprotein(a) to infarct artery patency in survivors of myocardial infarction. *Circulation*. 1993;88:935-940.
- [57] Ridker PM, Stampfer MJ, Rifai N. Novel risk factors for systemic atherosclerosis: a comparison of C-reactive protein, fibrinogen, homocysteine, lipoprotein(a), and standard cholesterol screening as predictors of peripheral arterial disease. *JAMA*. 2001;285:2481-2485.
- [58] Moliterno DJ, Jokinen EV, Miserez AR, Lange RA, Willard JE, Boerwinkle E, Hillis LD, Hobbs HH. No association between plasma lipoprotein(a) concentrations and the

- presence or absence of coronary atherosclerosis in African-Americans. *Arterioscler Thromb Vasc Biol.* 1995;15:850-855.
- [59] Akanji AO. Apo(a) isoforms do not predict risk for coronary heart disease in a Gulf Arab population. *Ann Clin Biochem.* 2000;37(Pt 3):360-366.
- [60] Smolders B, Lemmens R, Thijs V. Lipoprotein (a) and stroke: a meta-analysis of observational studies. *Stroke.* 2007;38:1959-1966.
- [61] Shintani S, Kikuchi S, Hamaguchi H, Shiigai T. High serum lipoprotein(a) levels are an independent risk factor for cerebral infarction. *Stroke.* 1993;24:965-969.
- [62] Ariyo AA, Thach C, Tracy R; Cardiovascular Health Study Investigators. Lp(a) lipoprotein, vascular disease, and mortality in the elderly. *N Engl J Med.* 2003;349:2108-2115.
- [63] Milionis HJ, Filippatos TD, Loukas T, Bairaktari ET, Tselepis AD, Elisaf MS. Serum lipoprotein(a) levels and apolipoprotein(a) isoform size and risk for first-ever acute ischaemic nonembolic stroke in elderly individuals. *Atherosclerosis.* 2006;187:170-176.
- [64] Okura H, Inoue H, Tomon M, Nishiyama S, Yoshikawa T. Increased plasma lipoprotein(a) level in cardioembolic stroke with non-valvular atrial fibrillation. *Intern Med.* 1998;37:995.
- [65] Wiberg B, Sundström J, Arnlöv J, Terént A, Vessby B, Zethelius B, Lind L. Metabolic risk factors for stroke and transient ischemic attacks in middle-aged men: a community-based study with long-term follow-up. *Stroke.* 2006;37:2898-2903.
- [66] Jamieson DG, Fu L, Usher DC, Lavi E. Detection of lipoprotein(a) in intraparenchymal cerebral vessels: correlation with vascular pathology and clinical history. *Exp Mol Pathol.* 2001;71:99-105.
- [67] Nagaraj SK, Pai P, Bhat G, Hemalatha A. Lipoprotein (a) and other Lipid Profile in Patients with Thrombotic Stroke: Is it a Reliable Marker? *J Lab Physicians.* 2011;3:28-32.
- [68] Coresh J, Longenecker JC, Miller ER 3rd, Young HJ, Klag MJ. Epidemiology of cardiovascular risk factors in chronic renal disease. *J Am Soc Nephrol.* 1998;9(12 Suppl):S24-30.
- [69] Díaz-Peromingo JA, Carbajal DG, Albán-Salgado A. Lipoprotein (a) in patients on hemodialysis. *Acta Med Austriaca.* 2004;31:73-75.
- [70] Dejanova B, Filipce V, Dejanov P, Sikole A, Grozdanovski R, Maleska V. Atherosclerosis risk factors related to hemodialysis duration and erythropoietin therapy. *Clin Chem Lab Med.* 2001;39:484-486.
- [71] Kaysen GA. Lipid and lipoprotein metabolism in chronic kidney disease. *J Ren Nutr.* 2009;19:73-77.
- [72] Hoover-Plow J, Hart E, Gong Y, Shchurin A, Schneeman T. A physiological function for apolipoprotein(a): a natural regulator of the inflammatory response. *Exp Biol Med (Maywood).* 2009;234:28-34.
- [73] Fleischmann EH, Bower JD, Salahudeen AK. Are conventional cardiovascular risk factors predictive of two-year mortality in hemodialysis patients? *Clin Nephrol.* 2001;56:221-230.
- [74] Ozdemir FN, Güz G, Sezer S, Arat Z, Turan M, Haberal M. Atherosclerosis risk is higher in continuous ambulatory peritoneal dialysis patients than in hemodialysis patients. *Artif Organs.* 2001;25:448-452.

- [75] Kronenberg F, Lingenhel A, Neyer U, Lhotta K, König P, Auinger M, Wiesholzer M, Andersson H, Dieplinger H. Prevalence of dyslipidemic risk factors in hemodialysis and CAPD patients. *Kidney Int Suppl.* 2003;84:113-116.
- [76] Igarashi Y, Yamaura M, Ito M, Inuzuka H, Ojima K, Aizawa Y. Elevated serum lipoprotein(a) is a risk factor for left atrial thrombus in patients with chronic atrial fibrillation: a transesophageal echocardiographic study. *Am Heart J.* 1998;136:965-971.
- [77] Celik S, Baykan M, Orem C, Kiliç K, Orem A, Erdöl C, Kaplan S. Serum lipoprotein(a) and its relation to left ventricular thrombus in patients with acute myocardial infarction. *Jpn Heart J.* 2001;42:5-14.
- [78] Díaz-Peromingo JA, Albán-Salgado A, García-Suárez F, Sánchez-Leira J, Saborido-Froján J, Iglesias-Gallego M. Lipoprotein(a) and lipid profile in patients with atrial fibrillation. *Med Sci Monit.* 2006;12:122-125.
- [79] McLean JW, Tomlinson JE, Kuang WJ, Eaton DL, Chen EY, Fless GM, Scanu AM, Lawn RM. cDNA sequence of human apolipoprotein(a) is homologous to plasminogen. *Nature.* 1987;330:132-137.
- [80] Loscalzo J, Weinfeld M, Fless GM, Scanu AM. Lipoprotein(a), fibrin binding, and plasminogen activation. *Arteriosclerosis.* 1990;10:240-245.
- [81] Caplice NM, Panetta C, Peterson TE, Kleppe LS, Mueske CS, Kostner GM, Broze GJ Jr, Simari RD. Lipoprotein (a) binds and inactivates tissue factor pathway inhibitor: a novel link between lipoproteins and thrombosis. *Blood.* 2001;98:2980-2987.
- [82] Buechler C, Ullrich H, Ritter M, Porsch-Oezcuemez M, Lackner KJ, Barlage S, Friedrich SO, Kostner GM, Schmitz G. Lipoprotein (a) up-regulates the expression of the plasminogen activator inhibitor 2 in human blood monocytes. *Blood.* 2001;97:981-986.
- [83] Hajjar KA, Gavish D, Breslow JL, Nachman RL. Lipoprotein(a) modulation of endothelial cell surface fibrinolysis and its potential role in atherosclerosis. *Nature.* 1989;339:303-305.
- [84] Rimm EB, Williams P, Fosher K, Criqui M, Stampfer MJ. Moderate alcohol intake and lower risk of coronary heart disease: meta-analysis of effects on lipids and haemostatic factors. *BMJ.* 1999;319:1523-1528.
- [85] Rehm J, Gmel G, Sempos CT, Trevisan M. Alcohol-related morbidity and mortality. *Alcohol Res Health.* 2003;27:39-51.
- [86] Kolovou GD, Salpea KD, Anagnostopoulou KK, Mikhailidis DP. Alcohol use, vascular disease, and lipid-lowering drugs. *J Pharmacol Exp Ther.* 2006;318:1-7.
- [87] Böhm M, Rosenkranz S, Laufs U. Alcohol and red wine: impact on cardiovascular risk. *Nephrol Dial Transplant.* 2004;19:11-6.
- [88.] Wang JA, Chen XF, Yu WF, Chen H, Lin XF, Xiang MJ, Fang CF, Du YX, Wang B. Relationship of heavy drinking, lipoprotein (a) and lipid profile to infrarenal aortic diameter. *Vasc Med.* 2009;14:323-329.
- [89] Sofi F, Marcucci R, Giusti B, Pratesi G, Lari B, Sestini I, Lo Sapio P, Pulli R, Pratesi C, Abbate R, Gensini GF. High levels of homocysteine, lipoprotein (a) and plasminogen activator inhibitor-1 are present in patients with abdominal aortic aneurysm. *Thromb Haemost.* 2005;94:1094-1098.
- [90] Petersen E, Wågborg F, Angquist KA. Does lipoprotein(a) inhibit elastolysis in abdominal aortic aneurysms? *Eur J Vasc Endovasc Surg.* 2003;26:423-428.

- [91] Parhofer KG. Lipoprotein(a): medical treatment options for an elusive molecule. *Curr Pharm Des.* 2011;17:871-876.
- [92] McKenney J. New perspectives on the use of niacin in the treatment of lipid disorders. *Arch Intern Med.* 2004;164:697-705.
- [93] Chasman DI, Shiffman D, Zee RY, Louie JZ, Luke MM, Rowland CM, Catanese JJ, Buring JE, Devlin JJ, Ridker PM. Polymorphism in the apolipoprotein(a) gene, plasma lipoprotein(a), cardiovascular disease, and low-dose aspirin therapy. *Atherosclerosis.* 2009;203:371-376.
- [94] Galvano F, Li Volti G, Malaguarnera M, Avitabile T, Antic T, Vacante M, Malaguarnera M. Effects of simvastatin and carnitine versus simvastatin on lipoprotein(a) and apoprotein(a) in type 2 diabetes mellitus. *Expert Opin Pharmacother.* 2009;10:1875-1882.
- [95] McKenney JM, Jones PH, Bays HE, Knopp RH, Kashyap ML, Ruoff GE, McGovern ME. Comparative effects on lipid levels of combination therapy with a statin and extended-release niacin or ezetimibe versus a statin alone (the COMPELL study). *Atherosclerosis.* 2007;192:432-437.
- [96] Gonbert S, Malinsky S, Sposito AC, Laouenan H, Doucet C, Chapman MJ, Thillet J. Atorvastatin lowers lipoprotein(a) but not apolipoprotein(a) fragment levels in hypercholesterolemic subjects at high cardiovascular risk. *Atherosclerosis.* 2002;164:305-311.
- [97] Pan J, Van JT, Chan E, Kesala RL, Lin M, Charles MA. Extended-release niacin treatment of the atherogenic lipid profile and lipoprotein(a) in diabetes. *Metabolism.* 2002;51:1120-1127.
- [98] Schaefer EJ, McNamara JR, Tayler T, Daly JA, Gleason JL, Seman LJ, Ferrari A, Rubenstein JJ. Comparisons of effects of statins (atorvastatin, fluvastatin, lovastatin, pravastatin, and simvastatin) on fasting and postprandial lipoproteins in patients with coronary heart disease versus control subjects. *Am J Cardiol.* 2004 Jan 1;93(1):31-9.
- [99] Bell DA, Hooper AJ, Burnett JR. Mipomersen, an antisense apolipoprotein B synthesis inhibitor. *Expert Opin Investig Drugs.* 2011;20:265-272.
- [100] Angelin B, Rudling M. Lipid lowering with thyroid hormone and thyromimetics. *Curr Opin Lipidol.* 2010;21:499-506.
- [101] Perrone G, Capri O, Galoppi P, Brunelli R, Bevilacqua E, Ceci F, Ciarla MV, Strom R. Effects of either tibolone or continuous combined transdermal estradiol with medroxyprogesterone acetate on coagulatory factors and lipoprotein(a) in menopause. *Gynecol Obstet Invest.* 2009;68:33-39.
- [102] Daher R, Al-Amin H, Beaini M, Usta I. Effect of tibolone therapy on lipids and coagulation indices. *Clin Chem Lab Med.* 2006;44:1498-1499.
- [103] Tselmin S, Schmitz G, Julius U, Bornstein SR, Barthel A, Graessler J. Acute effects of lipid apheresis on human serum lipidome. *Atheroscler Suppl.* 2009;10:27-33.
- [104] Jaeger BR, Richter Y, Nagel D, Heigl F, Vogt A, Roeseler E, Parhofer K, Ramlow W, Koch M, Utermann G, Labarrere CA, Seidel D; Group of Clinical Investigators. Longitudinal cohort study on the effectiveness of lipid apheresis treatment to reduce high lipoprotein(a) levels and prevent major adverse coronary events. *Nat Clin Pract Cardiovasc Med.* 2009;6:229-239.

Recent Advances in Cardiovascular Risk Factors

Edited by Prof. Mehnaz Atiq

ISBN 978-953-51-0321-9

Hard cover, 522 pages

Publisher InTech

Published online 21, March, 2012

Published in print edition March, 2012

Among the non-communicable diseases, cardiovascular disorders are the leading cause of morbidity and mortality in both the developed and the developing countries. The spectrum of risk factors is wide and their understanding is imperative to prevent the first and recurrent episodes of myocardial infarction, stroke or peripheral vascular disease which may prove fatal or disabling. This book has tried to present an update on risk factors incorporating new research which has thrown more light on the existing knowledge. It has also tried to highlight regional diversity addressing such issues. It will hopefully be resourceful to the cardiologists, general practitioners, family physicians, researchers, graduate students committed to cardiovascular risk prevention.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

José Antonio Díaz Peromingo (2012). Lipoprotein (a) and Cardiovascular Risk, Recent Advances in Cardiovascular Risk Factors, Prof. Mehnaz Atiq (Ed.), ISBN: 978-953-51-0321-9, InTech, Available from: <http://www.intechopen.com/books/recent-advances-in-cardiovascular-risk-factors/lp-a-and-cardiovascular-risk>

INTech
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](https://creativecommons.org/licenses/by/3.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen