We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Gamma Radiation

Richard Stalter and Dianella Howarth
St. John's University
USA

1. Introduction

The content of this chapter includes a brief history of gamma radiation, units of radiation measurement, ecological importance, tables including the half life of gamma emitting nuclides, comparative sensitivity of living organisms to gamma radiation, biological magnification of radioactive and nuclear materials, and brief descriptions of case studies of Woodwell (1962), Stalter and Kincaid 2009), and nuclear power plant disasters (Three Mile Island, USA, 1980, Chernobyl 1986, Japan 2011).

Gamma radiation is somewhat similar to x-rays in that both pass through living materials easily. Also referred to as "photons" they travel at the speed of light. Gamma rays have sufficient energy to ionize matter and therefore can damage living cells. The damage produced in the cell or tissue is proportional to the number of ionizing paths produced in the absorbing material. Isotopes of elements that are emitters are radionuclides important in fission products from nuclear testing, nuclear power plant disasters or waste.

The injurious affect of gamma rays depends on (1) their number (2) their energy and (3) their distance from the source of radiation. Radiation intensity decreases exponentially with increasing distance. Radiation damage on vascular plant species was demonstrated by Woodwell (1962) who subjected a mature pine oak forest at Brookhaven National Laboratory to gamma radiation from a cesium 137 source (Figure 1).

Fig. 1. Radiation dose and damage to a pine-oak forest, Brookhaven National Laboratory, 1961. Zones delineated by vertical lines (Woodwell 1962, Stalter and Kincaid 2009).

Gamma rays are external emitters that penetrate biological materials easily and produce their insidious effects without being taken internally. Alpha and beta particles are internal emitters; their damage to organisms is greatest when taken internally. Odum (1971) summarizes this concept best, "the alpha beta gamma series is one of increasing penetration but decreasing concentration of ionization and local damage." Alpha and beta radiation, unlike gamma radiation, are corpuscular in nature. While alpha particles travel but a few centimeters, and can be stopped by a layer of dead skin, they are dangerous because they produce a large amount of local ionization which can cause mutations disrupting cell processes. Beta particles are high speed electrons. While much smaller than alpha particles, they are able to travel up to a couple of centimeters in living tissue, giving up their energy over a large path. Beta particles, like alpha particles can damage tissue, and like alpha particles, can cause mutations that affect the functioning of cells.

2. The history of gamma radiation as applied to biological systems

Most are familiar with the discovery of x-radiation by Roentgen in 1895 and the isolation of radium by the Curies in 1898 (Goodspeed and Uber 1939). Researchers soon learned that both x-rays and radioactive substances such as radium produced similar effects on biological materials. Koernicke (1905) noted that cell division was delayed on x-ray and radium treated cells. Both Koernicke (1905) and Gager (1907) described "striking chromosomal disruptions" after cells were dosed with x-rays or exposed to radium, a gamma emitter. Gamma irradiated cells were also broken or fragmented by radiation treatment (Gager 1907, 1908). For additional historical work on radiation and plant cytogenetics the reader is directed to a review article by Goodspeed and Uber (1939). Smith (1958) compiled a paper on the use of radiation in the production of useful mutations based on papers presented in three symposia in the United States from August 1956 to January 1957. A more recent review article on ionizing radiation damage to plants was prepared by Klein and Klein (1971).

There are numerous studies applying gamma radiation to biological systems. Several investigations involving botanicals follow. Nuttall et al (1961) found that yellow sweet Spanish onions exposed to 4000 or 8000 rad prevented sprouting in 97% of their experimental group suggesting that irradiation might be a viable method of prolonging storage life for onions. This study, while intriguing, has not been generally accepted by a public concerned with the problems of radiation. A second article by Heeney and Rutherford (1964) examined the effects of gamma radiation on the storage life of fresh strawberries. A dose of 330,000 rad prevented fungal development of the redcoat strawberry variety stored at 40 degrees F for 26 days. The fungal free period was sharply reduced at lower radiation doses and/or at higher temperatures. Pritchard et al (1962) studied the effect of gamma radiation on the utilization of wheat straw by rumen microorganisms. They concluded that, "high levels of gamma radiation were needed to release nutrients trapped in wheat straw needed by microbes. However, the levels of gamma irradiation necessary for nutrient release were well above what was practical for commercial purposes."

Baumhover et al (1955) investigated the use of gamma irradiation on male sterilization on the control of screw-worm flies in the southern United States while Bushland (1960) Cutcomp (1967) and Lawson (1967) discussed this practice as a general way of controlling certain insect pests. Gambino and Lindberg (1964) examined the response of the pocket

mouse to ionizing radiation. McCormick and Golley (1966) presented data on irradiation of natural vegetation in the southeastern United States while Monk (1966) published a similar study on the effects of short-term gamma radiation on an old field. Witherspoon (1965, 1969) examined radiation damage to a forest surrounding an unshielded fast reactor in 1965, and followed this study with a report in 1969 on radiosensitivity of forest tree species to acute fast neutron radiation. Odum and Pigeon (1970) researched the effect of irradiation and ecology of a tropical rain forest in Puerto Rico.

3. Units of measurement

Three units, the gigabecquerel (GBq), gray (GY), and roentgen (R) are used to measure radiation. The GBq measures the number of gamma rays emitted from a source of radiation and is a unit of radioactivity that is defined as 1.37×10^{-12} atomic decays each second. The weight of the material comprising a GBq varies. One gram of radium is 37 GBq while 10^{-7} th of a gram of newly formed radio-sodium is also 37 GBq since both release 3.7×10^{10} disintegrations/second (Odum 1971). In dealing with biological systems, smaller units are generally used such as the millicurie microcurie and picocurie which are 10^{-3} , 10^{-6} and 10^{-12} respectively.

A second measurement of radiation is the GY. The absorbed dose of 1 GY means the absorption of 1 joule of radiation energy per kg of tissue. The third, the roentgen is nearly the same as the GY, and is used as a unit of measurement for exposure to gamma and x rays. Both are units of the total dose of radiation received by an organism. The dose rate is the amount of radiation received per unit time.

4. Ecological importance of radionuclides

There are different kinds of atoms of each element; these are referred to as isotopes. Some isotopes are radioactive, some not. Radioactive isotopes are unstable. These decay into other isotopes releasing radiation. Each radioactive isotope, radionuclide, have a specific rate of disintegration, its half life.

Radionuclides fall into well defined groups (Tables 1 and 2). Naturally occurring nuclides are included in Table 1 while those from fallout produced by fission or uranium and other elements are found in Table 2. Fission isotopes are produced from nuclear explosions which have for the most part been eliminated and from "controlled" reactions that produce nuclear power. While most of the aforementioned nuclides are not essential for the growth of organisms, they may be incorporated in biogeochemical cycles and become concentrated in food chains, especially strontium and cesium. Thus Woodwell (1962) used cesium as a gamma radiation emitter in his well published study of an irradiated pine oak forest at Brookhaven National Laboratory, Long Island, New York. More will be said about this research later in this paper.

5. Sensitivity of organisms to radioactivity

There is a wide range of sensitivity of organisms to radioactivity. Mammals are most sensitive while bacteria are most resistant especially as spores. Moreover there is a wide range of tolerance to radiation during the life cycle of an organism. Radiation sickness in

humans can be caused by as little as 0.35 Gy while a dose of 6-8 Gy is lethal to nearly 100% of individuals (Donnelly et al 2010). A dose of 2 Gy may kill some insect embryos while a dose of 100 Gy is necessary to kill all adult individuals (Odum 1971). Dividing cells are generally more susceptible to radiation than resting cells. The toxicity of radionuclides depends on the absorption, distribution in the body, half-life, elimination half-time, type of radiation emitted, and their energy.

Element	Half-Life	Radiations Emitted	
Uranium-235(²³⁵ U)	7 x 10 ⁸ yrs.	Alpha ³	Gamma ⁰
Radium-226 (²²⁶ Ra)	1620 yrs.	Alpha ³	Gamma ⁰
Potassium-40 (⁴⁰ K)	1.3 x 10 ⁹ yrs.	Beta ²	Gamma ²
Carbon-14 (See Table 3.)			

Table 1. Naturally occurring gamma emitting isotopes which contribute to background radiation (Odum 1971).

Element	Half-Life	Radiations Emitted	
The cesium group	33 yrs.	Beta ²	Gamma
Cesium-137 (¹³⁷ Cs) and	2.6 min	Beta	Gamma ¹
daughter barium-137 (¹³⁷ Ba)	2.3 yrs.	Beta ¹	Gamma ²
Cesium-134 (¹³⁴ Cs)			
The cerium group	285 days	Beta ¹	Gamma ⁰
Cerium-144 (¹⁴⁴ Ce) and	17 min.	Beta ²	Gamma ²
daughter praseodymium-144	33 days	Beta ¹	Gamma ¹
(¹³⁷ Pr)			
Cerium-141 (¹⁴¹ Ce)			
The ruthenium group	1 yr.	Beta ²	
Ruthenium-106 (¹⁰⁶ Ru) and	30 sec.	Beta ³	Gamma ²
daughter rhodium-106 (¹⁰⁶ Rh)	40 days	Beta ¹	Gamma ¹
Ruthenium-103 (¹⁰³ Ru)	65 days	Beta ¹	Gamma ¹
Zirconium-95 (⁹⁵ Zr) and daughter	35 days	Beta ⁰	Gamma ¹
niobium-95(⁹⁵ Nb)	12.8 days	Beta ¹	Gamma ¹
Barium-140 (¹⁴⁰ Ba) and daughter	40 hrs	Beta ²	Gamma ²
lanthanum-140(¹⁴⁰ La)	11.3 days	Beta ¹	Gamma ¹
Neodymium-147 (¹⁴⁷ Nd) and	2.6 yrs.	Beta ¹	Gamma
daughter promethium-147(¹⁴⁷ Pm)	61 days	Beta ²	Gamma ¹
Yttrium-91 (⁹¹ Y)	2.4 x 10 ⁴ yrs.	Alpha ³	Gamma ¹
Plutonium-239 (²³⁹ Pu)	8 days	Beta ¹	Gamma ¹
Iodine-131 (¹³¹ I)	7 x 10 ⁸ yrs.	Alpha ³	Gamma ⁰
Uranium-235 (²³⁵ U)			

Table 2. Elements important in fission products entering the environment through fallout or waste disposal.

Sparrow (1962), Sparrow and Evans (1961), Sparrow and Woodwell (1962), and Sparrow et al (1963) have demonstrated that sensitivity of ionizing radiation is directly proportional to the size of the cell nucleus or chromosome volume. The larger the chromosome volume the more sensitive the material is to radiation. There are also differences in radiation tolerance between wild and laboratory rodent populations. Gambino and Lindberg (1964) and Golley et al (1965) have reported that the lethal dose for 50% of some wild rodent populations is roughly twice that of laboratory white mice or white rats, likely due to the reduced variation in the latter.

Radioactivity has been successfully used to sterilize certain male insect pests. Sterile males are introduced to natural populations in large numbers which mate with females. A female mates only once, and once mated with a sterile male produces no young. Introducing radiated sterile male screw-worm flies in areas where they occur successfully reduced the number of screw-worm flies, a major pest in the southern United States. For those seeking more general information on this topic see Baumhover et al (1955) Bushland (1960), Cutcomp (1967), Knipling (1960, 1964, 1965, 1967) and Lawson (1967).

6. Radiation effects on ecosystems

Since the early 1960's there have been numerous studies on the effect of gamma radiation on ecosystems. These studies were fueled by the arms race between the Soviet Union and the United States (Stalter and Kincaid 2009). After lengthy negotiations between the two powers the SALT (Strategic Arms Limitation Treaty) was signed in 1971 and extended in 1977. With the signing of the treaty, less funding for irradiation studies was available (Stalter and Kincaid 2009). Thus most studies cited in this paper are those conducted prior to the SALT agreement of 1971. The gamma source that has been used has been either cesium 137 or cobalt 60. These include the studies of Woodwell (1962, 1965a) at Brookhaven National Laboratory, Long Island, New York, a tropical rain forest, Puerto Rico (Odum and Pigeon 1970) and the desert of Nevada (French 1965). Additional studies have been conducted in the fields and forests of Georgia (Odum and Kuenzler 1963) (Platt 1965), and Oak Ridge, Tennessee (Witherspoon 1965, 1969). Much additional work involving a portable gamma source on plant communities has been conducted at the Savanna River Ecology Laboratory, Aiken, South Carolina (McCormick and Platt 1962, McCormick and Golly 1966, Monk 1966, McCormick 1969).

Stalter and Kincaid (2009) investigated community development following gamma radiation at a pine-oak forest, Brookhaven National Laboratory, Long Island, New York. The objective of this study was to compare vascular plant community change at five vegetation zones the site of Woodwell's (1962) gamma irradiated forest (Figure 1). The zones were: the dead zone where all vegetation was killed; a graminoid *Carex pensylvanica* zone; an ericaceous zone; an oak dominated zone; and a control, the original oak pine forest. Radiation greater than 63,000 roentgens killed all vegetation. *Carex* dominated the zone receiving 27,000 to 63,000 roentgens, ericaceous shrubs, *Vaccinium* spp. and *Gaylussacia baccata* were dominant at the zone receiving 11,000 to 27,000 roentgens while oaks survived at the zone receiving 3600 to 11,000 roentgens. Upon completion of the Woodwell study in the 1970's, pitch pine (*Pinus rigida*) has invaded the total kill zone as bare mineral soil favors pine regeneration (Stalter and Kincaid 2009). *Carex* remained the dominant taxon in the

original *Carex* zone demonstrating again that different plant species vary in their tolerance of radiation.

Herbaceous plant communities may be more resistant to radiation than mature forests because many early successional species have small nuclei (Sparrow and Evans 1961) and also because herbaceous taxa like *Carex pensylvanica* have more below ground plant material which is shielded from gamma radiation. Sparrow (1962), Sparrow and Evans (1961), and Sparrow et al (1963) present detailed information on the relationship between nuclear volumes, chromosome numbers and relative radiosensitivity.

7. Biological magnification of radioactive material

Radioactive material may become concentrated or "biologically magnified" during food chain transfer. Numerous biology and ecology text books include information on how living organisms take up nutrients pesticides and radioactive material and concentrate them. Because this concept is well known, we direct the reader to several early studies involving the concentration of radioactive material (See the work of Foster and Rostenbach, 1954; Hanson and Kornberg 1956; Davis and Foster 1958). Ophel (1963) reported a concentration of strontium 90 in perch flesh as 5x that of lake water while that in perch bone was 3000x! Additional information on radioecological concentration can be found in Auberg and Crossley (1958), Auberg and Hungate (1967) and Polikarpov (1966).

8. Radioactive fallout

Radioactive particles that fall to the earth after above ground nuclear tests and nuclear power plant accidents are called radioactive fallout. Radioactive particles mix with the dust in the atmosphere and eventually fall to earth often thousands of miles from the initial explosion.

There are two types of nuclear weapons, the fission bomb and fusion bomb or thermonuclear weapon. In thermonuclear devices, deuterium fuses to form a heavier element with the release of energy and neutrons. A fission bomb is needed to trigger the fusion reaction. The thermonuclear weapon produces more neutrons which induce radioactivity in the environment than a fission device per unit of energy released. Roughly ten percent of the energy of a nuclear weapon is in residual radiation which may become dispersed in the atmosphere (Glasstone 1957). The amount of fallout produced depends on the type of weapon, size of the weapon and also on the amount of naturally occurring material that is mixed with the radioactive material released in the explosion. Fallout patterns and intensity depend upon the direction of the wind, speed and direction of the jet stream, presence and amount of precipitation.

Atomic explosions carry radioactive material high in the atmosphere where the radioactive material becomes fused with silica dust and other material present in the vicinity of the explosion. These particles are largely insoluble. The fallout particles may adhere to vegetation where they enter food chains at the primary consumer level. Fallout from Chernobyl in 1986 was deposited in Lappland (Sweden) where caribou consumed contaminated vegetation. Shifting winds also carried Chernobyl radiation particles to northern Italy where rabbit growers fed their rabbits vegetation contaminated with

radioactive fallout from Chernobyl. Ultimately the rabbits were destroyed because of the high concentration of radioactive material in their flesh.

There are differences in the kind of radionuclides that enter terrestrial and marine food chains. Soluble fission products, strontium 90 and cesium 137, are generally found in the highest amounts in land plants and animals. In marine systems fallout that forms strong complexes with organic matter such as cobalt 60, iron 59, zinc 65, and manganese 54 are most likely to be concentrated in marine organisms. In addition, those found in colloidal form such as cesium 134 and zirconium 95 are also found in high concentration in marine organisms. Cesium 134 is mostly from the fission products of a power reactor whereas cesium 137 can be formed during atomic power plant accidents or as a product of nuclear bomb explosions.

There are additional considerations/problems associated with concentrating radioactive material entering food chains as the concentration of radioactivity is also a function of nutrient richness, and the exchange and storage capacity of soils. Nutrient poor soils and thin soils such as those found on granite outcrops act as a nutrient trap providing more radionuclides to the vegetation. For example, sheep grazing on hill pastures in England accumulated 20x as much strontium 90 in their bones than sheep pastured in deep valleys where calcium content of the soil was higher and the grasses taller (Bryant et al 1957). For additional radiological work on tracers in food chains and trophic levels see Odum and Golley (1963), Odum and Kuenzler (1963), de la Cruz (1963), Ball and Hooper (1963), Foster (1958), and Foster and Davis (1956).

9. Nuclear power plant accidents

Brief descriptions of three power plant accidents in the United States the Soviet Union and Japan follow. The first nuclear power plant accident occurred at 4 am on March 28, 1979, near Harrisburg, Pennsylvania, USA, the state's capital. A malfunction in the cooling system resulted in a portion of the core to melt in the Number 2 reactor. The approximately 2 million people who lived near the plant had an average dose of 0.14 Gy (Rogovin 1980). Although some radioactive gas was released from the plant on the 29th and 30th of March there was, "not enough to cause any radiation dose above background levels in the neighborhood of the accident" (<http://www.world-nuclear.org/info/info/info/inf36.html>). Fortunately, there were no reported injuries or health issues emanating from the Three Mile Island accident.

A more serious nuclear accident occurred at the Chernobyl power plant located 80 miles north of the city of Chernobyl in the Ukraine, one of the original Soviet Republics. A "routine" shut down and test that began on the 25th of April, 1986, led to this disaster. At one in the morning, 26 April, the reactor's power source dropped and when the backup safety system failed, the reactor, Reactor Four, exploded. Shortly after the initial explosion at Chernobyl, the Swedish government reported high levels of radiation at their Forsmark nuclear power plant at Stockholm. When additional European nuclear power plants also experienced higher than normal levels of radiation, they contacted the USSR for an explanation. Although initially denying the nuclear disaster, on the 28th of April the USSR acknowledged that one of their reactors had been compromised.

Group A. Naturally occurring isotopes which contribute to background radiation.					
NUCLIDE		HALF-LIFE		RADIATIONS EMITTED	
Uranium-235 (²³⁵ U)		7 x 10 ⁸ yrs.		Alpha ³	Gamma ⁰
Radium-226 (²²⁶ Ra)		1620 yrs.		Alpha ³	Gamma ⁰
Potassium-40 (⁴⁰ K)		1.3 x 10 ⁹ yrs.		Beta ²	Gamma ²
Carbon-14 (¹⁴ C)		5568 yrs.		Beta ⁰	
⁰ Very low energy, less than 0.2 Mev; ¹ relatively low energy, 0.2-1 Mev; ² high energy, 1-3 Mev; ³ very high energy, over 3 Mev.					
Group B. Gamma emitting nuclides of elements which are essential constituents of organisms.					
Modified from Odum (1971).					
NUCLIDE		HALF-LIFE		RADIATIONS EMITTED	
Cobalt-60 (⁶⁰ Co)		5.27	yrs.	Beta ¹	Gamma ²
Copper-64 (⁶⁴ Cu)		12.8	hrs.	Beta ¹	Gamma ²
Iodine-131 (¹³¹ I)		8	days	Beta ¹	Gamma
Iron-59 (⁵⁹ Fe)		45	days	Beta ¹	Gamma ²
Manganese-54 (⁵⁴ Mn)		300	days	Beta ²	Gamma ²
Potassium-42 (⁴² K)		12.4	hrs.	Beta ³	Gamma ²
Sodium-22 (²² Na)		2.6	yrs.	Beta ¹	Gamma ²
Sodium-24 (²⁴ Na)		15.1	hrs.	Beta ²	Gamma ²
Zinc-65 (⁶⁵ Zn)		250	days	Beta ¹	Gamma ²
Also barium-140 (¹⁴⁰ Ba), bromine-82 (⁸² Br), molybdenum-99 (⁹⁹ Mo) and other trace elements.					
Group C. Nuclides important in fission products entering the environment through fallout or waste disposal.					
NUCLIDE		HALF-LIFE		RADIATIONS EMITTED	
The strontium group					
Strontium-90 (⁹⁰ Sr) and		28	yrs.	Beta ¹	
daughter yttrium-90 (⁹⁰ Y)		2.5	days	Beta ²	
Strontium-89 (⁸⁹ Sr)		53	days	Beta ²	
The cesium group					
Cesium-137 (¹³⁷ Cs) and		33	yrs.	Beta ²	Gamma
daughter barium-137 (¹³⁷ Ba)		2.6	min.	Beta	Gamma ¹
Cesium-134 (¹³⁴ Cs)		2.3	yrs.	Beta ¹	Gamma ²
The cerium group					
Cerium-144 (¹⁴⁴ Ce) and		285	days	Beta ¹	Gamma ⁰
daughter praseodymium-144 (¹⁴⁴ Pr)		17	min.	Beta ²	Gamma ²
Cerium-141 (¹⁴¹ Ce)		33	days	Beta ¹	Gamma ¹
The ruthenium group					
Ruthenium-106 (¹⁰⁶ Ru) and		1	yr.	Beta ⁰	
daughter rhodium-106 (¹⁰⁶ Rh)		30	sec.	Beta ³	Gamma ²
Ruthenium-103 (¹⁰³ Ru)		40	days	Beta ¹	Gamma ¹
Zirconium-95 (⁹⁵ Zr) and daughter		65	days	Beta ¹	Gamma ¹
niobium-95 (⁹⁵ Nb)		35	days	Beta ⁰	Gamma ¹
Barium-140 (¹⁴⁰ Ba) and daughter		12.8	days	Beta ¹	Gamma ¹

lanthanum-140 (¹⁴⁰ La)	40	hrs.	Beta ²	Gamma ²
Neodymium-147 (¹⁴⁷ Nd) and	11.3	days	Beta ¹	Gamma ¹
daughter promethium-147 (¹⁴⁷ Pm)	2.6	yrs.	Beta ¹	Gamma
Yttrium-91 (⁹¹ Y)	61	days	Beta ²	Gamma
Plutonium-239 (²³⁹ Pu)	2.4 x 10 ⁴	yrs.	Alpha ³	Gamma ¹
Iodine-131 (see Group B)				
Uranium (see Group A)				

Table 3. Radionuclides of Ecological Importance

Scientists estimate that the radiation from the Chernobyl accident was 100x that of the two atom bombs dropped on Hiroshima and Nagasaki. It is estimated that the total atomospheric release was 5200 PBq (petabecquerel, 10¹⁵ Bq). The immediate death toll was 31 individuals though many more may die from the long term effects of radiation. The Soviets battled blazes at the Chernobyl power plant for two weeks. Those battling the fires were heroes in this author’s eyes because they knew they were exposing themselves to dangerous levels of radiation. Ultimately the Soviet authorities encased the Chernobyl reactor in concrete. A second more stable sarcophagus is currently being constructed over the original; its scheduled completion date is 2013.

There may have been additional unreported nuclear power plant accidents in the Soviet Union. Radioactive monitoring stations in Europe have picked up higher levels of radiation at various times which may have been the result of other Soviet nuclear power plant accidents.

The third and most recent nuclear power plant crisis occurred at the Fukushima Daiichi power plant in Japan. The cause of this disaster was a severe earthquake and tsunami on the 11th of March, 2011. The earth quake, which registered approximately 9 on the Richter Scale, was the event that set this tragedy in motion. The earthquake and resulting tsunami damaged the power plant compromising the cooling systems to the reactors causing the fuel rods to overheat. This disaster was rated greater than that at Three Mile Island. As of June 2011, the Fukushima disaster has released approximately one tenth the total amount of radiation as was released at Chernobyl. Unfortunately, the damaged Japanese reactor continues to spew forth radiation so the ultimate amount of radiation released from the plant cannot be determined with certainty.

10. Acknowledgements

The authors extend their thanks to the following individuals who assisted in the preparation of this chapter: Huizhong Xu, Associate professor of Physics, St. John’s University, who reviewed the physics section, to Natacha Lamarre, undergraduate biology major, St. John’s University, who typed the paper, prepared the tables and checked the references.

11. References

Auberg B., Hungate, F. P. (eds.). 1967. *Radioecological Concentration Processes*. Pergamon Press, Oxford. 1040 pp.

- Auerbach, S. I., and Crossley, D. A. 1958. Strontium-90 and cesium-137 uptake under natural conditions. Proc. Int. Conf. Peaceful Uses Atomic Energy, Geneva Paper No. 401.
- Ball, R. C., and Hooper, F. F. 1963. Translocation of phosphorus in a trout stream ecosystem. In: *Radioecology* (V. Schultz and A. W. Klement, eds.). Reinhold Publishing Company, New York. pp. 217-228.
- Baumhover, A. H., Graham A. J., Hopkins, D. E., Dudley, F. H., New, W. D., and Bushland, R. C. 1955. Screw-worm control through release of sterilized flies. J. Econ. Entomol., 48:462-466.
- Bryant, F. J., Chamberlain, A. C., Morgan, A., and Spicer, C. S. 1957. Radiostrontium in soil, grass, milk and bone in U.K.; 1956 results. J. Nuc. En., 6:22-40.
- Bushland, R. C. 1960. Male sterilization for the control of insects. In: *Advances in Pest Control Research* (R. L. Metcalf, ed.), Vol. III. John Wiley & Sons, Inc., New York.
- de la Cruz, A. A., and Wiegert, R. G. 1967. 32-Phosphorus tracer studies of a horse weed aphid-ant food chain. Amer. Midl. Nat., 77:501-509.
- Cutcomp, L. K. 1967. Progress in insect control by irradiation induced sterility. Pans, 13:61-70.
- Davis, J. J., and Foster, R. F. 1958. Bioaccumulation of radioisotopes through aquatic food chains. Ecology, 39:530-535.
- Donnelly, E. H., Nemhauser, J. B., Smith, J. M., Kazzi, Z. N. Farfan, E. B., Chang, A. S., and Naeem, S. F. 2010. Acute radiation syndrome: assessment and management. Southern Medical Journal. 103(6):541-546.
- Foster, R. F. 1958. Radioactive tracing of the movement of an essential element through an aquatic community with specific reference to radiophosphorus. Publ. della Stazione Zool. di Napoli.
- Foster, R. F., Davis, J. J. 1956. The accumulation of radioactive substances in aquatic forms. Proc. Int. Conf. Peaceful Uses Atomic Energy, Geneva, 13:364-367.
- Foster, R. F., and Rostenbach, R. E. 1954. Distribution of radioisotopes in the Columbian River. J. Amer. Water Works Assoc., 46:663-640.
- French, N. R. 1965. Radiation and animal population: problems, progress and projections. Health Physics, 11:1557-1568.
- Gager, C. S. 1907. Some effects of radioactivity on plants. Science 25:264.
- Gager, C. S. 1908. Effects of the rays of radium on plants. Mem. N. Y. Bot. Gard. 4:1-278.
- Gambino, J. J., and Lindberg, R. G. 1964. Response of the pocket mouse to ionizing radiation. Read. Res., 22:586-597.
- Glasstone, S. 1957. *The Effects of Nuclear weapons*. U. S. Atomic Energy Commission, Washington, D. C.
- Golley, F. B., Gentry, J. B., Menhinick, E., and Carmon, J. L. 1965. Response of wild rodents to acute gamma radiation. Rad. Res., 24:350-356.
- Goodspeed, T. H., and Uber, F. M. 1939. Radiation and plant cytogenetics. Botanical Review 5(1):1-48.
- Hanson, W. C. and Kornberg, H. A. 1956. Radioactivity in terrestrial animals near an atomic energy site. Proc. Int. Conf. Peaceful Uses Atomic Energy, Geneva 13:385-388.
- Heeney, H. B., and Rutherford, W. M. 1964. Some effects of gamma radiation on the storage life of fresh strawberries. Canadian Journal of Plant Science 44:188-194.
- Hungate, R. E. 1966. *The Rumen and Its Microbes*. Academic Press, New York. 533 pp.

- Klein, R. M., and Klein, D. T. 1971. Post-irradiation modulation of ionizing radiation damage to plants. *Botanical Review* 34(4):397-436.
- Knipling, E. F. 1960. The eradication of the screwworm fly. *Scient. Amer.*, 203(4):54-61.
- Knipling, E. F. 1963. The sterility principle. *Agr. Sci. Rev.*, 1(1):2.
- Knipling, E. F. 1965. The sterility method of pest population control. In: *Research in Pesticides* (G. O. Chichester, ed.). Academic Press, New York. Pp. 233-249.
- Knipling, E. F. 1967. Sterile technique, principles involved, current application, limitations and future applications. In: *Genetics of Insect Vectors of Disease* (Wright and Pal, eds.). Elsevier Publishing Co., Amsterdam. Pp. 587-616.
- Koernicke, M. 1905. Über die Wirkung von Röntgen- und Radiumstrahlen auf pflanzliche Gewebe und Zellen. *Ber. Deut. Bot. Ges.* 23:404-415.
- Lawson, F. R. 1967. Theory of control of insect population by sexually sterile males. *Ann. Entomol. Soc. Amer.*, 60:713-722.
- McCormick, F. J. 1969. Effects of ionizing radiation on a pine forest. In: *2nd Nat. Sym. Radioecology* (D. Nelson and F. Evans, eds.). Clearinghouse Fed. Sci. Tech. Info., U. S. Dept. Commerce, Springfield, VA. Pp. 78-87.
- McCormick, F. J., and Platt, R. B. 1962. Effects of ionizing radiation on a natural plant community. *Rad. Bot.*, 2:161-204.
- McCormick, F. J., and Golley, F. B. 1966. Irradiation of natural vegetation-an experimental facility, procedures and dosimetry. *Health Physics*, 12:1467-1474.
- Monk, C. D. 1966. Effects of short-term gamma irradiation on an old field. *Rad. Bot.*, 6:329-335.
- Nuttall, V. W.; Lyall, L. H.; and McQueen, K. F. 1961. Some effects of gamma radiation on stored onions. *Canadian Journal of Plant Science*, 41:805-813.
- Odum, E. P. 1971. *Fundamentals of Ecology*. W. B. Saunders Company, Philadelphia. Pp. 451-467.
- Odum, E. P. and Golley, F. B. 1963. Radioactive tracers as an aid to the measurement of energy flow at the population level in nature. In: *Radio ecology* (V. Shultz and A.W. Klement, eds.). Reinhold Publishing Company, New York. Pp. 403-410.
- Odum, E. P., and Kuenzler, E. J. 1963. Experimental isolation of food chains in an old field ecosystem with use of phosphorus-32. In: *Radio ecology* (V. Shultz and A.W. Klement, eds.). Reinhold Publishing Company, New York. pp. 113-120.
- Odum, H. T., and Pigeon, R.F. (eds.). 1970. *A tropical rainforest. A study of irradiation and ecology at El Verde, Puerto Rico*. Nat. Tech. Info, Service, Springfield, VA. 1678 pp.
- Odum, H. T., and Pigeon, R.F. 1965. Ionizing radiation and homeostasis of ecosystems. In: *Ecological Effects of Nuclear War* (Woodwell, ed.). Brookhaven National Laboratories, Publ. No. 917. Pp. 39-60.
- Ophel, I. L. 1963. The fate of radiostrontium in a freshwater community. In: *Radioecology* (V. Scultz and W. Klement, eds.), Reinhold Publishing Company, New York. Pp. 213-216.
- Platt, R. B. 1965. Ionizing Radiation and homeostasis of ecosystems. In: *Ecological Effects of Nuclear War* (Woodwell, ed.) Brookhaven National Library, Publ. No. 917:39-60.
- Polikarpou, G. C. 1966. *Radioecology of aquatic organisms*. (Translated from Russian by S. Technica and edited by Schultz and Klement).

- Pritchard, G. I., W. J. Pigden and D. J. Minson. 1962. Effect of gamma radiation on the utilization of wheat straw by rumen microorganisms. *Canadian Journal of Animal Science* 42:215-217.
- Rogovin, M. 1980. Three Mile Island: A report to the Commissioners and to the Public, Volume I. Nuclear Regulatory Commission, Special Inquiry Group.
- Smith, H. H. 1958. Radiation in the production of useful mutations. *Botanical Review* 24(1): 1-24.
- Sparrow, A. H. 1962. The role of the cell nucleus in determining radiosensitivity. Brookhaven Lecture Series No. 17. Brookhaven Nat. Lab. Publ. No. 766.
- Sparrow, A. H., and Evans, H. J. 1961. Nuclear factors affecting radiosensitivity. 1. The influence of nuclear size and structure, chromosome complement and DNA content. In: *Fundamental Aspects of Radiosensitivity*. Brookhaven Nat. Lab. Pp. 76-100.
- Sparrow, A. H.; Shairer, L. A.; and Sparrow, R. C. 1963. Relationship between nuclear volumes, chromosome numbers, and relative radiosensitivity. *Science*, 141:163-166.
- Sparrow, A. H., and Woodwell, G. M. 1962. Prediction of the sensitivity of plants to chronic gamma irradiation. *Rad. Bot.*, 2:9-26.
- Stalter, R., and Kincaid, D. T. 2009. Community development following gamma radiation at a pine-oak forest, Brookhaven National Laboratory, Long Island, New York. *American Journal of Botany* 96(12):2206-2213.
- Witherspoon, J. P. 1965. Radiation Damage to forest surrounding an unshielded fast reactor. *Health Physics*, 11:1637-1642.
- Witherspoon, J. P. 1969. Radiosensitivity of forest tree species to acute fast neutron radiation. In: *Proc. 2nd Natl. Sym. Radioecology* (D. Nelson and F. Evans, eds.). Clearinghouse Fed. Sci. Tech. Info., Springfield VA. Pp. 120-126.
- Woodwell, G. M. 1962. Effects of ionizing radiation on terrestrial ecosystems. *Science*, 138:572-577.
- Woodwell, G. M. (ed.). 1965. *Ecological Effects of Nuclear War*. Brookhaven National Laboratory Publ. no. 917, 72 pp.
- Woodwell, G. M. 1965a. Effects of ionizing radiation on ecological systems. In: *Ecological Effects of Nuclear War* (Woodwell, ed.). Brookhaven National Laboratory Publ. no. 917. Pp. 20-38.

Gamma Radiation

Edited by Prof. Feriz Adrovic

ISBN 978-953-51-0316-5

Hard cover, 320 pages

Publisher InTech

Published online 21, March, 2012

Published in print edition March, 2012

This book brings new research insights on the properties and behavior of gamma radiation, studies from a wide range of options of gamma radiation applications in Nuclear Physics, industrial processes, Environmental Science, Radiation Biology, Radiation Chemistry, Agriculture and Forestry, sterilization, food industry, as well as the review of both advantages and problems that are present in these applications. The book is primarily intended for scientific workers who have contacts with gamma radiation, such as staff working in nuclear power plants, manufacturing industries and civil engineers, medical equipment manufacturers, oncologists, radiation therapists, dental professionals, universities and the military, as well as those who intend to enter the world of applications and problems of gamma radiation. Because of the global importance of gamma radiation, the content of this book will be interesting for the wider audience as well.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Richard Stalter and Dianella Howarth (2012). Gamma Radiation, Gamma Radiation, Prof. Feriz Adrovic (Ed.), ISBN: 978-953-51-0316-5, InTech, Available from: <http://www.intechopen.com/books/gamma-radiation/gamma-radiation>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](https://creativecommons.org/licenses/by/3.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen