We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Solar Radiation in Tidal Flat

M. Azizul Moqsud
Kyushu University
Japan

1. Introduction

The Ariake Sea, which is located in the north-western part of Kyushu Island, is one of the best-known semi-closed shallow seas in Japan. Many rivers flow into the eastern coast area of the Ariake Sea and carry 4.4×10^8 kg of sediments per year (Azad et al. 2005). Coarse sediments accumulate in the eastern coast, and fine grains brought by the residual current accumulate in the bay head to form vast tidal flats with fine sediments (Kato and Seguchi 2001). The vast tidal flat mud of the Ariake Sea, which is almost 40% of the total tidal flat area of Japan, is famous for its rich fishery products and *Porphyra* sp. (sea weed) cultivation. Different types of shells like *Sinonovacula constricta*, *Atrina pectinata* and *Crassostrea gigas* are important creatures in the Ariake tidal mud. However, a dramatic decrease in the catch of these shells is observed in the tidal flat area. From Fig. 1 it is seen that the catch of *Crassostrea gigas* usually living in the near surface mud, dropped from 7.99×10^5 kg in 1976 to only 1.26×10^5 kg in 1999; that of *Atrina pectinata*, living in the upper 0.10-0.15 m of the mud, declined from 1.3395×10^7 kg in 1976 to 7.9×10^4 kg in 1999, and the situation in the case of *Sinonovacula constricta*, living in the depth of 0-0.7 m of the mud, was even worse: 1.7×10^5 kg catch in 1976 dropped to practically nil by 1992.

The acid treatment practice for *Porphyra* sp. cultivation is one of the major causes for this declination of the shells as this practice has made the geo-environmental condition of the Ariake tidal mud unfavorable for the living creatures of the tidal mud (Hayashi and Du, 2005, Moqsud et al. 2007). During the period of the cultivation (December -March), the acid (which is mainly organic chemicals) is used as the disinfectant acid to treat the *Porphyra* sp. cultivated in the sea and also to provide some nutrient phosphorus to it.

This organic acid provides ample of foods for the sulphate reducing bacteria living in the mud and consequently increase the sulfide content in the mud. The generation of sulfide is also influenced by the seasonal temperature and shows a higher value during the summer and the late autumn as bacteria becomes more active in the higher temperature. The higher sulfide content created by acid treatment practice is the main reason for the unfavorable condition for the benthos in the Ariake Sea. Moreover, the activities of the benthos depend strongly on the thermal environment near the sediment surface. Photosynthetic capacity of micro phytobenthos on an intertidal flat was strongly influenced by mud surface temperature (Blanchard et al. 1997). The filtration rate of bivalves was dependent on the water temperature (Hosokawa et al., 1996). As a result, to evaluate geo-thermal environment is important especially for the acid contaminated Ariake Sea. Thermal properties dictate the storage and movement of heat in soils and as such influence the temperature and heat flux

in soils as a function of time and depth (Anandkumar et. al, 2001). In recent years, considerable efforts have gone into developing techniques to determine these properties (Ochsner et al, 2001). The propagation of heat in a soil is governed by its thermal characteristics (De Vries, 1963). Main factors influencing soil thermal properties are mineralogical composition, the organic content and water content (De Vries, 1952, Wierenga et. al, 1969). No study has been carried out before to get the information about the thermal properties as well as thermal environment of the Ariake sea mud. So the objective of this study is to assess the thermal environment of the tidal mud by getting the information of the temperature distribution in different depths and find a diurnal and seasonal profile of it in the tidal flat region, and finally thermal properties variation with respect to depth for the temperature distribution in different seasons. The thermal properties of the Ariake Sea mud collected from both tidal flat and inside the deep sea of the Ariake Sea were conducted as a part of thermal environmental studies of the Ariake Sea.

Fig. 1. The graph of catch vs year

2. Sampling sites

Two sampling sites from tidal flat areas, sample 1(S1) and sample 2 (S2) and three sampling sites (sample 3 (S3), sample 4 (S4) and sample 5 (S5)) inside the Ariake Sea were selected as the study areas. Figure 2 shows the locations of the two tidal flat areas (Higashiyoka and Iida) and the three different areas inside the sea, along with the two types (pillar type and float type) of *Porphyra* sp. cultivation areas.

The tidal currents sweep into the sea and move northwards along the eastern shoreline and create a counterclockwise water movement. This would sweep the finer suspended particles delivered by rivers on the east side towards the inland end, where sedimentation would occur. Sediments in the Ariake Sea tidal flats are medium sand to silty mud. Medium sand, which accounts for 71% of the total tidal flats, is located mainly in the east and south coast areas (Azad et al. 2005). The silty mud is mainly in the bay head. Higashiyoka tidal flat located in the bay head was chosen as a study area (S1) which is near to Chikugo River (the biggest river in Kyushu Island), Okinohota River as well as other rivers and thought to be affected by the river waters. Another study area in tidal flat was Iida (S2), which seems to be

the most affected by the acid treatment practice. The other three study areas are chosen inside the Ariake Sea where all the time they are under water. The sample 1 and sample 2 (Higashiyoka and Iida) were collected during the ebb tide and the tidal flat was exposed to the sun directly. The other three mud samples (S3, S4, and S5 in Figure 2) were collected from under the sea water at different depths in different locations in the Ariake Sea. The sample collection was done in the last week of April 2006. The typical values of basic physicochemical properties of the mud samples collected from five study areas are tabulated in the Table 1. The mud samples were collected from the 0-0.2 m in the Ariake Sea.

Fig. 2. Map of the Ariake Sea & study area

3. Materials and methods

The vast tidal flat area of the Ariake Sea, which is 40 % of the total tidal area of Japan, is mainly muddy with high water content. The percentage of clay content is much higher than the sand or silt. To evaluate the temperature variation in different depths of the tidal mud, 5 numbers of thermocouples (Tokyo sokki kenkyojo Co., Ltd. Model no. N004853) were installed at 0.10 m, 0.20 m, 0.50 m, 1.0 m and 2.0 m depth which were connected with data logger (TDS- 530) to store the continuous hourly data of the temperature at Higashiyoka tidal flat mud. The sensors were placed about 20 m away from the shore line. The data loggers were kept in a watertight box and put in a small ship which was tied with some anchor and moved upward and downward during the high tide and ebb tide, respectively. Every two days the automatically stored data was collected from the data logger in the ship. This field investigation was carried out from 1st April, 2006 to 8th April, 2006 at Higashiyoka tidal flat.

In order to measure the seasonal temperature variation, the data were collected from both Iida and Higashiyoka tidal flat, 20 m away from the shore line during the ebb tide once in every month. By inserting the thermocouple (3 m long and 0.96 cm diameter) vertically into the tidal flat upto 3.0 m depth and at each 0.10 m interval the data was measured. The thermocouple was connected with a battery and a digital display. The temperature data was displayed directly in degree celcius. The mud samples from tidal flat were collected during the ebb tide and about 20 m distance from the shore line. The sample was then sliced into specified layers in the laboratory to measure various properties in each layer. The sulfide content was measured following the standard method prescribed by the Japan fisheries resource conservation association. The instrument which was used to measure the sulfide content is the GASTEC 201L/H which was also used by Wu et al. (2003) to determine the sulfide content of the marine sediments.

In-situ samples were collected by inserting vertically a thin wall steel tube sampler with a diameter of 0.07 m and a length of 0.90 m at five sites. For sample collection from tidal flat region an amphibious ship was used. The mud samples from tidal flat were collected during the ebb tide and about 40 m distance from the shore line. For sample collection from inside the sea, a ship was used. The ship was stopped in the predetermined location which was fixed by the global positioning system (GPS). The diver dived into the sea and collected the mud samples by inserting the steel tube into the sea bed floor and capped the two openings of the tube. The sample was then sliced into 0.05 m layers in the laboratory to measure the thermal properties in each layer.

The thermal properties analyzer KD2 Decagon Devices, Inc. was used to measure the thermal properties. Thermal conductivity and thermal diffusivity were measured directly from the thermal properties analyzer.

Physicochemical Parameters	S1	S2	S3	S4	S5
Density (x10 ⁻³ kg m ⁻³)	2.71	2.69	2.68	2.69	2.64
Water content (%)	168	235	160	239	253
Liquid limit <i>w_L</i> (%)	130	150	107	149	142
Plasticity index <i>I_p</i>	73	87	67	89	88
Ignition loss (%)	11.9	13.3	14.4	12.6	13.7
pH	8.03	7.92	7.60	7.53	7.59
ORP (mV)	-40.7	-121.4	128	130	46.38
Acid volatile sulphide (x10 ⁻³ kg kg ⁻¹ dry-mud)	0.16	0.42	0.14	0.30	0.49
Salinity(kg m ⁻³)	17	16	20	21	22
Grain size analysis(%)					
Sand	9	7	11	6	6
Silt	36	30	49	46	45
Clay	55	63	36	47	47

Table 1. Basic physicochemical properties of the samples

The volumetric heat capacity was calculated by the relation: Volumetric heat capacity = Thermal conductivity/thermal diffusivity.

4. Results and discussion

4.1 Daily variation of temperature

Figure 3 shows the variation of the tidal mud temperature at different depths from 1st April, 2006 to 8th April; 2006. It is seen that at 0.10 m and 0.20 m depth, the fluctuation of temperature was more prominent. However, from 0.50 m to 2.0 m depth, the diurnal variation was not so prominent. In the sub-surface region, the solar radiation affected the soil temperature more than the deeper part of the tidal mud.

This type of diurnal profile of temperature also agrees with the findings in Baeksu tidal flat in Korea (Yan-k et al. 2005). At 2.0 m depth, the temperature shows higher value than 1.0 m depth. This is probably due to the volumetric heat capacity of the tidal mud and the time lag for absorbing and releasing the heat during the summer and the winter. The peak temperature reached at different times at different depths. During the ebb tide, the time lag to reach the peak at different depths, is more than that at the high tide due to infiltration of sea water in the deeper depth.

Fig. 3. Variation of diurnal temperature with depth in the Higashiyoka tidal flat

Figure 4 illustrates one day (24h) variation of tidal flat mud temperature influenced by the solar radiation. It is seen that at 0.1 m depth, the peak value was reached when the solar radiation was also at the peak. At night, the temperature did not show any variation both during the ebb tide and the high tide time. This proves that the tidal mud temperature is only influenced by the solar radiation in the subsurface region. The tidal mud temperature at subsequent depths reaches the peak at different times, with the time lag increasing with depth. The peak temperature was reached about 2:00 PM and the value was about 17 °C at 0.10 m. The temperature at 0.50 m, 1.0 m and 2.0 m remained almost constant around 12-13 °C. It is concluded from this Figure that time lag increased with increasing depth but

the rate of increasing decreased with the increasing depth. Thermal properties of the tidal flat mud govern this type of phenomenon.

Fig. 4. Effects of solar radiation on the soil temperature in different depths

Fig. 5. Seasonal variation of temperature at Higashiyoka tidal flat

Fig. 6. Seasonal variation of temperature and consequently expansion and contraction tendency and acid treatment practice effects

4.2 Seasonal variation of temperature in tidal flat

Figure 5 shows that the seasonal variation of temperature at different depths at Higashiyoka tidal mud in 2007. During the spring and summer the surface temperature shows a higher value than the subsequent depths. During this time, heat was absorbed by the tidal mud and heat was transferred from the surface to the deeper part of the tidal mud. On the other hand, during winter and autumn the surface temperature was lower than the subsequent depths. During this time heat is released at the surface. During April, the variation was not so prominent. It showed almost straight line graph. Iida site also showed the same trend as with Higashiyoka site during the summer and the winter.

The acid treatment practice started during the winter season (December-February). In winter, the temperature drop down about 5⁰ C. Due to the lowering of temperature the tidal flat mud showed a contraction. The acid treatment practice and the contraction tendency of the tidal mud occur during the same time. As a result, the chemicals used in the sea laver treatment agent entered into the tidal mud. On the other hand, during the summer the surface temperature reached about 38⁰ C. The high temperature results in an expansion of the tidal mud. The tidal mud expansion causes easy movement of some biogenic gases generated inside the tidal mud.

4.3 Conceptual image of seasonal temperature effects and acid treatment practice on tidal flat

Figure 6 shows the conceptual image of the seasonal temperature variation and the acid treatment practice in the Ariake Sea. The various chemicals which are inside the sea laver treatment medicine enter into the tidal mud during the winter season due to the contraction effect of tidal mud. The chemicals and organic acid supply a lot of foods to the sulfate reducing bacteria. With this ample of foods and the convenient temperature during the spring and summer the sulfate reducing bacteria becomes very active and consequently produces hydrogen sulfide, sulfur di oxide, and as a result, the acid volatile sulfide (AVS) content increased. The AVS content at the Iida tidal flat area shows much higher than the safe limit for the living creatures in the tidal mud. The AVS represents a complex and dynamic biogeochemical system which is not defined simply by analysis of acid volatile

sulfide materials (Richard and Morse, 2005). Actually there are many factors which are liable to produce AVS in some specific regions. However, the laboratory test showed that due to the acid treatment practice the AVS value increased in the tidal flat mud (Moqsud et al. 2007). So the conceptual image of acid treatment practice and the seasonal variation of temperature are thought to be rational.

4.4 Proposed mechanisms of pore water movement

Figure 7 illustrates the conceptual image of the pore water movement in the tidal mud due to the seasonal variation of temperature. During spring and summer, the temperature at the shallow depth of the Iida tidal mud of the Ariake sea was higher than that of deeper depth, whereas opposite phenomenon was found during autumn and winter. The temperature gradient in the mud causes pore water to move in the vapor phase from a higher temperature site to a lower temperature site. The vapor condenses at the lower temperature area and becomes water, which increases the total head and drives the water liquid phase from lower temperature site to the higher temperature site (Nassar et al. 2000). Aforementioned process is titled coupled heat-pore water vapor-pore water liquid flow, as shown in Fig. 7.

Fig. 7. Proposed concept of coupled heat-pore water vapor-pore water liquid flow in tidal flat

5. Thermal properties

5.1 Thermal conductivity variation with depth

Figure 8 shows the variation of thermal conductivity at different depths in the Ariake sea. In the samples of tidal flats (sample 1 and sample 2), the variation is more prominent than the other samples collected from deep sea. This is probably due to much turbulent in the tidal flat mud in the region and introduces various kinds of matter during the tidal water movement as well as the direct exposure to the sun light during the ebb tide. All the samples show great variations in the sub surface (0-0.20 m) region but less variation in deeper region. Thermal conductivity of mud varies with soil texture, water content and organic matter content (Hamdeh and Reeder, 2000). The water content of the Ariake mud is always over 130% in different depths, which indicates that the conductivity of the Ariake mud is not affected by the water content at different depths.

Fig. 8. Variation of thermal conductivity with depth in the Ariake sea

5.2 Thermal diffusivity variation with depth

Figure 9 shows the variation of thermal diffusivity with depth for all the Ariake mud. It is seen that in the tidal flats (sample 1 and sample 2); the thermal diffusivity varied much at the different depths. On the other hand in the case of deep sea mud sample (sample 3, sample 4 and sample 5) the thermal diffusivity was constant at different depths. This is due to a small chance in turbulence in the deep sea bed floor. However, in the tidal flat area, during the low tide, the tidal mud is exposed directly to the sunlight, and during the ebb tide, a lot of foreign matters come and disturb the homogeneity in the mud of the tidal mud layers. It is seen that in the deep sea mud, the value of thermal diffusivity is always in $0.12 \times 10^{-6} \text{ m}^2/\text{s}$. In the tidal flat, the peak was reached at $0.13 \times 10^{-6} \text{ m}^2/\text{s}$ at different depths.

Fig. 9. Variation of thermal diffusivity with depth

5.3 Volumetric heat capacity variation with depth

The volumetric heat capacity of the tidal mud refers to the value which indicates the ability to store heat. If the volumetric heat capacity of a soil is high then the soil is more stable in terms of temperature change or the thermal environment. Figure 10 illustrates the variation of volumetric heat capacity with depth of the various samples. Sample 2 shows a great variation in volumetric heat capacity. The peak shows at 0.35 m depth and value is about 6.3 MJ/m³ °C. Clay soil generally has higher volumetric heat capacity than sandy soil for the same water content and soil density (Hamed, 2003). Volumetric heat capacity is very important for the acid infected tidal mud. Sulphate reducing bacteria (SRB) plays an important role in the geo-environmental condition of the Ariake Sea. These Bacteria like the layer where the volumetric heat capacity is higher (Moqsud et al.2006). Because in that layer it shows the more stable condition which is liked by the bacteria.

Fig. 10. Variation of volumetric heat capacity with depth

The temperature of underground soil is affected mainly by the soil thermal properties (Nassar et al., 2006) and these properties play a significant role in the geo-environmental condition in the global environment. The thermal properties of the mud are also induced by the mineralogical matter presence in the mud. The effects of this mineral matter on the thermal properties of the Ariake sea mud needs further study.

6. Conclusions

The temperature profiles, which result primarily from the molecular diffusion of heat through the sediment, resemble those typical of field soils and their form is similarly dependent on the thermal properties of the mud and the ambient meteorological conditions. The diurnal temperature variation is more visible near the surface (0.10 m and 0.20 m). The temperature increase gradually from morning, peak at noon and gradually decrease at afternoon. However, at 1.0 m and 2.0 m depth, the variation of temperature was not so prominent. This is due to the volumetric heat capacity and the thermal conductivity of the tidal mud. From the seasonal variation of temperature, it is seen that during late summer,

the surface and subsurface temperature is always higher than the deeper depth of the mud while in the winter the opposite phenomenon occurs. The thermal properties of mud collected from tidal flat showed a different trend from the mud collected inside the sea due to the exposure to the sunlight and the tidal wave turbulation in the tidal flat areas. In this study an innovative idea has been adopted to explain the deterioration and the natural remediation of the tidal flat mud in the Ariake Sea. The proposed mechanisms for understanding the transient seepage of pore water liquid of the tidal mud which contributes to the transport of sea laver treatment acid in the tidal mud and also natural remediation of contaminated tidal mud of the Ariake Sea was described clearly. The seasonal variation of temperature and the volumetric heat capacity of the mud have played a significant role for the maintaining the deterioration and natural remediation in the Ariake sea mud.

7. References

- Azad, K.A., Ohira, S.I., Oda, M. and Toda, K. (2005). On-site measurements of hydrogen sulphide and sulfur dioxide emissions from tidal flat sediments of the Ariake Sea, Japan. *Atmospheric Environment*, 39: 6077-6087.
- Anandkumar K., Venkatesan R. and Prabha T. "Soil thermal properties at Kalpakkam in coastal south India", *Earth Planet Science*, Vol.110,No. 3,2001,pp. 239-245.
- Ben-Dor,E and Banin A. "Determination of organic matter content in arid-zone soils using a simple loss-on-ignition method. *Commun".Soil SciencePlant Analysis*, Vol. 20,1989, pp.1675-1695
- Blanchard G.F.,Guarini P. and Richard P. "Seasonal effect on the relationship between the photosynthetic capacity of intertidal microphytobenthos and temperature". *Journal of Phycol.*Vol.33, 1997, pp.723-728
- De Vries D A. The thermal conductivity of soil. *Meded. Landbouwhogeschool*, Wageningen, 1952.
- De Vries D A. Thermal properties of soils. In: *physics of plant Environment*. Amsterdam, Holland, 1963.
- Ekwue E., Stone R.,Maharaj V. and Bhagwat D.. "Thermal conductivity and diffusivity of four Trinidadian soils as affected by peat content." *Transactions of the ASAE*. Vol.48, No.5, 2005, pp.1803-1815.
- Hamdeh H.N. "Thermal properties of soils as affected by density and water content." *Biosystems engineering*, Vol.86,No. 1,2003, pp. 97-102.
- Hamdeh H.N and Reeder R C. "Soil thermal conductivity: effects of density, moisture, salt concentration, and organic matter". *Soil science society of America Journal*, Vol.64, 2000, pp. 1285-1290.
- Hayashi S. And Du Y.J. (2005). Effect of acid treatment agent of sea laver on geo-environmental properties of tidal flat muds in the Ariake Sea. *Journal of ASTM International*, 3 (7), 52-59.
- Holmer, M and P. Storkholm,. "Sulphate reduction and sulphur cycling in lake sediments: a review." *Freshwater Biology*, 46,2001,pp. 431-451.
- Hosokawa Y, Kiebe E.,Miyoshi Y.,Kuwaie T. and Furukawa K.. "Distribution of aerial filtration rate of short-necked clam in coastal tidal flat." *The technical note of the port and harbour research institute*, Vol. 844, 1996, pp.21 (in Japanese with abstract in English).

- Kato, O. and Seguchi, M. (2001). The flow features and forming process of tidal flat in Ariake Sea. A report on the fifth international conference on the environmental management of enclosed coastal seas. EMECS 2001, 6-368.
- Moqsud, M.A., Hayashi, S., Du, Y.J. and Suetsugu D. (2007). Impacts of acid treatment practice on geo-environmental conditions in the Ariake Sea, Japan. *Journal of Southeast Asian geotechnical society*, 38 (2): 79-86.
- Moqsud M.A., Hayashi S., Du Y.J., Suetsugu D., Ushihara Y., Tanaka S. and Okuzono K. "Effects of acid treatment on geo-environmental conditions in the Ariake Sea, Japan." The 21st ICSW, March 26-29., Philadelphia, USA. 2006, pp.436-445.
- Moqsud M. A., Hayashi S., Du Y. J., Suetsugu D. "Evaluation of thermal properties of the mud of the Ariake Sea." International symposium on lowland technology, September 14-16. (2006 a) Saga, Japan, pp. 263-268.
- Nassar IN, Horton R, Flerchinger GN (2000) Simultaneous heat and mass transfer in soil columns exposed to freezing/thawing conditions. *Soil Sci.* 165 (3): 208-216.
- Naidu A. and Singh D.. "A generalized procedure for determining thermal resistivity of soils." *International Journal of thermal Sciences*. Vol.43, 2004, pp. 43-51.
- Ochsner T E., Horton R., Ren T. "A new perspective on soil thermal properties." *Soil science society of America Journal*, Vol. 65, 2001, pp.1641-1647.
- Rickard, D. and Morse, J.W. (2005). Acid volatile sulfide (AVS). *Marine chemistry*, 97:141-197.
- Sorensen, J. and Jorgenson, B.B. (1987). Early diagnosis in sediments from Danish coastal water: microbial activity and Mn-Fe-S geochemistry. *Geochimica et Cosmochimica Acta*, 51: 1883-1890.
- Wierenga P J., Nielsen D.R and Hagan R M. "Thermal properties of soil based upon field and laboratory measurements". *Soil science society of America Journal*, Vol.33, 1969, pp. 354-360.
- Wu, S.S., Tsutsumi, H., Kita-Tsukamoto, K., Kogure, K., Ohwada, K. and Wada, M. (2003). Visualization of the respiring bacteria in sediments inhabited by *Capitella* sp.1. *Fisheries Research*, 69:170-175.
- Yadav M R and Saxena G S.. "Effect of compaction and moisture content on specific heat and thermal capacity of soils". *Journal of Indian society of soil science*, Vol.21, 1973, pp. 129
- Yadav M R and Saxena G S.. "Effect of compaction and moisture content on specific heat and thermal capacity of soils". *Journal of Indian society of soil science*, Vol.21, 1973, pp. 129
- Yang-Ki C., Kim T., You, K., Park, L., Moon, H., Lee, S., Yoon, Y., (2005). "Temporal and spatial variability in the sediment temperature on the Baeksu tidal flat, Korea." *Estuarine coastal and shelf science*, Vol. 65, pp. 302-308.

Solar Radiation

Edited by Prof. Elisha B. Babatunde

ISBN 978-953-51-0384-4

Hard cover, 484 pages

Publisher InTech

Published online 21, March, 2012

Published in print edition March, 2012

The book contains fundamentals of solar radiation, its ecological impacts, applications, especially in agriculture, architecture, thermal and electric energy. Chapters are written by numerous experienced scientists in the field from various parts of the world. Apart from chapter one which is the introductory chapter of the book, that gives a general topic insight of the book, there are 24 more chapters that cover various fields of solar radiation. These fields include: Measurements and Analysis of Solar Radiation, Agricultural Application / Bio-effect, Architectural Application, Electricity Generation Application and Thermal Energy Application. This book aims to provide a clear scientific insight on Solar Radiation to scientist and students.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

M. Azizul Moqsud (2012). Solar Radiation in Tidal Flat, Solar Radiation, Prof. Elisha B. Babatunde (Ed.), ISBN: 978-953-51-0384-4, InTech, Available from: <http://www.intechopen.com/books/solar-radiation/solar-radiation-in-tidal-flat>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](https://creativecommons.org/licenses/by/3.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen