We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Analysis of SNARE-Mediated Exocytosis Using a Cell Fusion Assay

Chuan Hu, Nazarul Hasan and Krista Riggs
*Department of Biochemistry and Molecular Biology,
 University of Louisville School of Medicine, Louisville, KY
 USA*

1. Introduction

Exocytosis is the fusion of transport vesicles with the plasma membrane. By exocytosis, eukaryotic cells secrete soluble proteins and endogenous chemicals to the extracellular space, and deliver new membrane proteins and lipids to the plasma membrane. A large body of work has demonstrated that the interactions of SNARE (soluble N-ethylmaleimide-sensitive factor attachment protein receptor) proteins on vesicles (v-SNAREs) and on target membranes (t-SNAREs) catalyze intracellular vesicle fusion events, including exocytosis (Bonifacino and Glick, 2004; Jahn et al., 2003; Rothman, 1994) (Fig. 1). The vesicle-associated membrane proteins (VAMPs), *i.e.*, VAMPs 1, 2, 3, 4, 5, 7 and 8, are v-SNAREs that reside in various post-Golgi vesicular compartments, and have been implicated in exocytosis. In this chapter, we review recent progress of using a novel cell fusion assay to analyze the specificity and membrane fusion activities of VAMPs (Hasan et al., 2010).

Fig. 1. Interactions of VAMPs and plasma membrane t-SNAREs drive exocytosis.

2. SNAREs – Core machinery of vesicle fusion

SNAREs are cytoplasmic oriented type I membrane proteins. SNAREs share one homologous domain, the 'SNARE motif,' which contains eight heptad repeats ready for coiled-coil formation. The SNARE proteins that mediate the fusion of synaptic vesicles with the presynaptic plasma membrane are well studied (Sollner et al., 1993b). In synapses, the v-

SNARE VAMP2 resides in synaptic vesicles, whereas t-SNAREs syntaxin1 and synaptosomal-associated protein of 25 kD (SNAP-25) are located in the plasma membrane. Syntaxin1 and SNAP-25 constitute an acceptor complex for VAMP2 (Fasshauer and Margittai, 2004). The cytoplasmic domains of VAMP2, syntaxin1 and SNAP-25 form an extremely stable complex that is resistant to sodium dodecyl sulfate (SDS) (Hayashi et al., 1994) and heat stable up to $\sim 90^{\circ}\text{C}$ (Yang et al., 1999), indicating that SNARE complex formation is thermodynamically favorable. One α -helix from VAMP2, one α -helix from syntaxin1 and two α -helices from SNAP-25 intertwine to form a four-helix bundle (Sutton et al., 1998). Assembly of SNARE complexes is initiated at the N-termini and proceeds to the transmembrane domains at the C-termini in a zipper-like fashion (Stein et al., 2009). When v- and t-SNARE proteins are incorporated into liposomes, they spontaneously drive liposome fusion (McNew et al., 2000b; Weber et al., 1998), demonstrating that SNAREs form the minimal machinery for membrane fusion. Using a cell fusion assay, we showed that v- and t-SNARE proteins ectopically expressed on the cell surface spontaneously drive cell-cell fusion (Hu et al., 2003; Hu et al., 2007), providing further proof that SNAREs form the core machinery for intracellular membrane fusion. After membrane fusion, the adapter protein SNAP (soluble NSF attachment protein) and the ATPase NSF (N-ethylmaleimide-sensitive factor) dissociate v-/t-SNARE complexes at the expense of ATP (Mayer et al., 1996; Sollner et al., 1993a) to free SNAREs for the next round of fusion.

Genomic analysis indicates that there are 36 SNAREs in humans (Bock et al., 2001). Individual members of the SNARE family localize to distinct subcellular organelles (Chen and Scheller, 2001), suggesting that each SNARE has a selective role in vesicle trafficking. Using yeast SNARE proteins as models, a series of experiments showed that to a remarkable degree the specificity of intracellular membrane fusion can be predicted from the pattern of liposome fusion mediated by isolated v- and t-SNARE proteins (McNew et al., 2000a; Parlati et al., 2002). However, membrane fusion by SNAREs in mammalian cells is more promiscuous (Brandhorst et al., 2006; Shen et al., 2007). Here we show that with the exception of VAMP5, VAMPs are essentially redundant in mediating membrane fusion with plasma membrane t-SNAREs (Hasan et al., 2010).

3. Roles of VAMPs in exocytosis

VAMPs have been implicated in vesicle fusion with the plasma membrane, the *trans*-Golgi network (TGN) and endosomes. VAMP1 (synaptobrevin 1) and VAMP2 (synaptobrevin 2) mediate regulated exocytosis in neurons and endocrine cells (Hanson et al., 1997; Kesavan et al., 2007; Morgenthaler et al., 2003). In addition, VAMP2 is involved in the exocytosis of the water channel aquaporin 2 (Procino et al., 2008) and $\alpha 5\beta 1$ integrin (Hasan and Hu, 2010), as well as insulin-stimulated translocation of the glucose transporter GLUT4 (Randhawa et al., 2000). Enriched in recycling endosomes and endosome-derived vesicles (Galli et al., 1994; McMahon et al., 1993), VAMP3 (cellubrevin) mediates the recycling of transferrin receptors to the cell surface (Galli et al., 1994), integrin trafficking (Luftman et al., 2009; Proux-Gillardeaux et al., 2005; Skalski and Coppelino, 2005), and the secretion of α -granules in platelets (Feng et al., 2002; Polgar et al., 2002). Present primarily in the TGN, VAMP4 participates in the transport between the TGN and endosomes (Mallard et al., 2002; Steegmaier et al., 1999), as well as in homotypic fusion of early endosomes (Brandhorst et al., 2006). Expressed in muscle cells, VAMP5 (myobrevin) is associated with the plasma

membrane and intracellular vesicles (Zeng et al., 1998). In addition to vesicular transport from endosomes to lysosomes (Advani et al., 1999), the tetanus neurotoxin-insensitive VAMP (VAMP7) is involved in apical exocytosis in polarized epithelial cells (Galli et al., 1998; Pocard et al., 2007). Associated with early endosomes (Advani et al., 1998; Wong et al., 1998), VAMP8 (endobrevin) is required in regulated exocytosis in pancreatic acinar cells (Wang et al., 2004).

VAMPs have high sequence homology in their SNARE motifs (Fig. 2). All VAMPs possess a conserved arginine residue at the center of SNARE motifs (Fig. 2), and have been classified as R-SNAREs based on crystal structures (Fasshauer et al., 1998). The N-terminal 51 residues of VAMP4 contain a dominant signal for targeting to the TGN, while the SNARE motif of VAMP5 is responsible for its targeting to the plasma membrane (Zeng et al., 2003). In VAMP7, the N-terminal ‘longin domain’ regulates subcellular targeting (Pryor et al., 2008).

VAMPs 3, 4, 7 and 8 have broad tissue distribution (Advani et al., 1998; McMahon et al., 1993). Originally identified in nervous tissues, VAMPs 1 and 2 are also detected in skeletal muscle, fat and other tissues (Jagadish et al., 1996; Martin et al., 1998; Procino et al., 2008;

Fig. 2. Sequence alignment of human VAMP proteins. The conserved arginine residues in the center of SNARE motifs are labeled red.

Randhawa et al., 2000; Veale et al., 2010). Therefore, multiple VAMPs are co-expressed in mammalian cells. However, it is not clear if the seven VAMPs have differential membrane fusion activities. Using a cell fusion assay, we compare the membrane fusion activities of VAMPs.

3. Cell fusion assays

3.1 Expression of flipped SNAREs at the cell surface

By fusing the pre-prolactin signal sequence, which specifies translocation across the endoplasmic reticulum, to N-termini of the neuronal SNAREs VAMP2, syntaxin1 and SNAP-25, the cell fusion assay was originally developed in Dr. James Rothman's lab (Hu et al., 2003). The engineered SNAREs are called 'flipped' SNAREs because the orientation of their SNARE motifs against cellular membranes is flipped. A Myc tag is inserted between the signal sequence and N-termini of SNAREs to detect flipped SNARE proteins (Fig. 3A). When COS-7 cells are transfected with flipped SNARE constructs, flipped SNARE proteins are expressed at the cell surface (Fig. 3B). To express plasma membrane t-SNAREs, flipped syntaxins 1 or 4 are cotransfected with flipped SNAP-25. SNAP-25 does not contain a transmembrane domain. After expression, flipped SNAP-25 proteins are anchored to the cell surface by assembling with flipped syntaxin proteins.

Flow cytometry is used to measure the expression levels of SNARE proteins at the cell surface (Figs. 3C and D). When flipped VAMPs 1, 3, 4, 5, 7 and 8, and syntaxins 1 and 4 plasmids are transfected at the same concentration, cell surface expression of VAMPs 5 and 8 is higher than VAMPs 1, 3, 4 and 7, and cell surface expression of syntaxin4 is higher than syntaxin1 (Hasan et al., 2010). To express the v- and t-SNAREs at the same level, we optimized the concentration of each flipped SNARE plasmid used in transfection (concentrations see Section 4.2), so that VAMPs 1, 3, 4, 5, 7 and 8, and syntaxins 1 and 4 are expressed at the same level at the cell surface, respectively (Fig. 3D).

3.2 Microscopic cell fusion assay

Multiple readout systems have been developed to detect fusion of the cells that express flipped v-SNAREs (v-cells) and the cells that express flipped t-SNARE proteins (t-cells) (Hasan et al., 2010; Hu et al., 2003; Hu et al., 2007). In the microscopic assay shown in Fig. 4A, flipped v-SNARE constructs are cotransfected with a plasmid that encodes the green fluorescent protein EGFP. In t-cells, the flipped t-SNARE constructs are cotransfected with a plasmid that encodes the red fluorescent protein DsRed2. Fusion of the v- and t-cells results in fused cells that contain both EGFP and DsRed2. In the merged channel, the cytoplasm of the fused cells is yellow (arrows, Fig. 4A).

3.3 Enzymatic cell fusion assay

The microscopic cell fusion assay becomes less efficient when used to analyze multiple v-/t-SNARE combinations quantitatively. To develop a quantitative cell fusion assay, we take advantage of the strong transcriptional activation by binding of the tetracycline-controlled transactivator (tTA) to the tetracycline-response element (TRE) (Gossen and Bujard, 1992). Two plasmids in CLONTECH's Tet-Off gene expression system are used (Hasan et al., 2010). The first plasmid pTet-Off encodes the transcriptional activator tTA, and the second

Fig. 3. Expression of flipped SNAREs at the cell surface. (A) Domain structure of flipped SNAREs. (B) Twenty-four hours after transfection with empty vector pcDNA3.1(+) or flipped VAMP5 plasmid, unpermeabilized COS-7 cells are stained with an anti-Myc antibody. Representative confocal images are shown. Scale bar, 50 μ m. (C and D) Twenty-four hours after transfection with empty vector or flipped SNARE plasmids, unpermeabilized cells are stained with the anti-Myc antibody and analyzed by flow cytometry. (C) Representative FACS profiles of the cells transfected with empty vector or flipped VAMP1. (D) To express VAMPs and syntaxins at the same level at the cell surface, flipped SNARE plasmids are transfected at titrated concentrations. The mean fluorescence intensity of staining is obtained using CellQuest Pro software.

Fig. 4. Cell fusion assays. (A) Microscopic cell fusion assay. Cells that express flipped v-SNAREs (v-cells) are labeled by the green fluorescent protein EGFP, whereas cells that express flipped t-SNAREs (t-cells) are labeled by the red fluorescent protein DsRed2. Fusion of v- and t-cells results in fused cells (arrows) whose cytoplasm is yellow under fluorescence microscope. Scale bar, 50 μ m. (B) Enzymatic cell fusion assay. The tetracycline-controlled transactivator (tTA) is expressed in v-cells, and a reporter plasmid that encodes β -galactosidase under control of the tetracycline-response element (TRE-LacZ) is transfected into t-cells. Fusion of the v- and t-cells leads to the binding of tTA to TRE and the expression of β -galactosidase, which is measured using a colorimetric method by absorbance at 420 nm. Only baseline β -galactosidase activity is detected when either flipped VAMP2 or SNAP-25 is not expressed.

plasmid pBI-G encodes the *LacZ* gene under control of the tetracycline-response element (TRE-*LacZ*). In the absence of tTA, transcription of the *LacZ* gene in TRE-*LacZ* is silent. When tTA is present, it binds to the TRE and activates the transcription of *LacZ*, resulting in the expression of β -galactosidase. We hypothesize that if tTA is located in v-cells and TRE-*LacZ* is located in t-cells, β -galactosidase will not be expressed. Fusion of the v- and t-cells would result in the binding of tTA to TRE and transcriptional activation of *LacZ* (Fig. 4B).

The neuronal SNAREs are used to test feasibility of the assay. VAMP2 is coexpressed with tTA in v-cells, and syntaxin1 and SNAP-25 are coexpressed with TRE-*LacZ* in t-cells. When the v- and t-cells are combined, robust β -galactosidase expression is indeed detected (Fig. 4B). However, when either VAMP2 or SNAP-25 is not expressed, only baseline β -galactosidase activity is detected, indicating that cell fusion and expression of β -galactosidase rely on interactions of v- and t-SNAREs.

3.4 Experimental procedures

3.4.1 Cell culture and reagents

COS-7 cells were obtained from the American Type Culture Collection, and cultured in Dulbecco Modified Eagle's Medium (DMEM) supplemented with 4.5 g/l glucose and 10% fetal bovine serum (FBS). The anti-Myc monoclonal antibody 9E10, developed by Dr. Bishop, was obtained from the Developmental Studies Hybridoma Bank maintained by the University of Iowa. pEGFP-N3, pDsRed2-N1, pTet-Off and pBI-G were obtained from CLONTECH. Plasmid transfection is done with Lipofectamine according to the manufacturer's instructions (Invitrogen).

3.4.2 FACS analysis

Expression levels of SNAREs at the cell surface are measured using immunostaining and flow cytometry as we previously reported (Hasan and Hu, 2010). Briefly, COS-7 cells are seeded in 6-well plates. Twenty-four hours after transfection with the flipped SNARE, pTet-Off and pBI-G plasmids, cells are fixed with 1% paraformaldehyde in PBS++ (PBS supplemented with 0.1 g/l CaCl_2 and 0.1 g/l MgCl_2). After labeling with the anti-Myc monoclonal antibody 9E10 and FITC-conjugated secondary antibodies, the cells are scraped off the plates with a cell scraper. 15,000 cells are analyzed using a FACSCalibur flow cytometer (BD Biosciences). The mean fluorescence intensity of each sample is obtained using CellQuest Pro software.

3.4.3 Microscopic cell fusion assay

The day before transfection, 1.2×10^6 COS-7 cells are seeded in each 100-mm cell culture dish, and 5×10^4 COS-7 cells are seeded on sterile 12-mm glass coverslips contained in 24-well plates. For v-cells, 5 μg of flipped v-SNARE and pEGFP-N3 are cotransfected into the cells grown in each 100-mm culture dish. For t-cells, 0.25 μg each of flipped syntaxin, SNAP-25 and pDsRed2-N1 are cotransfected into the cells seeded in the 24-well plates. Twenty-four hours after transfection, the v-cells are detached from culture dishes with EDTA (Enzyme-free Cell Dissociation Buffer (Invitrogen)). Detached cells are counted with a hemocytometer and resuspended in HEPES-buffered DMEM supplemented with 10% FBS,

6.7 $\mu\text{g/ml}$ tunicamycin and 0.67 mM DTT. v-Cells (1.2×10^5) are added to each coverslip already containing the t-cells. After 24 hours at 37°C in 5% CO_2 , the cells are gently washed once with PBS++, then fixed with 4% paraformaldehyde. Confocal images are collected on an Olympus laser scanning confocal microscope. The 488 nm argon laser line is used to excite EGFP and the 543 nm HeNe laser line is used to excite DsRed2. To prevent cross-contamination between EGFP and DsRed2, each channel is imaged sequentially before merging the images.

3.4.4 Enzymatic cell fusion assay

The day before transfection, 1.2×10^6 COS-7 cells are seeded in each 100-mm cell culture dish, and 2×10^5 COS-7 cells are seeded in each well of 6-well plates. For v-cells, 5 μg each of flipped v-SNARE plasmid are cotransfected with 5 μg of pTet-Off into the cells in each 100-mm culture dish. Control cells are cotransfected with empty vector pcDNA3.1(+) and pTet-Off. For t-cells, 1 μg each of flipped syntaxins 1 or 4, SNAP-25 and pBI-G are cotransfected into the cells in each well of the 6-well plates. There are putative N-glycosylation motifs (Asn-X-Ser/Thr) in VAMPs. Our previous studies (Hu et al., 2003; Hu et al., 2007) showed that N-glycosylation disrupts the function of flipped SNAREs, and that treatment of tunicamycin, an antibiotic that inhibits N-glycosylation (Elbein, 1984), restores the fusion activities of flipped SNAREs. To prevent N-glycosylation of VAMPs 1, 4, 5, 7 and 8, v-cells expressing these VAMP proteins and control cells are incubated in cell culture medium containing 10 $\mu\text{g/ml}$ of tunicamycin during transfection. Since we have mutated the putative N-glycosylation sites in flipped VAMP2 (Hu et al., 2003) and VAMP3 proteins (Hu et al., 2007), v-cells that express VAMPs 2 or 3 are not treated with tunicamycin. Since flipped syntaxins 1 or 4, SNAP-25 proteins are not N-glycosylated (Hu et al., 2003; Hu et al., 2007), the t-cells are not treated with tunicamycin.

Twenty-four hours after transfection, the v-cells are detached from the culture dishes with Enzyme-free Cell Dissociation Buffer, and added (4.8×10^5 cells) to each well already containing the t-cells. After 6, 12 or 24 hours at 37°C in 5% CO_2 , the expression of β -galactosidase is measured using the β -Galactosidase Enzyme Assay System with Reporter Lysis Buffer according to the manufacturer's instructions (Promega). The cells are washed twice with PBS, and then lysed in Reporter Lysis Buffer. Cell lysates are mixed with equal volume of Assay 2 $\times$ Buffer. As a blank control, Reporter Lysis Buffer is mixed with Assay 2 $\times$ Buffer. After 90 minutes, the colorimetric reaction is stopped by adding 1 M sodium carbonate, and absorbance at 420 nm is measured using a HITACHI 100-40 spectrophotometer.

4. Membrane fusion by VAMPs and plasma membrane t-SNAREs

4.1 Fusogenic interactions of VAMPs and plasma membrane t-SNAREs

Using the enzymatic fusion assay (Fig. 4B), we examine the fusogenic pairings between the seven VAMPs and two plasma membrane t-SNARE complexes, syntaxin1/SNAP-25 (Fasshauer and Margittai, 2004) and syntaxin4/SNAP-25 (Reed et al., 1999). Robust β -galactosidase expression is detected when VAMPs 1, 2, 3, 4, 7 or 8 are combined with syntaxin1/SNAP-25 (Fig. 5A) or syntaxin4/SNAP-25 (Fig. 5B), indicating that these VAMPs mediate membrane fusion with plasma membrane t-SNAREs. With syntaxin1/SNAP-25, the

six VAMPs drive fusion to a similar degree. With syntaxin4/SNAP-25, VAMP8 fuses less efficiently than VAMPs 1, 2, 3 and 4 (31% lower fusion activity and $P = 0.046$ vs. VAMP1, Fig. 5B). In contrast, when VAMP5 is combined with the t-SNAREs, only baseline β -galactosidase activity is detected (Figs. 5A and B), suggesting that VAMP5 does not drive membrane fusion with the t-SNAREs. The stronger fusion activities of syntaxin4/SNAP-25 than syntaxin1/SNAP-25 (compare Figs. 5A and B) are likely caused by higher cell surface expression of syntaxin4/SNAP-25 than syntaxin1/SNAP-25 and higher fusion activity of syntaxin4 than syntaxin1 (Hasan et al., 2010). Together, the data shown in Fig. 5 indicate that VAMPs 1, 2, 3, 4, 7 and 8, but not VAMP5, drive membrane fusion when partnering with plasma membrane t-SNAREs.

Fig. 5. Cell fusion by VAMPs and plasma membrane t-SNAREs. Twenty-four hours after combining v-cells that express VAMPs 1, 2, 3, 4, 5, 7 or 8 and t-cells that express (A) syntaxin1/SNAP-25 or (B) syntaxin4/SNAP-25, cell fusion is quantified using the enzymatic cell fusion assay. Control cells (-VAMP) are transfected with the empty vector. The flipped SNARE plasmids are transfected at the same concentration. Error bars represent standard deviation of three independent experiments. * $P < 0.05$ vs. VAMP1.

VAMPs 1, 2, 3, 7 and 8 are known to drive vesicle fusion with the plasma membrane. The above data provide additional evidence that vesicles that carry either one of these five VAMPs are capable of fusing with the plasma membrane. Since these VAMPs are functionally redundant, they may compensate each other in loss of function studies (Borisovska et al., 2005).

To rule out the possibility that the baseline β -galactosidase activity detected in the VAMP5 combinations (Fig. 5) is caused by residual membrane fusion activity of VAMP5, we perform the microscopic cell fusion assay, which analyzes individual cell fusion events and can detect rare fusion events. In multiple experiments, no cell fusion is observed in the microscopic assay when VAMP5 is combined with syntaxin1/SNAP-25 or syntaxin4/SNAP-25, whereas VAMP4 drives fusion efficiently with both t-SNARE complexes (Hasan et al., 2010) (data not shown). Together, the results using enzymatic and microscopic cell fusion assays suggest that VAMP5 is unable to mediate membrane fusion with the plasma membrane t-SNAREs.

VAMP5 is expressed in muscle cells, in which it is mainly associated with the plasma membrane as well as intracellular vesicles (Zeng et al., 1998; Zeng et al., 2003). It is still possible that VAMP5 forms fusogenic interactions with other plasma membrane t-SNAREs to mediate exocytosis in muscle cells. Indeed, SNAP-23, a ubiquitously expressed SNAP-25 homolog, is expressed in skeletal muscle (Bostrom et al., 2010). The role of VAMP5 in exocytosis needs further investigation.

4.2 Differential membrane fusion activities of VAMPs

Do VAMPs have differential membrane fusion capacities? An ideal experimental system to answer this question will require equal expression of VAMP proteins and a quantitative membrane fusion assay. Using anti-Myc staining and flow cytometry as measurement (Figs. 3C and D), we express VAMP proteins at the same level at the cell surface. Furthermore, using the quantitative enzymatic cell fusion assay, we compare their membrane fusion activities.

When COS-7 cells are transfected with the flipped SNARE plasmids at the same concentration, cell surface expression of VAMPs 5 and 8 is more than 2 fold higher than VAMPs 1, 3, 4 and 7, and cell surface expression of syntaxin4 is 1.8 fold higher than syntaxin1 (Hasan et al., 2010) (data not shown). To express the v- and t-SNAREs at the same level, we titrate and optimize the concentration of each flipped SNARE plasmid used in transfection. Flipped SNARE plasmids are transfected at the following concentrations (per 10 cm² growth area, *i.e.*, per well in 6-well plates): VAMP1, 0.2 μ g; VAMP3, 0.5 μ g; VAMP4, 0.5 μ g; VAMP5, 0.05 μ g; VAMP7, 1.0 μ g; VAMP8, 0.1 μ g; syntaxin1, 0.5 μ g; syntaxin4, 0.05 μ g. tTA, TRE-LacZ and flipped SNAP-25 are cotransfected at 1 μ g per 10 cm² growth area. Under such conditions, VAMPs 1, 3, 4, 5, 7 and 8 are expressed at same level at the cell surface, while syntaxins 1 and 4 are expressed at the same level (Fig. 3D). Because the flipped VAMP2 protein does not contain a Myc tag (Hu et al., 2003), we are not able to compare its expression level with the other VAMPs.

After expressing VAMPs 1, 3, 4, 5, 7 and 8 at the same level, we compare their membrane fusion activities using the enzymatic cell fusion assay. With syntaxin1/SNAP-25, VAMPs 1, 3, and 8 have comparable and the highest fusion activities, whereas VAMPs 4 and 7 have 50% and 30% lower fusion activities, respectively (Fig. 6A). With syntaxin4/SNAP-25,

VAMPs 1 and 3 have comparable and the highest fusion activities, whereas VAMPs 4, 7 and 8 have 36%, 26% and 54% lower fusion activities, respectively (Fig. 6B). As expected, only baseline β -galactosidase activity is detected when VAMP5 is paired with the t-SNAREs. These data indicate that VAMPs have differential membrane fusion activities with plasma membrane t-SNAREs. However, when expressed at higher levels, VAMP4 drives membrane fusion as efficiently as VAMPs 1 and 3 (Fig. 5). Since the differences of fusion activities among the fusogenic VAMPs are within a factor of 2, these results imply that with the exception of VAMP5, VAMPs are essentially redundant in mediating membrane fusion with plasma membrane t-SNAREs.

Fig. 6. Comparison of fusion activities of VAMPs. With optimized transfection conditions, VAMPs 1, 3, 4, 5, 7 and 8 are expressed at the same level at the cell surface, while syntaxin1/SNAP-25 (A) and syntaxin4/SNAP-25 (B) are expressed at the same level. Cell fusion is quantified using the enzymatic fusion assay. The fusion activities (OD_{420}) of control cells (-VAMP) and the v-cells expressing VAMPs 3, 4, 5, 7 or 8 are normalized to the fusion activity of the v-cells expressing VAMP1. Error bars represent standard deviation of four independent experiments.

4.3 Cooperativity of VAMP proteins in the cell fusion reaction

The number of SNARE complexes that cooperate to mediate vesicle fusion is under active investigation. To investigate the cooperativity of VAMPs in membrane fusion, we determine the dependence of cell fusion activity on cell surface expression level of VAMP1. We choose VAMP1 in this experiment because it has high membrane fusion activity (Fig. 6). In t-cells, the cell surface expression levels of syntaxin1 and SNAP-25 are kept constant. v-Cells are transfected with increasing concentrations of the flipped VAMP1 plasmid. At each concentration, we measure the cell surface expression level of VAMP1 proteins using flow cytometry, and determine cell fusion activity of VAMP1 with syntaxin1/SNAP-25 using the enzymatic fusion assay. Cell fusion activity is then plotted as a function of the mean fluorescence intensity of VAMP1 staining (Fig. 7A). The correlation is best fit with a polynomial regression. The hyperbolic instead of sigmoidal correlation (Fig. 7A) suggests that there is no cooperativity of VAMP1 proteins in driving cell fusion.

Fig. 7. Dependence of cell fusion activity on cell surface density of VAMP1. (A) v-cells expressing increasing amount of VAMP1 at the cell surface are combined with t-cells expressing syntaxin1/SNAP-25. Cell fusion activities are quantified and correlated with the mean fluorescence intensity of VAMP1 staining. (B) Log-log plot of cell fusion activity vs. mean fluorescence intensity of VAMP1 staining.

The log-log plot has been used to determine the cooperativity of viral fusion proteins in membrane fusion (Danieli et al., 1996). Using log-log plot, we further analyze the cooperativity of VAMP1 proteins. If two VAMP1 (V) proteins take part in the cell fusion reaction, *i.e.*, $V + V \rightarrow \text{Fusion}$, the rate of fusion (F) = $k [V]^2$. Therefore, $\log (F) = \log (k) + 2 \log [V]$, and the slope of the resulting log-log plot will be 2. A log-log analysis of the dependence of cell fusion activity on VAMP1 cell surface density is shown in Fig. 7B. Linear regression is performed to model the log-log correlation (Fig. 7B), and the resulting slope is 0.52. These results further suggest that the cell fusion reaction does not involve concerted action of VAMP1 proteins.

Previous studies have estimated that 1 to 11 SNARE complexes are needed for membrane fusion (Domanska et al., 2009; Hua and Scheller, 2001; Karatekin et al., 2010; van den Bogaart et al., 2010). In the cell fusion reaction, we do not observe cooperativity of VAMP1 proteins, suggesting that concerted action of multiple SNARE complexes is not required to fuse cellular membranes. However, to achieve fast exocytosis in intact cells, concerted action of multiple SNARE complexes is clearly needed (Mohrmann et al., 2010). Such cooperativity of SNARE complexes may be organized by the binding of regulatory proteins such as synaptotagmins and Munc18 (Chicka et al., 2008; Shen et al., 2007).

5. Conclusion

To examine v-/t-SNARE interactions quantitatively, we developed an enzymatic cell fusion assay that utilizes activated expression of β -galactosidase and spectrometric measurement. Using this assay, we show that VAMPs 1, 2, 3, 4, 7 and 8 mediate membrane fusion efficiently with plasma membrane t-SNAREs syntaxin1/SNAP-25 and syntaxin4/SNAP-25, whereas VAMP5 does not drive fusion with the t-SNAREs. By expressing VAMPs 1, 3, 4, 7 and 8 at the same level, we further compare their membrane fusion activities. VAMPs 1 and 3 exhibit comparable and the highest fusion activities, whereas VAMPs 4, 7 and 8 have 30 - 50% lower fusion activities. Collectively, these data indicate that VAMPs have differential membrane fusion activities, and imply that with the exception of VAMP5, VAMPs are essentially redundant in mediating membrane fusion with plasma membrane t-SNAREs. Furthermore, no cooperativity of VAMP1 proteins is observed in the cell fusion reaction, suggesting that concerted action of multiple SNARE complexes is not required to fuse cellular membranes.

6. Acknowledgement

We thank Adrienne Bushau and David Humphrey for critically reading the manuscript. This work was supported by startup funds from the University of Louisville and CA135123 from the National Institutes of Health (to C.H.).

7. References

- Advani, R.J., Bae, H.R., Bock, J.B., Chao, D.S., Doung, Y.C., Prekeris, R., Yoo, J.S., and Scheller, R.H. (1998). Seven novel mammalian SNARE proteins localize to distinct membrane compartments. *Journal of Biological Chemistry*, Vol.273, pp. 10317-10324.

- Advani, R.J., Yang, B., Prekeris, R., Lee, K.C., Klumperman, J., and Scheller, R.H. (1999). VAMP-7 mediates vesicular transport from endosomes to lysosomes. *Journal of Cell Biology*, Vol.146, pp. 765-776.
- Bock, J.B., Matern, H.T., Peden, A.A., and Scheller, R.H. (2001). A genomic perspective on membrane compartment organization. *Nature*, Vol.409, pp. 839-841.
- Bonifacino, J.S., and Glick, B.S. (2004). The mechanisms of vesicle budding and fusion. *Cell*, Vol.116, pp. 153-166.
- Borisovska, M., Zhao, Y., Tsytsyura, Y., Glyvuk, N., Takamori, S., Matti, U., Rettig, J., Sudhof, T., and Bruns, D. (2005). v-SNAREs control exocytosis of vesicles from priming to fusion. *EMBO Journal*, Vol.24, pp. 2114-2126.
- Bostrom, P., Andersson, L., Vind, B., Haversen, L., Rutberg, M., Wickstrom, Y., Larsson, E., Jansson, P.A., Svensson, M.K., Branemark, R., Ling, C., Beck-Nielsen, H., Boren, J., Hojlund, K., and Olofsson, S.O. (2010). The SNARE protein SNAP23 and the SNARE-interacting protein Munc18c in human skeletal muscle are implicated in insulin resistance/type 2 diabetes. *Diabetes*, Vol.59, pp. 1870-1878.
- Brandhorst, D., Zwillig, D., Rizzoli, S.O., Lippert, U., Lang, T., and Jahn, R. (2006). Homotypic fusion of early endosomes: SNAREs do not determine fusion specificity. *Proceedings of the National Academy of Sciences of the United States of America*, Vol.103, pp. 2701-2706.
- Chen, Y.A., and Scheller, R.H. (2001). SNARE-mediated membrane fusion. *Nature Reviews Molecular Cell Biology*, Vol.2, pp. 98-106.
- Chicka, M.C., Hui, E., Liu, H., and Chapman, E.R. (2008). Synaptotagmin arrests the SNARE complex before triggering fast, efficient membrane fusion in response to Ca^{2+} . *Nature Structural and Molecular Biology*, Vol.15, pp. 827-835.
- Danieli, T., Pelletier, S.L., Henis, Y.I., and White, J.M. (1996). Membrane fusion mediated by the influenza virus hemagglutinin requires the concerted action of at least three hemagglutinin trimers. *Journal of Cell Biology*, Vol.133, pp. 559-569.
- Domanska, M.K., Kiessling, V., Stein, A., Fasshauer, D., and Tamm, L.K. (2009). Single vesicle millisecond fusion kinetics reveals number of SNARE complexes optimal for fast SNARE-mediated membrane fusion. *Journal of Biological Chemistry*, Vol.284, pp. 32158-32166.
- Elbein, A.D. (1984). Inhibitors of the biosynthesis and processing of N-linked oligosaccharides. *CRC Critical Reviews in Biochemistry*, Vol.16, pp. 21-49.
- Fasshauer, D., and Margittai, M. (2004). A transient N-terminal interaction of SNAP-25 and syntaxin nucleates SNARE assembly. *Journal of Biological Chemistry*, Vol.279, pp. 7613-7621.
- Fasshauer, D., Sutton, R.B., Brunger, A.T., and Jahn, R. (1998). Conserved structural features of the synaptic fusion complex: SNARE proteins reclassified as Q- and R-SNAREs. *Proceedings of the National Academy of Sciences of the United States of America*, Vol.95, pp. 15781-15786.
- Feng, D., Crane, K., Rozenvayn, N., Dvorak, A.M., and Flaumenhaft, R. (2002). Subcellular distribution of 3 functional platelet SNARE proteins: human cellubrevin, SNAP-23, and syntaxin 2. *Blood*, Vol.99, pp. 4006-4014.
- Galli, T., Chilcote, T., Mundigl, O., Binz, T., Niemann, H., and De Camilli, P. (1994). Tetanus toxin-mediated cleavage of cellubrevin impairs exocytosis of transferrin receptor-containing vesicles in CHO cells. *Journal of Cell Biology*, Vol.125, pp. 1015-1024.

- Galli, T., Zahraoui, A., Vaidyanathan, V.V., Raposo, G., Tian, J.M., Karin, M., Niemann, H., and Louvard, D. (1998). A novel tetanus neurotoxin-insensitive vesicle-associated membrane protein in SNARE complexes of the apical plasma membrane of epithelial cells. *Molecular Biology of the Cell*, Vol.9, pp. 1437-1448.
- Gossen, M., and Bujard, H. (1992). Tight control of gene expression in mammalian cells by tetracycline-responsive promoters. *Proceedings of the National Academy of Sciences of the United States of America*, Vol.89, pp. 5547-5551.
- Hanson, P.I., Heuser, J.E., and Jahn, R. (1997). Neurotransmitter release - four years of SNARE complexes. *Current Opinion in Neurobiology*, Vol.7, pp. 310-315.
- Hasan, N., Corbin, D., and Hu, C. (2010). Fusogenic Pairings of Vesicle-Associated Membrane Proteins (VAMPs) and Plasma Membrane t-SNAREs - VAMP5 as the Exception. *PLoS One*, Vol.5, pp. e14238.
- Hasan, N., and Hu, C. (2010). Vesicle-associated membrane protein 2 mediates trafficking of alpha5beta1 integrin to the plasma membrane. *Experimental Cell Research*, Vol.316, pp. 12-23.
- Hayashi, T., McMahon, H., Yamasaki, S., Binz, T., Hata, Y., Sudhof, T.C., and Niemann, H. (1994). Synaptic vesicle membrane fusion complex: action of clostridial neurotoxins on assembly. *EMBO Journal*, Vol.13, pp. 5051-5061.
- Hu, C., Ahmed, M., Melia, T.J., Sollner, T.H., Mayer, T., and Rothman, J.E. (2003). Fusion of cells by flipped SNAREs. *Science*, Vol.300, pp. 1745-1749.
- Hu, C., Hardee, D., and Minnear, F. (2007). Membrane fusion by VAMP3 and plasma membrane t-SNAREs. *Experimental Cell Research*, Vol.313, pp. 3198-3209.
- Hua, Y., and Scheller, R.H. (2001). Three SNARE complexes cooperate to mediate membrane fusion. *Proceedings of the National Academy of Sciences of the United States of America*, Vol.98, pp. 8065-8070.
- Jagadish, M.N., Fernandez, C.S., Hewish, D.R., Macaulay, S.L., Gough, K.H., Grusovin, J., Verkuylen, A., Cosgrove, L., Alafaci, A., Frenkel, M.J., and Ward, C.W. (1996). Insulin-responsive tissues contain the core complex protein SNAP-25 (synaptosomal-associated protein 25) A and B isoforms in addition to syntaxin 4 and synaptobrevins 1 and 2. *Biochemical Journal*, Vol.317, pp. 945-954.
- Jahn, R., Lang, T., and Sudhof, T.C. (2003). Membrane fusion. *Cell*, Vol.112, pp. 519-533.
- Karatekin, E., Di Giovanni, J., Iborra, C., Coleman, J., O'Shaughnessy, B., Seagar, M., and Rothman, J.E. (2010). A fast, single-vesicle fusion assay mimics physiological SNARE requirements. *Proceedings of the National Academy of Sciences of the United States of America*, Vol.107, pp. 3517-3521.
- Kesavan, J., Borisovska, M., and Bruns, D. (2007). v-SNARE actions during Ca(2+)-triggered exocytosis. *Cell*, Vol.131, pp. 351-363.
- Luftman, K., Hasan, N., Day, P., Hardee, D., and Hu, C. (2009). Silencing of VAMP3 inhibits cell migration and integrin-mediated adhesion. *Biochemical and Biophysical Research Communications*, Vol.380, pp. 65-70.
- Mallard, F., Tang, B.L., Galli, T., Tenza, D., Saint-Pol, A., Yue, X., Antony, C., Hong, W., Goud, B., and Johannes, L. (2002). Early/recycling endosomes-to-TGN transport involves two SNARE complexes and a Rab6 isoform. *Journal of Cell Biology*, Vol.156, pp. 653-664.
- Martin, L.B., Shewan, A., Millar, C.A., Gould, G.W., and James, D.E. (1998). Vesicle-associated membrane protein 2 plays a specific role in the insulin-dependent

- trafficking of the facilitative glucose transporter GLUT4 in 3T3-L1 adipocytes. *Journal of Biological Chemistry*, Vol.273, pp. 1444-1452.
- Mayer, A., Wickner, W., and Haas, A. (1996). Sec18p (NSF)-driven release of Sec17p (alpha-SNAP) can precede docking and fusion of yeast vacuoles. *Cell*, Vol.85, pp. 83-94.
- McMahon, H.T., Ushkaryov, Y.A., Edelmann, L., Link, E., Binz, T., Niemann, H., Jahn, R., and Sudhof, T.C. (1993). Cellubrevin is a ubiquitous tetanus-toxin substrate homologous to a putative synaptic vesicle fusion protein. *Nature*, Vol.364, pp. 346-349.
- McNew, J.A., Parlati, F., Fukuda, R., Johnston, R.J., Paz, K., Paumet, F., Sollner, T.H., and Rothman, J.E. (2000a). Compartmental specificity of cellular membrane fusion encoded in SNARE proteins. *Nature*, Vol.407, pp. 153-159.
- McNew, J.A., Weber, T., Parlati, F., Johnston, R.J., Melia, T.J., Sollner, T.H., and Rothman, J.E. (2000b). Close is not enough: SNARE-dependent membrane fusion requires an active mechanism that transduces force to membrane anchors. *Journal of Cell Biology*, Vol.150, pp. 105-117.
- Mohrmann, R., de Wit, H., Verhage, M., Neher, E., and Sorensen, J.B. (2010). Fast Vesicle Fusion in Living Cells Requires at Least Three SNARE Complexes. *Science*, Vol.330, pp. 502-505.
- Morgenthaler, F.D., Knott, G.W., Floyd Sarria, J.C., Wang, X., Staple, J.K., Catsicas, S., and Hirling, H. (2003). Morphological and molecular heterogeneity in release sites of single neurons. *European Journal of Neuroscience*, Vol.17, pp. 1365-1374.
- Parlati, F., Varlamov, O., Paz, K., McNew, J.A., Hurtado, D., Sollner, T.H., and Rothman, J.E. (2002). Distinct SNARE complexes mediating membrane fusion in Golgi transport based on combinatorial specificity. *Proceedings of the National Academy of Sciences of the United States of America*, Vol.99, pp. 5424-5429.
- Pocard, T., Le Bivic, A., Galli, T., and Zurzolo, C. (2007). Distinct v-SNAREs regulate direct and indirect apical delivery in polarized epithelial cells. *Journal of Cell Science*, Vol.120, pp. 3309-3320.
- Polgar, J., Chung, S.H., and Reed, G.L. (2002). Vesicle-associated membrane protein 3 (VAMP-3) and VAMP-8 are present in human platelets and are required for granule secretion. *Blood*, Vol.100, pp. 1081-1083.
- Procino, G., Barbieri, C., Tamma, G., De Benedictis, L., Pessin, J.E., Svelto, M., and Valenti, G. (2008). AQP2 exocytosis in the renal collecting duct -- involvement of SNARE isoforms and the regulatory role of Munc18b. *Journal of Cell Science*, Vol.121, pp. 2097-2106.
- Proux-Gillardeaux, V., Gavard, J., Irinopoulou, T., Mege, R.M., and Galli, T. (2005). Tetanus neurotoxin-mediated cleavage of cellubrevin impairs epithelial cell migration and integrin-dependent cell adhesion. *Proceedings of the National Academy of Sciences of the United States of America*, Vol.102, pp. 6362-6367.
- Pryor, P.R., Jackson, L., Gray, S.R., Edeling, M.A., Thompson, A., Sanderson, C.M., Evans, P.R., Owen, D.J., and Luzio, J.P. (2008). Molecular basis for the sorting of the SNARE VAMP7 into endocytic clathrin-coated vesicles by the ArfGAP Hrb. *Cell*, Vol.134, pp. 817-827.
- Randhawa, V.K., Bilan, P.J., Khayat, Z.A., Daneman, N., Liu, Z., Ramlal, T., Volchuk, A., Peng, X.R., Coppola, T., Regazzi, R., Trimble, W.S., and Klip, A. (2000). VAMP2, but not VAMP3/cellubrevin, mediates insulin-dependent incorporation of GLUT4 into

- the plasma membrane of L6 myoblasts. *Molecular Biology of the Cell*, Vol.11, pp. 2403-2417.
- Reed, G.L., Hough, A.K., and Fitzgerald, M.L. (1999). Human platelets contain SNARE proteins and a Sec1p homologue that interacts with syntaxin 4 and is phosphorylated after thrombin activation: implications for platelet secretion. *Blood*, Vol.93, pp. 2617-2626.
- Rothman, J.E. (1994). Mechanisms of intracellular protein transport. *Nature*, Vol.372, pp. 55-63.
- Shen, J., Tareste, D.C., Paumet, F., Rothman, J.E., and Melia, T.J. (2007). Selective activation of cognate SNAREpins by Sec1/Munc18 proteins. *Cell*, Vol.128, pp. 183-195.
- Skalski, M., and Coppelino, M.G. (2005). SNARE-mediated trafficking of alpha5beta1 integrin is required for spreading in CHO cells. *Biochemical and Biophysical Research Communications*, Vol.335, pp. 1199-1210.
- Sollner, T., Bennett, M.K., Whiteheart, S.W., Scheller, R.H., and Rothman, J.E. (1993a). A protein assembly-disassembly pathway in vitro that may correspond to sequential steps of synaptic vesicle docking, activation, and fusion. *Cell*, Vol.75, pp. 409-418.
- Sollner, T., Whiteheart, S.W., Brunner, M., Erdjument-Bromage, H., Geromanos, S., Tempst, P., and Rothman, J.E. (1993b). SNAP receptors implicated in vesicle targeting and fusion. *Nature*, Vol.362, pp. 318-324.
- Steegmaier, M., Klumperman, J., Foletti, D.L., Yoo, J.S., and Scheller, R.H. (1999). Vesicle-associated membrane protein 4 is implicated in trans-Golgi network vesicle trafficking. *Molecular Biology of the Cell*, Vol.10, pp. 1957-1972.
- Stein, A., Weber, G., Wahl, M.C., and Jahn, R. (2009). Helical extension of the neuronal SNARE complex into the membrane. *Nature*, Vol.460, pp. 525-528.
- Sutton, R.B., Fasshauer, D., Jahn, R., and Brunger, A.T. (1998). Crystal structure of a SNARE complex involved in synaptic exocytosis at 2.4 Å resolution. *Nature*, Vol.395, pp. 347-353.
- van den Bogaart, G., Holt, M.G., Bunt, G., Riedel, D., Wouters, F.S., and Jahn, R. (2010). One SNARE complex is sufficient for membrane fusion. *Nature Structural and Molecular Biology*, Vol.17, pp. 358-364.
- Veale, K.J., Offenhauser, C., Whittaker, S.P., Estrella, R.P., and Murray, R.Z. (2010). Recycling endosome membrane incorporation into the leading edge regulates lamellipodia formation and macrophage migration. *Traffic*, Vol.11, pp. 1370-1379.
- Wang, C.C., Ng, C.P., Lu, L., Atlashkin, V., Zhang, W., Seet, L.F., and Hong, W. (2004). A role of VAMP8/endobrevin in regulated exocytosis of pancreatic acinar cells. *Developmental Cell*, Vol.7, pp. 359-371.
- Weber, T., Zemelman, B.V., McNew, J.A., Westermann, B., Gmachl, M., Parlati, F., Sollner, T.H., and Rothman, J.E. (1998). SNAREpins: minimal machinery for membrane fusion. *Cell*, Vol.92, pp. 759-772.
- Wong, S.H., Zhang, T., Xu, Y., Subramaniam, V.N., Griffiths, G., and Hong, W. (1998). Endobrevin, a novel synaptobrevin/VAMP-like protein preferentially associated with the early endosome. *Molecular Biology of the Cell*, Vol.9, pp. 1549-1563.
- Yang, B., Gonzalez, L., Jr., Prekeris, R., Steegmaier, M., Advani, R.J., and Scheller, R.H. (1999). SNARE interactions are not selective. Implications for membrane fusion specificity. *Journal of Biological Chemistry*, Vol.274, pp. 5649-5653.

- Zeng, Q., Subramaniam, V.N., Wong, S.H., Tang, B.L., Parton, R.G., Rea, S., James, D.E., and Hong, W. (1998). A novel synaptobrevin/VAMP homologous protein (VAMP5) is increased during in vitro myogenesis and present in the plasma membrane. *Molecular Biology of the Cell*, Vol.9, pp. 2423-2437.
- Zeng, Q., Tran, T.T., Tan, H.X., and Hong, W. (2003). The cytoplasmic domain of Vamp4 and Vamp5 is responsible for their correct subcellular targeting: the N-terminal extension of VAMP4 contains a dominant autonomous targeting signal for the trans-Golgi network. *Journal of Biological Chemistry*, Vol.278, pp. 23046-23054.

Crosstalk and Integration of Membrane Trafficking Pathways

Edited by Dr. Roberto Weigert

ISBN 978-953-51-0515-2

Hard cover, 246 pages

Publisher InTech

Published online 11, April, 2012

Published in print edition April, 2012

Membrane traffic is a broad field that studies the complex exchange of membranes that occurs inside the cell. Protein, lipids and other molecules traffic among intracellular organelles, and are delivered to, or transported from the cell surface by virtue of membranous carriers generally referred as "transport intermediates". These carriers have different shapes and sizes, and their biogenesis, modality of transport, and delivery to the final destination are regulated by a multitude of very complex molecular machineries. A concept that has clearly emerged in the last decade is that each membrane pathway does not represent a close system, but is fully integrated with all the other trafficking pathways. The aim of this book is to provide a general overview of the extent of this crosstalk.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Chuan Hu, Nazarul Hasan and Krista Riggs (2012). Analysis of SNARE-Mediated Exocytosis Using a Cell Fusion Assay, *Crosstalk and Integration of Membrane Trafficking Pathways*, Dr. Roberto Weigert (Ed.), ISBN: 978-953-51-0515-2, InTech, Available from: <http://www.intechopen.com/books/crosstalk-and-integration-of-membrane-trafficking-pathways/analysis-of-snare-mediated-exocytosis-using-a-cell-fusion-assay>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](https://creativecommons.org/licenses/by/3.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen