We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Introgression and Long-Term Naturalization of Archaeophytes into Native Plants – Underestimated Risk of Hybrids

Hiroyuki Iketani¹ and Hironori Katayama²

¹*Institute of Fruit Science, National Agriculture and Food Research Organization,
Food Resources Education and Research Center,*

²*Graduate School of Agricultural Science, Kobe University,
Japan*

1. Introduction

1.1 Difficulties in the study of archaeophytes

Alien plants pose problems in the conservation of biodiversity, especially by invasion and successive mal-effects on local ecosystem and biodiversity (Pyšek et al., 1995; Ellstrand, 2003; Nentwig, 2007). Even if they are only in cultivation, they could affect by hybridization, vector of diseases and pests and other factors. Since an alien plant is defined as one whose distribution has expanded out of its native range under human influence, the history of alien plants begins with human migration, especially in association with agriculture. As most have spread with active migration since the Age of Discovery, alien plants can be delineated into archaeophytes (introduced before the Age of Discovery) and neophytes (introduced since) (Pyšek et al., 2002; Preston et al., 2004).

Although this classification is commonly used, particularly in Europe, it is difficult to distinguish these two categories (Pyšek et al., 2004; Willis & Birks, 2006). It is practically impossible to prove the non-nativeness of a plant, especially from morphological, ecological, or phytogeographical data. For example, most archaeophytes grow in or near human-made environments, for example, as agricultural weeds or ruderal plants. They have already extended into that ecological system and have been held in equilibrium. Therefore, although research tries to distinguish archaeophytes from true native plants, the two are usually treated the same in the practice of biodiversity conservation. In fact, some archaeophytes are listed in Red Lists (Cheffings & Farrell, 2005; Ministry of Environment, Japan, 2007), since exclusion of archaeophytes from native plants without distinct evidence would make their conservation value lower (Willis & Birks, 2006).

1.2 Biodiversity in Japan and archaeophytes

The Japanese archipelago is rich in biodiversity (Ohwi, 1965; Iwatsuki et al., 1993–2011). The climate of this region ranges widely from subarctic to subtropical, with high humidity and precipitation throughout. It lies near the Asian continent and has been connected with it several times in geological history (continental islands). This placement may have promoted

migration from north and south and produced flora with mixed boreal and temperate elements. By contrast, the archipelago was not connected with the continent during the last glacial maximum, and this isolation may have stimulated the phylogenetic differentiation of species or intraspecific taxa from the continental mother taxa. In addition, its mountainous topography with about twenty 3000-m-class peaks provides refugia for many alpine plants that are relicts of the glacial period.

However, this biologically affluent archipelago is also one of the most populous regions in the world. Although preservation measures beginning in the early modern age have maintained a high percentage of forest (about 70% of land) and protected this “green archipelago” (Totman, 1987), population pressure still poses threats. Therefore Japan is nominated in one of the biodiversity hotspots (Mittermeier et al., 2004).

Humans arrived in Japan relatively recently, ca. 35 000 years BP or later (Keally, 2009). Agriculture is estimated to have begun in Japan 3000 years BP or later (Shōda, 2007). The introduction of alien plants may have begun from this period. Maekawa (1943) first proposed the concept of “prehistorically naturalized plants”, presumably introduced from the Asian continent before the beginning of written history (end of the 6th century CE). He listed about 120 ruderal plants, mainly agricultural weeds. Other plants, including trees, are now also included in this category (Shimizu, 2003). Of course, many plants were also introduced from the Asian continent in historic times before the Age of Discovery. However, since most of these plants are crops or ornamentals, only a few became completely naturalized. Therefore, most of the naturalized archaeophytes in Japan are prehistorically naturalized plants.

1.3 Archaeophytic naturalized useful trees in Japan

Several fruit trees and a few industrial crop trees were introduced into Japan prehistorically. Representative examples are pear (*Pyrus pyrifolia* Nakai), peach (*Prunus persica* L.), plum (*Prunus salicina* L.), apricot (*Prunus armeniaca* L.), persimmon (*Diospyros kaki* Thunb.), loquat (*Eriobotrya japonica* Lindl.), and princess tree (*Paulownia tomentosa* Steud.). These plants now grow more or less wild and can reproduce without human intervention. Some have extended their wild distribution into the upper temperate deciduous forest zone (1000–1600 m a.s.l. in central Japan), where human settlement has been rare and many native flagship species of Japanese biodiversity grow. Whereas some or all of these species were formerly regarded as native, archaeological and phytogeographical research now estimates that they are introduced; for example, the remains, usually seeds or stones, of these fruit trees are found only at archaeological sites dating after the beginning of agriculture, in contrast to native edible fruit species such as wild grape (*Vitis coignetiae* Planch.) and raspberries (*Rubus* spp.) (Kobayashi, 1990).

2. Introgression of archaeophytic *Pyrus pyrifolia* into native *Pyrus ussuriensis* in NE Japan

2.1 Problems in the classification of Japanese *Pyrus*

Among these naturalized useful trees, only *Pyrus pyrifolia* has intercrossable wild relatives in the flora of Japan, namely *Pyrus calleryana* and *Pyrus ussuriensis*. *Pyrus calleryana* is doubtlessly native because it has a very distinctive distribution in common with other

famous endangered plants such as *Magnolia stellata* (Ueda, 1989). On the other hand, the status of *P. ussuriensis* in the flora of Japan has been obscure. This plant was originally described in the early 20th century as two new native species from remote regions (Nakai, 1918): namely, *Pyrus hondoensis* in central Japan (Chubu region) and *Pyrus aromatica* in northeastern Japan (Kitakami Mountains; Fig. 1). However, since then, trees morphologically intermediate between these plants and *P. pyrifolia* were reported, and many of them were described as distinct species. As a result, more than 70 “species” of *Pyrus* were described throughout Japan. Since most of them exist only sporadically in human areas, modern taxonomists treat them as synonyms of either *P. pyrifolia* or *P. ussuriensis* (Ohwi, 1965; Kitamura, 1979; Iketani & Ohashi, 2001).

Solid ellipses indicate the distribution site of the populations in the Kitakami Mountains and the Chubu region. The dashed line indicates the region where wild trees other than the true native type are common. Prefectural boundaries are also shown.

Fig. 1. Localities of two native populations of *Pyrus ussuriensis* in Japan

This taxonomic confusion leads to doubt about the nativeness of wild populations of *P. ussuriensis* (Kitamura, 1979), which is heightened for other reasons. First, this species is edible and is cultivated in northeastern China and Korea. Therefore, *P. ussuriensis* could also have been introduced and naturalized. Second, species of *Pyrus* easily hybridize with each other. Third, morphological distinction between species is obscure: only the presence/absence of calyx lobes in mature fruits is the discriminative character in floristic and taxonomic works (Yü & Ku, 1974; Ku & Sponberg, 2003). Finally, the original sites of discovery are in the lower mountain zone, not far from human settlements.

2.2 Approaches from phytogeography and morphology

As one of the main reasons for this suspicion may be a lack of understanding of the true wild populations of *P. ussuriensis*, my group performed extensive field investigations (Iketani & Ohashi, 2003). We found this species both near the original places of discovery and in more elevated places, which correspond to the upper temperate deciduous forest zone. In contrast to the sporadic distribution of trees growing in human areas at lower elevation, the wild trees grew more densely at higher elevation. They were also morphologically distinct from other seemingly wild or cultivated trees. The fruits and leaf laminae are small in the former but vary in the latter from as small as in wild trees to as large as in edible cultivars (Fig. 2). In addition, the disjunctive geographical distribution at higher elevations of the Chubu region and the Kitakami Mountains is one of the distinctive patterns of endemic taxa in the flora of Japan (Ohashi, 1987). These findings support the indigenesness of the wild populations.

● *Pyrus ussuriensis* from northeastern Japan. ○ *P. ussuriensis* from central Japan. ▲ *P. ussuriensis* from Asian continent. △ Cultivars of *P. ussuriensis* and naturalized *Pyrus*. Vertical lines show the maximum and minimum values of fruit length. (From Iketani & Ohashi, 2003)

Fig. 2. Relation between fruit length and maximum lamina length in *Pyrus* accessions

2.3 Molecular approach reveals introgression between native and prehistorically naturalized plants

Traditional methods made the presence of true native populations plausible. However, the evidence was not conclusive, and the reason for the appearance of morphologically intermediate trees between *P. ussuriensis* and *P. pyrifolia* remained in doubt. Worse, many pear trees with wide morphological variations were discovered in the Kitakami Mountains and surrounding region (Katayama & Uematsu, 2006). To resolve these difficulties, we investigated population genetics using microsatellite loci and with Bayesian statistical inference (Iketani et al., 2010).

The analysis of 226 individuals from six regions implied five hypothetical ancestral populations (Fig. 3; Iketani et al., 2010). These results were at least partly predictable, but not entirely expected. The two true native populations of *P. ussuriensis* in Japan showed genetic distinctiveness, and both were much differentiated from wild plants of *P. ussuriensis* in the Asian continent. These results show that the Japanese populations of *P. ussuriensis* are truly native. Similarly, the introduction of *P. pyrifolia* from China was also supported, since the genetic structure of old Japanese cultivars and local landraces shows a partially common element with Chinese pear cultivars.

Five hypothetical ancestral populations are shown with different shadings. Modified based on Iketani et al., (2010)

Fig. 3. Bayesian statistical inference of population structure of *P. ussuriensis* and *P. pyrifolia*.

Populations from northeastern Japan were more or less genetically admixed with *P. pyrifolia*. This phenomenon was more conspicuous in local landraces and wild individuals collected from outside of the Kitakami Mountains, but was also evident in wild individuals in that region. This result and the ubiquity of intermediate trees clearly show introgression between native *P. ussuriensis* and prehistorically naturalized *P. pyrifolia* trees. In addition, truly native trees proved to be much rarer than introgressed trees and should be protected. *Pyrus ussuriensis* has now been added to the Japanese National Red List of Threatened Plants (Ministry of Environment, Japan, 2007).

3. Implications for biological conservation

3.1 Long-term hybridization and naturalization?

At present we have almost no empirical evidence to infer the historical introgression of *P. pyrifolia* into native *P. ussuriensis*. Although agriculture began later in northeastern Japan than in western Japan, the introduction of *P. pyrifolia* into this region would have also begun prehistorically, or early in historical times at the latest. Since *P. pyrifolia* itself is not widely escaped and naturalized even now, hybridization might have occurred first, and naturalization of hybrid offspring might have followed. Acceleration of naturalization due to hybridization between alien and native species, which is a well-known conservation problem (Ellstrand, 2003; Fitzpatrick & Shaffe, 2007), might have happened.

However, there are still many unclear points in the above scenario. Neither *P. ussuriensis* nor *P. pyrifolia* is very invasive. When assessed for invasiveness, they are judged as posing no or limited risk. For example, their weed risk factor (Food and Agriculture Organization of the United Nations [FAO], 2005) is 4 at most, against a critical score of 6. Even if the fitness of hybrids is better than that of the parents, it is unlikely that they would have expanded like other invasive plants. Therefore, we have to ascribe the cause of this limited expansion to the long passage of time, perhaps more than a thousand years. Kowarik (1995) and Pyšek & Jarošík (2005) studied the time-lag between the introduction and naturalization of alien plants in terms of centuries. The case of *Pyrus* could be appended as an example in terms of millennia.

A long residence time must raise the chance of successful naturalization (Pyšek & Jarošík, 2005). However, human activities such as repeated secondary release would promote invasion beyond the threshold of naturalization (Kowarik, 2003). As both *Pyrus* species bear edible fruits, ancient peoples in northeastern Japan might have grown and propagated hybrid pear trees, which would have become the source of secondary release, although there is no supportive evidence from history, ethnology, archaeology, or demography. Thus, research in these fields will be necessary.

3.2 Negative effect of archaeophytes through hybridization

How should we treat these hybrid pears? On one hand, they now grow widely in northeastern Japan but are not very invasive. The number of individuals is not very large and they are perhaps held in equilibrium in nature. Therefore, they could be treated similarly as native plants, as many other prehistorically naturalized plants (archaeophytes) are.

However, we cannot ignore these pears, because the plant that now grows widely is not an archaeophyte itself. Instead, an archaeophyte has hybridized with a native plant, and the hybrid offspring grow better than their native parent. In this case, hybrid plants should be controlled for the conservation of native plants, especially if the latter are threatened.

This case shows that archaeophytes that are themselves not invasive could still pose a risk to native plants in the long term. The negative effects of alien plants on natural ecosystems are already known. For example, there is much evidence that crops and other domesticated plants can hybridize with their wild relatives while still in cultivation (Ellstrand, 2003). Reproductive interference, which is the depression of fitness by interspecific pollination

interactions such as competition for limited pollinators and pollen loss by interspecific pollen transfer, has recently been recognized (Matsumoto et al., 2010; Takakura et al., 2011). In both phenomena, negative effects could become evident on the human timescale if a plant was cultivated extensively, such as agricultural crops. Even with ornamental use, the effect could appear on such a timescale.

Sukopp & Sukopp (1993) expressed concern about the naturalization (“becoming feral”) of cultivated plants (“cultigens”) and long-term ecological effects such as hybridization with related wild plants. They listed several pairs of cultivated and wild plants in Central Europe which could hybridize. Since then, several actual cases among these pairs have been reported; e.g., *Daucus carota* ssp. *sativus* and ssp. *carota* (Magnussen & Hause, 2007), *Beta vulgaris* ssp. *vulgaris* and ssp. *maritima* (Arnauld et al., 2003; Fénart et al., 2008), and *Brassica napus* and *Brassica rapa* (Andersen et al., 2009).

In conservation policy, introduced cultivated plants which prove not to be invasive are not usually regarded as dangerous. However, in the conservation of threatened plants, their crossable relatives should not be grown near natural habitat. The International Union for Conservation of Nature and Natural Resources [IUCN] (2000) recommended: “Since the impacts on biological diversity of many alien species are unpredictable, any intentional introductions and efforts to identify and prevent unintentional introductions should be based on the precautionary principle.” The case of *Pyrus* supports this policy. We have to control even archaeophytes if they could harm native plants, once they prove to be truly naturalized alien plants. This is why we should distinguish archaeophytes and true native plants more precisely.

3.3 Potential problems in the Pyrinae

The invasiveness and other negative ecological effects of *Pyrus* and related plants are also relevant to the case for distinguishing archaeophytes and true native plants more precisely. The genus *Pyrus* belongs to the subtribe Pyrinae of the tribe Pyreae (formerly subfamily Maloideae) in the family Rosaceae (Potter et al., 2007). Since the reproductive barriers between the members of this tribe are low, interspecific hybrids which are usually fertile are common in nature; even intergeneric hybrids are not rare. This is one of the most extreme cases in higher plants, even though plants produce hybrids much more easily than animals. Previously this character has been a problem only in taxonomy (Kovanda, 1965; Robertson et al., 1991), but it could also become a problem in conservation. Species of this subtribe naturalize commonly in the temperate zones of both hemispheres; e.g., *Malus pumila* (apple), *Pyrus communis* (common pear), *P. pyrifolia*, *Cotoneaster* spp., and *Crataegus* spp. (hawthorn).

So far, only a few species of the Pyrinae are nominated in national and other lists of invasive plants (Table 1). This list shows the obvious invasiveness of some species in Oceania and the Pacific islands, where no native species of the Pyrinae grow (except for *Osteomeles*). *Eriobotrya japonica*, *Pyracantha* spp., and *Sorbus* spp. are also naturalized in the temperate zone of Europe and Japan (Tutin et al., 1968; Shimizu, 2003), but they are not recognized as invasive. However, hybridization between introduced plants and their wild relatives would happen in the long term even without naturalization.

Recently, the invasiveness of *Pyrus calleryana*, which was introduced from East Asia and has been widely propagated in the USA as an ornamental during the past 50 years, has been

Species	Nominated list ^a
<i>Amelanchier spicata</i>	EMPPO
<i>Cotoneaster franchettii</i>	AWC, CIPC
<i>Cotoneaster glaucophyllus</i>	AWC, MoE NZ
<i>Cotoneaster lacteus</i>	CIPC
<i>Cotoneaster pannosus</i>	AWC, HER, CIPC
<i>Cotoneaster salicifolius</i>	AWC
<i>Cotoneaster simonsii</i>	AWC, MoE NZ
<i>Crataegus monogyna</i>	AWC, CIPC
<i>Crataegus sinaica</i>	AWC
<i>Eriobotrya japonica</i>	MoE NZ, HER
<i>Pyracantha angustifolia</i>	AWC, HER, MoE NZ
<i>Pyracantha coccinea</i>	AWC
<i>Pyracantha fortuneana</i>	AWC
<i>Sorbus aucuparia</i>	MoE NZ
<i>Sorbus</i> spp.	AWC

a: AWC, Australian Weeds Committee (2011). CIPC, California Invasive Plant Council (2006). EMPPO, European and Mediterranean Plant Protection Organization (2011). HER, Hawaiian Ecosystems at Risk (2011). MoE NZ, Ministry of Environment, New Zealand (2011).

Table 1. Species of the Pyrinae nominated in national and other lists of invasive plants.

recognized (Vincent, 2005). This is perhaps the first case of a species of *Pyrus* being recognized as invasive. This sudden realization of invasion is explained as the overcoming of self-incompatibility owing to the planting of different cultivars and the Allee effect (Culley & Hardiman, 2009; Hardiman & Culley, 2010). However, since there are no native species of *Pyrus* in North America, negative genetic effects on native relatives would not occur.

In Europe, in contrast, although Sukopp & Sukopp (1993) stated that *Pyrus communis* became feral in Europe in the absence of wild relatives, in fact there are many wild species in this continent (Tutin et al., 1968; Aldasoro et al., 1996), some possibly of hybrid origin, either wild × wild or wild × cultivated. Introgression between cultivated apple (*Malus pumila*, syn. *M. ×domestica*) and a wild relative (*Malus sylvestris*) in Europe is already known (Coart et al., 2003, 2006; Larsen et al., 2006). These cases are comparable to our case of *Pyrus ussuriensis* in terms of introgression between archaeophytic cultivated fruit trees and wild relatives.

The most crucial but incompletely understood situation occurs in East Asia, the center of differentiation of both *Pyrus* and *Malus*, as well as of many other genera of the Pyrinae. About 20 species of *Pyrus* and 30 of *Malus* are now recognized in this region (Iketani & Ohashi, 2001; Ohashi, 1993; Wu et al., 2003). Some species, especially those now found only in cultivation, might have a hybrid origin. These hybridizations might have occurred among native species, although some might have occurred between alien and native. For example, the Chinese pear cultivars we studied proved to be admixtures between *P. ussuriensis* and *P. pyrifolia* (Fig. 3). These cultivars originated in northern China (Yü, 1979), where *P. ussuriensis*

is native but *P. pyrifolia* is alien. Thus, the hybridization would have occurred between wild trees or local landraces of the former and introduced plants of the latter. The time of hybridization is uncertain, but is perhaps not recent. Floristic studies show that trees referred to as *P. ussuriensis* with fairly large fruits occur in nature (Yü & Ku, 1974; Ku & Sponberg, 2003). They are perhaps hybrids, as found in Japan.

4. Future research possibilities

Our case of *Pyrus* suggests that whether a plant is native or alien could be inferred from molecular data. However, studies based on this strategy are rare. In Japan, Sasanuma et al. (2002) and Hori et al. (2006) reported the genetic uniformity of *Elymus humidus* and *Lycoris radiata*, respectively, both of which are prehistorically naturalized plants, in comparison with a certain level of genetic diversity among their wild relatives. European studies have focused on cultivated plants and their relatives. In addition to the abovementioned studies of *Daucus*, *Beta*, *Brassica*, and *Malus*, there are several studies of *Triticum* and *Aegilops* (e.g., Zaharieva & Monneveux, 2006; Arrigo et al., 2011). However, investigations in Europe would be difficult, because Europe is contiguous with the true native habitats of many archaeophytes, and archaeophytes could have been introduced much earlier (from the 6th millennium BCE) than in Japan. Nevertheless, as there are many more archaeophytes in Europe than in Japan and as many putative hybrids between native and alien plants have already been reported (Vila et al., 2000; Pyšek et al., 2002), many opportunities for research remain. For example, English elm (*Ulmus minor* var. *vulgaris*) was proved to be a 2000-year-old Roman clone (Gil et al., 2004).

The importance of research in East Asia and other parts of Eurasia is obvious (Catri, 1989). In research on alien plants, basic biological data such as floristic and ecological status and phytogeography are necessary. Research in historical documents and archaeological data is important for the assessment of archaeophytes. Fortunately, East Asian countries are rich in these resources. Traditional Eastern herbalism, which originated in China and was also developed in Japan and Korea, may provide plant records. Although research in the East still has many constraints compared with that in the West, our long historical cultural heritage offers one important advantage.

5. References

- Aldasoro, J.J., Aedo, C., & Garmendia, F.M. (1996). The genus *Pyrus* L. (Rosaceae) in southwest Europe and North Africa, *Botanical Journal of the Linnean Society*, Vol. 121, No. 2, (June 1996), pp. 143–158, ISSN 1095-8339
- Andersen, N.S., Poulsen, G., Andersen, B. A., Kiær, L.P., D'Hertefeldt, T., Wilkinson, M.J., & Jørgensen, R.B. (2009). Processes affecting genetic structure and conservation: a case study of wild and cultivated *Brassica rapa*. *Genetic Resources and Crop Evolution*, Vol. 56, No. 2, (March 2009), pp. 189–200, ISSN 0925-9864
- Arnaud, J.F., Viard, F., Delescluse, M., & Cuguen, J. (2003). Evidence for gene flow via seed dispersal from crop to wild relatives in *Beta vulgaris* (Chenopodiaceae): consequences for the release of genetically modified crop species with weedy lineages. *Proceedings of the Royal Society of London, Series B*, Vol. 270, No. 1524 (August 2003), pp. 1565–1571, ISSN 0080-4649

- Arrigo, N., Guadagnuolo, R., Lappe, S., Pasche, S., Parisod, C., & Felber, F. (2011). Gene flow between wheat and wild relatives: empirical evidence from *Aegilops geniculata*, *Ae. neglecta* and *Ae. triuncialis*. *Evolutionary Applications*, online, ISSN 1752-4571
- Australian Weeds Committee (2011). Noxious weed list for Australian states and territories Ver.24.00, 15.09.2011, Available from <http://www.weeds.org.au/docs/weednet6.pdf>
- California Invasive Plant Council (2006). *California Invasive Plant Inventory*, Cal-IPC Publication, Retrieved from <http://www.cal-ipc.org/ip/inventory/pdf/Inventory2006.pdf>
- Castri, F. di (1989). History of Biological Invasions with Special Emphasis on the Old World, In: *Biological Invasions – A Global Perspective*. Drake, J.A., Mooney, H.A., Castri, F. di, Groves, R.H., Kruger, F.J., Rejmánek, M., & Williamson, M. (Eds.), ISBN 0471920851, John Wiley & Sons, New York, USA, pp. 1–30. Retrieved from <http://www.scopenvironment.org/downloadpubs/scope37/scope37-ch01.pdf>
- Cheffings, C. & Farrell, L. (Eds.) (2005). *The vascular plant Red Data List for Great Britain*. 15.09.2011, Available from <http://jncc.defra.gov.uk/page-3354>
- Coart, E., Vekemans, X., Smulders, M.J., Wagner, I., Van Huylenbroeck, J., Van Bockstaele, E., & Roldan-Ruiz, I. (2003). Genetic variation in the endangered wild apple (*Malus sylvestris* (L.) Mill.) in Belgium as revealed by amplified fragment length polymorphism and microsatellite markers. *Molecular Ecology*, Vol. 12, No. 4, (April 2003), pp. 845–57, ISSN 0962-1083
- Coart, E., Van Glabeke, S., De Loose, M., Larsen, A.S., & Roldán-Ruiz, I. (2006). Chloroplast diversity in the genus *Malus*: new insights into the relationship between the European wild apple (*Malus sylvestris* (L.) Mill.) and the domesticated apple (*Malus domestica* Borkh.). *Molecular Ecology*, Vol. 15, No. 8, (July 2006), pp. 2171–2182, ISSN 0962-1083
- Culley, T.M. & Hardiman, N.A. (2009). The role of intraspecific hybridization in the evolution of invasiveness: a case study of the ornamental pea tree *Pyrus calleryana*. *Biological Invasions*, Vol. 11, No. 5, (May 2009), pp. 1107–1119, ISSN 1387-3547
- Ellstrand, N.C. (2003). *Dangerous liaisons? When cultivated plants mate with their wild relatives*, Johns Hopkins University Press, ISBN 0-8018-74-5-X, Baltimore, MD, USA
- European and Mediterranean Plant Protection Organization (2011). Invasive alien plants – EPPO lists and documentation, 15.09.2011, Available from http://www.eppo.org/INVASIVE_PLANTS/ias_plants.htm
- Fénart, S., Arnaud, J.F., De Cauwer, I., & Cuguen, J. (2008). Nuclear and cytoplasmic genetic diversity in weed beet and sugar beet accessions compared to wild relatives: new insights into the genetic relationships within the *Beta vulgaris* complex species. *Theoretical and Applied Genetics*, Vol. 116, No. 8, (May 2008), pp. 1063–1077, ISSN 0040-5752
- Fitzpatrick, B.M., & Shaffe, H.B. (2007). Hybrid vigor between native and introduced salamanders raises new challenges for conservation. *Proceedings of the National Academy of Science of the United States of America*, Vol. 104, No. 40 (October 2007), pp. 15793–15798, ISSN 0027-8424
- Food and Agriculture Organization of the United Nations [FAO] (2005). *Procedures for Weed Risk Assessment*. 15.09.2011, Available from <ftp://ftp.fao.org/docrep/fao/009/y5885e/y5885e00.pdf>

- Gil, L., Fuentes-Utrilla, P., Soto, Á., Cervera, M.T., & Collada, C. (2004). English elm is a 2,000-year-old Roman clone. *Nature* Vol. 431, No. 7012 (October 2004), pp. 1053, ISSN 0028-0836
- Hardiman N.A., & Culley T.M. (2010). Reproductive success of cultivated *Pyrus calleryana* (Rosaceae) and establishment ability of invasive, hybrid progeny. *American Journal of Botany*, (October 2010) Vol. 97, No. 10, pp. 1698-1706, ISSN 0002-9122
- Hawaiian Ecosystems at Risk project (2011). Hawaii's Most Invasive Horticultural Plants. In: *Hawaiian Ecosystems at Risk project*, 15.09.2011, Available from <http://www.state.hi.us/dlnr/dofaw/hortweeds/specieslist.htm>
- Hori, T., Hayashi, A., Sasanuma, T., & Kurita, S. (2006). Genetic variations in the chloroplast genome and phylogenetic clustering of *Lycoris* species. *Genes & Genetic Systems*, Vol. 81, No. 4, (October 2006), pp. 243–253, ISSN 1341-7568
- Iketani, H. & Ohashi, H. (2001). Rosaceae Subfam. II. Maloideae (=Pomoideae). In: *Flora of Japan IIb. Angiospermae Dicotyledoneae Archichlamydeae*, Iwatsuki, K., Boufford, D.E., Ohba, H. (Eds.), 111–124, Kodansha, ISBN 4-0615-4605-8, Tokyo, Japan
- Iketani, H., & Ohashi, H. (2003). Taxonomy and distribution of Japanese populations of *Pyrus ussuriensis* Maxim. (Rosaceae). *Journal of Japanese Botany*, Vol. 78, No. 3, (June 2003), pp. 119–134, ISSN 0022-2062
- Iketani, H., Yamamoto, T., Katayama, H., Uematsu, C., Mase, N. & Sato., Y. (2010). Introgression between native and prehistorically naturalized (archaeophytic) wild pear (*Pyrus* spp.) populations in Northern Tohoku, Northeast Japan. *Conservation Genetics*, Vol. 11, No. 1, (February 2010), pp. 115–126, ISSN 1566-0621
- International Union for Conservation of Nature and Natural Resources [IUCN] (2000). *IUCN guidelines for the prevention of biodiversity loss caused by alien invasive species*. 15.09.2011, Available from http://www.issg.org/pdf/guidelines_iucn.pdf
- Iwatsuki, K., Yamazaki, T., Boufford, D.E. & Ohba, H. (Eds.). (1993–2011). *Flora of Japan Vol. 1–4*, Kodansha Scientific, Tokyo, Japan
- Katayama, H. & Uematsu, C. (2006). Pear (*Pyrus* species) genetic resources in Iwate, Japan. *Genetics Resources and Crop Evolution*, Vol. 53, No. 3, (May 2006), pp. 483–498, ISSN 0925-9864
- Keally, C.T. (2009). *Japanese Archaeology*, 15.09.2011, Available from <http://www.t-net.ne.jp/~keally/index.htm>
- Kitamura, S. (1979). *Pyrus*, In: *Coloured Illustrations of Woody Plants of Japan Vol. 2*, Kitamura, S. & Murata, G. (Eds.), pp. 42–47, Hoikusha, Osaka, Japan (in Japanese)
- Kobayashi, A. (1990). *Bunka to Kudamono (Culture and Fruits)*, Yokendo, ISBN 4-8425-9009-2, Tokyo, Japan (in Japanese)
- Kovanda, M. (1965). On the genetic concepts in the Pomoideae. *Preslia*, Vol. 37, pp. 27–34, ISSN 0032-7786
- Kowarik, I. (1995). Time lags in biological invasions with regard to the success and failure of alien species. In: *Plant Invasions: General Aspects and Special Problems*, Pyšek, P., Prach, K., Rejmánek, M., & Wade, M. (Eds.). pp. 15–38, SPB Academic Publishing, ISBN 90-5103-097-5, Amsterdam, the Netherlands
- Kowarik, I. (2003). Human Agency in Biological Invasions: Secondary Releases Foster Naturalisation and population expansion of alien plant species. *Biological Invasions*, Vol. 5, No. 4, (December 2003), pp. 293–312, ISSN 1387-3547

- Ku, T.C. & Sponberg, S.A. (2003). *Pyrus*. In: *Flora of China*, Vol. 9, *Pittosporaceae through Connaraceae*, Wu, Z.Y., Raven, P.H., & Hong, D.Y. (Eds.). pp. 173–179, Science Press, ISBN 7030111729, Beijing, China
- Larsen, A.S., Asmussen, C.B., Coart, E., Olrik, D.C., & Kjær, E.D. (2006). Hybridization and genetic variation in Danish populations of European crab apple (*Malus sylvestris*). *Tree Genetics & Genomes*, Vol. 2, No. 2, (April 2006), pp. 86–97, ISSN 1614-2942
- Maekawa, F. (1943). Prehistoric-naturalized plants to Japan proper. *Acta Phytotaxomica et Geobotanica*, Vol. 13, (November 1943), pp. 274–279, ISSN 0001-6799 (in Japanese)
- Magnussen, L.S. & Hauser T.P. (2007). Hybrids between cultivated and wild carrots in natural populations in Denmark. *Heredity*, Vol. 99, No. 2, (August 2007), pp. 185–192, ISSN 0018-067X
- Matsumoto, T., Takakura, K., & Nishida, T. (2010). Alien pollen grains interfere with the reproductive success of native congener. *Biological Invasions*, Vol. 12, No. 6, pp. 1617–1626, ISSN 1387-3547
- Ministry of the Environment, Japan (2007). The Updated Japanese Red Lists on Mammals, Brackish-water/Freshwater Fishes, Insects, Shellfish, and Plants I and II. 15.09.2011, Available from <http://www.env.go.jp/en/headline/headline.php?serial=503>
- Ministry of the Environment, New Zealand (2011). Alien plants which are considered weeds on conservation lands in New Zealand, 15.09.2011, Available from <http://www.mfe.govt.nz/publications/ser/ser1997/html/tables/table9.4.html>
- Mittermeier, R.A; Gil, P.R., Hoffman, M., Pilgrim, J., Brooks, T., Mittermeier, C.G., Lamoreux, J., & da Fonseca, G.A.B. (Eds.). (2004). *Hotspots Revisited: Earth's Biologically Richest and Most Threatened Terrestrial Ecoregions*, University of Chicago Press, ISBN 978-9686-3977-72, Chicago, USA
- Nakai, T. (1918). Notulae ad plantas japonicae et koreae XVI. *Botanical Magazine, Tokyo*, Vol. 32, No. 374, pp. 28–37, ISSN 0006-808X
- Nentwig, W. (Ed.). (2003). *Biological Invasions*. Springer, ISBN 978-3-540-46919-6, Berlin, Germany
- Ohashi, H. (1987). Floristic regions in the Tohoku District of Japan. *Journal of Japanese Botany*, Vol. 62, pp. 119–126, ISSN 0022-2062 (in Japanese)
- Ohashi, H. (1993). Rosaceae. In: *Flora of Taiwan, 2nd Ed., Vol. 3*, Editorial Committee of the Flora of Taiwan (Eds.) pp. 69–157, Editorial Committee of the Flora of Taiwan, ISBN 957-9019-41-X, Taipei, Taiwan
- Ohwi, J. (1965). *Flora of Japan*, Smithsonian Institute, Washington DC, USA
- Potter D., Eriksson, T., Evans, R.C., Oh, S., Smedmark, J.E.E., Morgan, D.R., Kerr, M., Robertson, K.R., Arsenault, M., Dickinson T.A., & Campbell, C.S. (2007). Phylogeny and classification of Rosaceae. *Plant Systematics and Evolution*, Vol. 266, No. 1–2, pp. 5–43, ISSN: 0378-2697
- Preston, C.D., Pearman, D.A., & Hall, A.R. (2004). Archaeophytes in Britain. *Botanical Journal of the Linnean Society*, Vol. 145, No. 3, (July 2004), pp. 257–294, ISSN 1095-8339
- Pyšek, P. & Jarošík V. (2005). Residence time determines the distribution of alien plants. In: *Invasive plants: ecological and agricultural aspects*, Inderjit, B.P.C. (Ed.). pp. 77–96, Birkhäuser Verlag, Basel, Switzerland
- Pyšek, P., Prach, K., Rejmánek, M., & Wade, M. (Eds.). (1995). *Plant Invasions: General Aspects and Special Problems*, SPB Academic Publishing, ISBN 90-5103-097-5, Amsterdam, the Netherlands

- Pyšek, P., Sádlo, J., & Mandák, B. (2002). Catalogue of alien plants of the Czech Republic. *Preslia*, Vol. 74, No. 2, pp. 97–186, ISSN 0032-7786
- Pyšek, P., Richardson, D.M., Rejmánek, M., Webster, G., Williamson, M., & Kirschner, J. (2004). Alien plants in checklists and floras: towards better communication between taxonomists and ecologists. *Taxon*, Vol. 53, No. 1, (February 2004), pp. 131–143, ISSN 0040-0262
- Robertson, K.R., Phipps, J.B., Rohrer, J.R., & Smith, P.G. (1991). A synopsis of genera in Maloideae (Rosaceae). *Systematic Botany*, Vol. 16, No. 2, (April–June 1991), pp. 376–394, ISSN 0363-6445
- Sasanuma, T., Endo, T.R., & Ban, T. (2002). Genetic diversity of three *Elymus* species indigenous to Japan and East Asia (*E. tsukushiensis*, *E. humidus* and *E. dahuricus*) detected by AFLP. *Genes & Genetic Systems*, Vol. 77, No. 6, pp. 429–438, ISSN 1341-7568
- Shimizu, T. (Ed.). (2003). *Naturalized Plants of Japan*, Heibonsha, ISBN 4-582-53508-9, Tokyo, Japan (in Japanese)
- Shōda, S. (2007). A Comment on the Yayoi Period Dating Controversy. *Bulletin of the Society for East Asian Archaeology*, Vol. 1, pp. 1–7, ISSN 1864-6026
- Sukopp, H. & Sukopp, U. (1993). Ecological long-term effects of cultigens becoming feral and of naturalization of non-native species. *Cellular and Molecular Life Sciences*, Vol. 49, No. 3, (March 1993), pp. 210–218, ISSN 1420-682X
- Takakura, K., Matsumoto, T., Nishida, T., & Nishida, S. (2011). Effective range of reproductive interference exerted by an alien dandelion, *Taraxacum officinale*, on a native congener. *Journal of Plant Research*, Vol. 124, No. 2, (March 2011), pp. 269–276, ISSN 0918-9440
- Totman, C. (1987). *The Green Archipelago: Forestry in Pre-Industrial Japan*, University of California Press, ISBN 978-0520-0631-29, Berkeley, USA
- Tutin, T.G., Heywood, V.H., Burges, N.A., Moore, D.M., Valentine, D.H., Walters, S.M., & Webb D.A. (Eds.) (1968). *Flora Europaea Vol. 2, Rosaceae to Umbelliferae*, Cambridge University Press, ISBN 052106662X, Cambridge, UK
- Ueda, K. (1989). Phytogeography of Tōkai Hilly Land Element I. Definition. *Acta Phytotaxonomica and Geobotanica* Vol. 40, No. 1–4, (July 1989), pp. 190–202, ISSN 0001-6799 (in Japanese)
- Vilà, M., Weber E., & D’Antonio, C.M. (2000). Conservation Implications of Invasion by Plant Hybridization. *Biological Invasions*, Vol. 2, No. 3 (September 2000), pp. 207–217, ISSN 1387-3547
- Vincent, M.A (2005). On the Spread and Current Distribution of *Pyrus calleryana* in the United States. *Castanea*, Vol. 70, No. 1, (Mar 2005), pp. 20–31, ISSN 0008-7475
- Willis, K.J. & Birks H.J.B. (2006). What is natural? The need for a longterm perspective in biodiversity conservation. *Science*, Vol. 314, No. 5803, (November 2006), pp. 1261–1265, ISSN 0036-8075
- Wu, Z. & Raven, P.H. & Hong, D.Y. (Eds.) (2003). *Flora of China Vol. 9, Pittosporaceae through Connaraceae*, Science Press, ISBN 7030111729, Beijing, China
- Yü, T.T. (1979). *Zhong guo guo shu fen lei xue (Classification of the fruit trees in China)*, Agriculture Press, Beijing (in Chinese)
- Yü, T.T. & Ku, T.C. (1974). *Pyrus*. In: *Flora Reipublicae Popularis Sinicae*, Vol. 36, Yü, T.T. (Ed.), pp. 354–372, Science Press, Beijing, China (in Chinese)

Zaharieva, M. & Monneveux, P. (2006). Spontaneous hybridization between bread wheat (*Triticum aestivum* L.) and its wild relatives in Europe. *Crop Science*, Vol. 46, No. 2, pp. 512-527, ISSN 0011-183X

IntechOpen

IntechOpen

Topics in Conservation Biology

Edited by Dr. Tony Povillitis

ISBN 978-953-51-0540-4

Hard cover, 110 pages

Publisher InTech

Published online 02, May, 2012

Published in print edition May, 2012

Conservation biology is called a "crisis discipline." In a world undergoing rapid change, this science informs us about research, technologies, management practices, and policies that can help protect the earth's naturally-occurring biological diversity. The six chapters of this book provide insightful analysis on managing protected areas (Middle East), conserving biochemical and genetic diversity of carob tree (Tunisia) and wild pear (Japan), determining the health status of Amazon manatee, manipulating sex ratios to benefit wildlife, and narrowing the gap between religion and conservation. The authors approach threats to biological diversity from varied angles, reflecting the interdisciplinary nature of the field. This book offers room for reflection on the definition and utility of the word 'natural' on a planet now overwhelmingly dominated by people.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Hiroyuki Iketani and Hironori Katayama (2012). Introgression and Long-Term Naturalization of Archaeophytes into Native Plants - Underestimated Risk of Hybrids, Topics in Conservation Biology, Dr. Tony Povillitis (Ed.), ISBN: 978-953-51-0540-4, InTech, Available from: <http://www.intechopen.com/books/topics-in-conservation-biology/introgression-and-long-term-naturalization-of-archaeophytes-into-native-plants-underestimated-risk-o>

INTeCH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](https://creativecommons.org/licenses/by/3.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen