We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Human Male Meiosis and Sperm Aneuploidies

María Vera¹, Vanessa Peinado¹, Nasser Al-Asmar^{1,2},
Jennifer Gruhn³, Lorena Rodrigo¹, Terry Hassold³ and Carmen Rubio^{1,2}

¹Iviomics

²Instituto Valenciano de Infertilidad (IVI Valencia)

³Center for Reproductive Biology

School of Molecular Biosciences

Washington State University

^{1,2}Spain

³USA

1. Introduction

Infertility affects 15% of couples of reproductive age (De Krester, 1997; reviewed by O'Flynn O'Brien et al., 2010). For years most reproductive problems were attributed to women, but recent research has shown that between 30 and 50% of infertility cases result from male factor (Lipshultz & Howards, 1997; reviewed by Stahl et al., 2011).

Reports indicate that infertile males, especially those with low sperm count, have an increase in chromosomally-abnormal spermatozoa, therefore the study of these individuals is critical for the field of assisted reproduction. Importantly, aneuploid sperm remain able to fertilize eggs, often resulting in repetitive intracytoplasmic sperm injection (ICSI) failure, recurrent miscarriage, and offspring with increased genetic risk. Most numerical chromosomal abnormalities, especially monosomies, are inviable and result in spontaneous abortion. However, a subset of chromosomal abnormalities can result in live birth, including trisomies 13, 18, and 21 and sex chromosome aneuploidies. Offspring with these aneuploidies typically display physical disabilities, mental retardation, infertility, etc. (reviewed by Martin, 2008). Thus, effective diagnostic methods to detect sperm aneuploidy are of great interest.

An estimated 5% (minimally) of pregnancies are aneuploid. However, in assisted reproductive technology (ART), estimates are as high as 25% or more. Aneuploid conceptions often derive from aneuploid gametes: the incidence of aneuploidy in human oocytes is about 20-25%, and is more than 2% in sperm (reviewed by Hassold & Hunt, 2001). Aneuploidy in gametes is intimately associated with meiotic errors. This chapter focuses on the etiology of aneuploidy in the human male gamete, particularly examining male meiosis, meiotic errors, and their possible relationship with aneuploidy. We also describe aneuploidy rates in fertile and infertile men and their reproductive outcomes.

2. Male meiosis

2.1 Generalities

Meiosis in humans varies considerably between males and females (Fig. 1). In females, meiosis begins somewhat synchronously in all oocytes during fetal development, but then arrests before birth. Resumption of meiosis occurs asynchronously at or after puberty, at the time of ovulation, but arrests again until fertilization. In contrast, male meiosis does not begin until puberty and occurs continuously and asynchronously throughout adulthood. Primary spermatocytes divide into two secondary spermatocytes through meiosis I (MI), and each secondary spermatocyte divides into two spermatids, which will differentiate and mature into sperm. The main difference between the male and female production of gametes is that oogenesis only leads to the production of one final ovum from each primary oocyte, in contrast in males four sperm are originated from each primary spermatocyte.

Fig. 1. Gamete formation in males (spermatogenesis) and females (oogenesis).

Meiosis is characterized by an extended prophase followed by two divisions that produce haploid gametes. Prior to meiosis, during S phase, DNA is replicated and sister chromatids remain connected through proteins called cohesins. Meiosis begins with prophase I, arguably the most critical stage thanks to two key events: synapsis and recombination

(explained in detail later). Prophase I is divided into four stages: leptotene, zygotene, pachytene, and diplotene. During leptotene, homologous chromosomes begin to condense and approach each other to align, but do not yet pair. With the aid of cohesins (e.g., REC8 and SMC1B), a chromosomal scaffold – the synaptonemal complex (SC) – begins to form via assembly of axial elements composed of SC-specific proteins [e.g., synaptonemal complex protein 2 (SYCP2) and SYCP3]. Additionally, DNA double-strand breaks (DSB) are induced by the protein SPO11; this is the initiating event of recombination, or DNA exchange between homologous chromosomes. During zygotene chromosomes continue condensing and homologues pair and begin synapsis (reviewed by Handel & Schimenti, 2010). Axial elements become lateral elements and, joined by the transverse filaments (e.g., SYCP1), form the SC to maintain synapsis. Subsequently, during pachytene – the longest stage of prophase I – synapsis is completed and DSB are repaired by DNA break repair machinery; a small subset of DSB are repaired as crossovers (chiasmata). Finally, during diplotene the SC is disassembled. Each pair of homologues (bivalents) begins to separate, but they remain temporarily joined at chiasmata (reviewed by Burgoyne et al., 2009).

Following prophase I, germ cells continue into the first meiotic metaphase, in which axial elements are disassembled and cohesins are removed, except those located at centromeres. In this phase, homologous pairs of chromosomes migrate to the equatorial plane of the spindle, with their centromeres facing different poles, forming the metaphase plate. In anaphase I, homologous chromosomes separate. Finally, cell division at telophase I results in two secondary spermatocytes with haploid chromosomes, but with two chromatids per chromosome.

The second meiotic division is an equational division similar to mitosis except that, in this case, the parent cells are haploid. Four spermatids are thus generated from one pre-meiotic germ cell; each should contain 23 chromosomes, with a single chromatid for each chromosome.

Three events can lead to aneuploidy during MI: failure to resolve chiasmata, resulting in a true non-disjunction; no chiasma formation or early disappearance of one, resulting in an achiasmate non-disjunction; and, finally, a premature separation of sister chromatids. In meiosis II (MII) the only cause of aneuploidy is non-disjunction of sister chromatids (reviewed by Hassold & Hunt, 2001).

The importance of MI versus MII errors varies among chromosomes. For example, trisomy 16 is caused by MI errors, while trisomy 18 is usually caused by non-disjunction in MII (reviewed by Hassold & Hunt, 2001). In male, the chromosome with the highest estimated frequency of non-disjunction in MI is the XY bivalent and in MII is the Y chromosome (reviewed by Hall et al., 2006). Further, autosomal trisomies are usually maternal in origin, while sex chromosome aneuploidies are more frequently paternal in origin (reviewed by Templado et al., 2011).

The critical events of meiosis are reflected in two series of specific processes: First, pairing, synapsis, and formation of the SC between homologues establish and regulate cohesion between sister chromatids. Second, recombination between homologous chromosomes and proper orientation of the two centromeres of each bivalent must be completed. Both sets of events are controlled by a pachytene checkpoint, which induces delay or arrest of cells that have not completed these steps to avoid aberrant chromosome segregation and formation of

defective gametes (reviewed by Roeder & Bailis, 2000). However, a failure in any of these steps together with a checkpoint failure can result in aneuploidy or segregation failure.

2.2 Synapsis

The SC is a protein structure that mediates pairing, synapsis, and recombination between homologous chromosomes during meiosis. The SC is formed by two parallel lateral elements and a central element that links them with transverse filaments. Three protein components of the SC have been identified: SYCP1, forming the central element, and SYCP2 and SYCP3, which are recruited to the chromosomal cores to form the axial elements (reviewed by Handel & Schimenti, 2010). These structures have been visualized using monoclonal antibodies, particularly against SYCP3, in immunocytochemical assays. As the first component to localize to the forming SC, SYCP3 can be used to monitor its assembly and disassembly during prophase I. Indeed, SYCP3 distribution varies greatly during prophase (Fig. 2): in leptotene SYCP3 appears as small linear fragments; in zygotene the axial elements begin to elongate and synapse; in pachytene synapsis is complete revealing 22 structures (one for each bivalent) and one sex vesicle (XY bivalent); and, finally, in diplotene each pair of sister chromatids starts to separate, visualized as forks in the SC.

It has been observed that the synaptic initiation occurs in subtelomeric regions and not in the telomeres as initially was thought. Furthermore it has been seen that the centromere avoids spreading the synapsis from q-arm to p-arm and vice versa (Brown et al., 2005).

Fig. 2. SYCP3, MLH1, and CREST (stained in orange, green, and blue respectively) distribution in prophase I stages (from left to right: leptotene, zygotene, pachytene, and diplotene).

Synapsis fidelity can be measured by observing the frequency of certain anomalies: gaps, splits, and lack of crossovers. Gaps are discontinuities in the SC and occur often in normal males. Splits are misaligned chromosomal regions that form looplike structures; these occur less commonly than gaps. Finally, SCs without crossovers are detected by absence of MutL protein homolog 1 (MLH1) foci on the bivalent. This anomaly is less common than gaps and even splits. The existence of gaps is reported to correlate with the location of the MLH1 protein. Further, recombination frequency is reduced on SCs with gaps, but not those with splits, which suggests that the nature of splits and gaps differs (Sun et al., 2005, 2007).

2.3 Recombination

Proper recombination between homologous chromosomes is required for normal meiosis and segregation of chromosomes. Chiasmata, physical structures corresponding to

crossover points between homologous chromosomes, keep homologues joined until anaphase I. The distribution of these points is not entirely random; rather, there are genomic regions that exhibit much higher rates of recombination, called hot spots, and other locations that rarely have recombination events, called cold spots. For example, centromeres seem to be cold spots, while subtelomeric regions seem to be hot spots with an excess of recombination points. Several studies of human chromosomes have shown that at least one crossover occurs in each arm, except in short arms of acrocentric chromosomes, where crossovers occur infrequently. Further, the existence of a crossover inhibits the formation of another in a nearby region, a phenomenon known as “interference” (Brown et al., 2005; reviewed by Lynn et al., 2004; Sun et al., 2004a).

Antibodies against MLH1, a DNA mismatch-repair protein involved in crossing-over, are used to localize recombination sites. Applying this method in human spermatocytes, groups have described inter-individual variations in MLH1 frequency (Hassold et al., 2004; Lynn et al., 2002; Sun et al., 2005, 2006). Additional findings demonstrated that patient age does not affect meiotic recombination frequency (Lynn et al., 2002). Further, a role for the SC in mediating recombination levels was observed, with a correlation between the number of MLH1 foci and chromosome arm lengths (Lynn et al., 2002; Sun et al., 2004a). Despite these advances, the basis of crossover formation remains incompletely understood, e.g., why crossovers form at a particular site. This indicates the importance of continued efforts toward revealing the mechanistic basis of recombination.

2.4 Methods for studying critical events of meiosis

Male meiosis can be studied using indirect methods, like linkage mapping, or direct assays such as cytogenetics and immunocytogenetics.

Linkage maps, also called genetic or meiotic maps, are derived from genotype data in families by examining heredity of short tandem repeat polymorphisms to detect all recombination events per meiosis. This type of assay provides high resolution, but has some disadvantages. First, at least three generations need to be examined, and second, only half of all recombination events can be observed since only two of the four chromatids are involved in the exchange process (reviewed by Lynn et al., 2004).

Direct assays, although they forfeit some resolution, can help combat these disadvantages. For example, analysis of diakinesis/metaphase I stage cells by conventional cytogenetic techniques is used to directly observe crossovers. However, because diakinesis and metaphase I have a short duration, finding these cells can be difficult. Additionally, material for this kind of study requires testicular biopsy. Finally, chromosomes are highly condensed, making it difficult to identify them correctly (reviewed by Lynn et al., 2004; reviewed by Vallente et al., 2006). This last problem can be solved by adding M-FISH (Multiplex-fluorescent *in situ* hybridization), which allows the identification of individual bivalents. Unfortunately, this technique is costly and quite laborious, prohibiting its application in clinical practice (Sarrate et al., 2004).

Another, more recently developed direct assay uses immunocytogenetics to study prophase I, particularly pachytene, germ cells. Antibodies against lateral element proteins (SYCP3) and transverse filament proteins (SYCP1) are used to visualize the SC, anti-MLH1 antibodies are used to detect recombination sites, and the centromere can be localized with

CREST sera (calcinosis-Raynaud's phenomenon-esophageal dysfunction-sclerodactyly-telangiectasia) (Fig. 2). This assay allows the study of meiotic progression, synaptic defects, recombination rates, etc. Additionally, immunocyto genetics can be combined with FISH to analyze recombination rates and crossover placement in individual chromosomes. While this assay also requires testicular biopsies to obtain material, pachytene has a long duration and nuclei are not as condensed as in diakinesis (Gonsalves et al., 2004, 2005; Judis et al., 2004; Lynn et al., 2002; Ma et al., 2006; Sun et al., 2004a, 2004b, 2006, 2007, 2008a, 2008b).

3. Male meiosis abnormalities

Different types of meiotic abnormalities have been described, and they are not mutually exclusive. One type of abnormality is meiotic arrest, which occurs when spermatogenesis stops at any stage of maturation of the germ line (spermatogonia, primary spermatocyte, secondary spermatocyte, or spermatid). Arrest can be partial, affecting only some germline cells and causing oligozoospermia, or complete, affecting all germline cells and causing azoospermia. In 74% of cases, meiotic arrest is caused by *synaptic defects*, which are due to a decrease in the number of exchanges (chiasmata) between homologous chromosomes in prophase I. This can induce abnormal chromosome segregation in MI, resulting in secondary spermatocytes with an altered chromosome complement (reviewed by Egozcue et al., 2005).

Importantly, a decrease in the total number of crossovers can result from homologues devoid of recombination sites. This, in turn, could cause misalignment of chromosomes on the metaphase plate and, subsequently, improper segregation (reviewed by Martin et al., 2008). The XY bivalent is particularly susceptible to this phenomenon, since only a small region of homology exists between the X and Y chromosomes, and, thus, the pair has only a single recombination site (Shi & Martin, 2000; Thomas & Hassold, 2003). In fact, several studies have demonstrated that sex chromosomes have the highest frequency of achiasmate bivalents (Sun et al., 2006).

In addition to altered frequency of recombination, abnormal crossover distribution can also occur. In a recent study was found an altered MLH1 distribution in one of four infertile males, these could have negative consequences as aneuploid sperm (Ferguson et al., 2009).

3.1 Abnormal meiosis in infertile males

Reports indicate that up to 8% of the general infertile population exhibit meiotic defects (reviewed by Egozcue et al., 2005). Studies of patients with obstructive azoospermia (OA) are typically used to define normal meiosis, since, in these cases, the absence of sperm in semen is due exclusively to a physical barrier (reviewed by Vallente et al., 2006). In fact, recent studies in testicular sperm from OA patients did not show differences in aneuploidy rates compared to testicular control subjects (Rodrigo et al., 2011). Further, recombination levels showed a similar pattern in all patients, with a mean recombination level of 49.5 ± 0.7 crossovers (MLH1 foci) (Al-Asmar et al., 2010, 2011).

On the other hand, in most non-obstructive azoospermic (NOA) patients the cause of testicular damage is idiopathic (Judis et al., 2004). In these individuals various meiotic abnormalities have been observed. Indeed, several studies have reported a significant decrease in recombination levels in NOA, with averages of 32.7-42.7 MLH1 foci versus 46.0-

48.5 for controls (Sun et al., 2004b, 2007). Additionally, one report described a NOA patient with a significant increase (73% vs. 4.5% in controls) in cells with at least one autosomal bivalent without an MLH1 focus (Sun et al., 2004b). Further, NOA samples may have altered distributions of germ cells in meiotic stages. In particular, leptotene and zygotene stage cells are more common in azoospermic patients than in controls (7.95% vs. 2.30% and 9.75% vs. 1.45% respectively), with corresponding decreases in pachytene stage cells (75.30% vs. 96.25% in controls). This phenomenon may reflect meiotic arrest in the earlier stages of prophase (Tassistro et al., 2009). In fact, complete meiotic arrest was observed in one case of NOA (Judis et al., 2004), in which the blockage was at zygotene/pachytene stage and no evidence was detected for synapsis or crossovers between homologues. Another study reported a partial or complete arrest during zygotene in 4 of 40 NOA patients (Gonsalves et al., 2004).

Other types of meiotic abnormalities have been observed in NOA patients. For example, SC discontinuities are more frequent in these patients than in controls (Sun et al., 2004b; Tassistro et al., 2009). In addition, differences have been detected in the frequency of asynapsis events between NOA and control populations (7.97% vs. 2.95%) (Tassistro et al., 2009).

Meiotic abnormalities have also been found in other patients. In a study with oligoasthenozoospermic patients, 17.5% of individuals had an increased incidence of meiotic anomalies (reviewed by Egozcue et al., 2000; Vendrell et al., 1999). Additionally, in a retrospective study of 500 patients with different types of infertility or sterility, the incidence of synaptic abnormalities was inversely proportional to the quantitative level of spermatogenesis. Specifically, altered or incomplete meiosis was found in 24% of azoospermic males, 33% of asthenoteratozoospermic males, 51.5% of oligoastheno-teratozoospermia (OAT) patients, and in 90% of NOA males (García et al., 2005).

Furthermore, genetic studies have identified mutations for proteins involved in meiosis in infertile males. For example, mutations in SCP3, MLH1, SPO11 –a protein involved in the formation of DSB– and RAD54 –a protein that participates in recombination and DNA repair– have been found (reviewed by Sanderson et al., 2008). Otherwise, variations in DNA sequence and their relationship with an increase or decrease in the recombination rate have been described (Kong et al., 2008).

3.2 Abnormal meiosis and reproductive outcomes

Alterations in meiosis do not necessarily lead to azoospermia. Indeed, meiotic errors can produce abnormal sperm that retain fertilization capabilities, resulting in abnormal embryos and either recurrent miscarriage or abnormal offspring. Aran et al. (2004) observed a high number of chromosomal abnormalities (42.5%) in embryos from ICSI cycles for males with meiotic abnormalities (Aran et al., 2004). Furthermore, it has been reported that approximately 50% of individuals with Klinefelter syndrome (47,XXY) have resulted from paternal non-disjunction. Indeed, paternal origin has been associated with a decrease in recombination for sex chromosomes (reviewed by Martin et al., 2008).

3.3 Correlation with sperm aneuploidy

Both aberrant meiosis and increased frequency of sperm aneuploidy have been described in infertile men. However, very few studies have examined meiosis and chromosomal

abnormalities in the same patients. One study to determine the relationship between frequency of recombination in pachytene cells and frequency of sperm aneuploidy in six fertile men did not find a significant correlation (Sun et al., 2008b). Other studies observed that recombination frequency in 24,XY spermatozoa was significantly lower than in normal sperm (25.3 vs. 38.3%) (Shi et al., 2001) and that 67.1% of those XY bivalents were achiasmate (Thomas & Hassold, 2003). Also, in one infertile male with extremely high sperm aneuploidy rates, a correlation was observed between low recombination frequency in pachytene cells for chromosomes 13, 21, and the sex chromosomes and a high aneuploidy frequency for the same chromosomes (Ma et al., 2006).

A study in testicular tissue of seven NOA patients reported an increase in the frequency of pachytene cells with at least one achiasmate bivalent compared with controls (12.4% vs. 4.2% respectively). The same patients exhibited an increase in the frequency of aneuploidies, specifically more sperm disomy than controls for chromosomes 21 (1.00% vs. 0.24%), X, (0.16% vs. 0.03%), and Y (0.12% vs. 0.03%). Further, a significant correlation was found between cells with sex vesicles lacking MLH1 foci and sex chromosome disomy in sperm (Sun et al., 2008a).

Another recent study by Peinado and colleagues (2011) analyzed testicular biopsies of 11 NOA patients and compared these with a control group of 10 patients with obstructive azoospermia post-vasectomy. NOA patients had a mean of 44.9 ± 3.6 MLH1 foci versus 48.2 ± 2.1 in controls. This decrease in recombination levels resulted from a significant increase in the percentage of SC without exchanges (1.1% vs. 0.4%) and with one exchange (19.4% vs. 14.6%) and a significant decrease in complexes with three (19.7% vs. 22.8%) and four exchanges (3.6% vs. 5.5%) compared to the control group. In the same study using FISH analysis, NOA patients displayed an increase in sex chromosome disomies (0.39% vs. 0.18%) and a three-fold increase in disomy for chromosomes 13 (0.4% vs. 0.09%) and 21 (0.3% vs. 0.09%) as well as in diploidy rates (0.13% vs. 0.05%). These findings corroborate the correlation between both parameters, recombination frequency and synapsis defects, and higher aneuploidy risk for offspring in NOA patients (Peinado et al., 2011).

4. Sperm aneuploidies in males

In recent years, the use of ICSI has significantly improved the fertility prognosis of infertile couples affected by severe oligozoospermia (Palermo et al., 1992; Van Steirteghem et al., 1993) or azoospermia, in the latter case using spermatozoa retrieved from the epididymis (Tournaye et al., 1994) or testicle (Devroey et al., 1995; Gil-Salom et al., 1995a, 1995b, 1996, 2000; Schoysman et al., 1993). However, prenatal diagnosis (i.e., preimplantation genetic diagnosis, PGD) following ICSI has shown statistically significant increases in *de novo* sex chromosome abnormalities and structural autosomal aberrations (Bonduelle et al., 2002; Van Steirteghem et al., 2002), most of which seem to have a paternal origin (Meschede et al., 1998; Van Opstal et al., 1997). This finding emphasizes the importance of a strict genetic evaluation of ICSI candidates.

4.1 Approaches to studying aneuploidy in sperm samples

The first chromosomal studies of spermatozoa were developed in 1970, with differential staining of concrete regions of chromosomes (Barlow & Vosa, 1970; Pearson & Bobrow, 1970). These showed that 1.4% of sperm have aneuploidies for sex chromosomes. Other

authors published an individual chromosomal aneuploidy rate of 2% and a global aneuploidy rate of 38% (Pawlowitzki & Pearson, 1972). However, this method was later demonstrated to be non-specific, resulting in over-reporting of aneuploidy. Thus, more reliable techniques were needed.

A later technique introduced spermatozoa to hamster oocytes (Yanagimachi et al., 1976) to directly study human sperm chromosomes (Rudak et al., 1978). After incubating spermatozoa with several hamster eggs, cells were fixed and stained using karyotyping methods to observe metaphase nuclei and analyze numerical and structural chromosome abnormalities. However, this method could not test samples from men with severe male factor because their sperm were not able to capacitate and penetrate zona-free oocytes. Additionally, this technique allowed analysis of only a limited number of sperm (reviewed by Carrell, 2008). Finally, the time and animals required limited the use of this technique in clinical practice (reviewed by Martin, 2008).

In the 1990s, the first FISH assays were developed, offering a faster, easier, and less costly method to detect aneuploidies in human spermatozoa (reviewed by Martin, 2008). This technique uses fluorescent nucleic acid probes complementary to DNA to visualize regions of interest. This approach permits analysis of a large number of sperm from the same sample. However, FISH cannot be performed for all chromosomes of a single sperm nucleus because the sperm head is too small and signals overlap. Therefore, FISH is typically performed for just five chromosomes (13, 18, 21, X, and Y—aneuploidies for these chromosomes can result in live offspring), and resulting aneuploidy rates refer only to those chromosomes analyzed (Fig. 3) (reviewed by Templado et al., 2011). Despite the drawbacks, FISH remains the most widely used technique to detect aneuploidies.

Fig. 3. FISH assays in human spermatozoa. a: diploid spermatozoon (13,13,21,21), b: disomy for chromosome 21 (13,21,21), c: disomy for chromosome 13 (13,13,21), d: diploid spermatozoon (X,Y,18,18), e: disomy for chromosome X (X,X,18), f: disomy for sex chromosomes (X,Y,18).

Recently, an automated tool for the analysis of sperm aneuploidy has been developed. This system has several advantages over manual FISH. First, automated FISH reduces the time required from 10-20 hours to 1 hour. Additionally, the software is able to store all images captured under microscopy. However, this system is costly (reviewed by Carrell, 2008; reviewed by Templado et al., 2011) and is less efficient in identifying diploid sperm than manual analysis (i.e., manual analysis found higher diploidy rates than automatic analysis, Molina et al., 2009).

4.2 Incidence of aneuploidy in fertile males

The detection of alterations in aneuploidy rates first requires establishment of normal parameters. Such values must be based on sperm from normozoospermic males with

proven fertility to establish the baseline levels of human sperm aneuploidy. Notably, baseline sperm aneuploidy rates may be subject to inter-individual and intra-individual variability. Studies published to date show considerable variability in aneuploidy rates in fertile men. However, it is unclear whether this variability is real or is due to different methodologies used in each laboratory (reviewed by Templado et al., 2011). One study on intra-individual and inter-individual variability indicates that unknown life events may sporadically or consistently affect sperm aneuploidy rates (Tempest et al., 2009). Despite these factors, a recent review analyzed 32 studies that described aneuploidy rates from normal men. The authors found that total aneuploidy frequency in sperm is about 4.5%. However, disomy rates for autosomes were highly variable, ranging from 0.03% for chromosome 8 to 0.47% for chromosome 22. Sex chromosomes, the most commonly analyzed, exhibited a disomy rate of 0.27% (reviewed by Templado et al., 2011).

4.3 Incidence of aneuploidy in infertile males

FISH studies reveal a significantly increased incidence of numerical chromosomal abnormalities in sperm from infertile males, mainly in sex chromosomes, and in sperm from OAT patients (Aran et al., 1999; Bernardini et al., 1998, 2000; Calogero et al., 2001a,b; Colombero et al., 1999; Martin et al., 2003; Moosani et al., 1995; Nishikawa et al., 2000; Pang et al., 1999; Pfeffer et al., 1999; Rubio et al., 2001; Ushijima1 et al., 2000; Vegetti et al., 2000). Indeed, Rubio et al. observed statistically significant differences in disomies for chromosome 21 and sex chromosomes and in diploidy in OAT males compared to normozoospermic (Normo), asthenozoospermic (Asthen), teratozoospermic (Terato), and asthenoteratozoospermic (AT) males (Table 1).

	Normo (n=14)	Asthen (n=14)	Terato (n=8)	AT (n=6)	OAT (n=21)
% sex chromosome disomy	0.44	0.36	0.42	0.29	0.68 ^a
% disomy 13	0.14	0.09	0.07	0.11	0.17
% disomy 18	0.02	0.04	0.03	0.02	0.05
% disomy 21	0.17	0.12	0.10	0.08	0.24 ^b
% diploidy	0.10	0.11	0.15	0.06	0.25 ^a

Table 1. Percentage of chromosome aneuploidies. ^ap<0.0001, ^bp=0.0423 (Rubio et al., 2001).

Additionally, at a lower sperm concentration aneuploidy rates were higher. The highest percentage of abnormal FISH results was found at concentrations <1x10⁶ sperm/mL (severe oligozoospermia) –57% of these patients presented some kind of chromosomal abnormality (Fig. 4).

Although some studies have found similar incidences of chromosomal abnormalities in testicular sperm from NOA patients (Martin et al., 2000) and OA patients (Viville et al., 2000) compared to ejaculated sperm from fertile males, most FISH studies report a higher incidence of chromosomal abnormalities in testicular spermatozoa, particularly in NOA patients (Bernardini et al., 2000; Burrello et al., 2002; Levron et al., 2001; Mateizel et al., 2002; Palermo et al., 2002) than in ejaculated spermatozoa from normozoospermic donors. This difference seems to be more appreciable in sex chromosomes (Rodrigo et al., 2004).

Fig. 4. Percentage of abnormal FISH for different sperm concentration (Rodrigo, L. Unpublished data).

A recent study examined sperm aneuploidy rates in NOA and OA patients (Rodrigo et al., 2011). Importantly, this study used two control groups: one group included ejaculated sperm samples from normozoospermic donors with proven fertility (ejaculated control group, EC); the other included testicular sperm samples of males with proven fertility, for whom a testicular sperm aspiration was performed at the time of vasectomy (testicular control group, TC) because the endocrine environment of the testis had not been altered. In control samples, testicular sperm showed higher incidences of aneuploidies than ejaculated sperm. For azoospermic patients (both NOA and OA), the differences were greater when compared to EC than to TC. Additionally, these differences were greater in NOA than in OA patients. Therefore, to better assess NOA patients, testicular sperm from controls should be used for statistical comparisons.

5. Sperm aneuploidies and reproductive outcomes

5.1 Sperm aneuploidies in ART/infertile populations

Sperm aneuploidies can impact reproductive outcome at different stages: fertilization, embryo development, pregnancy, or birth. Several studies have analyzed the clinical consequences of aneuploidies during *in vitro* fertilization (IVF) cycles.

Fertilization rate

Chromosomally abnormal sperm have been related to repetitive ICSI failures in a prospective study. The authors found a higher aneuploidy rate in those with unsuccessful ICSI outcome, especially for chromosome 18 and the sex chromosomes (Nicolopoulos et al., 2008).

Embryo development

Sperm aneuploidies can also result in abnormal embryos. A FISH study in couples with abnormal sperm detected a significant decrease in the number of normal embryos and a significant increase in mosaic embryos (Rodrigo et al., 2003). A similar study in couples with abnormal FISH and low sperm concentration (<5X10⁶ sperm/mL) also found a significant increase in abnormal embryos, especially in mosaic embryos –embryos with cells with different chromosome complement– and sex chromosome aneuploidies (Pehlivan et al.,

2003). Mosaicism rate as high as 53% has been reported in patients with NOA (Silber et al., 2003). These findings could be explained by fertilization with sperm carrying multiple chromosomal alterations or centrosome abnormalities. Sperm defective centrosomes impede the formation of asters or lead to an abnormal spindle, with an abnormal distribution of chromosomes, resulting in aneuploid embryos (Chatzimeletiou et al., 2008). In addition, an abnormal number of male centrioles in the centrosome has been related with the production of haploid, poliploid, or mosaic embryos (Munne et al., 2002; Silber et al., 2003).

Sanchez-Castro et al. (2009) studied sperm aneuploidy rates and embryo chromosomal abnormalities in couples with oocyte donation. They found more abnormal embryos in the study group compared to the control group. Further, oligozoospermic patients showed a higher proportion of abnormal embryos (Sanchez-Castro et al., 2009). Recently, an increase in sperm chromosomal abnormalities has been reported to directly affect chromosomal constitution of preimplantation embryos. This study showed that an increase in disomy for sex chromosomes is associated with an elevated risk of generating potentially viable embryos whose sex chromosomes are affected. Additionally, they observed that an increase in diploid sperm results in a higher incidence of triploid embryos, which are associated with more spontaneous abortions (Rodrigo et al., 2010).

Implantation rate

Oocyte fertilization by a chromosomally abnormal sperm is believed to cause implantation failure (Pang et al., 1999). A study in patients with three or more implantation failures reported an increase in sex chromosome disomies in 31.6% of males (Rubio et al., 2001). Further, later studies correlated abnormal FISH in spermatozoa with a decrease in pregnancy and implantation rates in ICSI cycles (Burrello et al., 2003; Nicopoullos et al., 2008).

Miscarriage

Abnormal sperm has also been related to recurrent miscarriage. A study in sperm samples from couples with recurrent miscarriage showed that sex chromosome disomy was significantly increased compared to internal controls. Further, in a subset of seven couples who underwent oocyte donation, mean frequencies for sex chromosome disomy were even higher and diploidy was also significantly increased (Rubio et al., 1999). These results suggest an implication of sperm chromosome abnormalities in some cases of recurrent pregnancy loss. Later, other studies corroborated this hypothesis, reporting an increase in sex chromosome disomies and diploid spermatozoa in couples with recurrent miscarriage (Al-Hassan et al., 2005; Bernardini et al., 2004; Giorlandino et al., 1998; Rubio et al., 2001). Finally, a recent study described that approximately 66% of abnormal karyotypes from miscarriages originate from male factor (Kim et al., 2010).

5.2 Sperm aneuploidies and abnormal offspring

Although most embryonic abnormalities end in implantation failure or spontaneous abortion, a variable percentage of abnormal offspring has been reported and associated with the presence of aneuploid spermatozoa in the father.

Down syndrome

Blanco et al. (1998) studied two fathers of children with Down syndrome who had a paternally-derived extra chromosome 21. FISH sperm studies showed elevated incidences of

spermatozoa with disomy 21. Further, in one patient, an increase in diploid sperm and disomy for sex chromosomes was observed (Blanco et al., 1998). Later chromosomes 4, 13, and 22 were analyzed in the same patients; an increase in disomies for chromosomes 13 and 22 was reported (Soares et al., 2001).

Sex chromosomes

Several studies of sperm samples in fathers of children with sex chromosome abnormalities have described higher rates of sex chromosome aneuploidy in sperm, especially related to Klinefelter syndrome (47,XXY) and Turner syndrome (X0). Moosani et al. (1999) performed FISH in sperm from males with normal karyotypes whose children had Klinefelter syndrome. They observed a higher percentage of XY disomy in sperm from these patients compared to fertile donors (Moosani et al., 1999). In another study the incidence of sperm with XY disomy was compared among males with children with Klinefelter syndrome of paternal or maternal origin. A significant increase in disomic sperm was detected in the paternal origin group (Eskenazi et al., 2002).

One study of males having children with Turner syndrome (45,X0) of paternal origin analyzed aneuploidy incidence for sex chromosomes. An increase in disomies and nullisomies for sex chromosomes was observed compared to the control group. Oocyte fertilization by sperm with nullisomy for sex chromosomes can produce a (45,X0) embryo (Martinez-Pasarell et al., 1999). Later, aneuploidy incidences for chromosomes 4, 13, 21, and 22 were assessed in ejaculated sperm from the same four males. A significant increase in chromosomes 13 and 22 disomies was observed in one male and in chromosome 21 in two other males (Soares et al., 2001). Another group studied the incidence of sex chromosomal aneuploidies in a male with a previous miscarriage with Turner syndrome of paternal origin. This study reported a significant increase in sex chromosomal disomies and nullisomies compared to the control group (Tang et al., 2004).

6. Conclusion

Many couples with infertility receive a diagnosis of “male origin”. Thus, the study of male gametogenesis and accurate evaluation of male gametes are extremely important.

As a key step of spermatogenesis, meiosis has been the focus of many recent studies. These studies often use immunocytogenetic assays to detect relevant proteins for recombination and synapsis. Thanks to this technique, meiotic abnormalities in infertile males have been well-described, especially for NOA males. Importantly, meiotic abnormalities can result in abnormal chromosome segregation; in fact, a correlation between meiotic abnormalities and sperm aneuploidy has already been described.

Chromosome abnormalities are increasingly found in sperm of many infertile men. This makes the direct analysis of sperm aneuploidy of clinical relevance, since male infertility is now treated by ICSI, which has the implicit risk of transmitting chromosomal aberrations from paternal side. Therefore, in IVF settings, the analysis of sperm chromosomal aneuploidies by FISH is of great interest. Using sperm FISH analysis, an increased incidence of sex chromosome disomies has been described in patients with impaired sperm parameters and in patients with recurrent spontaneous abortions. In this regard, an inverse correlation between sperm quality and sperm aneuploidy rates has been reported.

Another important issue is how sperm aneuploidy may influence ICSI outcome in infertile patients. The studies published so far suggest that sperm aneuploidy may be associated with implantation failure and/or early fetal loss. For these couples at risk, several treatment options have been postulated that range from regular ICSI cycles, to genetic diagnosis or donor sperm. Genetic screening studies in male factor infertility have reported a higher incidence of chromosome abnormalities in embryos from infertile men with altered sperm parameters. In azoospermic patients, higher rates of mosaic and chromosomally abnormal embryos have been reported, most frequently for sex chromosomes.

Therefore, accurate detection of sperm aneuploidies by FISH could be a useful tool for reproductive counseling in couples with male infertility.

7. Acknowledgment

Special thanks go to author's colleagues who contributed to the research and clinical work, presented in this chapter.

8. List of abbreviations

ART	Assisted reproductive technology
CREST	Calcinosis-Raynaud's phenomenon-esophageal dysfunction sclerodactyly-telangiectasia
DSB	Double-strand breaks
EC	Ejaculated control
FISH	Fluorescent in situ hybridization
ICSI	Intracytoplasmic sperm injection
IVF	In vitro fertilization
M-FISH	Multiplex-fluorescent in situ hybridization
MI	Meiosis I
MII	Meiosis II
MLH1	MutL protein homolog 1
NOA	Non-obstructive azoospermia
OA	Obstructive azoospermia
OAT	Oligoastheno-teratozoospermia
RAD54	DNA repair and recombination protein RAD54-like
REC8	Meiotic recombination protein REC8 homolog
SC	Synaptonemal complex
SMC1B	Structural maintenance of chromosomes protein 1B
SPO11	Meiotic recombination protein SPO11
SYCP	Synaptonemal complex protein
TC	Testicular control

9. References

- Al-Asmar, N., Peinado, V., Gruhn, J., Susiarjo, M., Gil-Salom, M., Martínez-Jabaloyas, J.M., Pellicer, A., Remohí, J., Rubio, C., Hassold, T. (2010). Sperm aneuploidy and meiotic recombination in patients with post-vasectomy obstructive azoospermia,

- Proceedings in 26th Annual Meeting of the European Society of Human Reproduction and Embryology*, Rome, Italy, June 27-30, 2010
- Al-Asmar, N., Peinado, V., Gruhn, J., Gil-Salom, M., Martínez-Jabaloyas, J.M., Pellicer, A., Remohí, J., Hassold, T., Rubio, C., (2011). Meiotic recombination, crossover placement and sperm aneuploidy in fertile post-vasectomy patients, *Proceedings in 8th European Cytogenetics Conference*, Porto, Portugal, July 2-5, 2011
- Al-Hassan, S., Hellani, A., Al-Shahrani, A., Al-Deery, M., Jaroudi, K. & Coskun, S. (2005). Sperm chromosomal abnormalities in patients with unexplained recurrent abortions. *Arch.Androl.*, Vol.51, No.1, pp. 69-76
- Aran, B., Blanco, J., Vidal, F., Vendrell, J. M., Egozcue, S., Barri, P. N., Egozcue, J. & Veiga, A. (1999). Screening for abnormalities of chromosomes X, Y, and 18 and for diploidy in spermatozoa from infertile men participating in an in vitro fertilization-intracytoplasmic sperm injection program. *Fertil.Steril.*, Vol.72, No.4, pp. 696-701
- Aran, B., Veiga, A., Vidal, F., Parriego, M., Vendrell, J. M., Santalo, J., Egozcue, J. & Barri, P. N. (2004). Preimplantation genetic diagnosis in patients with male meiotic abnormalities. *Reprod.Biomed.Online*, Vol.8, No.4, pp. 470-476
- Barlow, P. & Vosa, C. G. (1970). The Y chromosome in human spermatozoa. *Nature*, Vol.226, No.5249, pp. 961-962
- Bernardini, L., Borini, A., Preti, S., Conte, N., Flamigni, C., Capitanio, G. L. & Venturini, P. L. (1998). Study of aneuploidy in normal and abnormal germ cells from semen of fertile and infertile men. *Hum.Reprod.*, Vol.13, No.12, pp. 3406-3413
- Bernardini, L., Gianaroli, L., Fortini, D., Conte, N., Magli, C., Cavani, S., Gaggero, G., Tindiglia, C., Ragni, N. & Venturini, P. L. (2000). Frequency of hyper-, hypohaploidy and diploidy in ejaculate, epididymal and testicular germ cells of infertile patients. *Hum.Reprod.*, Vol.15, No.10, pp. 2165-2172
- Bernardini, L. M., Costa, M., Bottazzi, C., Gianaroli, L., Magli, M. C., Venturini, P. L., Francioso, R., Conte, N. & Ragni, N. (2004). Sperm aneuploidy and recurrent pregnancy loss. *Reprod.Biomed.Online*, Vol.9, No.3, pp. 312-320
- Blanco, J., Gabau, E., Gomez, D., Baena, N., Guitart, M., Egozcue, J. & Vidal, F. (1998). Chromosome 21 disomy in the spermatozoa of the fathers of children with trisomy 21, in a population with a high prevalence of Down syndrome: increased incidence in cases of paternal origin. *Am.J.Hum.Genet.*, Vol.63, No.4, pp. 1067-1072
- Bonduelle, M., Liebaers, I., Deketelaere, V., Derde, M. P., Camus, M., Devroey, P. & Van Steirteghem, A. (2002). Neonatal data on a cohort of 2889 infants born after ICSI (1991-1999) and of 2995 infants born after IVF (1983-1999). *Hum.Reprod.*, Vol.17, No.3, pp. 671-694
- Brown, P. W., Judis, L., Chan, E. R., Schwartz, S., Seftel, A., Thomas, A. & Hassold, T. J. (2005). Meiotic synapsis proceeds from a limited number of subtelomeric sites in the human male. *Am.J.Hum.Genet.*, Vol.77, No.4, pp. 556-566
- Burgoyne, P. S., Mahadevaiah, S. K. & Turner, J. M. (2009). The consequences of asynapsis for mammalian meiosis. *Nat.Rev.Genet.*, Vol.10, No.3, pp. 207-216
- Burrello, N., Calogero, A. E., De Palma, A., Grazioso, C., Torrisi, C., Barone, N., Pafumi, C., D'Agata, R. & Vicari, E. (2002). Chromosome analysis of epididymal and testicular spermatozoa in patients with azoospermia. *Eur.J.Hum.Genet.*, Vol.10, No.6, pp. 362-366

- Burrello, N., Vicari, E., Shin, P., Agarwal, A., De Palma, A., Grazioso, C., D'Agata, R. & Calogero, A. E. (2003). Lower sperm aneuploidy frequency is associated with high pregnancy rates in ICSI programmes. *Hum.Reprod.*, Vol.18, No.7, pp. 1371-1376
- Calogero, A. E., De Palma, A., Grazioso, C., Barone, N., Romeo, R., Rappazzo, G. & D'Agata, R. (2001a). Aneuploidy rate in spermatozoa of selected men with abnormal semen parameters. *Hum.Reprod.*, Vol.16, No.6, pp. 1172-1179
- Calogero, A. E., De Palma, A., Grazioso, C., Barone, N., Burrello, N., Palermo, I., Gulisano, A., Pafumi, C. & D'Agata, R. (2001b). High sperm aneuploidy rate in unselected infertile patients and its relationship with intracytoplasmic sperm injection outcome. *Hum.Reprod.*, Vol.16, No.7, pp. 1433-1439
- Carrell, D. T. (2008). The clinical implementation of sperm chromosome aneuploidy testing: pitfalls and promises. *J.Androl.*, Vol.29, No.2, pp. 124-133
- Chatzimeletiou, K., Morrison, E. E., Prapas, N., Prapas, Y. & Handyside, A. H. (2008). The centrosome and early embryogenesis: clinical insights. *Reprod.Biomed.Online*, Vol.16, No.4, pp. 485-491
- Colombero, L. T., Hariprasad, J. J., Tsai, M. C., Rosenwaks, Z. & Palermo, G. D. (1999). Incidence of sperm aneuploidy in relation to semen characteristics and assisted reproductive outcome. *Fertil.Steril.*, Vol.72, No.1, pp. 90-96
- De Kretser, D. M. (1997). Male infertility. *Lancet*, Vol.349, No.9054, pp. 787-790
- Devroey, P., Liu, J., Nagy, Z., Goossens, A., Tournaye, H., Camus, M., Van Steirteghem, A. & Silber, S. (1995). Pregnancies after testicular sperm extraction and intracytoplasmic sperm injection in non-obstructive azoospermia. *Hum.Reprod.*, Vol.10, No.6, pp. 1457-1460
- Egozcue, S., Vendrell, J. M., Garcia, F., Veiga, A., Aran, B., Barri, P. N. & Egozcue, J. (2000). Increased incidence of meiotic anomalies in oligoasthenozoospermic males preselected for intracytoplasmic sperm injection. *J.Assist.Reprod.Genet.*, Vol.17, No.6, pp. 307-30
- Egozcue, J., Sarrate, Z., Codina-Pascual, M., Egozcue, S., Oliver-Bonet, M., Blanco, J., Navarro, J., Benet, J. & Vidal, F. (2005). Meiotic abnormalities in infertile males. *Cytogenet.Genome Res.*, Vol.111, No.3-4, pp. 337-342
- Eskenazi, B., Wyrobek, A. J., Kidd, S. A., Lowe, X., Moore, D., 2nd, Weisiger, K. & Aylstock, M. (2002). Sperm aneuploidy in fathers of children with paternally and maternally inherited Klinefelter syndrome. *Hum.Reprod.*, Vol.17, No.3, pp. 576-583
- Ferguson, K. A., Leung, S., Jiang, D. & Ma, S. (2009). Distribution of MLH1 foci and inter-focal distances in spermatocytes of infertile men. *Hum.Reprod.*, Vol.24, No.6, pp. 1313-1321
- García, F., Egozuce, S., López Teijón, M.L., Olivares, R., Serra, O., Aura, M., Moragas, M., Rabanal, A., Egozuce, J. (2005). Meiosis en biopsia testicular y patrón espermatogenico, *Proceedings in XII Congreso Nacional de Andrologia*, A Coruña, España, April 14-16, 2005
- Gil-Salom, M., Minguez, Y., Rubio, C., De los Santos, M. J., Remohi, J. & Pellicer, A. (1995a). Efficacy of intracytoplasmic sperm injection using testicular spermatozoa. *Hum.Reprod.*, Vol.10, No.12, pp. 3166-3170
- Gil-Salom, M., Minguez, Y., Rubio, C., Remohi, J. & Pellicer, A. (1995b). Intracytoplasmic testicular sperm injection: an effective treatment for otherwise intractable obstructive azoospermia. *J.Urol.*, Vol.154, No.6, pp. 2074-2077

- Gil-Salom, M., Romero, J., Minguez, Y., Rubio, C., De los Santos, M. J., Remohi, J. & Pellicer, A. (1996). Pregnancies after intracytoplasmic sperm injection with cryopreserved testicular spermatozoa. *Hum.Reprod.*, Vol.11, No.6, pp. 1309-1313
- Gil-Salom, M., Romero, J., Rubio, C., Ruiz, A., Remohi, J. & Pellicer, A. (2000). Intracytoplasmic sperm injection with cryopreserved testicular spermatozoa. *Mol.Cell.Endocrinol.*, Vol.169, No.1-2, pp. 15-19
- Giorlandino, C., Calugi, G., Iaconianni, L., Santoro, M. L. & Lippa, A. (1998). Spermatozoa with chromosomal abnormalities may result in a higher rate of recurrent abortion. *Fertil.Steril.*, Vol.70, No.3, pp. 576-577
- Gonsalves, J., Sun, F., Schlegel, P. N., Turek, P. J., Hopps, C. V., Greene, C., Martin, R. H. & Pera, R. A. (2004). Defective recombination in infertile men. *Hum.Mol.Genet.*, Vol.13, No.22, pp. 2875-2883
- Gonsalves, J., Turek, P. J., Schlegel, P. N., Hopps, C. V., Weier, J. F. & Pera, R. A. (2005). Recombination in men with Klinefelter syndrome. *Reproduction*, Vol.130, No.2, pp. 223-229
- Hall, H., Hunt, P. & Hassold, T. (2006). Meiosis and sex chromosome aneuploidy: how meiotic errors cause aneuploidy; how aneuploidy causes meiotic errors. *Curr.Opin.Genet.Dev.*, Vol.16, No.3, pp. 323-329
- Handel, M. A. & Schimenti, J. C. (2010). Genetics of mammalian meiosis: regulation, dynamics and impact on fertility. *Nat.Rev.Genet.*, Vol.11, No.2, pp. 124-136
- Hassold, T. & Hunt, P. (2001). To err (meiotically) is human: the genesis of human aneuploidy. *Nat.Rev.Genet.*, Vol.2, No.4, pp. 280-291
- Hassold, T., Judis, L., Chan, E. R., Schwartz, S., Seftel, A. & Lynn, A. (2004). Cytological studies of meiotic recombination in human males. *Cytogenet.Genome Res.*, Vol.107, No.3-4, pp. 249-255
- Judis, L., Chan, E. R., Schwartz, S., Seftel, A. & Hassold, T. (2004). Meiosis I arrest and azoospermia in an infertile male explained by failure of formation of a component of the synaptonemal complex. *Fertil.Steril.*, Vol.81, No.1, pp. 205-209
- Kim, J. W., Lee, W. S., Yoon, T. K., Seok, H. H., Cho, J. H., Kim, Y. S., Lyu, S. W. & Shim, S. H. (2010). Chromosomal abnormalities in spontaneous abortion after assisted reproductive treatment. *BMC Med.Genet.*, Vol.11, pp. 153
- Kong, A., Thorleifsson, G., Stefansson, H., Masson, G., Helgason, A., Gudbjartsson, D. F., Jonsdottir, G. M., Gudjonsson, S. A., Sverrisson, S., Thorlacius, T., Jonasdottir, A., Hardarson, G. A., Palsson, S. T., Frigge, M. L., Gulcher, J. R., Thorsteinsdottir, U. & Stefansson, K. (2008). Sequence variants in the RNF212 gene associate with genome-wide recombination rate. *Science*, Vol.319, No.5868, pp. 1398-1401
- Levron, J., Aviram-Goldring, A., Madgar, I., Raviv, G., Barkai, G. & Dor, J. (2001). Studies on sperm chromosomes in patients with severe male factor infertility undergoing assisted reproductive technology treatment. *Mol.Cell.Endocrinol.*, Vol.183 Suppl 1, pp. S23-8
- Lipshultz L.I., Howards, S. S. (1997). *Infertility in the male* (3), Mosby, ISBN 0815155018, 9780815155010, USA
- Lynn, A., Koehler, K. E., Judis, L., Chan, E. R., Cherry, J. P., Schwartz, S., Seftel, A., Hunt, P. A. & Hassold, T. J. (2002). Covariation of synaptonemal complex length and mammalian meiotic exchange rates. *Science*, Vol.296, No.5576, pp. 2222-2225

- Lynn, A., Ashley, T. & Hassold, T. (2004). Variation in human meiotic recombination. *Annu.Rev.Genomics Hum.Genet.*, Vol.5, pp. 317-349
- Ma, S., Arsovska, S., Moens, P., Nigro, M. & Chow, V. (2006). Analysis of early meiotic events and aneuploidy in nonobstructive azoospermic men: a preliminary report. *Fertil.Steril.*, Vol.85, No.3, pp. 646-652
- Martin, R. H., Greene, C., Rademaker, A., Barclay, L., Ko, E. & Chernos, J. (2000). Chromosome analysis of spermatozoa extracted from testes of men with non-obstructive azoospermia. *Hum.Reprod.*, Vol.15, No.5, pp. 1121-1124
- Martin, R. H., Greene, C., Rademaker, A. W., Ko, E. & Chernos, J. (2003). Analysis of aneuploidy in spermatozoa from testicular biopsies from men with nonobstructive azoospermia. *J.Androl.*, Vol.24, No.1, pp. 100-103
- Martin, R. H. (2008). Meiotic errors in human oogenesis and spermatogenesis. *Reprod.Biomed.Online*, Vol.16, No.4, pp. 523-531
- Martinez-Pasarell, O., Nogues, C., Bosch, M., Egozcue, J. & Templado, C. (1999). Analysis of sex chromosome aneuploidy in sperm from fathers of Turner syndrome patients. *Hum.Genet.*, Vol.104, No.4, pp. 345-349
- Mateizel, I., Verheyen, G., Van Assche, E., Tournaye, H., Liebaers, I. & Van Steirteghem, A. (2002). FISH analysis of chromosome X, Y and 18 abnormalities in testicular sperm from azoospermic patients. *Hum.Reprod.*, Vol.17, No.9, pp. 2249-2257
- Meschede, D., Lemcke, B., Exeler, J. R., De Geyter, C., Behre, H. M., Nieschlag, E. & Horst, J. (1998). Chromosome abnormalities in 447 couples undergoing intracytoplasmic sperm injection--prevalence, types, sex distribution and reproductive relevance. *Hum.Reprod.*, Vol.13, No.3, pp. 576-582
- Molina, O., Sarrate, Z., Vidal, F. & Blanco, J. (2009). FISH on sperm: spot-counting to stop counting? Not yet. *Fertil.Steril.*, Vol.92, No.4, pp. 1474-1480
- Moosani, N., Pattinson, H. A., Carter, M. D., Cox, D. M., Rademaker, A. W. & Martin, R. H. (1995). Chromosomal analysis of sperm from men with idiopathic infertility using sperm karyotyping and fluorescence in situ hybridization. *Fertil.Steril.*, Vol.64, No.4, pp. 811-817
- Moosani, N., Chernos, J., Lowry, R. B., Martin, R. H. & Rademaker, A. (1999). A 47,XXY fetus resulting from ICSI in a man with an elevated frequency of 24,XY spermatozoa. *Hum.Reprod.*, Vol.14, No.4, pp. 1137-1138
- Munne, S., Sandalinas, M., Escudero, T., Marquez, C. & Cohen, J. (2002). Chromosome mosaicism in cleavage-stage human embryos: evidence of a maternal age effect. *Reprod.Biomed.Online*, Vol.4, No.3, pp. 223-232
- Nicopoullou, J. D., Gilling-Smith, C., Almeida, P. A., Homa, S., Nice, L., Tempest, H. & Ramsay, J. W. (2008). The role of sperm aneuploidy as a predictor of the success of intracytoplasmic sperm injection? *Hum.Reprod.*, Vol.23, No.2, pp. 240-250
- Nishikawa, N., Murakami, I., Ikuta, K. & Suzumori, K. (2000). Sex chromosomal analysis of spermatozoa from infertile men using fluorescence in situ hybridization. *J.Assist.Reprod.Genet.*, Vol.17, No.2, pp. 97-102
- O'Flynn O'Brien, K. L., Varghese, A. C. & Agarwal, A. (2010). The genetic causes of male factor infertility: a review. *Fertil.Steril.*, Vol.93, No.1, pp. 1-12
- Palermo, G., Joris, H., Devroey, P. & Van Steirteghem, A. C. (1992). Pregnancies after intracytoplasmic injection of single spermatozoon into an oocyte. *Lancet*, Vol.340, No.8810, pp. 17-18

- Palermo, G. D., Colombero, L. T., Hariprashad, J. J., Schlegel, P. N. & Rosenwaks, Z. (2002). Chromosome analysis of epididymal and testicular sperm in azoospermic patients undergoing ICSI. *Hum.Reprod.*, Vol.17, No.3, pp. 570-575
- Pang, M. G., Hoegerman, S. F., Cuticchia, A. J., Moon, S. Y., Doncel, G. F., Acosta, A. A. & Kearns, W. G. (1999). Detection of aneuploidy for chromosomes 4, 6, 7, 8, 9, 10, 11, 12, 13, 17, 18, 21, X and Y by fluorescence in-situ hybridization in spermatozoa from nine patients with oligoasthenoteratozoospermia undergoing intracytoplasmic sperm injection. *Hum.Reprod.*, Vol.14, No.5, pp. 1266-1273
- Pawlowitzki, I. H. & Pearson, P. L. (1972). Chromosomal aneuploidy in human spermatozoa. *Humangenetik*, Vol.16, No.1, pp. 119-122
- Pearson, P. L. & Bobrow, M. (1970). Fluorescent staining of the Y chromosome in meiotic stages of the human male. *J.Reprod.Fertil.*, Vol.22, No.1, pp. 177-179
- Pehlivan, T.S., Rodrigo, L., Simón, C., Remohí, J., Pellicer, A., Rubio, C. (2003). Chromosomal abnormalities in day 3 embryos from severe oligozoospermic patients, *Proceedings in 59th Annual Meeting of the American Society for Reproductive Medicine*, Texas, USA, October, 2003
- Peinado, V., Al-Asmar, N., Gruhn, J., Rodrigo, L., Gil-Salom, M., Martinez, J.M., Pellicer, A., Remohi, J., Hassold, T.J., Rubio, C. Relationship between meiotic recombination and sperm aneuploidy and diploidy in non-obstructive azoospermic males, *Proceedings in 8th European Cytogenetics Conference*, Porto, Portugal, July 2-5, 2011
- Pfeffer, J., Pang, M. G., Hoegerman, S. F., Osgood, C. J., Stacey, M. W., Mayer, J., Oehninger, S. & Kearns, W. G. (1999). Aneuploidy frequencies in semen fractions from ten oligoasthenoteratozoospermic patients donating sperm for intracytoplasmic sperm injection. *Fertil.Steril.*, Vol.72, No.3, pp. 472-478
- Rodrigo, L., Rubio, C., Gil-Salom, M., Mateu, E., Pérez-Cano, I., Herrero, R., Simon, C., Remohí, J., Pellicer, A. (2003). Sperm chromosomal abnormalities: a new indication for preimplantational genetic diagnosis?, *Proceedings in 19th Annual Meeting of the European Society of Human Reproduction and Embryology*, Madrid, Spain, June-July, 2003
- Rodrigo, L., Rubio, C., Mateu, E., Simon, C., Remohi, J., Pellicer, A. & Gil-Salom, M. (2004). Analysis of chromosomal abnormalities in testicular and epididymal spermatozoa from azoospermic ICSI patients by fluorescence in-situ hybridization. *Hum.Reprod.*, Vol.19, No.1, pp. 118-123
- Rodrigo, L., Peinado, V., Mateu, E., Remohi, J., Pellicer, A., Simon, C., Gil-Salom, M. & Rubio, C. (2010). Impact of different patterns of sperm chromosomal abnormalities on the chromosomal constitution of preimplantation embryos. *Fertil.Steril.*, Vol.94, No.4, pp. 1380-1386
- Rodrigo, L., Rubio, C., Peinado, V., Villamon, R., Al-Asmar, N., Remohi, J., Pellicer, A., Simon, C. & Gil-Salom, M. (2011). Testicular sperm from patients with obstructive and nonobstructive azoospermia: aneuploidy risk and reproductive prognosis using testicular sperm from fertile donors as control samples. *Fertil.Steril.*, Vol.95, No.3, pp. 1005-1012
- Roeder, G. S. & Bailis, J. M. (2000). The pachytene checkpoint. *Trends Genet.*, Vol.16, No.9, pp. 395-403

- Rubio, C., Simon, C., Blanco, J., Vidal, F., Minguez, Y., Egozcue, J., Crespo, J., Remohi, J. & Pellicer, A. (1999). Implications of sperm chromosome abnormalities in recurrent miscarriage. *J.Assist.Reprod.Genet.*, Vol.16, No.5, pp. 253-258
- Rubio, C., Gil-Salom, M., Simon, C., Vidal, F., Rodrigo, L., Minguez, Y., Remohi, J. & Pellicer, A. (2001). Incidence of sperm chromosomal abnormalities in a risk population: relationship with sperm quality and ICSI outcome. *Hum.Reprod.*, Vol.16, No.10, pp. 2084-2092
- Rudak, E., Jacobs, P. A. & Yanagimachi, R. (1978). Direct analysis of the chromosome constitution of human spermatozoa. *Nature*, Vol.274, No.5674, pp. 911-913
- Sanchez-Castro, M., Jimenez-Macedo, A. R., Sandalinas, M. & Blanco, J. (2009). Prognostic value of sperm fluorescence in situ hybridization analysis over PGD. *Hum.Reprod.*, Vol.24, No.6, pp. 1516-1521
- Sanderson, M. L., Hassold, T. J. & Carrell, D. T. (2008). Proteins involved in meiotic recombination: a role in male infertility? *Syst.Biol.Reprod.Med.*, Vol.54, No.2, pp. 57-74
- Sarrate, Z., Blanco, J., Egozcue, S., Vidal, F. & Egozcue, J. (2004). Identification of meiotic anomalies with multiplex fluorescence in situ hybridization: Preliminary results. *Fertil.Steril.*, Vol.82, No.3, pp. 712-717
- Schoysman, R., Vanderzwalm, P., Nijs, M., Segal, L., Segal-Bertin, G., Geerts, L., van Roosendaal, E. & Schoysman, D. (1993). Pregnancy after fertilisation with human testicular spermatozoa. *Lancet*, Vol.342, No.8881, pp. 1237
- Shi, Q. & Martin, R. H. (2000). Aneuploidy in human sperm: a review of the frequency and distribution of aneuploidy, effects of donor age and lifestyle factors. *Cytogenet.Cell Genet.*, Vol.90, No.3-4, pp. 219-226
- Shi, Q., Spriggs, E., Field, L. L., Ko, E., Barclay, L. & Martin, R. H. (2001). Single sperm typing demonstrates that reduced recombination is associated with the production of aneuploid 24,XY human sperm. *Am.J.Med.Genet.*, Vol.99, No.1, pp. 34-38
- Silber, S., Escudero, T., Lenahan, K., Abdelhadi, I., Kilani, Z. & Munne, S. (2003). Chromosomal abnormalities in embryos derived from testicular sperm extraction. *Fertil.Steril.*, Vol.79, No.1, pp. 30-38
- Soares, S. R., Templado, C., Blanco, J., Egozcue, J. & Vidal, F. (2001). Numerical chromosome abnormalities in the spermatozoa of the fathers of children with trisomy 21 of paternal origin: generalised tendency to meiotic non-disjunction. *Hum.Genet.*, Vol.108, No.2, pp. 134-139
- Stahl, P. J., Sember, D. S. & Goldstein, M. (2011). Contemporary Management of Male Infertility. *Annu.Rev.Med.* Vol 63, No.18, pp. 1-16
- Sun, F., Oliver-Bonet, M., Liehr, T., Starke, H., Ko, E., Rademaker, A., Navarro, J., Benet, J. & Martin, R. H. (2004a). Human male recombination maps for individual chromosomes. *Am.J.Hum.Genet.*, Vol.74, No.3, pp. 521-531
- Sun, F., Kozak, G., Scott, S., Trpkov, K., Ko, E., Mikhaail-Philips, M., Bestor, T. H., Moens, P. & Martin, R. H. (2004b). Meiotic defects in a man with non-obstructive azoospermia: case report. *Hum.Reprod.*, Vol.19, No.8, pp. 1770-1773
- Sun, F., Oliver-Bonet, M., Liehr, T., Starke, H., Trpkov, K., Ko, E., Rademaker, A. & Martin, R. H. (2005). Discontinuities and unsynapsed regions in meiotic chromosomes have a cis effect on meiotic recombination patterns in normal human males. *Hum.Mol.Genet.*, Vol.14, No.20, pp. 3013-3018

- Sun, F., Oliver-Bonet, M., Liehr, T., Starke, H., Turek, P., Ko, E., Rademaker, A. & Martin, R. H. (2006). Variation in MLH1 distribution in recombination maps for individual chromosomes from human males. *Hum.Mol.Genet.*, Vol.15, No.15, pp. 2376-2391
- Sun, F., Turek, P., Greene, C., Ko, E., Rademaker, A. & Martin, R. H. (2007). Abnormal progression through meiosis in men with nonobstructive azoospermia. *Fertil.Steril.*, Vol.87, No.3, pp. 565-571
- Sun, F., Mikhaail-Philips, M., Oliver-Bonet, M., Ko, E., Rademaker, A., Turek, P. & Martin, R. H. (2008a). Reduced meiotic recombination on the XY bivalent is correlated with an increased incidence of sex chromosome aneuploidy in men with non-obstructive azoospermia. *Mol.Hum.Reprod.*, Vol.14, No.7, pp. 399-404
- Sun, F., Mikhaail-Philips, M., Oliver-Bonet, M., Ko, E., Rademaker, A., Turek, P. & Martin, R. H. (2008b). The relationship between meiotic recombination in human spermatocytes and aneuploidy in sperm. *Hum.Reprod.*, Vol.23, No.8, pp. 1691-1697
- Tang, S. S., Gao, H., Robinson, W. P., Ho Yuen, B. & Ma, S. (2004). An association between sex chromosomal aneuploidy in sperm and an abortus with 45,X of paternal origin: possible transmission of chromosomal abnormalities through ICSI. *Hum.Reprod.*, Vol.19, No.1, pp. 147-151
- Tassistro, V., Ghalamoun-Slami, R., Saias-Magnan, J. & Guichaoua, M. R. (2009). Chronology of meiosis & synaptonemal complex abnormalities in normal & abnormal spermatogenesis. *Indian J.Med.Res.*, Vol.129, No.3, pp. 268-278
- Tempest, H. G., Ko, E., Rademaker, A., Chan, P., Robaire, B. & Martin, R. H. (2009). Intra-individual and inter-individual variations in sperm aneuploidy frequencies in normal men. *Fertil.Steril.*, Vol.91, No.1, pp. 185-192
- Templado, C., Vidal, F. & Estop, A. (2011). Aneuploidy in human spermatozoa. *Cytogenet.Genome Res.*, Vol.133, No.2-4, pp. 91-99
- Thomas, N. S. & Hassold, T. J. (2003). Aberrant recombination and the origin of Klinefelter syndrome. *Hum.Reprod.Update*, Vol.9, No.4, pp. 309-317
- Tournaye, H., Devroey, P., Liu, J., Nagy, Z., Lissens, W. & Van Steirteghem, A. (1994). Microsurgical epididymal sperm aspiration and intracytoplasmic sperm injection: a new effective approach to infertility as a result of congenital bilateral absence of the vas deferens. *Fertil.Steril.*, Vol.61, No.6, pp. 1045-1051
- Ushijima, C., Kumasako, Y., Kihale, P. E., Hirotsuru, K. & Utsunomiya, T. (2000). Analysis of chromosomal abnormalities in human spermatozoa using multi-colour fluorescence in-situ hybridization. *Hum.Reprod.*, Vol.15, No.5, pp. 1107-1111
- Vallente, R. U., Cheng, E. Y. & Hassold, T. J. (2006). The synaptonemal complex and meiotic recombination in humans: new approaches to old questions. *Chromosoma*, Vol.115, No.3, pp. 241-249
- Van Opstal, D., Los, F. J., Ramlakhan, S., Van Hemel, J. O., Van Den Ouweland, A. M., Brandenburg, H., Pieters, M. H., Verhoeff, A., Vermeer, M. C., Dhont, M. & In't Veld, P. A. (1997). Determination of the parent of origin in nine cases of prenatally detected chromosome aberrations found after intracytoplasmic sperm injection. *Hum.Reprod.*, Vol.12, No.4, pp. 682-686
- Van Steirteghem, A. C., Nagy, Z., Joris, H., Liu, J., Staessen, C., Smits, J., Wisanto, A. & Devroey, P. (1993). High fertilization and implantation rates after intracytoplasmic sperm injection. *Hum.Reprod.*, Vol.8, No.7, pp. 1061-1066

- Van Steirteghem, A., Bonduelle, M., Devroey, P. & Liebaers, I. (2002). Follow-up of children born after ICSI. *Hum.Reprod.Update*, Vol.8, No.2, pp. 111-116
- Vegetti, W., Van Assche, E., Frias, A., Verheyen, G., Bianchi, M. M., Bonduelle, M., Liebaers, I. & Van Steirteghem, A. (2000). Correlation between semen parameters and sperm aneuploidy rates investigated by fluorescence in-situ hybridization in infertile men. *Hum.Reprod.*, Vol.15, No.2, pp. 351-365
- Vendrell, J. M., Garcia, F., Veiga, A., Calderon, G., Egozcue, S., Egozcue, J. & Barri, P. N. (1999). Meiotic abnormalities and spermatogenic parameters in severe oligoasthenozoospermia. *Hum.Reprod.*, Vol.14, No.2, pp. 375-378
- Viville, S., Warter, S., Meyer, J. M., Wittemer, C., Lorient, M., Mollard, R. & Jacqmin, D. (2000). Histological and genetic analysis and risk assessment for chromosomal aberration after ICSI for patients presenting with CBAVD. *Hum.Reprod.*, Vol.15, No.7, pp. 1613-1618
- Yanagimachi, R., Yanagimachi, H. & Rogers, B. J. (1976). The use of zona-free animal ova as a test-system for the assessment of the fertilizing capacity of human spermatozoa. *Biol.Reprod.*, Vol.15, No.4, pp. 471-476

IntechOpen

Aneuploidy in Health and Disease

Edited by Dr Zuzana Storchova

ISBN 978-953-51-0608-1

Hard cover, 244 pages

Publisher InTech

Published online 16, May, 2012

Published in print edition May, 2012

Aneuploidy means any karyotype that is not euploid, anything that stands outside the norm. Two particular characteristics make the research of aneuploidy challenging. First, it is often hard to distinguish what is a cause and what is a consequence. Secondly, aneuploidy is often associated with a persistent defect in maintenance of genome stability. Thus, working with aneuploid, unstable cells means analyzing an ever changing creature and capturing the features that persist. In the book *Aneuploidy in Health and Disease* we summarize the recent advances in understanding the causes and consequences of aneuploidy and its link to human pathologies.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Maria Vera, Vanessa Peinado, Nasser Al-Asmar, Jennifer Gruhn, Lorena Rodrigo, Terry Hassold and Carmen Rubio (2012). Human Male Meiosis and Sperm Aneuploidies, *Aneuploidy in Health and Disease*, Dr Zuzana Storchova (Ed.), ISBN: 978-953-51-0608-1, InTech, Available from:

<http://www.intechopen.com/books/aneuploidy-in-health-and-disease/human-male-meiosis-and-sperm-aneuploidies>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](https://creativecommons.org/licenses/by/3.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen